

The Jacobian Conjecture after the Three-Dimensional Counterexample: Exact Validation, a General Seed Family, the Geometry of Non-Properness, and Two-Dimensional Equivariant Rigidity

Felipe Santibañez-Leal 

CAOS Research program, github.com/fsantibanezleal/CAOS_RESEARCH

2026-07-24

Version 0.10

PREPRINT · not peer reviewed · CAOS Research programme
Paper A of a three-paper record · version 0.10 · 2026-07-24
DOI [10.5281/zenodo.21503365](https://doi.org/10.5281/zenodo.21503365) (concept DOI, resolves to the latest version)
Licensed CC BY 4.0 · sources and machine records:
github.com/fsantibanezleal/CAOS_RESEARCH

Abstract

On 2026-07-19, Levent Alpöge announced (crediting the language model Claude Fable 5 and a question posed by Akhil) an explicit polynomial map $F : \mathbb{C}^3 \rightarrow \mathbb{C}^3$ with constant Jacobian determinant -2 and a repeated value, refuting the Jacobian conjecture for every $N \geq 3$. We report the results of an exact-arithmetic experimental program built on this event. First, we validate the counterexample independently and compute the full fiber over the announced target: it consists of exactly the three announced points. Second, we reverse-engineer the structure of F : it is a weighted skew-product over the invariants $v = xy$, $t = x^2z$, its Keller property reduces to a two-variable identity, and its fibers are governed by a cubic in a single composite variable. Third, we derive and verify a general constructor: every admissible triple (seed polynomial p , scale k , section q) yields a Keller map with determinant $-kp(1)^2$ and generic fiber degree $\deg p + 1$; we exhibit new explicit counterexamples with rational collision certificates in fiber degrees 4, 5 and 6, plus a section-tail instance beyond the announced family shape, prove the degree law $(5d - 3, 5d - 4, 4)$ within the family, and show the announced map is its smallest member. Fourth, we make the failure geometry exact: a preimage escapes to infinity precisely when its fiber root is a multiple root, so the asymptotic variety of F is the plane $C = 0$ together with the explicit discriminant surface $27A^2C^2 - 18ABC + 16A + B^3C - B^2 = 0$; over $\mathbb{R}$ this wall separates a three-sheet from a one-sheet region, and F restricts to a surjective, non-injective real Keller map. The counterexamples reduce at suitable primes to non-injective Keller maps over $\mathbb{F}_\ell$ of degree below the characteristic. Fifth, we prove that the mechanism cannot reach the still-open two-variable case: for arbitrary weights, every $\mathbb{G}_m$ -equivariant Keller map of $\mathbb{C}^2$ is linear (a classification with a positive-factor kill at the top degree), so a planar counterexample, if any, must be genuinely non-equivariant. All computations are exact (rational or symbolic) and reproducible from the repository; every claim below is labeled as machine-verified, derived, or conjectural.

1 Introduction

Let k be a field of characteristic 0. A polynomial map $F = (F_1, \dots, F_N) : k^N \rightarrow k^N$ is a *Keller map* if $\det JF$ is a nonzero constant. The Jacobian conjecture $JC(N)$, stated in two variables by Kraus in 1884 (with a flawed proof whose gap, control of ramification at infinity, remained the subject's central obstacle [6]) and popularized by Keller in 1939 [4], asserted that every Keller map has a polynomial inverse. Over $\mathbb{C}$, invertibility is equivalent to injectivity, so a refutation requires exactly one repeated value. The classical positive ladder includes Wang's degree-2 theorem, Moh's analysis discarding planar counterexamples of degree up to 100 (with complete published detail only for the largest exceptional case; the modern audited floor is discussed in the planar companion), the Bass–Connell–Wright reduction to cubic homogeneous form [2], and Drużkowski's cubic-linear form [3]; Pinchuk [5] refuted the strong real variant by a non-proper local diffeomorphism with non-constant Jacobian. The conjecture is Problem 16 of Smale's list [7].

This is the *foundational* paper of a three-manuscript record, split so each audience reads a self-contained document while the repository log preserves the full history: (A) this paper, the three-dimensional aftermath (validation, structure, the seed family, the geometry of non-properness, the real census, characteristic p , the weighted landscape, and two-dimensional equivariant rigidity as the bridge out); (B) the planar-program paper (`manuscript-planar/`: the $JC(2)$ machine, the unconditional Newton-polygon exclusion theorems, the staircase transport, and the recalibrated frontier); (C) the cascade companion (`manuscript-cascade/`: the source-verified consequence cascade and the explicit dimension-48 symmetric witness).

On 2026-07-19/20, Alpöge announced the map, with $u = 1 + xy$,

$$F = (u^3z + y^2u(4 + 3xy), \quad y + 3xu^2z + 3xy^2(4 + 3xy), \quad 2x - 3x^2y - x^3z), \quad (1)$$

with $\det JF \equiv -2$ and $F(0, 0, -\frac{1}{4}) = F(1, -\frac{3}{2}, \frac{13}{2}) = F(-1, \frac{3}{2}, \frac{13}{2}) = (-\frac{1}{4}, 0, 0)$ [1]. Appending identity coordinates refutes $JC(N)$ for all $N \geq 3$; the case $N = 2$ remains open. At the time of writing no formal paper exists; the announcement is a machine-checkable certificate, which makes an exact experimental program both possible and necessary. This manuscript maps the findings of that program (repository: CAOS_RESEARCH, problem `algebraic-geometry/jacobian-conjecture`; experiments cited as EXP-*nnn*). Each claim is labeled: [MV] machine-verified exact identity, [D] derived analytically and certified on instances, [C] conjecture.

2 Exact validation and the full fiber (EXP-001)

Theorem 2.1 ([MV]). *For F as in (1): $\det JF = -2$ identically; the three announced points are pairwise distinct with the common image $(-\frac{1}{4}, 0, 0)$; and the fiber $F^{-1}(-\frac{1}{4}, 0, 0)$ consists of exactly those three points. In particular F is a non-injective Keller map and $JC(N)$ is false for every $N \geq 3$.*

The fiber computation is a lex Gröbner basis over $\mathbb{Q}$: the eliminant in z is $\frac{1}{8}(2z - 13)(4z + 1)$, and on the fiber $3x + 2y = 0$, $12y^2 = 4z + 1$. All coefficients of (1) are rational, so non-injectivity holds equally over $\mathbb{R}$.

3 The structure of the counterexample (EXP-002)

Theorem 3.1 ([MV]). *F is equivariant for the $\mathbb{G}_m$ -action with source weights $(1, -1, -2)$ and target weights $(-2, -1, 1)$. Writing $v = xy$, $t = x^2z$, $u = 1 + v$, the reduced data*

$$a_1 = x^2P, \quad b_1 = xQ, \quad c_1 = R/x = 2 - 3v - t$$

are polynomials in (v, t) ; each component of F is affine in z with coefficient vector $(u^3, 3xu^2, -x^3)$; and the generating identity $a_1 = \frac{ub_1}{2} - u^2 + \frac{u^3c_1}{2}$ holds.

Lemma 3.2 (Skew-product determinant lemma, [MV] with generic symbolic coefficients). *For any polynomials a_1, b_1, c_1 in (v, t) , the map $(a_1/x^2, b_1/x, xc_1)|_{v=xy, t=x^2z}$ satisfies*

$$\det JF \cdot c_1^2 = -J_2, \quad J_2 = \det \frac{\partial(V', T')}{\partial(v, t)}, \quad V' = b_1c_1, \quad T' = a_1c_1^2.$$

For F , $J_2 = 2c_1^2$ [MV]: the Keller property is a *two-variable* identity; the third dimension only carries weights. Fibers are governed by a single cubic: with $w = uc_1/2$ and $\Phi(w) = w^2 - w^3$,

$$\Phi(w) = \frac{1}{4}(wQR - PR^2) \quad \text{identically [MV]}, \quad (2)$$

so preimages of (A, B, C) correspond to roots of $4\Phi(w) - wBC + AC^2 = 0$; at the announced target this degenerates to $\Phi(w) = 0$ with roots $\{0, 0, 1\}$, matching the three preimages' w -values $(1, 0, 0)$ [MV]. Injectivity dies by *non-properness*: a Keller map is a local diffeomorphism, so sheets merge only at infinity; this is Pinchuk's loophole and the modern face of Kraus's gap.

4 The seed family: a general counterexample constructor (EXP-003/004)

The identification $c_1 = h(v) - t$ with $h = p/w$ is a degree-2 coincidence: an initial constructor built on it is *provably unsolvable* for $\deg p \geq 3$ (EXP-003, an exact unsolvability statement). The failure analysis yields the correct *potential form*.

Theorem 4.1 (Constructor v2, [D], certified on instances [MV]). *Let $p(w) = \sum_{i=1}^d p_i w^i$ satisfy*

$$p(0) = 0, \quad \int_0^1 p = 0, \quad p(1) \neq 0, \quad p'(1) \neq 2p(1),$$

let $k \neq 0$, $m = -kp(1)$, and let $q(v)$ be any polynomial with $q(0) = k$ and $q'(0) = k(p(1) - p'(1))/(p'(1) - 2p(1))$, with arbitrary higher-order tail. Set $\Phi(w) = \int_0^w p$, $s = c_1 = q(v) - t$, $w = us/k$, and define

$$V' = k^2p(w) + ms, \quad T' = wV' - k^2\Phi(w), \quad b_1 = V'/s, \quad a_1 = T'/s^2.$$

Then a_1, b_1 are polynomials in (v, t) , and $F_{p,k,q} = (a_1/x^2, b_1/x, xc_1)$ is a polynomial Keller map of $\mathbb{C}^3$ with

$$\det JF_{p,k,q} = -\frac{m^2}{k} = -kp(1)^2 \quad (\text{constant, for every admissible } (p, k, q)),$$

generic fiber degree $d + 1$, and the exact reconstruction inverse: from a root w of the fiber polynomial $k^2\Phi(w) - (wBC - AC^2)$,

$$s = \frac{BC - k^2p(w)}{m}, \quad u = \frac{kw}{s}, \quad x = \frac{C}{s}, \quad v = u - 1, \quad t = q(v) - s, \quad y = \frac{v}{x}, \quad z = \frac{t}{x^2}.$$

The announced map (1) is the instance $p = 2w - 3w^2$, $k = 2$, q affine; at $d = 2$ this shape is rigid up to the scale k [MV].

The three admissibility conditions are exactly the low-weight polynomiality constraints at the origin; the potential form makes $J_2 = m^2s^2/k$ automatic, so the Keller property costs nothing: this is the structural reason the conjecture is false in dimension ≥ 3 .

Theorem 4.2 (New explicit counterexamples, [MV]). *The following maps are non-injective polynomial Keller maps of $\mathbb{C}^3$, each certified by an engineered rational collision:*

Seed p	k	det	Component degrees	Generic fiber degree
$w - 2w^3$	3	-3	(12, 11, 4)	4
$8w - 12w^2 + 4w^3 - 5w^4$	14	-350	(17, 16, 4)	5
$2w - 3w^2$, section tail v^2	2	-2	(8, 7, 5)	3

For the first: $(-\frac{1}{15}, 18, 5130)$ and $(\frac{1}{24}, -18, -10368)$ both map exactly to $(-54, -54, 1)$. Collisions are engineered by prescribing two rational roots $w_1 \neq w_2$ and solving the linear system $w_i\tilde{V} - \tilde{T} = k^2\Phi(w_i)$ for the target invariants.

The section-tail instance changes the component-degree pattern at fixed fiber degree and determinant: the family is strictly larger than the seed-only family circulating informally. Every negative rational number is realized as a Keller determinant within the family.

Theorem 4.3 (Degree laws and family-internal minimality, [MV] for $d \leq 5$). *For affine sections q the component total degrees of $F_{p,k,q}$ are exactly $(5d - 3, 5d - 4, 4)$, $d = \deg p$ (verified $d = 2, \dots, 5$; the $d = 5$ instance, seed $w - 3w^5$, $k = 5$, $\det = -20$, degrees $(22, 21, 4)$, fiber degree 6, carries its own rational collision certificate). The scale k never changes degrees; section tails strictly increase them; degree-1 seeds are impossible (the integral condition forces $p = 0$), so the fiber-degree floor is 3. Consequently the minimal degree vector over the family is $(7, 6, 4)$, attained exactly at the announced map: the July 2026 counterexample is the smallest instance of its own family. Globally, only Wang's degree-2 theorem constrains $\mathbb{C}^3$; the existence of a counterexample of total degree 3 to 6 remains open.*

Remark 4.4 (Characteristic p , [MV]). The $d = 3$ instance reduces mod $\ell = 13$ and $\ell = 101$ to explicit non-injective Keller maps over $\mathbb{F}_\ell$ of total degree $12 < \ell$ (the rational collision reduces to distinct points with a common image, e.g. mod 13: $(6, 5, 8)$ and $(6, 8, 6)$ both to $(11, 11, 1)$). Unlike the classical char- p example $x - x^p$ (degree $= p$), the degree sits strictly below the characteristic; expected from standard integer-model reductions, now explicitly certified.

5 The asymptotic variety: where sheets merge

Theorem 5.1 (Escape = multiple fiber root, [MV] on F and the $d = 3$ instance). *In the potential form, the fiber polynomial $W(w) = k^2\Phi(w) - wBC + AC^2$ satisfies $W'(w) = k^2p(w) - BC$*

identically, and the reconstruction scale is $s = -W'(w)/m$. Hence a fiber root fails to reconstruct a finite preimage (the preimage escapes to infinity) precisely when it is a multiple root of W : over $C \neq 0$ the non-properness locus is the discriminant locus of W . For the announced F ,

$$A(F) = \{C = 0\} \cup \{27A^2C^2 - 18ABC + 16A + B^3C - B^2 = 0\}.$$

Verified end to end: the target $(0, 1, 1)$ lies on the surface and its exact fiber is the single point $(2, -\frac{1}{2}, \frac{9}{8})$ (two sheets escaped); over generic $(A, B, 0)$ the fiber is the single flat-sheet point $(0, B, A - 4B^2)$; the collision target $(-16, -16, 1)$ lies off the surface with a full 3-point rational fiber. Collisions are generic; escapes are codimension 1. Kraus's "ramification at infinity" is, for this family, the discriminant of a univariate polynomial.

6 The two-dimensional frontier: obstruction, rigidity, real census (EXP-005/006/010/011)

Theorem 6.1 ([MV] with generic symbolic f, g). *Let $\mathbb{G}_m$ act on $\mathbb{C}^2$ with weights $(1, -a)$, $a \geq 1$, and let $F = (x^{b_1}f(v), x^{b_2}g(v))$, $v = x^ay$, be an equivariant polynomial map. Then*

$$\det JF = x^{b_1+b_2+a-1} (b_1fg' - b_2f'g),$$

so the Keller condition forces $b_1 + b_2 = 1 - a$ and the single univariate identity $b_1fg' - b_2f'g = \text{const} \neq 0$.

Proposition 6.2 (One-invariant collapse). *If the reduced data (V', T') of a weighted skew-product depend on a single variable, then $dV' \wedge dT' = 0$, hence $J_2 \equiv 0$ and $\det JF \equiv 0$: no Keller map arises. Since a nontrivial $\mathbb{G}_m$ -action on $\mathbb{C}^2$ has one independent weight-0 invariant (against two in $\mathbb{C}^3$), the potential-form mechanism of Theorem 4.1 is structurally unavailable in dimension 2.*

Additionally [MV]: no coordinate slice of (1) (each variable frozen at rational test values, any output pair) is a 2D Keller map.

Theorem 6.3 (2D equivariant rigidity, [D], every mechanical step [MV] on sweeps). *For any $\mathbb{G}_m$ -action on $\mathbb{C}^2$ with weights $(w_1, -w_2)$, $w_1 \geq 1$, $w_2 \geq 0$, every equivariant polynomial Keller map is linear: (f_0x, g_0y) or (f_0y, g_0x) .*

Proof shape. Equivariant components are $P = x^{i_1}y^{j_1}f(v)$, $Q = x^{i_2}y^{j_2}g(v)$ with $v = x^{w_2}y^{w_1}$, and the determinant factors as

$$\det JF = x^{i_1+i_2-1} y^{j_1+j_2-1} [(i_1j_2 - i_2j_1)fg + v(\beta_1fg' - \beta_2f'g)], \quad \beta_r = w_1i_r - w_2j_r$$

(certified on 648 weight-times-exponent cases with generic symbolic f, g). Constancy forces $i_1 + i_2 = j_1 + j_2 = 1$, and the mixed shapes ($P = xyf$, $Q = g$ and its swap) admit no Keller solution (their bracket is divisible by v ; empty solves). The surviving shapes reduce to $fg + v(w_1fg' + w_2f'g) = c$ (or its swap), whose coefficient at the top bidegree $N = d_f + d_g$ is $(1 + w_1d_g + w_2d_f)f_{d_f}g_{d_g}$ with a strictly positive factor: the leading coefficients die and by descent f, g are constant (certified: empty constrained solves for all exact bidegrees ≤ 5 over eight weight pairs). $\square$

The independent enumeration EXP-006 (216 polynomial Keller instances over a bounded window, all 648 exactly-computed in-image fibers single-point, every instance linear [MV]) is the experimental companion of Theorem 6.3. Consequence: a two-variable counterexample, if one exists, must be genuinely non-equivariant: the symmetry class behind the entire three-dimensional family is completely closed in dimension 2.

Theorem 6.4 (Real fiber census of F , [MV]). *Over $\mathbb{R}$, the announced F restricts to a surjective, orientation-reversing ($\det = -2$), non-proper, non-injective real Keller map. For real targets off the discriminant wall the number of real preimages is 3 (region $D > 0$, which contains the collision targets) or 1 (region $D < 0$); two independent exact routes (Sturm isolation of the fiber cubic plus reconstruction, versus Gröbner elimination) agree at all samples, and no empty real fiber exists on a 36-target rational grid (an odd-degree argument makes this structural). In particular the constant-Jacobian real statement in dimension 3 falls with the same example; Pinchuk’s planar counterexample (1994) concerned the non-constant case.*

The honest map of remaining routes to a hypothetical 2D counterexample: no symmetry at all; non-reductive ($\mathbb{G}_a$) structures, which face the same one-invariant collapse in naive form; or gradings at infinity in the Newton-polygon/Abhyankar sense. Conversely, a proof of $JC(2)$ in this program’s terms is a properness statement: the reconstruction inverse of Theorem 4.1 shows exactly how asymptotic values are produced in dimension 3 (denominators $BC = k^2p(w)$), and whether one-invariant geometry leaves room for such escape is, in our view, the sharpest transferable open question.

7 The planar program (moved to the companion)

The planar arc of this program (the $JC(2)$ machine built on the linearity of the Keller condition in one component; the uniform closure of the $\min \deg \leq 2$ columns with one closed inverse; the four unconditional Newton-polygon exclusion theorems and the annihilation lemma; the diagonal-corner reduction and the staircase-transport instrument; and the source-audited recalibration of the gcd and degree-floor landscape) now lives in the companion manuscript `manuscript-planar/`, where it can grow with the active program without destabilizing this foundational record. Every result there keeps its experiment label (EXP-013 through EXP-043 at the time of this revision), and the repository history log remains the single chronological record for both papers.

8 Discussion and program

Why the conjecture is false. In the equivariant class, being Keller is a two-variable potential identity, satisfiable for free at any fiber degree ≥ 3 ; nothing in the determinant sees properness. **What is essential.** Two independent weight-0 invariants; the potential form; valuation conditions at the origin. **Consequence cascade (verified from primary sources, EXP-016).** The Mathieu conjecture is false for $SU(3)$ (Mathieu 1997: Mathieu for $SU(N)$ implies $JC(N)$; abelian case Duistermaat–van der Kallen 1998); the Gaussian moments conjecture is false (GMC implies JC, Derksen–van den Essen–Zhao 2017); Zhao’s vanishing conjecture is false (equivalent to JC); the Image conjecture is false in some dimension (implies the vanishing conjecture); and the full Dixmier conjecture is false (stable equivalence $JC(2n) \Leftrightarrow \text{Dixmier}(n)$),

Belov-Kanel–Kontsevich and Tsuchimoto: $JC(4)$ false gives Dixmier(2) false; Dixmier(1) remains open). The cascade is documented, with the explicit dimension-48 symmetric witness (a gradient Keller map $\text{id} + \nabla f$ with Hessian-nilpotent quartic potential and an exact two-point collision over $\mathbb{Q}(i)$, record EXP-041), in the cascade companion. **Standing work** lives in the planar companion: the transport closure of the fifth exclusion, the $B = 16$ normal-form attack, and the (72, 108) system; on this paper’s side, the global-minimality search in degrees 3 to 6 remains the one open experimental thread.

The weighted landscape: uniqueness of the mechanism (EXP-012)

Generalizing to source weights $(1, -1, -m)$, the determinant lemma $\det JF = x^{\sigma+m}D$ and the pairing reductions $J_2 = \pm c_1^{r_a+r_b-1}D$ hold generically [MV], and the reduced Jacobian vanishes to order $m - 1$ along the section. The survey outcome: at $m = 1$ the reduced pair is itself a planar Keller map, so that class is a *bridge* to $JC(2)$ (93 scanned instances, all injective; a collision there would refute $JC(2)$); at $m = 3$ the class $o = (-2, -2, 1)$ is *empty* (a one-line valuation proof, machine-certified) and $o = (-3, -1, 1)$ is rigid in scans (60 instances); at $m = 4$ the potential-form family does *not* exist: with seed degree up to 5 the admissibility system has Gröbner basis [1] under $\mu \neq 0$ [MV]. Our design hypothesis (a parity law producing families at every even m) was thereby refuted by its own experiment; the finding is stronger: *within the scanned weighted landscape, the announced counterexample’s weight system is the unique collision-capable mechanism*. Scope: sparse bounded scans, quantified in the experiment record; wider weight systems and higher-degree seeds are queued.

Reproducibility

Every theorem labeled [MV] is an exact symplectic computation over $\mathbb{Q}$ with a persisted script and output in the repository (experiments EXP-001 through EXP-012 under `problems/algebraic-geometry/jacobian`; each experiment declares its hypothesis before the run and records an adversarial-validation route with its verdict.

References

- [1] L. Alpöge, announcement of the counterexample, X (@_alpöge_), 2026-07-20; question posed by Akhil, map produced with Claude Fable 5. Independent press coverage: M. Sparkes, *New Scientist*, 2026-07-20.
- [2] H. Bass, E. H. Connell, D. Wright, The Jacobian conjecture: reduction of degree and formal expansion of the inverse, *Bull. Amer. Math. Soc. (N.S.)* **7** (1982), 287–330.
- [3] L. M. Drużkowski, An effective approach to Keller’s Jacobian conjecture, *Math. Ann.* **264** (1983), 303–313.
- [4] O.-H. Keller, Ganze Cremona-Transformationen, *Monatsh. Math. Phys.* **47** (1939), 299–306. doi:10.1007/BF01695502.
- [5] S. Pinchuk, A counterexample to the strong real Jacobian conjecture, *Math. Z.* **217** (1994), 1–4.

- [6] L. O. Rodríguez Díaz, On the origin of the Jacobian conjecture, *C. R. Math.* **364** (2026), 363–370. arXiv:2512.23614.
- [7] S. Smale, Mathematical problems for the next century, *Math. Intelligencer* **20** (1998), 7–15.
- [8] A. van den Essen, *Polynomial Automorphisms and the Jacobian Conjecture*, Progress in Mathematics 190, Birkhäuser, 2000.