

PRIMARY SYPHILIS



Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.

Primary Chancre—Male This dry-based, painless ulcer with indurated borders is typical for a primary chancre in a male patient. (Courtesy of A. Wisdom: *Sexually Transmitted Diseases*. London: Mosby-Wolfe; 1992.)



Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.

Primary Chancre—Female A solitary, painless genital chancre with a clean base in a patient with primary syphilis. (Courtesy of the Department of Dermatology, Naval Medical Center, Portsmouth, VA.)






Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.

Secondary Syphilis—Trunk Rash on trunk in secondary syphilis. (Courtesy of A. Wisdom: *Sexually Transmitted Diseases*. London: Mosby-Wolfe; 1992.)



Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.

Secondary Syphilis—Palms Papulosquamous rash of secondary syphilis. Note the annular appearance of the palmar rash. (Courtesy of H. Hunter Handsfield: *Atlas of Sexually Transmitted Diseases*. New York: McGraw-Hill; 1992.)






Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.

Secondary Syphilis—Soles Hyperkeratotic plantar rash in a patient with secondary syphilis. (Courtesy of H. Hunter Handsfield: *Atlas of Sexually Transmitted Diseases*. New York: McGraw-Hill; 1992.)



Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.

Mucous Patches Oral involvement in secondary syphilis manifest by mucous patches. These lesions are very infectious, and dark-field examination is often positive for spirochetes. (Courtesy of Morse, Moreland, Thompson: *Atlas of Sexually Transmitted Diseases*. London: Mosby-Wolfe; 1990.)



Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.

Condyloma Lata Typical appearance of the verrucous, heaped up lesions of condyloma lata, a manifestation of secondary syphilis. (Courtesy of H. Hunter Handsfield: *Atlas of Sexually Transmitted Diseases*. New York: McGraw-Hill; 1992.)



GONORRHEA



Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.

Male Urethritis Purulent, copious urethral discharge in a patient with gonococcal urethritis. (Courtesy of H. Hunter Handsfield: *Atlas of Sexually Transmitted Diseases*. New York: McGraw-Hill; 1992.)



Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.


Female Urethritis Gonococcal urethritis in a female patient. Note the purulent urethral discharge. (Courtesy of Morse, Moreland, Thompson: *Atlas of Sexually Transmitted Diseases*. London: Mosby-Wolfe; 1990.)






Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.

Cervicitis Endocervical purulent exudate in an asymptomatic patient with gonococcal cervicitis. The cervix is very friable. (Courtesy of King K Holmes, MD, from H. Hunter Handsfield: *Atlas of Sexually Transmitted Diseases*. New York: McGraw-Hill; 1992.)



Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.

Conjunctivitis Chemotic conjunctiva and copious purulent exudate in a patient with gonococcal conjunctivitis. (Courtesy of H. Hunter Handsfield: *Atlas of Sexually Transmitted Diseases*. New York: McGraw-Hill; 1992.)



Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.

Skin Lesions Small pustules with hemorrhage suggestive of the skin lesions of disseminated gonococcal infection. (Courtesy of H. Hunter Handsfield: *Atlas of Sexually Transmitted Diseases*. New York: McGraw-Hill; 1992.)



Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.

Bartholin's Cyst Enlarged, fluctuant, tender Bartholin's abscess of the labia, usually but not always a result of gonorrhea. (Courtesy of A. Wisdom: *Sexually Transmitted Diseases*. London: Mosby-Wolfe; 1992.)




CHLAMYDIAL INFECTION



Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.

Male Urethritis Thin urethral discharge of chlamydial urethritis.
(Courtesy of Walter Stamm, MD, from H. Hunter Handsfield: *Atlas of Sexually Transmitted Diseases*. New York: McGraw-Hill; 1992.)



Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.

Cervicitis Mucopurulent cervicitis from chlamydial infection. (Courtesy of H. Hunter Handsfield: *Atlas of Sexually Transmitted Diseases*. New York: McGraw-Hill; 1992.)

LYMPHOGRANULOMA VENEREUM





Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.

Lymphogranuloma Venereum Unilateral left lymphadenopathy in a patient with lymphogranuloma venereum. (Courtesy of Lawrence B. Stack, MD.)





Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.

Primary Lesions—Female Multiple coalescing superficial ulcerations of primary genital herpes. (Courtesy of Lawrence B. Stack, MD.)



Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.

Primary Lesions—Male Multiple genital vesicles of primary genital herpes. (Courtesy of H. Hunter Handsfield: *Atlas of Sexually Transmitted Diseases*. New York: McGraw-Hill; 1992.)



Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.

Recurrent Lesions—Female Solitary, minimally painful lesion of recurrent genital herpes. (Courtesy of H. Hunter Handsfield: *Atlas of Sexually Transmitted Diseases*. New York: McGraw-Hill; 1992.)



Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.

Herpes Simplex Virus—Cervix Erosive ulcerations of the cervix in a patient with genital herpes infection. This patient may be completely asymptomatic and may transmit the disease. (Courtesy of A. Wisdom: *Sexually Transmitted Diseases*. London: Mosby-Wolfe; 1992.)

CHANCROID



Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.

Chancroid Lesions Multiple painful, deep ulcerations of chancroid.
(Courtesy of H. Hunter Handsfield: *Atlas of Sexually Transmitted Diseases*. New York: McGraw-Hill; 1992.)



Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.


Chancroid Lesions and Inguinal Nodes Chancroid lesions with an enlarged lymph node. On examination, this node is tender and fluctuant. (Courtesy of H. Hunter Handsfield: *Atlas of Sexually Transmitted Diseases*. New York: McGraw-Hill; 1992.)





Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.

Pediculosis Pubis—on Hairs *Phthirus pubis*, or the crab louse, in the pubic hair of a patient complaining of itching. Note also the nits attached to the hairs. (Courtesy of Morse, Moreland, Thompson: *Atlas of Sexually Transmitted Diseases*. London: Mosby-Wolfe; 1990.)



Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.


Pediculosis Pubis—on Eyelashes *Phthirus pubis* lice noted in the eyelashes. (Courtesy of Spalton, Hitchings, Hunter: *Atlas of Clinical Ophthalmology*, 2d ed. London: Mosby-Year Book Europe; 1994.)

CONDYLOMA ACUMINATA (GENITAL WARTS)



Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.

Genital Warts—Female Verrucous lesions of the posterior fourchette in a patient with condyloma acuminata. (Used with permission from H. Hunter Handsfield: *Atlas of Sexually Transmitted Diseases*. New York: McGraw-Hill; 1992.)



Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.

Genital Warts—Male Typical appearance of condyloma acuminata of the glans penis. (Courtesy of Morse, Moreland, Thompson: *Atlas of Sexually Transmitted Diseases*. London: Mosby-Wolfe; 1990.)



Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.

Genital Warts—Female Verrucous lesions of the posterior fourchette in a patient with condyloma acuminata. (Used with permission from H. Hunter Handsfield: *Atlas of Sexually Transmitted Diseases*. New York: McGraw-Hill; 1992.)



Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.

Giant Warts—Female Giant warts of a female patient with extensive condyloma acuminata. (Courtesy of Morse, Moreland, Thompson; *Atlas of Sexually Transmitted Diseases*. London: Mosby-Wolfe; 1990.)



Copyright ©2006 by The McGraw-Hill Companies, Inc.
All rights reserved.

Giant Warts—Male Giant warts in a male patient with extensive condyloma acuminata. (Courtesy of A. Wisdom; *Sexually Transmitted Diseases*. London: Mosby-Wolfe; 1992.)

<http://nurseRD.blogspot.com>

www.authorstream.com/reynel89/Nursing

www.slideshare.net/reynel89/slideshows



THANK YOU!
Have a nice day :)

-
RDG

