


# X-Ray Grating Tomography without Phase-Retrieval using Spherically Symmetric Basis Functions

Florian Schiffrers<sup>1,2</sup>, Sebastian Kaeppler<sup>1</sup>, Shiyang Hu<sup>1</sup>, Jens Rieger<sup>2</sup>, Georg Pelzer<sup>2</sup>, Thilo Michel<sup>2</sup>, Christian Riess<sup>1</sup> and Gisela Anton<sup>2</sup>



**Figure:** *Illustration of a blob-based forward projection. A sample blob has a support given by the black circle and its energy - as indicated in green - is distributed around the center (red). The traditional (blue) and differential (orange) footprint depends only on the forward projection of the blob's center onto the detector.*

Traditional Talbot-Lau X-ray Phase-Contrast tomography requires prior phase-retrieval based on - for instance

- G2 stepping. Since several exposures per projection angle are required, high dose needs to be applied and simple integration into a continuously rotating gantry is cumbersome.

By modelling the phase-contrast image formation process using the object's attenuation, refractive and small-angle scattering properties, a maximum-likelihood-based algorithm is applied for tomographic reconstruction without the need of prior phase-retrieval [1]. Previous research has shown that a tomographic acquisition with only one phase-step per projection angle suffices using the proposed algorithm [2].

The choice of radial symmetrical image basis functions for iterative reconstruction has drawn particular interest to the Phase-Contrast community due to the analytical solution of the Radon transform and its differential footprint [3]. Thus discretization artifacts due to numerical differentiation are avoided implying a positive impact on the reconstruction result.

In this work we show reconstruction results of Talbot-Lau grating tomography using Kaiser-Bessel functions [4].

## REFERENCES

- [1] Ritter, André, et al. "Simultaneous maximum-likelihood reconstruction for x-ray grating based phase-contrast tomography avoiding intermediate phase retrieval." arXiv preprint arXiv:1307.7912 (2013).
- [2] Brendel, Bernhard, et al. "Penalized maximum likelihood reconstruction for x-ray differential phase-contrast tomography." Medical physics 43.1 (2016): 188-194.
- [3] Köhler, Thomas, Bernhard Brendel, and Ewald Roessl. "Iterative reconstruction for differential phase contrast imaging using spherically symmetric basis functions." Medical physics 38.8 (2011): 4542-4545.
- [4] Lewitt, Robert M. "Multidimensional digital image representations using generalized Kaiser-Bessel window functions." JOSA A 7.10 (1990): 1834-1846.

<sup>1</sup> Pattern Recognition Lab, Friedrich-Alexander-Universität Erlangen-Nürnberg

<sup>2</sup> Erlangen Centre for Astroparticle Physics, Friedrich-Alexander-Universität Erlangen-Nürnberg

✉ Florian Schiffrers, florian.schiffrers@fau.de