# Planetary nebulae: observing-list comparison
1,143 catalogue objects; 796 in main panel

195 unknown/limited radius

24
unavailable
both axes

$\log_{10}$ observed Hβ flux [erg s$^{-1}$ cm$^{-2}$]

−10
−11
−12
−13
−14

Additional objects: log Hβ > −12

128 without this Hβ quantity

$10^{-2}$ $10^{-1}$ $10^{0}$ $10^{1}$

Optical angular radius (arcmin; half the Acker diameter)

Observed flux; no extinction correction. Surface brightness assumes a circular optical area and is a diagnostic proxy.

First list by priority
(main-panel counts)

- Messier (4)
- Caldwell (13)
- SAC Urban (3)
- RASC Finest NGC (12)
- SAC Best NGC (1)
- Herschel 400 South (15)
- Herschel 400 (3)
- Herschel II (7)
- SAC Beyond NGC (23)
- AL Planetary Nebulae (40)
- AL Planetary Nebulae South (26)
- AL Northern Alternates (17)
- Other catalogue objects (632)

All memberships remain in the table.
Hollow symbols: uncertain diameter.
Diameter limits stay outside main panel.
Missing data are not faint detections.

Hβ is not visual magnitude or [O III].
No visibility cutoff has been applied.

Dashed: additional-object Hβ cut.
Reference-list members bypass it.