Planetary nebulae: inclusion criteria

Catalogue: planetary_nebulae_selection_v1_selected.ecsv

Include every resolved reference-list member OR finite observed log H-beta flux > -12. The cut is strict; list members bypass it.

319 selected of 1143: 178 reference-list members and 141 additional flux-only objects.

Reference lists: messier, caldwell, sac_urban, rasc_finest_ngc, sac_best_ngc, herschel400_south, herschel400, herschel2, sac_beyond_ngc, al_planetary_nebulae, al_planetary_nebulae_south, al_planetary_nebulae_alternates

Observed H-beta flux in erg s-1 cm-2; surface proxy per arcsec2. Not V or OIII; no extinction correction, visual-selection threshold or imputation.

log10(S_Hbeta) = log_hbeta_flux - log10(pi*(60*radius_arcmin)^2)

Upper/lower limits retained as limits; excluded from measured-radius main panels. Uncertain diameters are hollow. No radius or flux imputation.

Highest surface brightness first for review, with angular size and quality; no automatic public-interest approval.

Plots retain the full comparison population and unavailable-data margins. A surface-brightness ranking is a review aid; it is not an approved public-interest selection.

Scientific source SHA-256: 949d05e8bd41d05345024df52da69f1030b7818656798c8471815a3b6be939e3