



Adaptive Immune Markers for Predicting Cognitive Decline in PD

Kathleen L. Poston MD, MS ^{a,b}, Li Zhu MD ^a, Marian Shahid MSc ^a,
Marion Buckwalter MD, PhD ^{a,b}

^a Stanford University, Department of Neurology and Neurological Sciences

^b Stanford University, Department of Neurosurgery

PPMI Project ID: 181

Summary (or Abstract)

PD patients are at high risk of developing cognitive impairment and dementia. This could be because PD-associated neuronal death triggers a slow autoimmune process. Elevated IgG index may be a biomarker indicating this abnormal adaptive immune response.

Method

ELISA assays of CSF or plasma samples were conducted in duplicate according to manufacturer's guidelines, and between-plate controls utilized to adjust for differences between plates. If between-plate controls were more than 1.5-fold different, assays were repeated. For IgG ELISAs (Abcam #ab192515), CSF was diluted 1:4000 and plasma 1:2x10⁶. For albumin ELISAs (Abcam #ab179887), CSF was diluted 1:6000 and plasma 1: 1:4x10⁶. IgG index was calculated as [(CSF IgG/CSF albumin)÷(plasma IgG/plasma albumin)].

References

1. Standard Citation Format





PARKINSON'S
PROGRESSION
MARKERS
INITIATIVE

Play a Part in Parkinson's Research

About the Authors

This document was prepared by Kathleen Poston, Li Zhu, Marian Shahid, and Marion Buckwalter, Stanford University, Department of Neurology and Neurological Sciences. For more information please contact Marion Buckwalter at (650) 724-9098 or by email at marionb@stanford.edu or Kathleen Poston at (650) 723-2116 or by email at klposton@stanford.edu.

Notice: This document is presented by the author(s) as a service to PPMI data users. However, users should be aware that no formal review process has vetted this document and that PPMI cannot guarantee the accuracy or utility of this document.

