



1	Introduction to Derivatices Markets	5
1.1	Financial Markets, Assets	5
1.2	Present Value of Future Payments	5
1.3	Derivatives	5
1.4	Arbitrage	5
1.5	Forwards and Futures	5
1.6	Options	6
2	Discrete Time Models	
2.1	One-Period Binomial Model	7
2.2	Multi-Period Binomial Model	7
2.3	Option Pricing in the Binomial World	7
2.4	Dividends	7
2.5	Exotic Options	7
2.6	General Discrete-Time Market Models	7
3	Basic Stochastic Processes	9
3.1	Information and Filtration	9
3.2	Brownian Motion	11
3.3	The Ito Integral and the Ito-Doeblin Lemma	14
3 /	Arithmetic and Geometric Brownian Motion Models	19

4	Continuous-Time Financial Models	23
4.1	The Black-Scholes Model	23
4.2	Risk-Neutral Pricing and Girsanov Theorem	29
4.3	Monte-Carlo Method for Pricing	35
4.4	Implies Volatility	35
4.5	The Greeks	35
4.6	Hedging	35
5	Continuous-Time Interest Rate Models	37
5.1	Bonds and Interest Rates	37
5.2	Zero-Coupon Bond Pricing and the Term Structure Equation	40
5.3	Martingale Models for the Short Rate	43
	Index	49

1. Introduction to Derivatices Markets

- 1.1 Financial Markets, Assets
- 1.2 Present Value of Future Payments
- 1.3 Derivatives
- 1.4 Arbitrage
- 1.5 Forwards and Futures

Definition 1.5.1 A **forward** contract is a non-standardised agreement to buy or sell an asset at a certain future time T at a certain price K, where K is known as the **delivery price** or the **forward price**.

Remark 1.5.2 The delivery price is determined such that the value of the contract at inception is 0, i.e. no money is exchanged at the inception of a forward.

The part who agrees to buy the underlying has a long position.

Remark 1.5.3 — **Forwards Payoff.** Suppose a forward contract has expiration date T, forward price K, then the payoff to the long position in the forward at time T is $S_T - K$ and to the short position at time T is $K - S_T$.

Proposition 1.5.4 Suppose a forward contract has underlying asset to be a non-dividend stock with spot S_0 at inception, and the forward has maturity at T. If the annual effective rate is r compounded continuously, then for any $t \in [0, T]$ then no-arbitrage forward price is

$$K = S_t e^{r(T-t)}$$

Proof. Fix $t \in [0,T)$. At time t, suppose we enter a long forward agreement with forward price K and short one share of the underlying stock at S_t , and invest S_t at r, then the cash proceeds at time t is $0 + S_t - S_t = 0$.

At time T, the payoff of this strategy is

$$S_T - K - S_T + S_t e^{r(T-t)} = S_t e^{r(T-t)} - K.$$

If $S_t e^{r(T-t)} - K > 0$, then we have an arbitrage opportunity since the initial cost was 0. If $S_t e^{r(T-t)} < K$ then we could have done the reverse (i.e. short the forward and borrow S_t to buy one share of the stock, and this would have led to 0 initial cost but positive final payoff.

By the Law of One Price, we have

$$K = S_t e^{r(T-t)}$$
.

Theorem 1.5.5 Suppose a forward contract has a dividend-paying stock as underlying which has dividends D_1, \ldots, D_n at time t_1, \ldots, t_n before expiration date T, and that the interest rate is r compounded continuously, then the forward price K at time $t \in [0, T]$ is

$$K_t = S_t e^{r(T-t)} - \sum_{i=1}^n D_i e^{r(T-t_i)}$$

where S_t is the stock price at time t and $t_n > t_{n-1} > \cdots > t_1 \ge t$.

Proof. Fix time $t \in [0, T]$ and consider two portfolios at t.

Portfolio A has long forward and cash amount of $Ke^{r(T-t)} + \sum_{i=1}^{n} D_i e^{-r(t_i-t)}$ invested at r.

Portfolio B longs one share of the stock and reinvests dividends at rate r.

At time T, portfolio A has payoff

$$S_T - K + Ke^{r(T-t)}e^{r(T-t)} + \sum_{i=1}^n D_i e^{-r(t_i-t)}e^{r(T-t)}$$

$$= S_T - K + K + \sum_{i=1}^n D_i e^{r(T-t_i)}$$

$$= S_T + \sum_{i=1}^n D_i e^{r(T-t_i)}$$

and portfolio B has payoff

$$S_T + \sum_{i=1}^n D_i e^{r(T-t_i)}.$$

Because A and B have the same payoffs at T, the initial investments must be the same by the Law of One Price, so

$$Ke^{r(T-t)} + \sum_{i=1}^{n} D_i e^{-r(t_i-t)} = S_t$$

i.e.

$$K = S_t e^{r(T-t)} - \sum_{i=1}^n D_i e^{r(T-t_i)}.$$

1.6 Options



- 2.1 One-Period Binomial Model
- 2.2 Multi-Period Binomial Model
- 2.3 Option Pricing in the Binomial World
- 2.4 Dividends
- 2.5 Exotic Options
- 2.6 General Discrete-Time Market Models



3.1 Information and Filtration


Definition 3.1.1 Let $\{X_t : t \in [0, \infty)\}$ be a continuous-time stochastic process over probability space (Ω, \mathcal{F}, P) (STAT330 (S) Remark 1.1.6). The **information set** at time t, denoted \mathcal{F}_t , represents everything we know about X_t 's sample path.

We assume that

$$\mathcal{F}_s \subseteq \mathcal{F}_t$$

for all $0 \le s \le t$, i.e. no information is forgotten.

■ Example 3.1.2 Consider the following 2-period binomial model.



Here $\Omega = \{\omega_i : 1 \le i \le 4\}$ where ω_1 is the *uu* path, ω_2 is the *ud* path, ω_3 is the *du* path and ω_4

is the dd path. We also have

$$egin{aligned} \mathcal{F}_0 &= \{\Omega,\emptyset\} \ \\ \mathcal{F}_1 &= \{\Omega,\{\omega_1,\omega_2\},\{\omega_3,\omega_4\},\emptyset\} \ \\ \mathcal{F}_2 &= \mathcal{P}(\Omega), \text{ the power set of } \Omega \end{aligned}$$

Definition 3.1.3 Let (Ω, \mathcal{F}, P) be a probability space. A collection $\{\mathcal{F}_t : t \in [0, \infty)\}$ of σ -algebras over Ω is called a **filtration** when $\mathcal{F}_s \subseteq \mathcal{F}_t \subseteq \mathcal{F}$ for all $0 \le s \le t < \infty$. A probability space with such a filtration is a **filtered probability space** and is denoted $(\Omega, \{\mathcal{F}_t\}_t, \mathcal{F}, P)$. A continuous time stochastic process over $(\Omega, \{\mathcal{F}_t\}_t, \mathcal{F}, P)$ is **adapted** to $\{\mathcal{F}_t\}_t$ if every X_t (from the underlying stochastic process) is \mathcal{F}_t -measurable, i.e. for every possible values $r \in \mathbb{R}$, $X_t^{-1}(\{x \in \mathbb{R} : x \le r\}) \in \mathcal{F}_t$.

Definition 3.1.4 A stochastic process $X = \{X_t : t \in [0, \infty)\}$ defined on a filtered probability space $(\Omega, \{\mathcal{F}_t\}_t, \mathcal{F}, P)$ is called a **martingale** with respect to $\{\mathcal{F}_t\}_t$ if

- 1. *X* is adapted to $\{\mathcal{F}_t\}_t$.
- 2. $E(|X_t|) < \infty$ for all $t \in [0, \infty)$.
- 3. (martingale property) $E(X_t | \mathcal{F}_s) = X_s$ almost surely for all $0 \le s < t < \infty$.

Remark 3.1.5 In measure theory terms, we have the concept "almost everywhere" with respect to a particular measure.

So
$$E(X_t|\mathcal{F}_s) = X_s$$
 almost surely means

$$\Pr(E(X_t|\mathcal{F}_s) \neq X_s) = 0.$$

■ Example 3.1.6 Let $X := \{X_t : t \in [0, \infty)\}$ be a stochastic process based on a filtered probability space $(\Omega, \{\mathcal{F}_t\}_t, \mathcal{F}, P)$. Define random variable

$$Z_t := E(X|\mathcal{F}_t).$$

Assume $E(|X|) < \infty$. Consider the stochastic process

$$Z := \{Z_t : t \in [0, \infty)\}.$$

Because each \mathcal{F}_t is a σ -algebra in Ω , each Z_t is \mathcal{F}_t -measurable, hence Z is adapted to $\{\mathcal{F}_t\}_t$. Next,

$$E(|Z_t|) = E(|E(X|\mathcal{F}_t)|) \le E(E(|X||\mathcal{F}_t)) = E(|X|)$$

by Law of Total Expectation. By assumption,

$$E(|Z_t|) \leq E(|X|) < \infty.$$

Finally if s < t, then

$$E(Z_t|\mathcal{F}_s) = E(E(X|\mathcal{F}_t)|\mathcal{F}_s) = E(X|\mathcal{F}_s) = Z_s$$

by another application of Law of Total Expectation.

Hence Z is a martingale.

Remark 3.1.7 Intuitively, a stochastic process behaves like a martingale if it follows no discernable pattern, i.e. the best forecast of a future value is the currently observed value.

Formally, for an arbitrary u > 0, if $\{X_t\}_{t>0}$ is a martingale, then

$$E(X_{t+u}-X_t|\mathcal{F}_t)=E(X_{t+u}|\mathcal{F}_t)-E(X_t|\mathcal{F}_t)=E(X_t|\mathcal{F}_t)-E(X_t|\mathcal{F}_t)=0.$$

A martingale is defined with respect to a filtration and a probability measure. A non-martingale process may be converted into a martingale through a change of measure.

3.2 Brownian Motion

3.2 Brownian Motion

Definition 3.2.1 A continuous-time stochastic process $\{W_t : t \ge 0\}$ on a probability space (Ω, \mathcal{F}, P) is a **standard one-dimensional Brownian motion** if

- 1. $W_0(\omega) = 0$ for all $\omega \in \Omega$.
- 2. The sample paths $t \mapsto W(t, \omega)$ are continuous for all $\omega \in \Omega$.
- 3. For all $0 \le s < t$, $W_t W_s \sim N(0, t s)$.
- 4. For all $0 = t_0 < t_1 < t_2 < \ldots < t_n < \infty$, we have the random variables $W_{t_1} W_{t_0}$, $W_{t_2} W_{t_1}$, ..., $W_{t_n} W_{t_{n-1}}$ to be independent.

Remark 3.2.2 Stock price movements are often modelled using Brownian motion due to the latter's frantal nature.

Definition 3.2.3 A Brownian motion with **drift** μ and **diffusion** coefficient σ is

$$X_t = \mu t + \sigma W_t, t \geq 0$$

where $\{W_t\}_t$ is a standard Brownian motion.

Proposition 3.2.4 A Brownian motion $\{X_t\}_t$ with drift μ and diffusion coefficient σ satisfies

$$X_t - X_s \sim N(\mu(t-s), \sigma^2(t-s))$$

for all $0 \le s \le t$.

Proof. We have $X_t - X_s = \mu t + \sigma W_t - \mu s - \sigma W_s = \mu (t - s) + \sigma (W_t - W_s)$ where $W_t - W_s \sim N(0, t - s)$. Hence

$$E(X_t - X_s) = \mu(t - s) + \sigma E(W_t - W_s) = \mu(t - s)$$

and $Var(X_t - X_s) = \sigma^2 Var(W_t - W_s) = \sigma^2(s - t)$ and follows a normal distribution.

Definition 3.2.5 A random variable X on a probability space Ω, \mathcal{F}, P) is **independent of a** σ -algebra $\mathcal{F}_0 \subseteq \mathcal{P}(\Omega)$ if for any event $A \in \Omega$ corresponding to $X \in \mathcal{B}$ where \mathcal{B} is a Borel set in \mathbb{R} , and any $C \in \mathcal{F}_0$, we have

$$Pr(A \cap C) = Pr(A) Pr(C)$$
.

Definition 3.2.6 Let $\{W_t\}_{t\geq 0}$ be a 1-dimensional standard Brownian motion on (Σ, \mathcal{F}, P) , then a **filtration for** $\{W_t\}_{t\geq 0}$ is a filtration $\{\mathcal{F}_t\}_{t\geq 0}$ on (Σ, \mathcal{F}, P) such that

- 1. $\{W_t\}_{t>0}$ is adapted to $\{\mathcal{F}_t\}_{t>0}$.
- 2. For all $0 \le s < t$, the increment $W_t W_s$ is independent of the σ -algebra \mathcal{F}_s

Remark 3.2.7 A filtration on a standard Brownian motion has the property that future increments do not depend on the information today.

Definition 3.2.8 Let $X_1, ..., X_n$ be rv's on (Ω, \mathcal{F}, P) . A **filtration generated by** $X_1, ..., X_n$ is the σ -algebra over Ω generated by (Ω, \mathcal{F}, P) , i.e. the collection of the inverse images of the Borel sets of \mathbb{R} :

$$\{X_i^{-1}(S): S \in \mathcal{B}(\mathbb{R}), 1 \le i \le n\}.$$

Definition 3.2.9 For a standard Brownian motion $\{W_t\}_{t\geq 0}$ over (Ω, \mathcal{F}, P) , the filtration generated by $\{W_t\}_{t\geq 0}$ is a filtration for $\{W_t\}_{t\geq 0}$.

Proof. By Def. 3.2.8, clearly each W_t is measurable because the σ -algebras are generated with the inverse images of W_t , so the filtration is adapted to $\{\mathcal{F}_t\}_{t>0}$.

For the second property, let $A \in \Omega$ such that A corresponds to $W_t - W_s \in \mathcal{B}(\mathbb{R})$, and let $C \in \mathcal{F}_s$, where \mathcal{F}_s is the σ -algebra generated by W_s . Note that $W_s = W_s - W_0$ and by Def. 3.2.1(3), $W_t - W_s$ is independent of $W_s - W_0$. Hence A and C are independent.

Remark 3.2.10 Another way of saying Def. 3.2.5, given Def. 3.2.8, is to say that the σ -algebra generated by X is independent of \mathcal{F} .

Definition 3.2.11 Let Ω , \mathcal{F} , P) be a probability space and \mathcal{F}_1 , \mathcal{F}_2 be sub- σ -algebras of F. F_1 , F_2 are **independent** σ -algebras if

$$F_1 \perp F_2$$

for all $F_1 \in \mathcal{F}_1$ and $F_2 \in \mathcal{F}_2$.

Proposition 3.2.12 Let $\{W_t\}_{t\geq 0}$ be a standard Brownian motion on (Ω, \mathcal{F}, P) and $\{\mathcal{F}_t\}_{t\geq 0}$ be a filtration for $\{W_t\}_{t\geq 0}$, then

- 1. $E(W_t) = 0$ and $Var(W_t) = t$ for all $t \ge 0$.
- 2. For all $0 \le s \le t$:
 - $(2.1) E(W_t | \mathcal{F}_s) = W_s.$
 - $(2.2) \operatorname{Var}(W_t | \mathcal{F}_s) = t s.$
 - (2.3) $Corr(W_t, W_s) = min(s, t) = s$.
- 3. $\{W_t\}_{t>0}$ is a martingale with respect to $\{F_t\}_{t>0}$.

Proof. (1)By Def. 3.2.1(3), $W_t - W_0 \sim N(0,t)$, so

$$E(W_t) = E(W_t - W_0) = E(W_t) - E(W_0) = 0 - 0 = 0$$

where $Var(W_t) = Var(W_t - W_0) = t$.

(2.1)

$$E(W_t|\mathcal{F}_s) = E(W_t - W_s + W_s|\mathcal{F}_s)$$

$$= E(W_t - W_s|\mathcal{F}_s) + E(W_s|\mathcal{F}_s)$$

$$= 0 + W_s$$

$$= W_s.$$

(2.2)

$$Var(W_t | \mathcal{F}_s) = Var(W_t - W_s + W_s | \mathcal{F}_s)$$

$$= Var(W_t - W_s | \mathcal{F}_s) + Var(W_s | \mathcal{F}_s) \text{ by independence}$$

$$= t - s + 0$$

$$= t - s.$$

(2.3)

$$Corr(W_t, W_s) = E(W_t W_s) - E(W_t)E(W_s)$$

$$= E(W_t W_s)$$

$$= E((W_t - W_s + W_s)W_s)$$

$$= E((W_t - W_s)W_s) + E(W_s^2)$$

$$= E((W_t - W_s)(W_s - W_0)) + Var(W_s)$$

$$= 0 + Var(W_s) \text{ by independence}$$

$$= s.$$

3.2 Brownian Motion 13

(3) Clearly $\{W_t\}_{t\geq 0}$ is adapted to $\{\mathcal{F}_t\}_{t\geq 0}$ because $\{\mathcal{F}_t\}_{t\geq 0}$ is generated by $\{W_t\}_{t\geq 0}$. Moreover, equation 2.1 proves the martingale property, so it suffices to show that each

$$E(|W_t|) < \infty$$
.

By the Cauchy-Schwarz Inequality we have

$$E(|W_t|) \le (E(|W_t|^2))^{\frac{1}{2}} = \sqrt{t} < \infty.$$

This completes the proof.

Proposition 3.2.13 The sample paths $t \mapsto W_t(\omega)$ for a fixed $\omega \in \Omega$ are continuous but nowhere differentiable.

Proof. We omit the proof.

Definition 3.2.14 Let $f:[0,T] \to \mathbb{R}$ be a function and $\Pi = \{t_i : 0 \le i \le n, 0 = t_0 < t_1 < \cdots < t_n <$ $t_n = T$ } be a partition of [0, T]. The **total variation** of f is

$$TV(f) = \lim_{\|\Pi\| \to 0} \sum_{i=1}^{n} |f(t_i) - f(t_{i-1})|,$$

the quadratic variation of f is

$$QV(f) = [f, f]_T = \lim_{\|\Pi\| \to 0} \sum_{i=1}^{n} (f(t_i) - f(t_{i-1}))^2$$

where

$$||\Pi|| = \max_{1 \le i \le n} t_i - t_{i-1}$$

is the mesh of the partition.

f is of bounded variation if $TV(f) < \infty$, and of unbounded variation if otherwise.

Theorem 3.2.15 The sample paths of a Brownian motion $\{W_t\}_{t\geq 0}$, $t\mapsto W_t(\omega)$, are of unbounded variation but

$$QV(W) = (W, W)_T = T$$

for all $T \ge 0$ with probability 1.

Proof. We omit the proof.

Theorem 3.2.16 Continuously differentiable functions have quadratic variations of 0.

Proof. We omit the proof.

Theorem 3.2.17 Let $\{W_t\}_{t>0}$ be a Brownian motion, then

- 1. $[W,t]_T = \lim_{||\Pi|| \to 0} \sum_{i=1}^n (W_{t_i} W_{t_{i-1}})(t_i t_{i-1}) = 0.$ 2. $[t,t]_T = \lim_{||\Pi|| \to 0} \sum_{i=1}^n (t_i t_{i-1})^2 = 0.$

Proof. We omit the proof.

Remark 3.2.18 Informally, we write the above results to be:

Thm. 3.2.15: $d(W,W)_t = dW_t dW_t = (dW_t)^2 = dt$.

Thm. 3.2.17(1): $d(W,T)_t = dW_t dt = 0$.

Thm. 3.2.17(2): $d(T,T)_t = dtdt = 0$.

Proposition 3.2.19 Let $\{W_t\}_{t\geq 0}$ be a standard Brownian motion, then the process

$$\{W_t^2 - t\}_{t>0}$$

is a martingale with respect to the filtration generated by $\{W_t\}_{t\geq 0}$.

Proof. Since $\{W_t\}_{t\geq 0}$ is adapted, so is $\{W_t\}_{t\geq 0}$ and therefore so is $\{W_t - t\}_{t\geq 0}$. Next, $E(|W_t^2 - t|) \leq E(|W_t^2|) + t$ by the triangle inequality, and so in turn

$$E(|W_t^2 - t|) \le E(W_t^2) + t = t + t = 2t < \infty.$$

Finally,

$$E(W_t^2 - t | \mathcal{F}_s) \text{ for } 0 \le s < t$$

$$= E((W_t - W_s)^2 + 2(W_t - W_s)W_s + W_s^2 - t | \mathcal{F}_s)$$

$$= E((W_t - W_s)^2 | \mathcal{F}_s) + 2W_s E(W_t - W_s | \mathcal{F}_s) + E(W_s^2 | \mathcal{F}_s) - t$$

$$= Var(W_t - W_s) + 2W_s \cdot 0 + W_s^2 - t$$

$$= t - s + W_s^2 - t$$

$$= W_s^2 - s$$

which satisfies the martingale property.

3.3 The Ito Integral and the Ito-Doeblin Lemma

Definition 3.3.1 Let $(\Omega, \{\mathcal{F}_t\}_t, \mathcal{F}, P)$ be a filtered probability space such that $\{\mathcal{F}_t\}_t$ is a filtration for a standard Brownian motion $\{W_t\}_t$. Let f_t be a function of random variables depending on t such that

$$E\left(\int_0^t f_u^2 du\right) < \infty.$$

Then we say $\{f_t\}_t$ is a **square-integrable process** and we define the **Ito integral of** $\{f_t\}_t$ to be the random variable

$$I_t = \int_0^t f_u dW_u = \lim_{n \to \infty} \sum_{[t_{i-1}, t_i] \in \Pi_n} H_{t_{i-1}}(W_{t_i} - W_{t_{i-1}})$$

where $\Pi_n = \{0 = t_0 < t_1 < \dots < t_n = t\}$ is an *n*-partition of the interval [0, t] and the convergence of the limit is convergence in probability.

Proposition 3.3.2 Let $\{I_t\}_t$ be the Ito integral for $\{f_t\}_t$, then

- 1. I_t is a continuous function of t.
- 2. For each t, I_t is \mathcal{F}_t -measurable, i.e. for every Borel set $B \in \mathcal{B}(\mathbb{R})$, $I_t^{-1}(B) \in \mathcal{F}_t$.

3. For any constant $c \in \mathbb{R}$,

$$c\int_0^t f_u dW_u = \int_0^t cf_u dW_u.$$

4. For square-integrable processes $(f_t)_t$ and $(g_t)_t$,

$$\int_0^t f_u dW_u + \int_0^t g_u dW_u = \int_0^t (f_u + g_u) dW_u.$$

Proof. We omit the proof.

Theorem 3.3.3 Let $\{I_t\}_t$ be the Ito integral on a filtered probability space $(\Omega, \{\mathcal{F}_t\}_t, \mathcal{F}, P)$. $\{I_t\}_t$ is a martingale process with respect to $\{\mathcal{F}_t\}_t$, where $\{\mathcal{F}_t\}_t$ is adapted to the underlying standard Brownian motion of $\{I_t\}_t$.

Proof. We omit the proof.

Corollary 3.3.4 With the same setup as Thm. 3.3.3, we have

- 1. $E(I_t) = 0$ for all t.
- 2. $E\left(\int_{s}^{t} f_{u}dW_{u}|\mathcal{F}_{s}\right) = 0$ for all $0 \le s < t$.

Proof. 1. By the martingale property

$$E(I_t) = E(I_t | \mathcal{F}_0) = I_0 = \int_0^0 f_u dW_u = 0.$$

2. We have

$$E\left(\int_{s}^{t} f_{u}dW_{u}|\mathcal{F}_{s}\right)$$

$$=E\left(\int_{0}^{t} f_{u}dW_{u}|\mathcal{F}_{s}\right) - E\left(\int_{0}^{s} f_{u}dW_{u}|\mathcal{F}_{s}\right)$$

$$=E(I_{t}|\mathcal{F}_{s}) - I_{s}$$

$$=I_{s} - I_{s} \text{ by part (1)}$$

$$=0$$

Theorem 3.3.5 — Ito Isometry. Let $\{I_t\}_t$ be the Ito integral with respect to a process $\{f_t\}_t$, then

$$E(I_t^2) = E\left(\int_0^t f_u^2 du\right) < \infty.$$

Proof. The $< \infty$ part is a consequence of the fact that $\{f_t\}_t$ is square integrable. The proof of the first part, i.e.

$$E\left(\left(\int_0^t f_u dW_u\right)^2\right) = E\left(\int_0^t f_u^2 du\right),\,$$

is omitted.

Corollary 3.3.6 With the same setup as Thm. 3.3.5, we have

- 1. $Var(I_t) = E(I_t^2) = E(\int_0^t f_u^2 du)$.
- 2. $\operatorname{Var}(I_t | \mathcal{F}_s) = E\left(\int_s^t f_u^2 du | \mathcal{F}_s\right)$ for $0 \le s < t$.

Proof. 1.

$$Var(I_t) = E(I_t^2) - E(I_t)^2 = E\left(\int_0^t f_u^2 du\right) - 0^2 = E\left(\int_0^t f_u^2 du\right)$$

where the second step is by Thm. 3.3.5 and Corollary 3.3.4.

2.

$$Var(I_{t}|\mathcal{F}_{s})$$

$$= Var\left(I_{s} + \int_{s}^{t} f_{u}dW_{u}|\mathcal{F}_{s}\right)$$

$$= E\left(\left(I_{s} + \int_{s}^{t} f_{u}dW_{u}\right)^{2}|\mathcal{F}_{s}\right) - E\left(I_{s} + \int_{s}^{t} f_{u}dW_{u}|\mathcal{F}_{s}\right)^{2}$$

$$= E(I_{s}^{2}|\mathcal{F}_{s}) + 2E\left(I_{s}\int_{s}^{t} f_{u}dW_{u}|\mathcal{F}_{s}\right) + E\left(\left(\int_{s}^{t} f_{u}dW_{u}\right)^{2}|\mathcal{F}_{s}\right) - E(I_{t}|\mathcal{F}_{s})^{2}$$

$$= 0 + 2(0) + E\left(\left(\int_{s}^{t} f_{u}dW_{u}\right)^{2}|\mathcal{F}_{s}\right)$$

$$= E\left(\int_{s}^{t} f_{u}dW_{u}|\mathcal{F}_{s}\right)$$

as required.

Definition 3.3.7 Let $(\Omega, \{\mathcal{F}_t\}_t, \mathcal{F}, P)$ be a filtered probability space where $\{\mathcal{F}_t\}_t$ is adapted to a standard Brownian motion $\{W_t\}_t$, then a stochastic process $\{X_t\}_t$ is an **Ito process** if X_t has the form

$$dX_t = \alpha_t dt + \sigma_t dW_t$$

where:

 dX_t , dt, dW_t denote the infinitismal change of X_t , time and W_t respectively, α_t is a process depending on t such that it is adapted to $\{\mathcal{F}_t\}_t$ and

$$E\left(\int_0^t |\alpha_u|du\right) < \infty,$$

 σ_t is a stochastic process adaptd to $\{\mathcal{F}_t\}_t$.

We may write the Ito process in integral form

$$X_t - x_0 = \int_0^t \alpha_u du + \int_0^t \sigma_u dW_u$$

where x_0 is a non-random constant.

We call α_t the **drift**, and σ_t the **diffusion** or **volatility** of the Ito process.

We call

$$dX_t = \alpha_t dt + \sigma_t dW_t$$

to be the **differential form** of the Ito process.

Lemma 3.3.8 Let $\{X_t\}_t$ be an Ito process with

$$dX_t = \alpha_t dt + \sigma_t dW_t,$$

then

$$(dX_t)^2 = \sigma_t^2 dt.$$

Proof. We have

$$(dX_t)^2 = \alpha_t^2 (dt)^2 + 2\alpha_t \sigma^2 dt dW_t + \sigma_t^2 (dW_t)^2 = \alpha_t^2 \cdot 0 + 2\alpha_t \sigma^2 \cdot 0 + \sigma_t^2 dt$$

by Remark 3.2.18. Consequently

$$(dX_t)^2 = \sigma_t^2 dt.$$

Theorem 3.3.9 — Ito-Doeblin Lemma. Let $\{X_t\}_t$ be an Ito process defined on $(\Omega, \{\mathcal{F}_t\}_t, \mathcal{F}, P)$ with $dX_t = \alpha_t dt + \sigma_t dW_t$ and

$$f: [0,T] \times \mathbb{R} \to \mathbb{R}$$

be a function such that

$$f_t := \frac{\partial f}{\partial t}, f_x := \frac{\partial f}{\partial x}, f_{xx} := \frac{\partial f^2}{\partial x \partial x}$$

are well-defined and continuous, then

$$Y_t := f(t, X_t)$$

is also an Ito process with drift equal to

$$f_t(t,X_t) + f_x(t,X_t)\alpha_t + \frac{1}{2}f_{xx}(t,X_t)\sigma_t^2$$

and diffusion

$$f_X(t,X_t)\sigma_t$$
.

In other words

$$dY_t = df(t, X_t)$$

$$= \left(f_t(t, X_t) + f_x(t, X_t) \alpha_t + \frac{1}{2} f_{xx}(t, X_t) \sigma_t^2 \right) dt + f_x(t, X_t) dt \sigma_t dW_t$$

$$= f_t(t, X_t) dt + f_x(t, X_t) dX_t + \frac{1}{2} f_{xx}(t, X_t) (dX_t)^2 \text{ where } (dX_t)^2 = \sigma_t^2 dt \text{ by Lemma 3.3.8}$$

Proof. We omit the proof.

■ Example 3.3.10 Suppose we would like to compute

$$\int_0^T W_t dW_t$$

where $\{W_t\}_t$ is a standard Brownian motion.

Using Ito-Doeblin's Lemma, take $X_t = W_t$ and define

$$f: [0,T] \times \mathbb{R} \to \mathbb{R}$$
$$(t,x) \mapsto x^2$$

and $Y_t = f(t, X_t) = X_t^2$. It follows that

$$f_t(t,x) = 0, f_x(t,x) = 2x, f_{xx}(t,x) = 2.$$

By Ito-Doeblin's lemma,

$$dY_t = d(W_t)^2 = 0dt + 2W_t dW_t + \frac{1}{2}(2)(1)dt$$

since $\sigma_t = 1$ ($\{W_t\}_t$ is the standard Brownian motion).

Therefore

$$\int_{0}^{T} d(W_{t})^{2} = \int_{0}^{T} 2W_{t}dW_{t} + \int_{0}^{T} dt$$

and so

$$W_T^2 - W_0^2 = 2 \int_0^T W_t dW_t + T$$
$$\Leftrightarrow W_T^2 - T = 2 \int_0^T W_t dW_t$$
$$\Leftrightarrow \int_0^T W_t dW_t = \frac{1}{2} (W_T^2 - T)$$

3.4 Arithmetic and Geometric Brownian Motion Models

Definition 3.4.1 A stochastic process $\{X_t\}_t$ satisfying the stochastic differential equation

$$dX_t = \alpha dt + \sigma dW_t$$

where α, σ are constants, and $\{W_t\}_t$ is a standard Brownian motion, is called an **arithmetic Brownian motion (ABM)**.

Proposition 3.4.2 Let $\{X_t\}_t$ ba an ABM, then

- 1. $X_t = X_0 + \alpha t + \sigma W_t$.
- 2. $E(X_t) = X_0 + \alpha t$.
- 3. $Var(X_t) = \sigma^2 t$.
- 4. $X_t \sim N(X_0 + \alpha t, \sigma^2 t)$.
- 5. For $s \le t$, $E(X_t | \mathcal{F}_s) = X_s + \alpha(t s)$.

Proof. 1. In differential form we have

$$dX_t = \alpha dt + \sigma dW_t$$

and integrating both sides yields

$$\int_0^t X_u du = \int_0^t \alpha dt + \int_0^t \sigma dW_u$$

where $\int_0^t \sigma dW_u$ is an Ito integral. This yields

$$X_t - X_0 = \alpha t + \sigma W_t$$

as required.

- 2. $E(W_t) = 0$ by Proposition 3.2.12(1).
- 3. Proposition 3.2.12(1) has $Var(W_t) = t$.
- 4. We have $W_t \sim N(0,t)$ by Def. 3.2.1. The rest follows.

5.

$$E(X_t|\mathcal{F}_s) = E\left(X_s + \int_s^t \alpha du + \int_s^t \sigma dW_u|\mathcal{F}_s\right)$$

= $E(X_s) + \alpha(t-s) + \sigma E(W_t - W_s|\mathcal{F}_s)$
= $X_s + \alpha(t-s)$

Corollary 3.4.3 It an ABM $\{X_t\}_t$ has drift 0, then $\{X_t\}_t$ is a martingale.

Proof. By Proposition 3.4.2(5), we have

$$E(X_t|\mathcal{F}_s) = X_s$$

if the drift $\alpha = 0$. By definition, $\{X_t\}_t$ is a martingale.

Remark 3.4.4 If we use ABM to model asset returns, we get

$$\frac{dS_t}{S_t} = \mu dt + \sigma dW_t$$

where S_t is the price of the asset at time t. This motivates the following definition.

Definition 3.4.5 Suppose $\{S_t\}_t$ is a stochastic process that follows

$$dS_t = \mu S_t dt + \sigma S_t dW_t$$

where $\{W_t\}_t$ is the standard Brownian motion, then $\{S_t\}_t$ is said to follow a **geometric Brownian** motion (GBM).

Theorem 3.4.6 Let $\{S_t\}_t$ follow a geometric Brownian motion

$$dS_t = \mu S_t dt + \sigma S_t dW_t,$$

then the unique solution to this stochastic differential equation is

$$S_t = S_0 \exp\left(\left(\mu - \frac{\sigma^2}{2}\right)t + \sigma W_t\right),$$

and furthermore for $0 \le t \le T$,

$$S_T = S_t \exp\left(\left(\mu - \frac{\sigma^2}{2}\right)(T - t) + \sigma(W_T - W_t)\right).$$

Proof. Define function $f(t, S_t) = \log(S_t)$ and $X_t = S_t$, then

$$f_t = 0, f_s = \frac{1}{s}, f_{ss} = -\frac{1}{s^2}.$$

By the Ito-Doeblin Lemma, we get

$$d\log(S_t) = f_t dt + f_s dS_t + \frac{1}{2} f_{ss} (dS_t)^2$$

$$= \frac{1}{S_t} dS_t - \frac{1}{2S_t^2} (dS_t)^2$$

$$= \frac{1}{s} (\mu S_t dt + \sigma S_t dW_t) - \frac{1}{2S_t^2} (\mu S_t dt + \sigma S_t dW_t)^2$$

$$= \mu dt + \sigma dW_t - \frac{1}{2S_t^2} (\mu^2 S_t^2 (dt)^2 + 2\mu S_t^2 \sigma dt dW_t + \sigma^2 S_t^2 (dW_t)^2)$$

$$= \mu dt + \sigma dW_t - \frac{1}{2S_t^2} (0 + 0 + \sigma^2 S_t^2 dt) \text{ by Remark 3.2.18}$$

$$= \left(\mu - \frac{\sigma^2}{2}\right) dt + \sigma dW_t.$$

Integrating both sides gives

$$\int_0^t d\log(S_u) = \int_0^t \mu - \frac{\sigma^2}{2} du + \int_0^t \sigma dW_u$$

and consequently

$$\log(S_t) - \log(S_0) = \left(\mu - \frac{\sigma^2}{2}\right)t + \sigma W_t$$

which gives

$$S_t = S_0 \exp\left(\left(\mu - \frac{\sigma^2}{2}\right)t + \sigma W_t\right), t \geq 0.$$

Integrating $d \log(S_t)$ above from t to T gives

$$\int_{t}^{T} d\log(S_{u}) = \int_{t}^{T} \mu - \frac{\sigma^{2}}{2} du + \int_{t}^{T} \sigma dW_{u},$$

and similar to before,

$$\log(S_T) - \log(S_t) = \left(\mu - \frac{\sigma^2}{2}\right)(T - t) + \sigma(W_T - W_t)$$

and

$$S_T = S_t \exp\left(\left(\mu - \frac{\sigma^2}{2}\right)(T - t) + \sigma(W_T - W_t)\right), 0 \le t \le T,$$

which completes the proof.

Definition 3.4.7 If *X* is a random variable such that

$$\log(X) \sim N(\mu, \sigma^2)$$

for some μ , σ^2 , then X is said to follow a **log-normal distribution** of mean mu and standard deviation σ . We write $X \sim \text{LogN}(\mu, \sigma^2)$.

Proposition 3.4.8 If $X \sim \text{LogN}(\mu, \sigma^2)$, then

$$E(X) = e^{\mu + \frac{1}{2}\sigma^2}$$

and

$$Var(X) = (e^{\sigma^2} - 1)e^{2\mu + \sigma^2}.$$

Proof. We omit the proof.

Proposition 3.4.9 If $\{S_t\}_t$ is a GBM with

$$dS_t = \mu S_t dt + \sigma S_t dW_t$$

then for all $t \ge 0$,

$$E(S_t) = S_0 e^{-\mu t}$$

and for all $0 \le t \le T$,

$$E(S_T|\mathcal{F}_t) = S_t e^{\mu(T-t)}.$$

Proof. By Thm. 3.4.6

$$S_t = S_0 e^{\left(\mu - \frac{\sigma^2}{2}\right)t + \sigma W_t}$$
.

Let

$$Z_t = \log(S_0) + \left(\mu - \frac{\sigma^2}{2}\right)t + \sigma W_t,$$

then, since

$$W_t \sim N(0,t)$$
,

we have

$$Z_t \sim N\left(\log(S_0) + \left(\mu - \frac{\sigma^2}{2}\right)t, \sigma^2 t\right).$$

Note that by construction, $S_t = e^{Z_t}$, so

$$S_t \sim \text{LogN}\left(\log(S_0) + \left(\mu - \frac{\sigma^2}{2}\right)t, \sigma^2 t\right),$$

and by Proposition 3.4.8,

$$E(S_t) = S_0 e^{\mu t - \frac{\sigma^2}{2}t + \frac{1}{2}\sigma^2 t} = S_0 e^{\mu t}$$

as desired.

On the other hand, Thm. 3.4.6 also has that

$$S_T = S_t e^{\left(\mu - \frac{\sigma^2}{2}\right)(T - t) + \sigma(W_T - W_t)}.$$

Given \mathcal{F}_t , $t \leq T$, we have

$$\log(S_T) = \log(S_t) + \left(\mu - \frac{\sigma^2}{2}\right)(T - t) + \sigma Z,$$

where $Z \sim N(0, T - t)$ by the property of Brownian motion. It follows that

$$S_T | \mathcal{F}_t \sim \text{LogN}\left(\log(S_t) + \left(\mu - \frac{\sigma^2}{2}\right)(T - t), \sigma^2(T - t)\right).$$

and subsequently

$$E(S_T|\mathcal{F}_t) = S_t e^{\mu(T-t) - \frac{\sigma^2}{2}(T-t) + \frac{1}{2}\sigma^2(T-t)} = S_t e^{\mu(T-t)}$$

as desired.



4.1 The Black-Scholes Model

Definition 4.1.1 The **Black-Scholes model** has 2 assets, a risk-free asset with price process $\{B_t\}_t$, a risky asset with price process $\{S_t\}_t$ which does not pay dividend, from time $t \in [0,T]$, in a filtered probability space $(\Omega, \{\mathcal{F}_t\}_t, \mathcal{F}, P)$, and a corresponding Brownian motion $\{W_t^P\}_t$ with respect to P, where P, called the **physical probability measure**, is such that

$$dB_t = rB_t dt$$

for a fixed risk-free rate r, and $\{S_t\}_t$ follows a geometric Brownian motion

$$dS_t = \alpha S_t dt + \sigma S_t dW_t^P$$

for some constants α and σ .

A **trading strategy** is a process $\{h_t\}_t$ where each $h_t := (h_t^B, h_t^S)$ represents h_t^B units of the risk-free asset held in interval $[t, t + \Delta t)$ and h_t^S units of the risky asset held in interval $[t, t + \Delta t)$. Each h_t has **valuation**

$$V_t^h = h_t^B B_t + h_t^S S_t.$$

The process $\{V_t^h\}_t$ of trading strategy $\{h_t\}_t$ is the **valuation process** of the trading strategy.

Definition 4.1.2 A trading strategy $\{h_t\}_t$ in the Black-Scholes model is **self-financing** if

$$dV_t^h = h_t^S dS_t + h_t^B dB_t,$$

and it is an **arbitrage opportunity** if it is self-financing with $V_0^h \le 0$, $P(V_T^h \ge 0) = 1$, and $P(V_T^h > 0) > 0$.

The model is **arbitrage free** if there does not exist arbitrage opportunities.

Lemma 4.1.3 Suppose a contingency claim in the Black-Scholes model has pricing process $\{\Pi_t\}_t$ where each Π_t is a function of t and the risky asset price S_t : $\Pi_t = F(t, S_t)$, and that there exists a replicating portfolio $\{h_t\}_t$ that is also self-financing, then we have

1.

$$\alpha h_t^S S_t + r h_t^B B_t = \frac{\partial F}{\partial t} + \alpha S_t \frac{\partial F}{\partial S} + \frac{1}{2} \frac{\partial^2 F}{\partial S^2} \sigma^2 S_t^2$$

2.

$$\sigma h_t^S S_t = \frac{\partial F}{\partial S} \sigma S_t$$

and consequently $h_t^S = \frac{\partial F}{\partial S}$.

Proof. Since $\{h_t\}_t$ is self-financing, we have

$$dV_t^h = h_t^S dS_t + h_t^B dB_t$$

where $dS_t = \alpha S_t dt + \sigma S_t dW_t^P$ and $dB_t = rB_t dt$ by definition. Expand to get

$$dV_t^h = h_t^S(\alpha S_t dt + \sigma S_t dW_t^P) + h_t^B r B_t dt$$

= $(\alpha h_t^S S_t dt + h_t^B r B_t) dt + \sigma S_t h_t^S dW_t^P$.

On the other hand, $\{S_t\}_t$ is an Ito process, so by the Ito-Doeblin Lemma, $\{\Pi_t\}_t = \{F(t, S_t)\}_t$ is an Ito process with

$$d\Pi_t = dF(t, S_t) = \frac{\partial F}{\partial t}dt + \frac{\partial F}{\partial S}dS_t + \frac{1}{2}\frac{\partial^2 F}{\partial S^2}(dS_t)^2.$$

Substitute in $dS_t = \alpha S_t dt + \sigma S_t dW_t^P$ to get

$$d\Pi_{t} = \frac{\partial F}{\partial t}dt + \frac{\partial F}{\partial S}(\alpha S_{t}dt + \sigma S_{t}dW_{t}^{P}) + \frac{1}{2}\frac{\partial^{2} F}{\partial S^{2}}(\sigma^{2}S_{t}^{2}dt)$$

where the last term above is a consequence of Remark 3.2.18. Simplify this further to get

$$d\Pi_t = \left(\frac{\partial F}{\partial t} + \alpha S_t \frac{\partial F}{\partial S} + \frac{1}{2} \frac{\partial^2 F}{\partial S^2} \sigma^2 S_t^2\right) dt + \frac{\partial F}{\partial S} \sigma S_t dW_t^P.$$

Now, $\{h_t\}_t$ replicates $\{\Pi_t\}_t$, so $dV_t^h=d\Pi_t$, and we equate their expanded forms to get

$$(\alpha h_t^S S_t dt + h_t^B r B_t) dt + \sigma S_t h_t^S dW_t^P = dV_t^h$$

= $d\Pi_t = \left(\frac{\partial F}{\partial t} + \alpha S_t \frac{\partial F}{\partial S} + \frac{1}{2} \frac{\partial^2 F}{\partial S^2} \sigma^2 S_t^2\right) dt + \frac{\partial F}{\partial S} \sigma S_t dW_t^P.$

Equating drift yields

$$\alpha h_t^S S_t + r h_t^B B_t = \frac{\partial F}{\partial t} + \alpha S_t \frac{\partial F}{\partial S} + \frac{1}{2} \frac{\partial^2 F}{\partial S^2} \sigma^2 S_t^2$$

which is equation 1, and equating diffusion yields

$$\sigma S_t h_t^S = \frac{\partial F}{\partial S} \sigma S_t,$$

which is equation 2. This completes the proof.

Theorem 4.1.4 — The Black-Scholes Partial Differential Equation. Suppose a contingency claim in the Black-Scholes model has pricing process $\{\Pi_t\}_t$ where each Π_t is a function of t and the risky asset price S_t : $\Pi_t = F(t, S_t)$, and that there exists a replicating portfolio $\{h_t\}_t$ that is also self-financing, then we have

$$\frac{\partial F}{\partial t} + rS\frac{\partial F}{\partial S} + \frac{1}{2}\frac{\partial^2 F}{\partial S^2}\sigma^2 S^2 = rF$$

where $F(T, S_T) = \Pi_T$ is the payoff of the contingency at time T.

Proof. Lemma 4.1.3(1) gives $h_t^S = \frac{\partial F}{\partial S}$, but via the definition of the valuation process we also have

$$h_t^B B_t = V_t^h - h_t^S S_t.$$

Hence $h_t^B B_t = V_t^h - \frac{\partial F}{\partial S} S_t$. Moreover, because $\{h_t\}_t$ replicates $\{\Pi_t\}_t$, we have

$$h_t^B B_t = \Pi_t - \frac{\partial F}{\partial S} S_t = F(t, S_t) - \frac{\partial F}{\partial S} S_t.$$

Take Lemma 4.1.3(1) and substitute $h_t^B B_t = \Pi_t - \frac{\partial F}{\partial S} S_t$ and $h_t^S = \frac{\partial F}{\partial S}$ to get

$$\alpha \frac{\partial F}{\partial S} S_t + rF - r \frac{\partial F}{\partial S} S_t = \frac{\partial F}{\partial t} + \alpha S_t \frac{\partial F}{\partial S} + \frac{1}{2} \frac{\partial^2 F}{\partial S^2} \sigma^2 S_t^2.$$

This in turn yields

$$rF = \frac{\partial F}{\partial t} + rS\frac{\partial F}{\partial S} + \frac{1}{2}\frac{\partial^2 F}{\partial S^2}\sigma^2 S^2$$

as required.

Corollary 4.1.5 In a replicating portfolio $\{h_t\}_t$ of a contingency claim $\{\Pi_t\}_t$ in a Black-Scholes model, we have

$$h_t = (h_t^B, h_t^S) = \left(\frac{1}{B_t} \left(F - \frac{\partial F}{\partial S} S_t\right), \frac{\partial F}{\partial S}\right)$$

where $F(t, S_t) = \Pi_t$ for all $t \in [0, T]$.

Proof. This is in the proof of Thm. 4.1.4.

Theorem 4.1.6 — Feynman-Kac Theorem. Consider the partial differential equation

$$\frac{\partial}{\partial t}F(t,x) + \mu(t,x)\frac{\partial}{\partial t}F(t,x) + \frac{1}{2}\sigma^{2}(t,x)\frac{\partial^{2}}{\partial x^{2}}F(t,x) = V(t,x)F(t,x) - f(t,x)$$

defined for all $x \in \mathbb{R}$ and $t \in [0, T]$ subject to the boundary condition

$$F(T,x) = \Phi(x)$$

where μ, σ, Φ, V, f are known functions, $T \in \mathbb{R}^+$ is known, and

$$F: [0,T] \times \mathbb{R} \to \mathbb{R}$$

is the unknown function, then the solution F^* can be written as the conditional expectation

$$F^*(t,x) = E^P\left(\left.\int_t^T e^{-\int_t^r V(u,X_u)du} f(r,X_r) dr + e^{-\int_t^T V(u,X_u)du} \Phi(X_T)\right| X_t = x\right)$$

under the probability measure P such that X is an Ito process driven by

$$dX = \mu(X,t)dt + \sigma(t,X)dW_t^P$$

where $\{W_t^P\}$ is a Brownian motion under P and the initial condition X(t) is X(t) = x.

Proof. We omit the proof.

Corollary 4.1.7 If F is the solution to the partial differential equation

$$\frac{\partial F}{\partial t}(t,x) + \mu(t,x)\frac{\partial F}{\partial x}(t,x) + \frac{1}{2}\sigma^{2}(t,x)\frac{\partial^{2}}{\partial x^{2}}F(t,x) = rF(t,x)$$

subjected to the boundary condition

$$F(T,x) = \Phi(x)$$

where μ, Φ, σ are known functions, $r, T \in \mathbb{R}^+$, and $F : [0, T] \times \mathbb{R} \to \mathbb{R}$, then F has representation

$$F(t,x) = e^{-r(T-t)}E^{P}(\Phi(X_T)|X_t = x)$$

where X satisfies the partial differential equation

$$dX_s = \mu(s, X_s)ds + \sigma(s, X_s)dW_s^P, X_t = x$$

where $\{W_s\}_s$ is a Brownian motion under P.

Proof. Let f(t,x) = 0 and V(t,x) = r in the Feynman-Kac Theorem, and we yield the result.

Corollary 4.1.8 The solution to the Black-Scholes PDE in Thm. 4.1.4 has the form

$$\Pi_t = F(t, S_t) = e^{-r(T-t)} E^{\mathcal{Q}}(\Phi(S_T) | \mathcal{F}_t)$$

where S_t is the spot price at time t that follows the Ito process

$$dS_t = rS_t dt + \sigma S_t dW_t^Q$$

and $\{\mathcal{F}_t\}_t$ is the filtration adapted to the Brownian motion $\{W_t\}_t$ with respect to probability measure Q.

Proof. Define functions $\mu(t,x) = rx$, $\sigma(t,x) = \sigma x$, and fix S_t to be a price at a certain fixed time t. Re-write the Black-Scholes PDE as

$$\frac{\partial F}{\partial t}(t, S_t) + \mu(t, S_t) \frac{\partial F}{\partial S}(t, S_t) + \frac{1}{2}\sigma^2(t, S_t) \frac{\partial^2}{\partial S^2} F(t, S_t) = rF(t, S_t)$$

with boundary condition

$$F(T, S_T) = \Pi_T = \Phi(S_T)$$

and then apply Corollary 4.1.7 to get

$$F(t, S_t) = e^{-r(T-t)} E^Q(\Phi(S_T)|S_t = S_t).$$

Note that the condition in the conditional expectation above is simply \mathcal{F}_t , the information set at time t, so

$$F(t, S_t) = e^{-r(T-t)} E^{\mathcal{Q}}(\Phi(S_T) | \mathcal{F}_t)$$

as required, where

$$dS_t = rS_t dt + \sigma S_t W_t^Q.$$

Theorem 4.1.9 — Solution to Black-Scholes PDE. Under the Black-Scholes model, the arbitrage-free price at time t of the derivative instrument ξ with maturity T and payoff $\xi_T = \Phi(S_T)$ is given by

$$\Pi_t = e^{-r(T-t)} \int_{-\infty}^{\infty} \Phi(e^y) f_{Y_T}(y) dy,$$

where

$$Y_T | \mathcal{F}_t \sim N\left(\log(S_t) + \left(r - \frac{\sigma^2}{2}\right)(T - t), \sigma^2(T - t)\right).$$

Proof. From Corollary 4.1.8 we inferred the Ito process of the spot price $\{S_t\}_t$ if $F(t,S_t)$ is a solution to the Black-Scholes PDE. By Def. 3.4.5, $\{S_t\}_t$ follows a geometric Brownian motion with drift rS_t and diffusion σS_t . By Thm. 3.4.6 we have

$$S_T = S_t e^{\left(r - \frac{\sigma^2}{2}\right)(T - t) + \sigma(W_T - W_t)}$$

Write $Y_T = \log(S_t) + \left(r - \frac{\sigma^2}{2}\right)(T - t) + \sigma(W_T - W_t)$ and consequently $S_T = e^{Y_T}$. Moreover because $W_T - W_t \sim N(0, T - t)$, we have

$$Y_T | \mathcal{F}_t \sim N \left(\log(S_t) + \left(r - \frac{\sigma^2}{2} \right) (T - t), \sigma^2(T - t) \right).$$

For convenience, denote

$$\tilde{\mu} = \log(S_t) + \left(r - \frac{\sigma^2}{2}\right)(T - t), \tilde{\sigma}^2 = \sigma^2(T - t).$$

It follows that

$$S_T | \mathcal{F}_t \sim \text{LogN}(\tilde{\mu}, \tilde{\sigma}^2).$$

With the distribution of $S_T | \mathcal{F}_t$ known, we can evaluate the expression in Corollary 4.1.8 to get

$$\Pi_{t} = e^{-r(T-t)} E^{\mathcal{Q}}(\Phi(S_{T})|\mathcal{F}_{t})$$

$$= e^{-r(T-t)} E^{\mathcal{Q}}(\Phi(e^{Y_{T}})|\mathcal{F}_{t})$$

$$= e^{-r(T-t)} E^{\mathcal{Q}}(\Phi(e^{Y_{T}})|S_{t})$$

$$= e^{-r(T-t)} \int_{-\infty}^{\infty} \Phi(e^{Y}) f_{Y_{T}}(y) dy$$

by the definition of expectation, $f_{Y_T}(y)$ being the density function of Y_T . This completes the proof.

Lemma 4.1.10 If $X \sim \text{LogN}(\mu, \sigma^2)$, then for all K > 0,

$$E(X\mathbf{1}_{\{X>K\}}) = E(X)N\left(\frac{\mu + \sigma^2 - \log(K)}{\sigma}\right)$$

where $N(\cdot)$ is the distribution function of the standard normal distribution.

Proof. Write $X = e^Y$ where $Y \sim N(\mu, \sigma^2)$. We have

$$f_Y(y) = \frac{1}{\sigma\sqrt{2\pi}}e^{-\frac{1}{2}\left(\frac{y-\mu}{\sigma}\right)^2}, y \in \mathbb{R},$$

to be the density function of Y. It follows that

$$E(X\mathbf{1}_{\{X>K\}}) = E(e^{Y}\mathbf{1}_{\{Y>\log(K)\}}) = \int_{\log(K)}^{\infty} e^{y} \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{y-\mu}{\sigma}\right)^{2}} dy.$$

Let $z = \frac{y-\mu}{\sigma}$. It follows that $dz = \frac{1}{\sigma}dy$ and $y = \sigma z + \mu$. Re-write the above to be

$$\begin{split} E(X\mathbf{1}_{\{X>K\}}) &= \int_{\frac{\log(K) - \mu}{\sigma}}^{\infty} e^{\mu + \sigma z} \frac{1}{\sigma \sqrt{2\pi}} e^{-\frac{1}{2}z^{2}} \sigma dz \\ &= e^{\mu} \int_{\frac{\log(K) - \mu}{\sigma}}^{\infty} \frac{1}{\sqrt{2\pi}} e^{\frac{2\sigma z - z^{2}}{2}} dz \\ &= e^{\mu} \int_{\frac{\log(K) - \mu}{\sigma}}^{\infty} \frac{1}{\sqrt{2\pi}} e^{\frac{\sigma^{2} - (z - \sigma)^{2}}{2}} dz \\ &= e^{\mu + \frac{\sigma^{2}}{2}} \int_{\frac{\log(K) - \mu}{\sigma}}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-\frac{(z - \sigma)^{2}}{2}} dz. \end{split}$$

Let $u = z - \sigma$, and thus du = dz, $z = u + \sigma$, we get

$$E(X\mathbf{1}_{\{X>K\}}) = e^{\mu + \frac{\sigma^2}{2}} \int_{\frac{\log(K) - \mu}{2} - \sigma}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-\frac{u^2}{2}} du.$$

Note that $\frac{1}{\sqrt{2\pi}} \exp\left(-\frac{u^2}{2}\right)$ is the density function of N(0,1), so

$$\begin{split} E(X\mathbf{1}_{\{X>K\}}) &= e^{\mu + \frac{\sigma^2}{2}} \left(1 - N \left(\frac{\log(K) - \mu}{\sigma} - \sigma \right) \right) \\ &= e^{\mu + \frac{\sigma^2}{2}} N \left(\sigma - \frac{\log(K) - \mu}{\sigma} \right) \\ &= e^{\mu + \frac{\sigma^2}{2}} N \left(\frac{\mu + \sigma^2 - \log(K)}{\sigma} \right) \end{split}$$

as required.

Lemma 4.1.11 If $X \sim \text{LogN}(\mu, \sigma^2)$, then for all K > 0,

$$E(X\mathbf{1}_{\{X < K\}}) = E(X)N\left(\frac{\log(K) - \mu - \sigma^2}{\sigma}\right)$$

where $N(\cdot)$ is the distribution function of the standard normal distribution.

Proof. To be completed.

Lemma 4.1.12 If $X \sim \text{LogN}(\mu, \sigma^2)$, then for all K > 0,

1.

$$E(K1_{\{X>K\}}) = KN\left(\frac{\mu - \log(K)}{\sigma}\right)$$

2.

$$E(K\mathbf{1}_{\{X < K\}}) = KN\left(\frac{\log(K) - \mu}{\sigma}\right)$$

where $N(\cdot)$ is the distribution function of the standard normal distribution.

Proof. To be completed.

Theorem 4.1.13 In the Black-Scholes model, the price of a European call option with strike K and maturity T at time t when spot is S_t is

$$c(t, S_t, K, T) = S_t N(d_1(t, S_t)) - e^{-r(T-t)} K N(d_2(t, S_t))$$

where $N(\cdot)$ is the distribution function of the standard normal distribution, r is the risk-free rate, and

$$d_1(t,S_t) = \frac{\log\left(\frac{S_t}{K}\right) + \left(r + \frac{\sigma^2}{2}\right)(T-t)}{\sigma\sqrt{T-t}},$$

$$d_2(t,S_t) = d_1(t,S_t) - \sigma\sqrt{T-t}.$$

Proof. To be completed.

Corollary 4.1.14 In the Black-Scholes model, the price of a European put option with strike K and maturity T at time t when spot is S_t is

$$p(t, S_t, K, T) = e^{-r(T-t)}KN(-d_2(t, S_t)) - S_tN(-d_1(t, S_t))$$

where $N(\cdot)$ is the distribution function of the standard normal distribution, r is the risk-free rate, and

$$d_1(t,S_t) = \frac{\log\left(\frac{S_t}{K}\right) + \left(r + \frac{\sigma^2}{2}\right)(T-t)}{\sigma\sqrt{T-t}},$$

$$d_2(t,S_t) = d_1(t,S_t) - \sigma\sqrt{T-t}.$$

Proof. To be completed.

4.2 Risk-Neutral Pricing and Girsanov Theorem

Proposition 4.2.1 Suppose the risky asset in the Black-Scholes model $\{S_t\}_t$ satisfies

$$dS_t = rS_t dt + \sigma S_t dW_t^Q$$

where Q is the probability measure in the solution of the Black-Scholes PDE in Corollary 4.1.8,

then the series

$$\{S_t e^{-rt}, t \ge 0\}$$

is a martingale under Q.

Proof. Define $Y_t = f(t, S_t) = e^{-rt}S_t$ where $f(t, x) \mapsto e^{-rt}x$ is a function, then

$$f_t(x) = \frac{\partial f}{\partial t}(x) = -re^{-rt}x, f_x(t) = \frac{\partial f}{\partial x}(t) = e^{-rt}, f_{xx}(t) = \frac{\partial^2 f}{\partial x^2}(t) = 0.$$

Since $\{S_t\}_t$ is an Ito process (Corollary 4.1.8), we have, by the Ito-Doeblin Lemma,

$$dY_t = -re^{-rt}xdt + e^{-rt}dS_t + \frac{1}{2}(0)$$

$$= -re^{-rt}xdt + e^{-rt}dS_t$$

$$= -re^{-rt}xdt + e^{-rt}(rS_tdt + \sigma S_tdW_t^Q)$$

$$= \sigma S_t e^{-rt}dW_t^Q$$

to be an Ito process as well. Put $\{g_t\}_t$ to be

$$g_t = \sigma S_t e^{-rt}$$

we note that

$$E\left(\int_0^t g_u^2 du\right) = E\left(\int_0^t \sigma^2 S_u^2 e^{-2ru} du\right) < \infty,$$

so $\{Y_t\}_t$ is a well-defined Ito process.

By Theorem 3.3.3, $\{Y_t\}_t$ is a martingale with respect to $\{\mathcal{F}_t\}_t$, which is adapted to (generated by, in fact) the Brownian motion $\{W_t^Q\}_t$. This completes the proof.

Proposition 4.2.2 If $\{V_t\}_t$ is the value process of a self-financing portfolio, then the discounted portfolio values

$$\{e^{-rt}V_t, t \ge 0\}$$

is a martingale under Q, the probability measure in the solution to to the Black-Scholes PDE.

Proof. Because $\{V_t\}_t$ is self-financing, we have

$$dV_t = h_t^B dB_t + h_t^S dS_t$$

where $dB_t = re^{rt}dt$.

Let $Y_t = e^{-rt}V_t$ and define function $f:(t,x) \mapsto e^{-rt}x$ and get f_t , f_x , and f_{xx} in similar fashion to Proposition 4.2.1. By the Ito-Doeblin Lemma, we get

$$dY_t = -re^{-rt}V_tdt + e^{-rt}dV_t$$

where $V_t = h_t^B B_t + h_t^S S_t$. Substitute this and dV_t in yields

$$dY_{t} = -re^{-rt}(h_{t}^{B}B_{t} + h_{t}^{S}S_{t})dt + e^{-rt}(h_{t}^{B}dB_{t} + h_{t}^{S}dS_{t})$$

$$= -re^{-rt}(h_{t}^{B}B_{t} + h_{t}^{S}S_{t})dt + e^{-rt}(h_{t}^{B}re^{rt}dt + h_{t}^{S}dS_{t})$$

$$= -re^{-rt}h_{t}^{S}S_{t}dt + e^{-rt}h_{t}^{S}dS_{t}$$

$$= -re^{-rt}h_{t}^{S}S_{t}dt + e^{-rt}h_{t}^{S}(rS_{t}dt + \sigma S_{t}dW_{t}^{Q})$$

$$= h_{t}^{S}e^{-rt}\sigma S_{t}dW_{t}^{Q}$$

which is, using similar reasoning as Proposition 4.2.1, an Ito process and consequently a martingale with respect to Q.

Definition 4.2.3 In a continuous-time financial model, a contingent claim with payoff $\Phi(S_T)$ at time T is **attainable** if there exists a self-financing portfolio strategy $\{h_t\}_{t\geq 0} = \{(h_t^B, h_t^S): t\geq 0\}$ with valuation process

$$V_t = h_t^B B_t + h_t^S S_t$$

such that $S_T = \Phi(S_T)$.

If all contingency claims are attainable, the market is said to be **complete**.

Remark 4.2.4 Note that we already used the content of Def. 4.2.3 in the statement of Lemma 4.1.3.

Definition 4.2.5 In a model with a riskless asset and a risky asset, a **martingale measure** is a probability measure under which the discounted expectation of the risky asset price is equal to the current risky asset price.

Theorem 4.2.6 — First Fundamental Theorem of Asset Pricing. A market model in continuous-time is arbitrage-free if and only if there exists a martingale measure.

Proof. We omit the proof.

Theorem 4.2.7 — Second Fundamental Theorem of Asset Pricing. A continuous-time arbitrage-free market model is complete if and only if the martingale measure is complete (in a measure space (X, \mathcal{B}, μ) , μ is a complete measure if for all $E \in \mathcal{B}$ such that $\mu(E) = 0$, and $F \subset E$, then $F \in \mathcal{B}$).

Proof. We omit the proof.

Theorem 4.2.8 — Risk-Neutral Valuation. Suppose we have riskless asset with price process $\{B_t\}_t$ and a risk asset with price process $\{S_t\}_t$, such that

$$dS_t = \alpha(t, S_t)dt + \sigma S_t dW_t$$

$$dB_t = r_t B_t dt, B_0 = 1$$

in an arbitrage-free model with martingale measure Q, then the price of an attainable contingency claim at time t is given by

$$F(t,S_t) = E^Q \left(\Phi(S_T) \frac{B_t}{B_T} \middle| \mathcal{F}_t \right).$$

Proof. Since the contingency claim is attainable, let $\{V_t\}_t$ be the valuation process of the replicating portfolio. Because the model is arbitrage-free, we have

$$V_t = F(t, S_t)$$

for all $t \in [0, T]$.

Because Q is a martingale measure, the process

$$\left\{ Z_t = \frac{S_t}{B_t} : t \in [0, T] \right\}$$

is a martingale, and in particular

$$E(Z_t|\mathcal{F}_s) = E\left(\frac{S_t}{B_t}\middle|\mathcal{F}_s\right) = \frac{S_s}{B_s} = Z_s = \frac{1}{B_t}E(S_t|\mathcal{F}_s).$$

Re-write $\{Z_t\}_t$ in its differential form

$$dZ_t = g_t dW_t^Q$$

for some function g_t .

Define a function $f:(t,x)\mapsto \frac{x}{B_t}$ and we get

$$f_t(x) = \frac{\partial f}{\partial t}(t, x) = \left(-\frac{x}{B_t^2}\right) \frac{\partial}{\partial t} B_t = \left(\frac{-x}{B_t^2}\right) r_t B_t$$

$$f_x(t) = \frac{\partial f}{\partial x}(t, x) = \frac{1}{B_t}$$

$$f_{xx}(t) = \frac{\partial^2 f}{\partial x^2}(t, x) = 0.$$

Applying the Ito-Doeblin lemma, we have

$$dZ_t = \frac{-1}{B_t} S_t r_t B_t dt + \frac{1}{B_t} dS_t = \frac{-r_t}{B_t} S_t dt + \frac{1}{B_t} dS_t.$$

On the other hand, if we consider $Y_t := V_t/B_t$ for $t \ge 0$, using the same f, we apply the Ito-Doeblin Lemma again to get

$$dY_t = f_t(t, Y_t)dt + f_x(t, V_t)dV_t$$

$$= \frac{-V_t}{B_t^2} r_t B_t dt + \frac{1}{B_t} dV_t$$

$$= -\frac{r_t}{B_t} (h_t^B B_t + h_t^S S_t) dt + \frac{1}{B_t} (h_t^S dS_t + h_t^B dB_t)$$

$$= -\frac{r_t}{B_t} h_t^S S_t dt + \frac{1}{B_t} h_t^S dS_t - r_t h_t^B dt + \frac{1}{B_t} h_t^B (r_t B_t dt)$$

$$= \frac{-r_t}{B_t} h_t^S S_t dt + \frac{1}{B_t} h_t^S dS_t$$

$$= h_t^S \left(\frac{-r_t}{B_t} S_t dt + \frac{1}{B_t} dS_t \right)$$

$$= h_t^S dZ_t \text{ by the previous identity}$$

$$= h_t^S q_t dW_t^Q.$$

Hence $\{Y_t\}_t$ is a martingale under Q, and by the martingale property (Def. 3.1.4(3), we have

$$E(Y_t|\mathcal{F}_s) = Y_s \text{ for all } s \leq t$$

and in particular $Y_t = E(Y_T | \mathcal{F}_t)$ for any $t \in [0, T]$. Thus

$$\frac{V_t}{B_t} = E(Y_T | \mathcal{F}_t) = \frac{F(t, S_t)}{B_t} \Rightarrow F(t, S_t) = B_t E(Y_T | \mathcal{F}_t).$$

Now $Y_T = \frac{1}{B_T}\Phi(S_T)$ because $V_T = \Phi(S_T)$, so substitution yields

$$F(t,S_t) = B_t E\left(\left.\frac{\Phi(S_T)}{B_T}\right| \mathcal{F}_t\right) = E\left(\left.\Phi(S_T)\frac{B_t}{B_T}\right| \mathcal{F}_t\right)$$

as required.

Theorem 4.2.9 — Girsanov's Theorem. Let $\{W_t\}_t$ be a standard Brownian motion on the filtered probability space $(\Omega, \{\mathcal{F}_t\}_t, \mathcal{F}, P)$ and $\{\Psi_t\}_t$ be a stochastic process such that

$$E^P\left(e^{rac{1}{2}\int_0^T \Psi_t^2 dt}
ight) < \infty$$

for some fixed T > 0. Define a process $\{L_t\}_t$ on $t \in [0,T]$ where

$$L_0 = 1, L_t = e^{\int_0^t \Phi_s dW_s^P - \frac{1}{2} \int_0^t \Phi_s^2 ds},$$

i.e. $dL_t = \Phi_t L_t dW_t^P$, then there exists a probability measure Q on Ω such that

$$L_T = \frac{dQ}{dP}$$

and

$$dW_t^P = \Phi_t dt + dW_t^Q,$$

where $\frac{dQ}{dP}$ is the Radon-Nikodym derivative between measures Q and P and $\{W_t^Q\}_t$ is the Brownian motion under Q.

Proof. We omit the proof.

Remark 4.2.10 Since $dL_t = \Psi_t L_t dW_t^P$ is a Brownian motion with no drift, it is a martingale with respect to P, and

$$L_t = E^P \left(\left. \frac{dQ}{dP} \right| \mathcal{F}_t \right).$$

We sometimes write the process as

$$L_t = \left(\frac{dQ}{dP}\right)_t$$

i.e. the Radon-Nikodym derivative is a random variable.

Corollary 4.2.11 Consider a Black-Scholes model with price processes $dB_t = rB_t dt$ and $dS_t = \mu S_t dt + \sigma S_t dW_t^P$ for the riskless and the risky assets respectively over a filtered probability space $(\Omega, (\mathcal{F}_t)_t, \mathcal{F}, P)$, then there exists a probability measure Q such that the Brownian motion with respect to Q satisfies

$$dW_t^Q = dW_t^P - \frac{r - \mu}{\sigma}.$$

Proof. By Corollary 4.1.8, the solution to the Black-Scholes PDE has the spot price following the process

$$dS_t = rS_t dt + \sigma S_t dW_t^Q$$

where Q is a result of the Feynman-Kac Theorem.

Define the process $(\Psi_t)_t$ such that $\Psi_t = \frac{r-\mu}{\sigma}$ for all t > 0. Note that

$$E^P\left(e^{\frac{1}{2}\int_0^T\frac{(r-\mu)^2}{\sigma^2}dt}\right) = E^P\left(e^{\frac{1}{2}T(t-\mu)^2/\sigma^2}\right) < \infty$$

as long as $\sigma > 0$, which is true in any risky asset prices, so by Girsanov's Theorem, there exists a probability measure Q such that

$$\frac{dQ}{dP} = e^{\int_0^t \frac{r-\mu}{\sigma} dW_s^P - \frac{1}{2} \int_0^t \frac{(r-\mu)^2}{\sigma^2} ds}$$

and $dW_t^P = \frac{r-\mu}{\sigma}dt + dW_t^Q$, as required.

Corollary 4.2.12 The Black-Scholes model is arbitrage-free.

Proof. Direct consequence of Corollary 4.2.11, Proposition 4.2.1, and the First Fundamental Theorem of Asset Pricing.

Theorem 4.2.13 The Black-Scholes model is complete.

Proof. We omit the proof.

Proposition 4.2.14 Let P be the physical probability measure in the Black-Scholes model, Q be the risk-neutral probability obtained in Corollary 4.2.11, then the expectation of S_T given S_0 and riskless rate r under Q is

$$E^Q(S_T) = S_0 e^{rT}$$
.

Proof. From the proof of Corollary 4.2.11, we let $(\Psi_t)_t$ be

$$\Psi_t = \frac{r-\mu}{\sigma}$$
.

From the statement of Girsanov's Theorem, we define $(L_t)_t = (dQ/dP)_t$, where

$$\left(\frac{dQ}{dP}\right)_T = L_T = \exp\left(\int_0^T \Psi_s dW_s^P - \frac{1}{2}\int_0^T \Psi_s^2 ds\right).$$

Now

$$E^{Q}(S_{T}) = E^{P}\left(\left(\frac{dQ}{dP}\right)_{T} S_{T}\right)$$

where $E^P(S_T) = S_0 e^{\mu T}$ since dS_t follows a geometric Brownian motion and we invoke Thm. 3.4.6 to get

$$E^{P}(S_T) = E^{P}\left(S_0 e^{\left(\mu - \frac{\sigma^2}{2}\right)T + \sigma(W_T - W_t)}\right)$$
$$= S_0 e^{\left(\mu - \frac{\sigma^2}{2}\right)T} E^{P}(e^{\sigma(W_T - W_t)})$$

where $W_T - W_0 \sim N(0,T)$ and the result follows from the moment-generating function of the normal distribution.

Substitute in L_T to get

$$\begin{split} E^{\mathcal{Q}}(S_T) &= E^P \left(\left(\frac{d\mathcal{Q}}{dP} \right)_T S_T \right) \\ &= E^P \left(\exp \left(\int_0^T \frac{r - \mu}{\sigma} dW_s^P - \frac{1}{2} \int_0^T \left(\frac{r - \mu}{\sigma} \right)^2 ds \right) \cdot S_T \right) \\ &= E^P \left(\exp \left(\frac{r - \mu}{\sigma} W_T^P - \frac{1}{2} \left(\frac{r - \mu}{\sigma} \right)^2 T \right) \cdot S_0 \exp \left(\left(\mu - \frac{\sigma^2}{2} \right) T + \sigma W_T^P \right) \right) \\ &= e^{-\frac{1}{2} \left(\frac{r - \mu}{\sigma} \right)^2 T} \cdot S_0 e^{\left(\mu - \frac{\sigma^2}{2} \right)^T} E^P \left(e^{\left(\frac{r - \mu}{\sigma} + \sigma \right) W_T^P} \right) \\ &= e^{-\frac{1}{2} \left(\frac{r - \mu}{\sigma} \right)^2 T} S_0 e^{\left(\mu - \frac{\sigma^2}{2} \right) T} \cdot e^{\frac{1}{2} T \left(\frac{r - \mu}{\sigma} + \sigma \right)^2} \text{ by moment-generating function of } N(0, T) \\ &= S_0 e^{rT} \end{split}$$

as required.

■ Example 4.2.15 Consider a continuous-time model and a forward contract with forward price K on an asset which has price process $(S_t)_{t=0}^T$.

The payoff at time T is $S_T - K$ and by definition of a forward, the value of the forward at time 0 is 0. Suppose the riskless asset has price B_T at time T, then a risk-neutral measure Q must satisfy, by Thm. 4.2.8,

$$0 = E^Q \left(S_T - \frac{K}{B_T} \right).$$

Now $\frac{S}{B}$ is a martingale under Q by the proof of Thm. 4.2.8, thus

$$0 = E^{\mathcal{Q}}\left(S_T - \frac{K}{B_T}\right) = S_0 - \frac{K}{B_T}$$

and so $K = S_0 B_T$.

If there is a riskless rate r such that $B_t = e^{rt}$, then $K = S_0 e^{rt}$, which is the same result as Proposition 1.5.4.

On the other hand, the value of the contract itself is given by

$$f_{t,T} = E^{Q} \left((S_{T} - K) \frac{B_{t}}{B_{T}} \middle| \mathcal{F}_{t} \right)$$

$$= B_{t} E^{Q} \left(\frac{S_{T}}{B_{T}} \middle| \mathcal{F}_{t} \right) - E^{Q} \left(K \frac{B_{t}}{B_{T}} \right)$$

$$= B_{t} \frac{S_{t}}{B_{t}} - S_{0} B_{T} \frac{B_{t}}{B_{T}} \text{ by martingale property}$$

$$= S_{t} - S_{0} B_{t}$$

at time $t \in [0, T]$.

- 4.3 Monte-Carlo Method for Pricing
- 4.4 Implies Volatility
- 4.5 The Greeks
- 4.6 Hedging

5. Continuous-Time Interest Rate Models

5.1 Bonds and Interest Rates

Definition 5.1.1 A zero-coupon bond with maturity T is called a T-bond. The price of the T-bond at time t is denoted p(t,T), $0 \le t \le T$. We assume p(t,T) is a differentiable function with respect to t.

Definition 5.1.2 The **short rate** at time $t \in [0,T]$, denoted r(t), is a random variable which represents the continuously compounded interest rate at which one can borrow or lend for an infinitesimal amount of time Δt at time t.

Definition 5.1.3 At time $t \in [0, T]$, the [S, T] **LIBOR forward rate** for some $S \in [t, T]$, denoted L(t; S, T), is the simple interest on \$1 invested at time S till time T.

Proposition 5.1.4 Suppose $0 \le t \le S \le T$, then the LIBOR forward rate is

$$L(t; S, T) = -\frac{p(t, T) - p(t, S)}{(T - S)p(t, T)}$$

and in particular

$$L(S;S,T) = -\frac{p(t,T)-1}{(T-S)p(t,T)}.$$

Proof. By the definition of L(t; S, T) we have the payoff of \$1, invested at time S to be

$$1 + (T - S)L(t; S, T)$$

at time T. On the other hand, we can pay p(t,S) at time t to enter into a T-bond. This yields \$1 at time S, and becomes p(t,S)/p(t,T) at time T. Thus we have the equation

$$1 + (T - S)L(t; S, T) = \frac{p(t, S)}{p(t, T)}$$

and equivalently

$$L(t; S, T) = \frac{\frac{p(t, S)}{p(t, T)} - 1}{T - S} = -\frac{p(t, T) - p(t, S)}{(T - S)p(t, T)}.$$

For the second result, simply substitute in p(S,S) = 1 and get

$$L(S;S,T) = -\frac{p(t,T)-1}{(T-S)p(t,T)}.$$

Definition 5.1.5 The **simple spot rate** or the **LIBOR spot rate** for [S,T] is the L(S;S,T) in Proposition 5.1.4 above.

Definition 5.1.6 At time $t \in [0,T]$, the **continuously compounded forward rate for** [S,T] **contracted at** $t \leq S$ is the interest on \$1 invested at S till time T, assuming that interest is continuously compounded, denoted R(t;S,T).

The **continuously compounded spot rate**, or the **zero-coupon yield**, is R(S;S,T), or denoted R(S,T).

Proposition 5.1.7 We have

$$R(t; S, T) = -\frac{\log(p(t, T)) - \log(p(t, S))}{T - S}$$

and

$$R(S;S,T) = R(S,T) = -\frac{\log(p(t,T))}{T-S}.$$

Proof. Consider two strategies at time t. The first strategy enters a contract that invests \$1 at time S. By definition, the \$1 will grow at rate R(t;S,T) for the period of T-S. The second strategy invests p(t,S) at time t to enter a T-bond. This yields \$1 at time S, and becomes p(t,S)/p(t,T) at time T. This these two strategies both effectively have investment of \$1 at time S, their payoff at time T are equal:

$$e^{R(t;S,T)(T-S)} = \frac{p(t,S)}{p(t,T)}.$$

It follows that

$$\begin{split} R(t;S,T) &= \frac{1}{T-S} \log \left(\frac{p(t,S)}{p(t,T)} \right) \\ &= \frac{1}{T-S} (\log(p(t,S)) - \log(p(t,T))) \\ &= -\frac{\log(p(t,T)) - \log(p(t,S))}{T-S} \end{split}$$

as required. If t = S, then p(S,S) = 1 and $\log p(S,S) = 0$, and the rest is immediate.

Definition 5.1.8 The instantaneous forward rate with maturity T contracted at time t is

$$f(t,T) = -\frac{\partial}{\partial T} \log p(t,T),$$

while the **instantaneous short rate** at time $t \in [0, T]$ is

$$r(t) = f(t,t) = -\lim_{\Delta \to 0} \frac{\log(p(t,t+\Delta)) - \log(p(t,t))}{\Delta}.$$

Remark 5.1.9 For an infinitismal Δ we have

$$e^{r(t)\Delta} \approx \frac{p(t,t)}{p(t,t+\Delta)} = \frac{1}{p(t,t+\Delta)}.$$

Definition 5.1.10 The money market process, or the risk-free asset, for $t \in [0,T]$, is

$$\beta(t) = \exp\left(\int_0^t r(s)ds\right)$$

where $r(\cdot)$ is the instantaneous short rate function.

Proposition 5.1.11 A zero-coupon *T*-bond at time *t* has price

$$p(t,T) = p(t,S) \exp\left(-\int_S^T f(t,u) du\right)$$
 for all $S \le T$

and

$$p(t,T) = \exp\left(-\int_{t}^{T} f(t,s)ds\right).$$

Proof. The instantaneous forward rate f(t, u) can be written as

$$f(t, u) = -\frac{\partial}{\partial u} \log p(t, u).$$

Hence, by the Fundamental Theorem of Calculus,

$$\int_{S}^{T} f(t, u) du = -\log p(t, T) + \log p(t, S)$$

$$\Rightarrow \int_{S}^{T} f(t, u) du = -\log \left(\frac{p(t, T)}{p(t, S)}\right)$$

and we have the result.

The second equation is the consequence of the fact that p(t,t) = 1.

Definition 5.1.12 Define a function

$$y: [0, \infty) \times \mathbb{R} \to \mathbb{R}$$

 $(t, T) \mapsto R(t, T)$

where

$$R(t,T) = -\frac{\log p(t,T)}{T-t}.$$

R(t,T) is the **zero-coupon bond yield**. The map (t,y(t,T)) is the **yield curve of zero-coupon bonds**, and the function y is the **yield function**.

Remark 5.1.13 Note that

$$p(t,T) = e^{-y(t,T)(T-t)}$$

and the payoff, at time T, of \$1 invested in a zero-coupon bond with maturity T, is

$$\frac{1}{p(t,T)} = e^{y(t,T)(T-t)}.$$

This shows that $y(\cdot)$ is the rate of guaranteed return on the zero-coupon bond.

5.2 Zero-Coupon Bond Pricing and the Term Structure Equation

Definition 5.2.1 The **short rate model** has the following setup:

1. The short rate at time t follows the process

$$dr(t) = \mu(t, r(t))dt + \sigma(t, r(t))dW_t^P$$

for some drift function μ , diffusion function σ , and W_t^P , the standard Brownian motion with respect to a physical probability measure P.

2. There is a risk-free asset whose price follows

$$d\beta(t) = r(t)\beta(t)dt$$

where $\beta(t)$ is a money-market process.

- 3. For all T > 0, there exists a market for the T-bond.
- 4. For all T > 0, the price of the T-bond at time t is of the form

$$p(t,T) = F(t,r(t),T),$$

which can also be written as

$$p(t,T) = F^{T}(t,r), t \in [0,T]$$

where F is a smooth function and satisfies the boundary condition $F^{T}(T,r) = 1$.

Proposition 5.2.2 Under the short rate model, the dynamics of the *T*-bond price is

$$dF^{T} = F^{T} \alpha_{T}(t) dt + F^{T} \sigma_{T}(t) dW_{t}^{P}$$

where

$$\begin{split} \alpha_T(t) &= \frac{1}{F^T} \left(F_t^T + \mu F_r^T + \frac{1}{2} \sigma^2 F_{rr}^T \right) \\ F_t^T &= \frac{\partial F^T}{\partial t}, F_r^T = \frac{\partial F^T}{\partial r}, F_{rr}^T = \frac{\partial^2 F^T}{\partial r \partial r}, \\ \sigma_T(t) &= \frac{\sigma F_r^T}{F^T}, \end{split}$$

 μ and σ being the drift and diffusion of the short rate process r(t) respectively.

Proof. Write $Y_t = F^T(t, r(t))$ where

$$dr(t) = \mu dt + \sigma dW_t^P$$

as defined in the short rate model and use the Ito-Doeblin Lemma to get

$$dY_t = dF^T(t, r(t))$$

$$= \left(F_t^T + F_r^T \mu + \frac{1}{2} F_{rr}^T \sigma^2\right) dt + F_r^T \sigma dW_t^P$$

$$= F^T \left(\frac{1}{F^T}\right) \left(F_t^T + F_r^T \mu + \frac{1}{2} F_{rr}^T \sigma^2\right) dt + F^T \left(\frac{F_r^T \sigma}{F^T}\right) dW_t^P$$

$$= F^T \alpha_T(t) dt + F^T \sigma_T(t) dW_t^P$$

as required.

Proposition 5.2.3 Let α_T and σ_T be the drift and diffusion of the T-bond price process in the short rate model, then, if the bond market is arbitrage-free, then there exists a stochastic process $\{\lambda(t): t \geq 0\}$ such that

$$\lambda(t) = \frac{\alpha_T(t) - r(t)}{\sigma_T(t)}$$
 for all $t \ge 0$.

Proof. We omit the proof.

Definition 5.2.4 In an arbitrage-free bond market under the short rate model, the stochastic process

$$\lambda(t) = \frac{\alpha_T(t) - r(t)}{\sigma_T(t)}$$
 for all $t \ge 0$

where $\alpha_T(t)$ and $\sigma_T(t)$ are the drift and diffusion of the *T*-bond price process, is called the **market price of risk**.

Remark 5.2.5 In an arbitrage-free bond market under the short rate model, T-bonds of all maturities should have the same market price of risk. The substitution of $\alpha_T(t)$ and $\sigma_T(t)$ from Proposition 5.2.2 into Def. 5.2.4 yields the following formal definition.

Definition 5.2.6 Let $F^T(t, r(t))$ be a T-bond price process in a short rate model with the usual definitions of μ and σ , then the **term structure equation** is as follows:

$$\begin{cases} F_t^T + (\mu - \lambda \sigma) F_r^T + \frac{1}{2} \sigma^2 F_{rr}^T - r F^T &= 0 \\ F^T(T, r) &= 1 \end{cases}$$

for some real-valued process $\{\lambda(t): t \geq 0\}$. Note that both μ and σ are also functions of t.

Theorem 5.2.7 In an arbitrage-free bond market under the short rate model, the zero-coupon bond prices p(t,T) are given by

$$p(t,T) = F^{T}(t,r) = E^{Q}\left(\exp\left(-\int_{t}^{T} r(s)ds\right)\middle|\mathcal{F}_{t}\right)$$

where Q is a probability measure such that

$$dr(s) = (\mu - \sigma \lambda)ds + \sigma dW_s^Q$$

with initial condition $r(t) \in \mathbb{R}$.

Proof. Direct application of Feynman-Kac Theorem.

Corollary 5.2.8 The time-*t*-risk-neutral price of an interest-rate-contingent claim X paying $X(T) = \Phi(r(T))$ at maturity T is given by

$$X(t) = E^{Q}\left(\Phi(r(T))\exp\left(-\int_{t}^{T} r(s)ds\right) \middle| \mathcal{F}_{t}\right)$$

where the Q-dynamic of the short rate is given by

$$dr(s) = (\mu = \lambda \sigma)ds + \sigma dW_s^Q$$

$$r(t) = r \in \mathbb{R}.$$

Proof. We omit the proof.

Proposition 5.2.9 The relationship between probability measures P and Q in the short rate model and the risk-neutral valuation formula is

$$dW^{Q}(t) = dW^{P}(t) + \lambda dt$$

where λ is the market price of risk.

Proof. Under *P* we have

$$dr(t) = \mu dt + \sigma dW_t^P$$

by Def. 5.2.1, while under Q we have

$$dr(t) = (\mu - \lambda \sigma)dt + \sigma dW_t^Q.$$

By equating the two expressions, we have

$$dW_t^P + \lambda dt = dW_t^Q$$

as required.

Corollary 5.2.10 Under the probability measure Q yielded in Corollary 5.2.8, the price of T-bonds under the short rate model follows the dynamic

$$dF^{T} = r(t)F^{T}dt + F^{T}\sigma_{T}(t)dW_{t}^{Q}.$$

Proof. By Proposition 5.2.2 we have

$$dF^{T} = \alpha_{T}(t)F^{T}dt + F^{T}\sigma_{T}(t)dW_{t}^{P}.$$

Substitute in Proposition 5.2.9 to get

$$dF^{T} = F^{T} \alpha_{T}(t) dt + F^{T} \sigma_{T}(t) (dW_{t}^{Q} - \lambda dt).$$

Now
$$\lambda(t) = (\alpha_T(t) - r(t))/\sigma_T(t)$$
, so

$$dF^{T} = F^{T} \alpha_{T}(t)dt + F^{T} \sigma_{T}(t) \left(dW_{t}^{Q} - \frac{\alpha_{T}(t) - r(t)}{\sigma_{T}(t)} dt \right)$$

$$= F^{T} \alpha_{T}(t)dt + F^{T} \sigma_{T}(t)dW_{t}^{Q} - F^{T}(\alpha_{T}(t) - r(t))dt$$

$$= F^{T} r(t)dt + F^{T} \sigma_{T}(t)dW_{t}^{Q}$$

as required.

Corollary 5.2.11 Under the risk-neutral probability Q, the process

$$\left\{ \frac{F^T(t)}{\beta(t)}, t \ge 0 \right\}$$

in the short rate model is a martingale.

Proof. Define a function

$$f: [0,T] \times \mathbb{R} \to \mathbb{R}$$

 $(t,x) \mapsto \frac{x}{t}$

and define $Y_t := f(t, F^T)$.

It follows that

$$f_t = (-1)(\beta(t))^{-2} \left(\frac{d\beta}{dt}\right) (F^T)$$

where $d\beta(t) = r(t)\beta(t)dt$ by the assumption of the short rate model, so

$$f_t(t, F^T) = \frac{-1}{\beta(t)^2} r(t) \beta(t) F^T = -\frac{r(t)}{\beta(t)} F^T.$$

Also,

$$f_x(t, F^T) = \frac{1}{\beta(t)}, f_{xx}(t, F^T) = 0.$$

By the Ito-Doeblin Lemma,

$$dY_t = f_t(t, F^T)dt + f_x(t, F^T)dF^T$$

$$= -\frac{r(t)}{\beta(t)}F^Tdt + \frac{1}{\beta(t)}dF^T$$

$$= -\frac{r(t)}{\beta(t)}F^Tdt + \frac{1}{\beta(t)}(r(t)F^Tdt + F^T\sigma_T(t)dW_t^Q)$$

$$= F^T\sigma_T(t)dW_t^Q$$

 dY_t has no drift term, so by Thm. 3.3.3, Y_t is a martingale. By the construction of $\{Y_t\}_t$, this completes the proof.

5.3 Martingale Models for the Short Rate

Definition 5.3.1 Let Q be the risk-neutral measure under a short rate model. The following are models for the short rate under Q:

1. The Varsicek model states

$$dr = (b - ar)dt + \sigma dW_t^Q, a > 0.$$

2. The Cox-Ingersoll-Ross (CIR) model states

$$dr = a(b-r)dt + \sigma \sqrt{r}dW_t^Q.$$

3. The **Dothan model** states

$$dr = ardt + \sigma r dW_t^Q$$
.

4. The **Ho-Lee model** states

$$dr = \Omega(t)dt + \sigma dW_t^Q$$

where Ω is a function of time t chosen such that the model fits the original term structure.

5. The **Black-Derman-Toy model** states

$$dr = \Omega(t)rdt + \sigma(t)rdW_t^Q$$

where Ω and σ are both functions of time t.

6. The Hull-White one-factor model states

$$dr = (\Omega(t) - a(t)r)dt + \sigma(t)dW_t^Q, a(t) > 0,$$

where Ω , a, and σ are functions of time t.

7. The Hull-White two-factor model states

$$dr = (\Omega(t) - a(t)r)dt + \sigma(t)\sqrt{r}dW_t^Q, a(t) > 0,$$

where Ω , a, and σ are functions of time t. In general, these models specify the dynamic for

$$dr = \tilde{\mu}(t, r(t); \alpha)dt + \sigma(t, r(t); \alpha)dW_t^Q$$

under probability measure Q for some functions $\tilde{\mu}$ and σ of time t, where α can be a vector or a function, and contains additional parameters for $\tilde{\mu}$.

Definition 5.3.2 Given a particular model for the short rate *r*:

$$dr = \tilde{\mu}(t, r(t); \alpha)dt + \sigma(t, r(t); \alpha)dW^Q$$

the **"inverting the yield curve" procedure** for estimating α is as follows:

1. Solve, for every maturity T, the term structure equation

$$\begin{cases} F_t^T + \tilde{\mu}F_r^T + \frac{1}{2}F_{rr}^T - rF^T &= 0 \\ F^T(T, r) &= 1 \end{cases}$$

to obtain the zero-coupon prices $p(0,T;\alpha)$ for $T \ge 0$. Denote the results as $\{T,p(0,T;\alpha)\}$, the **theoretical term structure**.

- 2. Collect empirical data from the bond market for all maturities T. Denote this $\{T, p^*(0,T)\}$, the **empirical term structure**.
- 3. Choose α such that $\{T, p(0,T;\alpha)\}$ fits $\{T, p^*(0,T)\}$ based on some fitting criteria. Denote the optimal parameter to be α^* .
- 4. Insert α^* into $\tilde{\mu}$ and σ , and denote the estimated functions to be μ^* and σ^* .
- 5. Use

$$dr = \mu^*(t, r(t); \alpha^*)dt + \sigma^*(t, r(t); \alpha^*)dW^Q$$

to compute prices of interest rate derivatives with the methodology of Corollary 5.2.8.

Definition 5.3.3 Suppose $\{p(t,T): t \in [0,T], T \ge 0\}$ is a term structure of zero-coupon bond prices and each p(t,T) has the form

$$p(t,T) = F(t,r(t),T)$$

where

$$F(t,r(t),T) = e^{A(t,T)-B(t,T)r}$$

and A and B are deterministic functions, then the model of the bond prices is said to possess an **affine term structure (ATS)**.

Definition 5.3.4 For a short rate model

$$dr = \tilde{\mu}dt + \sigma dW_t^Q$$

under some probability measure Q, the model is **mean-reverting** is the mean of r tends to a constant level in the long run.

Proposition 5.3.5 Suppose the short rate follows a dynamic

$$dr = \tilde{\mu}(t, r(t); \alpha)dt + \sigma(t, r(t); \alpha)dW^Q$$

where $\tilde{\mu}$ and σ have the form

$$\begin{cases} \tilde{\mu}(t,r(t)) &= \alpha(t)r + \beta(t) \\ \sigma(t,r(t)) &= \sqrt{\gamma(t)r + \delta(t)}, \end{cases}$$

then the model admits an affine term structure where A and B satisfy the system

$$\begin{cases} B_t(t,T) + \alpha(t)B(t,T) - \frac{1}{2}\gamma(t)B^2(t,T) &= -1\\ B(T,T) &= 0, \end{cases}$$

$$\begin{cases} A_t(t,T) &= \beta(t)B(t,T) - \frac{1}{2}\delta(t)B^2(t,T) \\ A(T,T) &= 0. \end{cases}$$

Proof. We omit the proof.

Corollary 5.3.6 The Vasicek model for short rate

$$dr = (b - ar)dt + \sigma dW_t^Q$$

possess affine term structure

$$p(t,T) = e^{A(t,T) - B(t,T)r(t)}$$

where

$$A(t,T) = \frac{(B(t,T) - T + t) \left(ab - \frac{1}{2}\sigma^2\right)}{a^2} - \frac{\sigma^2 B^2(t,T)}{4a}$$
$$B(t,T) = \frac{1}{a}(1 - e^{-a(T-t)}.$$

Proof. By Proposition 5.3.5, we solve for A and B in systems

$$\begin{cases} B_t(t,T) + (-a)B(t,T) &= -1 \\ B(T,T) &= 0, \end{cases}$$

$$\begin{cases} A_t(t,T) &= bB(t,T) - \frac{1}{2}\sigma^2 B^2(t,T) \\ A(T,T) &= 0. \end{cases}$$

The first system easily yields

$$B(t,T) = \frac{1}{a}(1 - e^{-a(T-t)}),$$

and integrating the second system yields

$$A(t,T) = \frac{\sigma^2}{2} \int_t^T B^2(s,T) ds - b \int_t^T B(s,T) ds,$$

and substitution yields

$$A(t,T) = \frac{\sigma^2}{2} \int_t^T \frac{1}{a^2} (1 - e^{-a(T-s)})^2 ds - b \int_t^T \frac{1}{a} (1 - e^{-a(T-s)}) ds.$$

Integrating this yields the result.

Remark 5.3.7 1. The Vasicek model is mean-reverting with mean of r to be $\frac{b}{a}$.

- 2. In the Vasicek, Ho-Lee, and Hull-White one-factor models, the integral $\int r(s) ds$ has a normal distribution
- 3. Hence, the short rate *r* in the Vasicek model might become negative.

Corollary 5.3.8 Suppose the short rate follows a Ho-Lee model

$$dr = \Omega(t)dt + \sigma dW_t^Q$$

where, at t = 0, $\Omega(t)$ follows

$$p(0,T) = p^*(0,T)$$
, the observed T-bond price,

and

$$\Omega(t) = \frac{\partial f^*}{\partial T}(0, t) + \sigma^2 t$$

where $f^*(0,t)$ denotes the observed forward rates. Then, the T-bond price has closed-form expression

$$p(t,T) = \frac{p^*(0,T)}{p(0,t)} \exp\left((T-t)f^*(0,t) - \frac{\sigma^2}{2}t(T-t)^2 - (T-t)r(t) \right).$$

Proof. We omit the proof.

Corollary 5.3.9 Suppose the short rate follows a CIR model

$$dr = a(b-r)dt + \sigma \sqrt{r}dW_t^Q,$$

then the bond prices are given by

$$F^{T}(t,r) = A_0(T-t)e^{-B(T-t)r},$$

where

$$B(x) = \frac{2(e^{\gamma x} - 1)}{(\gamma + \alpha)(e^{\gamma x} - 1) + 2\gamma}$$

$$A_0(x) = \left(\frac{2\gamma e^{(a+\gamma)(x/2)}}{(\gamma + a)(e^{\gamma x} - 1) + 2\gamma}\right)^{2ab/\sigma^2}$$

$$\gamma = \sqrt{a^2 + 2\sigma^2}.$$

Proof. We omit the proof.

Remark 5.3.10 In the CIR model, there is mean reversion for p(t,T), r is always positive, the volatility of r depends on r, and r follows a chi-squared distribution, but we are unable to fir the time-0 observed prices of bonds.

Definition 5.3.11 A **European bond call option** with strike K and expiration S on a T-bond where S < T gives the option holder the right but not the obligation to buy a T-bond at K at time S.

A **European bond put option** with strike K and expiration S on a T-bond where S < T gives the option holder the right but not the obligation to sell a T-bond at K at time S.

Proposition 5.3.12 Suppose the short rate follows the Vasicek model

$$dr = (b - ar)dt + \sigma dW_t^Q, a > 0,$$

then the price of a European bond call option with strike K, expiration S, on a T-bond at time $t \in [0, S]$ is

$$c(t, S, K, T) = p(t, T)N(d) - p(t, S)KN(d - \sigma_p)$$

where

$$\begin{split} d &= \frac{1}{\sigma_p} \log \left(\frac{p(t,T)}{p(t,S)K} \right) + \frac{1}{2} \sigma_p, \\ \sigma_p &= \frac{1}{a} (1 - e^{-a(T-S)}) \sqrt{\frac{\sigma^2}{2a} (1 - e^{-2a(T-t)})}, \end{split}$$

and $N(\cdot)$ is the distribution function of the standard normal distribution.

Proof. We omit the proof.

Proposition 5.3.13 Suppose the short rate follows the Ho-Lee model

$$dr = \Omega(t)dt + \sigma dW_t^Q,$$

where

$$\Omega(t) = \frac{\partial f^*}{\partial T}(0, t) + \sigma^2 t,$$

then the price of a European call option with strike K, expiration S on a T-bond at time $t \in [0,S]$ is

$$c(t, S, K, T) = p(t, T)N(d) - p(t, S)KN(d - \sigma_p),$$

where

$$d = \frac{1}{\sigma_p} \log \left(\frac{p(t, T)}{p(t, S)K} \right) + \frac{1}{2} \sigma_p,$$

$$\sigma_p = \sigma(T - S) \sqrt{S - t},$$

and $N(\cdot)$ is the distribution function of the standard normal distribution.

Proof. We omit the proof.



	A	Continuously compounded spot rate 38
	see Term structure	
	В	Diffusion 16 Dothan model 44
Black-Derman-Toy model44Black-Scholes model23Arbitrage opportunity23Arbitrage-free34Call option price29Partial differential equation25Physical probability measure23Put option price29Solution27Valuation process23		Fernman-Kac Theorem 25 Filtered probability space 10 Adapted 10 Filtration 10 Generated by random variables 11 On standard Brownian motion 11 Forward 5
Bounded Brownia Arit	otions 47 d variation 13 an motion 11 thmetic 18 fusion 11	Forward Price
	ft	
	С	Girsanov's Theorem
Complet	del see Cox-Ingersoll-Ross model te market	Ho-Lee model
('Ontinu	ously compounded forward rate 38	Hull-White model 44

50 INDEX

1	Term structure
Independent σ-algebras 12 Information set 9 Instantaneous forward rate 39 Instantaneous short rate 39 Inverting the yield curve 44 Ito integral 14 Expectation 15 Martingale property 15 Ito isometry 15 Ito process 16 Differential form 16 Integral form 16 Ito-Doeblin Lemma 17	Affine 45 Empirical 44 Theoretical 44 Term structure equation 41 Total variation 13 U U Unbounded variation 13 V Vasicek model Y 44
LIBOR forward rate	Yield Curve of zero-coupon bonds
Market price of risk	Zero-coupon yield
Quadratic variation	
R	
Risk-free asset	
S	
Short rate 37 Mean-reverting 45 Model 40 Simple spot rate 38 Square-integrable process 14	
T	
T-bond	