



Covariation-derived residue contacts in *ab initio* modelling and Molecular Replacement

Felix Simkovic

Thesis submitted in accordance with the requirements of the
University of Liverpool
for the degree of
Doctor in Philosophy

September 2018

Institute of Integrative Biology
University of Liverpool
United Kingdom

Contents

List of Figures	iii
List of Tables	iv
List of Equations	v
List of Abbreviations	vii
1 Introduction	1
2 Materials & Methods	3
3 Evolutionary covariance in <i>ab initio</i> structure prediction-based Molecular Replacement	5
4 Evaluation of ROSETTA distance-restraint energy functions on contact-guided <i>ab initio</i> structure prediction	7
5 Alternative <i>ab initio</i> structure prediction algorithms for AMPLE	9
6 Decoy subselection using contact information to enhance MR search model creation	11
7 Protein fragments as search models in Molecular Replacement	13
8 Conclusion & Outlook	15
8.1 Conclusion	16
8.2 Outlook	17
A Appendix	19
Bibliography	21

List of Figures

List of Tables

List of Equations

List of Abbreviations

MR Molecular Replacement

Chapter 1

Introduction

Chapter 2

Materials & Methods

Chapter 3

Evolutionary covariance in *ab initio* structure prediction-based Molecular Replacement

Chapter 4

Evaluation of ROSETTA distance-restraint energy functions on contact-guided *ab initio* structure prediction

Chapter 5

Alternative *ab initio* structure prediction algorithms for AMPLE

Chapter 6

Decoy subselection using contact information to enhance MR search model creation

Chapter 7

Protein fragments as search models in Molecular Replacement

Chapter 8

Conclusion & Outlook

8.1 Conclusion

The successful disentanglement of direct and indirect residue contacts in contact prediction revolutionised many aspects of Structural Bioinformatics research [1]. Successful applications of contact information range from accurately defining domain boundaries [2] to identifying druggable protein-protein interfaces [3]. Although many such applications have been highlighted over the last few years [1], few concerned the topic of Molecular Replacement (MR) in X-ray crystallography. In this thesis, work was presented that made first attempts to apply contact information to explore some of its application spectrum in MR.

The use of contact information in *ab initio* protein structure prediction allowed researchers to predict the structure of many previously unknown protein folds based on their sequence alone [e.g., 4–12]. The major benefit of adding such information was to reduce the conformational search space, which allowed more challenging folds to be sampled correctly. Work presented in Chapters 3 to 5 further confirm such findings. More importantly, the presented results highlight that the modelling algorithm ROSETTA is very sensitive to the way contact information is introduced into the ROSETTA folding protocol. Two important examples include the up-weighting of β -strand contacts and the choice of energy function used to “reward” satisfied contacts. Furthermore, work in Chapter 5 highlights that fragment-based structure prediction algorithms may no longer be essential for accurate structure prediction. CONFOLD2, a fragment-independent algorithm, predicts the protein structure using secondary structure and contact information alone, which provided models of comparable accuracy to the state-of-the-art ROSETTA.

Beyond the prediction of protein structures, a major focus of the presented research centred on the benefit of such improved structure predictions in unconventional MR. In-line with prior expectations, better structure predictions yield more MR structure solutions. In particular, previous weaknesses of the AMPLE routine — a target’s chain length and fold — can be partially addressed with contact-guided structure predictions. Some examples for which structure solutions were obtained exceed 200 residues in chain length, whilst many others contain large proportions of β -structure. Nevertheless, simply adding contact information to *ab initio* protein structure prediction is not sufficient to solve all trialled targets. In part, such limitation is incurred through the lack of precision of contact information, which depends significantly on the availability of divergent homologous sequences. Furthermore, further research is required to be address new limitations in AMPLE by means of suboptimal processing of much more accurate structure predictions. One such example, outlined in Chapter 6, explored the incorporation of contact information in the AMPLE processing pipeline to address the latter issue. Contact information was used to estimate the similarity of a predicted decoy to its native structure, by means of scoring its long-range contact satisfaction. Exclusion of the worst decoys by this metric prior to clustering allowed more fine-grain

sampling in AMPLE, which turned unsuccessful decoy sets into ones from which the native structure can be elucidated. However, key examples presented in Chapters 3 to 5 also highlight that further developments in MR-related software are required to enable the automatic detection and subsequent processing of AMPLE ensemble search models that are correctly placed but by current metrics cannot be identified as correct structure solutions.

A further topic of research concerned the use of supersecondary structure elements or subfolds as MR search models. The default mode in AMPLE currently relies on computationally expensive *ab initio* structure predictions. Since contact predictions have reached sufficient quality for protein families with many known sequences, such information could be used to identify matching subfolds in other, unrelated protein structures. In Chapter 7, a new hybrid approach demonstrated the successful implementation of such an idea. Although imperfect at this stage, several examples highlighted the successful identification of such subfolds and subsequently successful MR structure solution. Tied to this idea may also be recent research that attempts to identify subfolds by means of matching a predicted contact map to those extracted from protein structures [11, 13].

8.2 Outlook

In this thesis first applications of predicted contact information in MR were presented. Despite the already promising results, this area of research is still in its infancy and a great number of potentially promising routes remain unexplored [1].

Earlier studies by Rigden [14] and Sadowski [2] demonstrated the successful application of residue contacts to identify domain boundaries. Although unexplored to-date, precise domain boundary predictions could be applied for better domain boundary definitions in *ab initio* structure prediction to avoid sampling of terminal loops and linkers, and thus improve protein structure prediction quality. Furthermore, contact information was used to improve the AMPLE ensemble-generation pipeline with respect to identifying poorly predicted decoys. However, the AMPLE ensemble-generation pipeline might additionally benefit from contact information to aid the driving of the truncation procedure. For example, contact data could be used to rank individual residues by their contribution to a contact network, similar to [15], and truncation driven by the rank order. Additionally, contact prediction might be used in the context of identifying alternative conformational states [16–20], which AMPLE could exploit to identify conserved residues between both states and truncate to this conserved core, or attempt remodelling after successful disentanglement of state-dependent contact pairs and try both conformations separately as ensemble search models.

Simkovic et al. [1] outlined many further such applications of predicted contact

information in the field of Structural Biology. Ultimately, the precision of contact information improves daily with the increasing depth of sequence databases, thus enabling further and further applications with more precise outcomes. Furthermore, many more research groups start to identify the value in using contact information in their own studies, and by means of pushing the boundaries new tools and applications are most likely going to emerge. Therefore, the work presented in this thesis may enable other researchers to already answer questions, most likely in terms of MR structure solutions by unconventional MR, that were previously intractable.

Appendix A

Appendix

Bibliography

- [1] F. Simkovic, S. Ovchinnikov, D. Baker, D. J. Rigden, *IUCrJ* **May 2017**, *4*, 291–300.
- [2] M. I. Sadowski, en, *Proteins: Struct. Funct. Bioinf.* **Feb. 2013**, *81*, 253–260.
- [3] F. Bai, F. Morcos, R. R. Cheng, H. Jiang, J. N. Onuchic, en, *Proceedings of the National Academy of Sciences* **Dec. 2016**, *113*, E8051–E8058.
- [4] D. S. Marks, L. J. Colwell, R. Sheridan, T. A. Hopf, A. Pagnani, R. Zecchina, C. Sander, en, *PLoS One* **Dec. 2011**, *6*, e28766.
- [5] M. Michel, S. Hayat, M. J. Skwark, C. Sander, D. S. Marks, A. Elofsson, en, *Bioinformatics* **Sept. 2014**, *30*, i482–8.
- [6] T. Kosciolk, D. T. Jones, en, *PLoS One* **Mar. 2014**, *9*, e92197.
- [7] S. Ovchinnikov, L. Kinch, H. Park, Y. Liao, J. Pei, D. E. Kim, H. Kamisetty, N. V. Grishin, D. Baker, en, *Elife* **Sept. 2015**, *4*, e09248.
- [8] S. Ovchinnikov, D. E. Kim, R. Y.-R. Wang, Y. Liu, F. DiMaio, D. Baker, en, *Proteins* **Sept. 2016**, *84 Suppl 1*, 67–75.
- [9] M. Michel, D. Menéndez Hurtado, K. Uziela, A. Elofsson, *Bioinformatics* **2017**, *33*, i23–i29.
- [10] S. H. P. de Oliveira, E. C. Law, J. Shi, C. M. Deane, en, *Bioinformatics* **Nov. 2017**, DOI 10.1093/bioinformatics/btx722.
- [11] S. Ovchinnikov, H. Park, N. Varghese, P.-S. Huang, G. A. Pavlopoulos, D. E. Kim, H. Kamisetty, N. C. Kyrpides, D. Baker, en, *Science* **Jan. 2017**, *355*, 294–298.
- [12] S. Wang, S. Sun, Z. Li, R. Zhang, J. Xu, en, *PLoS Comput. Biol.* **Jan. 2017**, *13*, e1005324.
- [13] D. W. A. Buchan, D. T. Jones, *Bioinformatics* **Sept. 2017**, *33*, 2684–2690.
- [14] D. J. Rigden, en, *Protein Eng.* **Feb. 2002**, *15*, 65–77.
- [15] D. J. Parente, J. C. J. Ray, L. Swint-Kruse, en, *Proteins* **Dec. 2015**, *83*, 2293–2306.
- [16] T. A. Hopf, L. J. Colwell, R. Sheridan, B. Rost, C. Sander, D. S. Marks, en, *Cell* **June 2012**, *149*, 1607–1621.
- [17] B. Jana, F. Morcos, J. N. Onuchic, *Phys. Chem. Chem. Phys.* **2014**, *16*, 6496–6507.
- [18] P. Sfriso, M. Duran-Frigola, R. Mosca, A. Emperador, P. Aloy, M. Orozco, en, *Structure* **Jan. 2016**, *24*, 116–126.

- [19] F. Morcos, B. Jana, T. Hwa, J. N. Onuchic, en, *Proc. Natl. Acad. Sci. U. S. A.* **Dec.** **2013**, *110*, 20533–20538.
- [20] L. Sutto, S. Marsili, A. Valencia, F. L. Gervasio, en, *Proc. Natl. Acad. Sci. U. S. A.* **Nov.** **2015**, *112*, 13567–13572.