

Reconstruct Glacier Surface Velocity from Landsat Imagery 1984-2018



Surge Type Glacier Identification on Northeast Spitsbergen, Svalbard

NEGRIBREEN, SPITSBERGEN

<http://www.geonova.no/diaries/norway-from-above/norway-from-above-preview-spitsbergen/>

Supervisor: Rickard Pettersson
Presenter: Shunan Feng


Surge Type Glacier Identification on Northeast Spitsbergen, Svalbard from Landsat Imagery 1984-2018

Shunan Feng and Rickard Pettersson

Uppsala University, Department of Earth Sciences, Uppsala, Sweden (fsn.1995@gmail.com)

Svalbard archipelago is known as the “surge hot spot” for its high occurrence of glacial surge. Previous attempts of identifying surge events over the whole region are relying on the geomorphological evidence. Meanwhile, direct observation is sparse, especially on Northeast (NE) Spitsbergen. Landsat satellites provide us the longest space-borne earth observatory history record. This study utilizes all the available Landsat images (1984-2018) of 40 major maritime and valley glaciers on NE Spitsbergen, Svalbard to reconstruct the glacier velocity based on the Coregistration of Optically Sensed Images and Correlation (COSI-Corr) tool. Clouds and cloud shadows, together with the ETM+ Scan Line Corrector (SLC)-off Error are detected and masked by Fmask (Function of mask) which is one of the most effective and accurate cloud detection algorithm. It identifies clouds based on a generated cloud probability layer and the shadows are calculated from the similarity and the corresponding solar-sensor geometry. A principle component analysis is performed to the band 2-4 (green, red and near infrared bands respectively) images of Landsat 4 and 5. The first principal component is then re-gridded to obtain the 15 m reduced images with enhanced ice topography and improved surface feature, which match the panchromatic channel in Landsat 7 and 8. All the images are georeferenced at the precision of 0.01 pixels (0.15 m) by a single step discrete fourier transform. The surface texture of glaciers is enhanced by a high-pass filter transformation. The identification of surge events is done by the peak analysis of the reconstructed glacier surface velocity at the front. The identified timing of surge events of Tunabreen (2003-2005, winter of 2016) and Negribreen (from July 2017) had a good agreement with other reports of surges of these glaciers. In total 11 surge events are identified on 10 glaciers.


Global Distribution



(Sevestre, H., Benn, D.I., 2015. Journal of Glaciology 61, 646–662.)




Surge Type Glacier “hot spot”




- Lack of direct observation
- Can be derived from
 - geomorphology evidence of crevasse squeeze ridges in glacier forelands
 - the subtraction of digital terrain models (DTM)
 - SAR/Optic Satellite


Glacier Surge Mechanism



(Murray, T., Strozzi, T., Luckman, A., Jiskoot, H., Christakos, P., 2003. Journal of Geophysical Research: Solid Earth 108, 1–15.)



- high spatial resolution (15/30 m)
- accurate radiometric calibration
- high geometric precision
- the longest earth observatory history record





UPPSALA
UNIVERSITET

Challenge

- Low ice flow rate (long quiescent phase)
- Spatial and temporal resolution limitations of imagery




Technical challenge

- Cloud, cloud shadow mask



Fmask/MFmask





Clouds:

- Identification based on a generated cloud probability layer

Shadows

- calculated from the similarity and the corresponding solar-sensor geometry





UPPSALA
UNIVERSITET

Technical challenge

- Data integration





UPPSALA
UNIVERSITET

Landsat data

Cloud/Cloud
shadow
detection

Re-grid and
Principal
Component
Analysis

Georeferencing

COSI-Corr

Landsat 4-5

Bands

Band 2-Green

Band 3-Red

Band 4-Near Infrared
(NIR)

Wavelength (μm)

0.52-0.60

0.63-0.69

0.76-0.90

Resolution (m)

30

Landsat 7

Landsat 8

Bands

Band 8 -
Panchromatic

Wavelength (μm)

0.52-0.90

0.503-0.676

Resolution (m)

15

Band 2-Green

Band 3-Red

Band 4-Near
Infrared (NIR)

Principal
Component
Analysis


Re-grid

Improved
Landsat 4-5
images

Technical challenge

- Imagine co-registration







- Landsat systematic product accuracy requirement is better than **0.44** pixels
- Some of earlier archive of Landsat data were poorly geolocated (~100-1000 m)
- Single Step Discrete Fourier Transform

$$\rho_{XY} = \mathfrak{F}^{-1}(\mathfrak{F}^*(x) \cdot \mathfrak{F}(y))$$

- DFT increases the resolution of the original paired image in the spectral domain by a factor of k
- the displacement can be measured at a subpixel accuracy with the maximum peak located at a higher resolution
- The k value is set as 100 to obtain a **precision of 0.01(1/k) pixels.**



- Coregistration of Optically Sensed Images and Correlation (COSI-Corr)
- It matches the paired images by estimating the phase difference in the Fourier domain





UPPSALA
UNIVERSITET


Technical challenge

- Search window
- Time window






Results



- Tunabreen advanced in 1970 and remains in quiescent stage since then.
- It is observed to have surged in 2003-2005.





Results



- Tunabreen is observed to speed up and surge during the winter of 2016
- the timing coincides with a period of **record high temperatures** and **precipitation** in October and November.

Results


Negribreen: ice flows faster and start to surge in July 2017.




Discussion

- Identified 12 glaciers may have surged during 2003-2005.
- A systematic error? is detected around 2010 due to the scan line corrector error in Landsat 7.
- Future:
 - Glacier surge mechanism and climate?

Experiment result



Reference velocity map



Schellenberger, T., Dunse, T., Kääb, A., Schuler, T.V., Hagen, J.O. and Reijmer, C.H., 2017. Multi-year surface velocities and sea-level rise contribution of the Basin-3 and Basin-2 surges, Austfonna, Svalbard. *The Cryosphere Discuss.*, <https://doi.org/10.5194/tc-2017-5>.