

$$MB: \quad p(p) = \sqrt{\frac{2\beta}{\pi m}} e^{-\beta p^2 / 2m}, \quad \beta = \frac{1}{k_B T}$$

$$\text{Show: } \frac{\sigma_p^2}{\langle \sigma^2 \rangle^2} = 1 \quad \text{and} \quad \frac{\sigma_T^2}{\langle T \rangle_{nvt}^2} = \frac{1}{n}$$

1) calculate  $\langle p^2 \rangle$ ,  $\langle p^4 \rangle$

$$2) \sigma_{p^2}^2 = \langle p^4 \rangle - \langle p^2 \rangle^2$$

$$\langle p^2 \rangle = \int_0^\infty p^2 P(p) dp$$

$$= \int_0^{\infty} p^2 \sqrt{\frac{2\beta}{\pi m}} \exp\left(-\frac{\beta p^2}{2m}\right) dp$$

$$\text{Let } u = \frac{\beta p^2}{2m} \quad (\Rightarrow) \quad p^2 = \frac{u 2m}{\beta}$$

with  $u$  and  $p^2$

$$\Rightarrow u \rightarrow 0, p \rightarrow 0$$

$$u \rightarrow \infty, p \rightarrow \infty$$

$$\frac{du}{dp} = \frac{\beta}{\alpha u_m} \neq p \quad (=) \quad dp = du \frac{u}{\beta p} = du \frac{u}{\beta} \sqrt{\frac{\beta}{u u_m}}$$

$$= \int_0^{\infty} \frac{u^2 m}{\beta} \sqrt{\frac{2\beta}{\pi m}} \exp(-u) du \frac{m}{\beta} \sqrt{\frac{\beta}{m u^2}}$$

$$= \frac{2\mu}{\beta} \sqrt{\frac{2\beta}{\pi\ln}} \frac{\mu}{\beta} \sqrt{\frac{\beta}{2\mu}} \int_{-\infty}^{\infty} \sqrt{u} e^{+\rho(-u)} du$$

$$= \frac{m}{B}$$

$$\begin{aligned}\langle p^4 \rangle &= \int_0^\infty p^4 \cdot p(p) dp \\ &= \int_0^\infty p^4 \sqrt{\frac{2p}{\pi m}} \exp\left(-\frac{p^2}{2m}\right) dp\end{aligned}$$

$$\text{Let } u = \frac{\beta p^2}{2m} \quad (\Rightarrow) \quad p^2 = \frac{u}{\beta m}, \quad p^4 = \left(\frac{u}{\beta m}\right)^2$$

$$\frac{du}{dp} = \frac{\beta}{2m} \quad (\Rightarrow) \quad dp = \frac{du}{\frac{u}{\beta m}} = du \frac{m}{\beta} \sqrt{\frac{\beta}{u/2m}}$$

$$= \left(\frac{2m}{\beta}\right)^2 \sqrt{\frac{2\beta}{\pi m}} \int_0^\infty u^2 \exp(-u) du \frac{m}{\beta} \sqrt{\frac{\beta}{u/2m}}$$

$$= \frac{2m^2}{\beta^2} \sqrt{\frac{2\beta}{\pi m}} \frac{m}{\beta} \sqrt{\frac{\beta}{2m}} \underbrace{\int_0^\infty \frac{1}{\sqrt{u}} u^2 \exp(-u) du}_{\frac{3}{4} \sqrt{\frac{\pi}{m}}}$$

$$= \frac{4m^2}{\beta^2} \frac{\sqrt{2\beta}}{\sqrt{\pi} \sqrt{m}} \frac{m}{\beta} \frac{3}{4} \sqrt{\frac{\pi}{m}}$$

$$\underline{\underline{= \frac{3m^2}{\beta^2}}}$$

$$\sigma_{p^2}^2 = \langle p^4 \rangle - \langle p^2 \rangle^2$$

$$= \frac{3m^2}{\beta^2} - \left(\frac{m}{\beta}\right)^2 = \underline{\underline{\frac{2m^2}{\beta^2}}}$$

$$\frac{\sigma_p^2}{\langle p^2 \rangle^2} = \frac{2 \frac{m^2}{\hbar^2}}{\frac{m^2}{\hbar^2}} = 2$$

U  
 $2 \times \frac{1}{2} = \underline{\underline{1}}$

in 2D, due to the degrees of freedom a factor  $\times \frac{1}{2}$  has to be applied.

Show

$$\frac{\sigma_T^2}{\langle T \rangle^2} = \frac{1}{N}$$

without going into too much thermodynamical details:

For small temperature fluctuations  $\sigma_T^2$  may be expressed as follows:

$$\sigma_T^2 \approx \frac{\sigma_E^2}{(N k_B)^2}$$

some correction factors [like 1.25 are ignored]

with the energy fluctuation  $\sigma_E^2$

The variance in Energy  $\sigma_E^2 = \langle E^2 \rangle - \langle E \rangle^2$  is given from thermal physics as

$$(1) \quad \sigma_E^2 = k_B T^2 C_V \quad \text{with the heat capacity } C_V.$$

$$\text{with } C_V = \frac{\partial E}{\partial T} \Big|_V \quad \text{and the } \langle E \rangle = \frac{1}{2} k_B T N$$

$$\Rightarrow C_V = N k_B \quad (2)$$

for  $N$  particles with 2 degrees of freedom in 2D

$\Rightarrow (2)$  in (1)

$$\sigma_E^2 = k_B T^2 N k_B = k_B^2 T^2 N^2$$

$$\Rightarrow \sigma_T^2 \approx \frac{k_B^2 T^2 N}{N^2 k_B^2} = \frac{T^2}{N}$$

and with  $\langle T \rangle^2 = T^2$

$$\Rightarrow \frac{\sigma_T^2}{\langle T \rangle^2} = \frac{T^2}{N T^2} = \frac{1}{N}$$

□

Consequences

$\frac{\sigma_T^2}{\langle T \rangle^2} = \frac{1}{N} \Rightarrow$  for large  $N$  the temperature fluctuates very little.

$\Rightarrow$  for large systems ( $N$  big) the temperature is stable, for smaller systems it can vary more

Since  $T \propto \langle U_E \rangle \propto p$ : for large systems individual momenta of the particles fluctuate less

