श्री BalaJi

Narendra Avasthi

PROBLEMS IN PHYSICAL CHEMISTRY

for JEE Main & Advanced



Free 20 YEARS PREVIOUS PAPERS (WITH SOLUTIONS) WITH THIS BOOK WORTH ₹300/-

Purchase Genuine Books only

A genuine copy of this book is covered by 3D holographic film printed with the logo of Shri Balaji Publications. As you tilt the book, the logo disappears at a certain angle.

15th edition

For More Material Join: @JEEAdvanced_2024

श्री BalaJi

Problems in PHYSICAL CHEMISTRY

for **JEE**
Main & Advanced

by:
Narendra Avasthi
Director,
Vibrant Academy India (P) Ltd.
KOTA

SHRI BALAJI PUBLICATIONS
(EDUCATIONAL PUBLISHERS & DISTRIBUTORS)
AN ISO 9001-2008 CERTIFIED ORGANIZATION
Muzaffarnagar (U.P.) - 251001

For More Material Join: @JEEAdvanced_2024

Contents

1.	Stoichiometry	1-72
2.	Atomic Structure	73-129
3.	Gaseous State	130-193
4.	Thermodynamics	194-257
5.	Chemical Equilibrium	258-309
6.	Ionic Equilibrium	310-363
7.	Chemical Kinetics and Nuclear Chemistry	364-426
8.	Electrochemistry	427-478
9.	Dilute Solution	479-520
10.	Solid State	521-561
11.	Surface Chemistry	562-600

For More Material Join: @JEEAdvanced_2024



1

STOICHIOMETRY

Laws of Chemical Combination

Chemical reactions take place according to certain laws. These laws are called the Laws of Chemical Combination. "These are no longer useful in chemical calculations now but gives an idea of earlier methods of analysing and relating compounds by mass."

- **Law of Conservation of Mass [Lavoisier (1774)]**

  During any physical or chemical change, the sum of masses of all substances present in reactions vessel remain conserved.

- **Law of Constant Composition or Definite Proportions [Proust (1799)]**

  In a given chemical compound, the elements are always combined in the same proportions by mass.

- **Law of Multiple Proportions [Dalton (1803)]**

  Whenever two elements form more than one compound, the different masses of one element that combine with the same mass of the other element are in the ratio of small whole numbers.

- **Law of Reciprocal Proportions [Richter (1792)]**

  When two elements combine separately with a fixed mass of a third elements then the ratio of their masses in which they do so is either same or some whole number multiple of the ratio in which they combine with each other.

- **Gay-Lussac's Law of combining volumes**

  According to Gay-Lussac's law of combining volume, when gases react together, they always do so in volumes which bear a simple ratio to one another and to the volumes of the products, if these are also gases, provided all measurements of volumes are done under similar conditions of temperature and pressure.

- **Avogadro's Law**

  The volume of a gas (at fixed pressure and temperature) is proportional to the number of moles (or molecules of gas present). Mathematically we can write

  $$V \quad n$$

  Or Equal volumes of all gases under similar conditions of temperature and pressure contain equal number of molecules.

For More Material Join: @JEEAdvanced_2024

Dalton's Atomic Theory

- Matter consists of tiny particles called atoms.
- Atoms are indestructible. In chemical reactions, the atoms rearrange but they do not themselves break apart.
- In any sample of a pure element, all the atoms are identical in mass and other properties.
- The atoms of different elements differ in mass and other properties.
- When atoms of different elements combine to form compounds, new and more complex particles are formed. However, in a given compound the constituent atoms are always present in the same fixed numerical ratio.

Modern Atomic Theory

- **Atom is no longer considered to be indivisible :** It has been found that an atom has a complex structure and is composed of sub-atomic particles such as electrons, protons and neutrons.
- **Atoms of the same element may not be similar in all respects :**
  Ex. Isotopes ($^{23}_{11}Na$, $^{24}_{11}Na$)
- **Atoms of different elements may be similar in one or more respects :**
  Ex. Isobars. ($^{40}_{20}Ca$, $^{40}_{18}Ar$)
- **Atom is the smallest unit which takes part in chemical reactions :** Although atom is composed of sub-atomic particles, yet it is the smallest particle which takes part in chemical reactions.
- **The ratio in which atoms unite may be fixed and integral but may not be simple :** For example, in sugar molecule ($C_{12}H_{22}O_{11}$), the ratio of C, H and O atoms is 12 : 22 : 11 which is not simple.
- **Atoms of one element can be changed into atoms of other element :**
  Ex. Artificial Nuclear Reactions
- **The mass of atom can be changed into energy :** According to Einstein's equation $E \quad mc^2$ (E Energy, m mass, c the speed of light, *i.e.,* 3 10 cm sec ^{1}), mass and energy are inter-convertible.

Mole Concept

(a) Definition of one mole : One mole is a collection of that many entities as there are number of atoms exactly in 12 gm of C-12 isotope.

(b) $1u$ 1 amu $\frac{1}{12}$ of mass of 1 atom of C^{12} $\frac{1\text{g}}{N_A}$ 1.66 10^{-24} g

(c) For Elements :

- 1 g atom 1 mole of atoms N_A atoms.
- g atomic mass (GAM) mass of N_A atoms in g.
- Mole of atoms $\frac{\text{Mass } (g)}{\text{GAM or molar mass}}$

For More Material Join: @JEEAdvanced_2024

(d) For molecules :

- 1 g molecule 1 mole of molecule N_A molecule.
- g molecular mass (GMM) mass of N_A molecule in g.
- Moles of molecules $\dfrac{\text{Mass } (g)}{\text{GMM or molar mass}}$

(e) For ionic compounds :

- 1 g formula unit 1 mole of formula unit N_A formula unit.
- g formula mass (GFM) mass of N_A formula unit in g.
- Mole of formula unit $\dfrac{\text{Mass } (g)}{\text{GFM or molar mass}}$

(f) 1 mole of a substance →
- Contains 6.022×10^{23} particles
- Mass as much as molecular mass/atomic mass/ionic mass in grams
- If it is a gas, one mole occupies a volume of 22.4 L at 1 atm & 273 K

(g) Average or mean atomic mass : Average atomic mass of element

$$A_{\text{Avg.}} \quad \frac{A_1x_1 \quad A_2x_2 \quad \ldots\ldots}{x_1 \quad x_2 \quad \ldots\ldots}$$

Here A_1, A_2 are isotopic mass of element and x_1, x_2 are natural abundance of isotopes.

(h) Average or mean molar mass : The average molar mass of the different substances present in the container

$$M_{\text{Avg.}} \quad \frac{M_1n_1 \quad M_2n_2 \quad \ldots\ldots}{n_1 \quad n_2 \quad \ldots\ldots}$$

Here M_1, M_2 are molar mass of substances and n_1, n_2 are mole of substances present in the container.

Empirical Formula, Molecular Formula

(a) Empirical formula : Formula depicting constituent atom in their simplest ratio.

Molecular formula : Formula depicting actual number of atoms in one molecule of the compound.

(b) Relation between molecular formula and empirical formula :

$$n \quad \frac{\text{Molecular mass}}{\text{Empirical Formula mass}}$$

(c) Densities :

- Density $\dfrac{\text{Mass}}{\text{Volume}}$, Unit: g/cc
- Relative density $\dfrac{\text{Density of any substance}}{\text{Density of reference substance}}$
- Specific gravity $\dfrac{\text{Density of any substance}}{\text{Density of water at 4 C}}$
- Vapour density : Ratio of density of vapours to the density of hydrogen at similar pressure and temperature.

For More Material Join: @JEEAdvanced_2024

Vapour density $= \dfrac{\text{Density of vapours at some temperature and pressure}}{\text{Density of } H_2 \text{ gas at same temperature and pressure}}$

Vapour density $= \dfrac{\text{Molecular mass}}{2}$

Stoichiometry

Stoichiometry pronounced ("stoy – key – om – e – tree") is the calculations of the quantities of reactants and products involved in a chemical reaction. Following methods can be used for solving problems.

(a) Mole Method (Balancing is required)

For Ex. : $2KClO_3 \quad 2KCl \quad 3O_2$

$$\frac{\text{Moles of } KClO_3}{2} \quad \frac{\text{Moles of } KCl}{2} \quad \frac{\text{Moles of } O_2}{3}$$

(b) Principle of Atom Conservation (P.O.A.C.) method (Balancing is not required)

For Ex. : $KClO_3 \quad KCl \quad O_2$

POAC for K : 1 mole of $KClO_3$ 1 mole of KCl

POAC for Cl : 1 mole of $KClO_3$ 1 mole of KCl

POAC for O : 3 mole of $KClO_3$ 2 mole of O_2

Concept of Limiting Reagent

(a) Limiting Reagent : It is very important concept in chemical calculation. It refers to reactant which is present in minimum stoichiometry quantity for a chemical reaction. It is reactant consumed fully in a chemical reaction. So all calculations related to various products or in sequence of reactions are made on the basis of limiting reagent.

(b) Calculation of Limiting Reagent : Divide given moles of each reactant by their stoichiometric coefficient, the one with least ratio is limiting reagent.

Percentage Yield

The percentage yield of product $= \dfrac{\text{Actual yield}}{\text{theoretical maximum yield}} \quad 100$

Concentration Terms

(a) For solutions (*i.e.*, homogeneous mixtures) :

❖ If the mixture is not homogeneous, then none of them is applicable.

(i) % by mass $\dfrac{w}{W} \quad \dfrac{\text{Mass of solute}}{\text{Mass of solution}} \quad 100$

[X% by mass means 100 gm solution contains X gm solute ; (100 X) gm solvent]

(ii) % $\dfrac{w}{V} \quad \dfrac{\text{Mass of solute}}{\text{Volume of solution}} \quad 100$

[X% $\dfrac{w}{V}$ means 100 mL solution contains X gm solute]

For More Material Join: @JEEAdvanced_2024

(iii) % $\frac{v}{V}$ $\frac{\text{Volume of solute}}{\text{Volume of solution}}$ 100

❖ For gases % by volume is same as mole %

(iv) Mole % $\frac{\text{Moles of solute}}{\text{Total moles}}$ 100

(v) Mole fraction (X) $\frac{\text{Moles of solute}}{\text{Total moles}}$

(vi) Molarity (M) $\frac{\text{Moles of solute}}{\text{Volume of solution (in litre)}}$

(vii) Molality (m) $\frac{\text{Moles of solute}}{\text{Mass of solvent (in kg)}}$

(viii) Parts per million (ppm) $\frac{\text{Mass of solute}}{\text{Mass of solvent}}$ $10^6 \sim \frac{\text{Mass of solute}}{\text{Mass of solution}}$ 10^6

(ix) Formality (F) $\frac{\text{No. of formula unit}}{\text{Volume of solution (in litre)}}$

(b) (i) On adding solvent in a solution (dilution) : Number of mole of solute remains constant

$M_f V_f \quad M_i V_i$

(ii) Mixing of two solutions of same solute

$M_f V_f \quad M_1 V_1 \quad M_2 V_2 \quad \ldots.$

(c) Volume strength of H_2O_2:

H_2O_2 (aq) solution labelled as 'x V' 'volume H_2O_2 (for *e.g.,* '20 V' H_2O_2), it means x volume of O_2 (in litre) at 1 atm and 273K that can be obtained from 1 litre of such a sample when it decomposes according to

$$H_2O_2 \qquad H_2O \quad \frac{1}{2}O_2$$

$$\text{Molarity of } H_2O_2 \quad \frac{\text{Volume strength of } H_2O_2}{11.2}$$

Eudiometry

(For reactions involving gaseous reactants and products)

❖ Eudiometry or gas analysis involves the calculations based on gaseous reactions or the reactions in which at least two components are gaseous, in which the amounts of gases are represented by their volumes, measured at the same pressure and temperature.

❖ Gay-Lussac's law of volume combination holds good.

❖ Problem may be solved directly is terms of volume, in place of mole.

The volume of gases produced is often given by mentioning certain solvent which absorb certain gases.

For More Material Join: @JEEAdvanced_2024

Solvent	gas (es) absorbed
KOH (*aq.*)	CO_2, SO_2, Cl_2
Ammoniacal Cu_2Cl_2	CO
Turpentine oil	O_3
Alkaline pyrogallol	O_2
Water	NH_3, HCl
Anhydrous $CuSO_4/CaCl_2$	H_2O

Redox Processes

(a) Oxidation Number : It is the charge (real or imaginary) which an atom appears to have when it is in combined state. It may be a whole no. or fractional. An element may have different values of oxidation number depending . It depends on nature of compound in which it is present. There are some operational rules to determine oxidation number.

(b) Definition of Oxidation and Reduction :

- **Oxidation :** Addition of oxygen , removal of hydrogen , addition of electronegative element, removal of electropositive element , loss of electrons , increase in oxidation number (de-electronation).
- **Reduction :** Removal of oxygen, addition of hydrogen, removal of electronegative element, addition of electropositive element, gain of electrons, decrease in oxid. no. (electronation).
- **Redox Reaction :** A reaction in which oxidation & reduction occur simultaneously.

(c) Oxidising and Reducing Agents :

- **Oxidising Agents (Oxidants, Oxidisers) :** They oxidise others, themselves are reduced & gain electrons. *e.g.,* $O_2, O_3, HNO_3, MnO_2, H_2O_2$, halogens, $KMnO_4, K_2Cr_2O_7, KIO_3, Cl(SO_4)_3, FeCl_3$, NaOCl, hydrogen ions.(Atoms present in their higher oxidation state.)
  Compounds is which reacting atom is present in its highest oxidation state may act as oxidant only.
- **Reducing Agents (Reductants or Reducers) :** They reduce others, themselves get oxidised & lose electrons. Also called reductants or reducers. H_2 molecular form is weak reducing agent but Nascent hydrogen is strong reducing agent *e.g.*, C, CO, H_2S, SO_2, $SnCl_2$, Sodium thio Sulphate ($Na_2S_2O_3$), Al, Na, CaH_2, $NaBH_4$, $LiAlH_4$ etc.
  Compounds in which reacting atom is present in its lowest oxidation state may act as reductant only.
  Both Oxidising & Reducing Agents : Compounds having reactive atom in its intermediate (between highest and lowest oxidation states) oxidation state may act as both oxidant or reductant. *e.g.*, SO_2, H_2O_2, O_3, NO_2, etc.

(d) Balancing of redox reactions :

- Ion - electron method
- Oxidation number method

[Concept involved is that, in any chemical reaction electrons (e) cannot be produced so no. of e *s* in L.H.S. & R.H.S. should be the same]

For More Material Join: @JEEAdvanced_2024

Equivalent Concept

(a) Law of Chemical equivalence : It states that in any chemical reaction the equivalents of all the reactants and products must be same.

$$2A \quad 3B \qquad 4C$$

Equivalents of 'A' Equivalents of 'B' Equivalents of 'C'

(b) Terms used in equivalent concept :

- Equivalent mass of A $\dfrac{\text{Molar mass of } A}{\text{Valency factor or } n \text{ factor}}$
- Equivalents of 'A' $\dfrac{\text{Mass of '}A\text{' (in g)}}{\text{Equivalent mass of '}A\text{'}}$
- Numbers of equivalents of 'A' no. of moles of 'A' n-factor
- **Normality (N) :** For solutions, concentration term **normality (N)** is used, which can be defined as "the number of equivalent of solute present in one litre (1000 mL) solution".

$$N \quad \frac{\text{Number of } g\text{ - equivalents of solute}}{\text{Volume of solution (in L)}}$$

$$N \quad \frac{W(g) \quad 1000}{E \quad V \text{ (in mL)}}$$

$$N \quad \text{Molarity} \times \text{Valence factor}$$

milli-equivalents of solute $N \quad V$ (in mL) $M \quad V$ (in mL) n factor

(c) Valence factor (n-factor) calculation : n-factor here we mean a conversion factor by which we divide molar mass of substance to get equivalent mass and it depends on nature of substance which vary from one condition to another condition. We can divide n-factor calculations in two category.

- **In case of non-redox reaction.**

  n-factor moles of charge displaced per mole of species
- **In case of redox reaction.**

  n-factor of oxidizing or reducing agent mole of electrons gain or lost per mol of species.

(d) Volumetric analysis (Titration) : Titration is a procedure for determining the concentration of a solution by allowing a carefully measured volume to react with a standard solution of another substance, whose concentration is known.

- **Primary standard :** A substance available in a pure form or state of known purity which is used in standardizing a solution.
- **Standardization :** The process by which the concentration of a solution is accurately ascertained.
- **Standard solution :** A solution whose concentration has been accurately determined.
- **Titrant :** The reagent (a standard solution) which is added from a burette standard term to react with the analyte.
- **Titrate :** This mainly involves titrations based chemistry. It can be divided into two major category.

  (I) Non-redox reactions (II) Redox reactions

For More Material Join: @JEEAdvanced_2024

(e) Type of reactions :

(i) Non-redox reactions : This involve following kind of titrations :

1. Acid-Base titrations
2. Double indicator acid-base titration
3. Precipitation titration
4. Back titration

(ii) Redox reactions : This involve following kind of titrations :

1. Iodimetry titrations
2. Iodometry titrations
3. Back titration

(f) Titrations :

(I) Non-redox titrations

(1) Acid-Base titration : To find out strength or concentration of unknown acid or base, it is titrated against base or acid of known strength. At the equivalence point, equivalents of titrate and titrant solution are equal. So by using law of chemical equivalence strength of unknown solution in determined.

At equivalence point, milliequivalents of acid milliequivalents of base.

(2) Double indicator acid-base titration : In the acid-base titration the equivalence point is known with the help of indicator which changes its colour at the end point. In the titration of polyacidic base or polybasic acid there are more than one end point. Some times one indicator is not able to give colour change at every end point. So to find out each end point we have to use more than one indicator. For example, in the titration of Na_2CO_3 against HCl there are two end points.

Na_2CO_3 HCl $NaHCO_3$ NaCl At first equivalence point

$NaHCO_3$ HCl H_2CO_3 NaCl At second equivalence point

When we use phenolphthalein in the above titration it changes its colour at first end point when $NaHCO_3$ is formed and with it we can not determine second end point. Similarly with methyl orange it changes its colour at second end point only and we can not determine first end point. It is because all indicator changes colour on the basis of pH of titration mixture. So in titration of $NaHCO_3$, $KHCO_3$ against acid, phenolphthalein can not be used.

Titration (against acid)	**Indicator**	**pH Range**	***n*-factor**
Na_2CO_3	Phenolphthalein	8.3 – 10	1
Na_2CO_3	Methyl orange	3.1 – 4.4	2

Note : When we carry out dilution of solution, milliequivalent, equivalent, milli mole or mole of substance does not change because they represent amount of substance, however molar concentration may change.

(3) Precipitation titration : In precipitation titrations involving ionic reactions, we determine the strength of unknown solution of salt by titrating it against a reagent with which it can form precipitate. For example NaCl strength can be determine by titrating it against $AgNO_3$ solution with which it form white ppt. of AgCl.

For More Material Join: @JEEAdvanced_2024

At equivalence point,

Milliequivalent of NaCl = milliequivalent of $AgNO_3$ used = milliequivalent of AgCl formed

(II) Redox Titrations

At equivalence point,

milliequivalent of Oxidizing agent used = milliequivalent of reducing agent reacted.

S.No.	Reagent	Half Reaction	*n*-factor of reagent
1.	FAS (Mohr's salt) $[FeSO_4(NH_4)_2SO_4\ 6H_2O]$	$Fe^{2} \quad Fe^{3} \quad e$	1
2.	MnO_4 (Permanganate ion) (in acidic medium)	$MnO_4 \quad 8H \quad 5e \quad Mn^{2} \quad 4H_2O$	5
3.	MnO_4 (in basic medium)	$Mn^{7} \quad e \quad Mn^{6}$	1
4.	MnO_4 (in mild basic or neutral medium)	$Mn^{7} \quad e \quad Mn^{4}$	3
5.	$Cr_2O_7^{2}$ (dichromate ion)	$Cr_2O_7^{2} \quad 14H \quad 6e \quad 2Cr^{3} \quad 7H_2O$	6
6.	$C_2O_4^{2}$ (Oxalate ion)	$C_2O_4^{2} \quad 2CO_2 \quad 2e$	2
7.	As_2O_3	$As_2O_3 \quad 5H_2O \quad 2AsO_4^{3} \quad 10H \quad 4e$	4
8.	$CaOCl_2$ (Bleaching powder)	$CaOCl_2 \quad H_2O \quad 2KI \quad Ca(OH)_2 \quad I_2 \quad 2KCl$	2
9.	MnO_2	$MnO_2 \quad 4HCl\,(Conc.) \quad MnCl_2 \quad Cl_2 \quad 2H_2O$	2
10.	IO_3	$IO_3 \quad 5I \quad 6H \quad 3I_2 \quad 3H_2O$	5
11.	H_2O_2 (act as oxidizing agent)	$H_2O_2 \quad 2e \quad 2H_2O$	2
12.	H_2O_2 (act as reducing agent)	$H_2O_2 \quad O_2 \quad 2e$	2
13.	H_2O_2 (disproportion)	$H_2O_2 \quad H_2O \quad 1/2\,O_2$	1
14.	Cl_2 (disproportion)	$3Cl_2 \quad 6OH$ (strong) $\quad ClO_3 \quad 5Cl \quad 3H_2O$	5/3
15.	H_2S (in acidic medium)	$H_2S \quad I_2 \quad S \quad 2I \quad 2H$	2
16.	Sn^{2} (in acidic medium)	$Sn^{2} \quad I_2 \quad Sn^{4} \quad 2I$	2
17.	N_2H_4	$N_2H_4 \quad N_2 \quad 4H \quad 4e$	4
18.	SO_3^{2} (in acidic medium)	$SO_3^{2} \quad H_2O \quad SO_4^{2} \quad 2e \quad 2H$	2
19.	$Na_2S_2O_3$ (Sodium thiosulphate or Hypo)	$2Na_2S_2O_3 \quad I_2 \quad Na_2S_4O_6 \quad 2NaI$	1
20.	I_2	$I_2 \quad 2e \quad 2I$	2

(1) Iodimetric Titration or Iodimetry : In such titrations iodine solution is used as an oxidant and iodine is directly titrated against a reducing agent. This type of titrations are used for the determination of strength of reducing agents like sulphides, arsenides, thiosulphates etc., by titrating them against a standard solution of iodine.

For More Material Join: @JEEAdvanced_2024

This type of titration involves free iodine, here iodine solution is treated with known sodium thiosulphate solution.

$$I_2 \quad 2Na_2S_2O_3 \qquad 2NaI \quad Na_2S_4O_6$$

Equivalents of I_2 Equivalents of $Na_2S_2O_3$ used

(2) Iodometric Titration or Iodometry: It is an indirect method of estimation of iodine. In this titration an oxidizing agent is used to liberate iodine from solution containing iodide ion (*e. g.*, KI) and the liberated iodine is treated with a standard solution of a reducing agent added from a burette. Here a neutral or an acidic solution of an oxidizing agent is used and the amount of liberated I_2 is equal to the equivalents of this oxidizing agent.

These titrations are used to determine the concentration of $K_2Cr_2O_7$, $KMnO_4$, $CuSO_4$, Ferric ions, H_2O_2 etc.

These titrations are carried out in following two steps :

- **Step-1 :** Oxidizing agent (X) KI (excess) I_2 reduced state of oxidant
  Equivalents of (X) Equivalents of I_2
- **Step-2 :** Liberated I_2 $2Na_2S_2O_3$ $2NaI$ $Na_2S_4O_6$
  Equivalents of I_2 Equivalents of $Na_2S_2O_3$ used

(g) Back titration : Back titration is used in volumetric analysis to find out excess amount of reagent added by titrating it with suitable reagent. It is also used to find out percentage purity of sample.

For example in an acid-base titration suppose we have added excess base $[B(OH)_q]$ in acid (H_nA) solution.

To find excess base it is back titrated with another acid (H_mB) of known concentration.

Equivalents of base equivalents of $(H_nA \quad H_mB)$

q moles of base taken n moles of H_nA reacted m moles of H_mB reacted

Hardness of Water

(a) Definition of hard water : Hard water is having soluble salts of calcium and magnessium ions.

(b) Degree of hardness : Degree of hardness defined as number of parts by mass of $CaCO_3$ (or its equivalent quantities of other substance) present in million parts of mass of water.

$$\text{Hardness of water} \quad \frac{\text{Mass of } CaCO_3}{\text{Mass of water}} \quad 10^6 \quad \text{ppm.}$$

For More Material Join: @JEEAdvanced_2024



Level 1

1. Calculate number of neutrons present in 12 10^{25} atoms of oxygen ($_8O^{17}$) :
(Given: N_A 6 10^{23})
(a) 1800 (b) 1600 (c) 1800 N_A (d) 3200 N_A

2. If mass of one atom is 3.32 10 ^{23}g, then calculate number of nucleons (neutrons and protons) present in 2 atoms of the element :
(a) 40 (b) 20 (c) 10 (d) 40 N_A

3. Calculate number of electrons present in 9.5 g of PO_4^3 :
(a) 6 (b) 5 N_A (c) 0.1 N_A (d) 4.7 N_A

4. What is the number of moles of O-atom in 126 amu of HNO_3 ?
(a) 2 (b) $\frac{2}{N_A}$ (c) 6 (d) $\frac{6}{N_A}$

5. What is the charge of 96 amu of S^2 ?
(a) 2 C (b) 3.2 10 ^{19}C (c) 9.6 10 ^{19}C (d) 6 C

6. A sample of sodium has a mass of 46 g. What is the mass of the same number of calcium atoms as sodium atoms present in given sample ?
(a) 46 g (b) 20 g (c) 40 g (d) 80 g

7. The total number of neutrons present in 54 mL H_2O (l) are:
(a) 3 N_A (b) 30 N_A (c) 24 N_A (d) none of these

8. Total number of electrons present in 48 g Mg^{2+} are:
(a) 24 N_A (b) 2 N_A (c) 20 N_A (d) none of these

9. The number of neutrons in 5 g of D_2O (D is ^2_1H) are:
(a) 0.25 N_A (b) 2 5 N_A (c) 1 1 N_A (d) none of these

10. Cisplatin, an anticancer drug, has the molecular formula $Pt(NH_3)_2Cl_2$. What is the mass (in gram) of one molecule ? (Atomic masses : Pt = 195, H = 1.0, N = 14, Cl = 35.5)
(a) 4.98 10 ^{21} (b) 4.98 10 ^{22} (c) 6.55 10 ^{21} (d) 3.85 10 ^{22}

11. Aspirin has the formula $C_9H_8O_4$. How many atoms of oxygen are there in a tablet weighing 360 mg?
(a) 1.204 10^{23} (b) 1.08 10^{22} (c) 1.204 10^{24} (d) 4.81 10^{21}

12. 20 g of an ideal gas contains only atoms of S and O occupies 5.6 L at 1 atm and 273 K. What is the molecular mass of gas ?
(a) 64 (b) 80 (c) 96 (d) None of these

13. A sample of ammonium phosphate, $(NH_4)_3PO_4$, contains 6 moles of hydrogen atoms. The number of moles of oxygen atoms in the sample is:
(a) 1 (b) 2 (c) 4 (d) 6

14. Total number of moles of oxygen atoms in 3 litre $O_3(g)$ at 27°C and 8.21 atm are:
(a) 3 (b) 1 (c) 1 (d) none of these

15. 3.011 10^{22} atoms of an element weighs 1.15 gm. The atomic mass of the element is:
(a) 10 amu (b) 2.3 amu (c) 35.5 amu (d) 23 amu

For More Material Join: @JEEAdvanced_2024

16. One atom of an element x weighs $6.643 \quad 10^{-23}$ g. Number of moles of atoms in its 20 kg is:
(a) 4 (b) 40 (c) 100 (d) 500

17. Mass of one atom of the element A is $3.9854 \quad 10^{-23}$ g. How many atoms are contained in 1 g of the element A?
(a) $2.509 \quad 10^{23}$ (b) $6.022 \quad 10^{23}$ (c) $12.044 \quad 10^{23}$ (d) None of these

18. Which of the following contains the largest mass of hydrogen atoms?
(a) 5.0 moles $C_2H_2O_4$ (b) 1.1 moles $C_3H_8O_3$
(c) 1.5 moles $C_6H_8O_6$ (d) 4.0 moles $C_2H_4O_2$

19. Which has minimum number of oxygen atoms ?
(a) 10 mL H_2O (l) (b) 0.1 mole of V_2O_5 (s)
(c) 12 gm O_3(g) (d) $12.044 \quad 10^{22}$ molecules of CO_2

20. Rearrange the following (I to IV) in the order of increasing masses:
(I) 0.5 mole of O_3 (II) 0.5 gm atom of oxygen
(III) $3.011 \quad 10^{23}$ molecules of O_2 (IV) 5.6 litre of CO_2 at 1 atm and 273 K
(a) II < IV < III < I (b) II < I < IV < III (c) IV < II < III < I (d) I < II < III < IV

21. If the volume of a drop of water is 0.0018 mL then the number of water molecules present in two drop of water at room temperature is:
(a) $12.046 \quad 10^{19}$ (b) $1.084 \quad 10^{18}$ (c) $4.84 \quad 10^{17}$ (d) $6.023 \quad 10^{23}$

22. It is known that atom contain protons, neutrons and electrons. If the mass of neutron is assumed to half of its original value whereas that of proton is assumed to be twice of its original value then the atomic mass of $^{14}_{6}C$ will be :
(a) same (b) 14.28% less (c) 14.28% more (d) 28.56% less

23. Common salt obtained from sea-water contains 8.775% NaCl by mass. The number of formula units of NaCl present in 25 g of this salt is :
(a) $3.367 \quad 10^{23}$ formula units (b) $2.258 \quad 10^{22}$ formula units
(c) $3.176 \quad 10^{23}$ formula units (d) $4.73 \quad 10^{25}$ formula units

24. The number of hydrogen atoms present in 25.6 g of sucrose ($C_{12}H_{22}O_{11}$) which has a molar mass of 342.3 g is :
(a) $22 \quad 10^{23}$ (b) $9.91 \quad 10^{23}$ (c) $11 \quad 10^{23}$ (d) $44 \quad 10^{23}$

25. Caffiene has a molecular mass of 194. If it contains 28.9% by mass of nitrogen, number of atoms of nitrogen in one molecule of caffeine is :
(a) 4 (b) 6 (c) 2 (d) 3

26. The density of water is 1g/mL. What is the volume occupied by 1 molecule of water ?
(a) $1.44 \quad 10^{-23}$mL (b) 1mL (c) 18 mL (d) $2.88 \quad 10^{-23}$mL

27. A 25.0 mm 40.0 mm piece of gold foil is 0.25 mm thick. The density of gold is 19.32 g/cm^3. How many gold atoms are in the sheet ? (Atomic weight : Au = 197.0)
(a) $7.7 \quad 10^{23}$ (b) $1.5 \quad 10^{23}$ (c) $4.3 \quad 10^{21}$ (d) $1.47 \quad 10^{22}$

28. If average molecular mass of air is 29, then assuming N_2 and O_2 gases are there, which options are correct regarding composition of air ?
(i) 75% by mass of N_2 (ii) 75% by moles N_2 (iii) 72.41% by mass of N_2
(a) only (i) is correct (b) only (ii) is correct
(c) both (ii) and (iii) are correct (d) both (i) and (ii) are correct

For More Material Join: @JEEAdvanced_2024

29. Density of dry air containing only N_2 and O_2 is 1.15 g/L at 740 mm of Hg and 300 K. What is % composition of N_2 by mass in the air ?
(a) 78% (b) 85.5% (c) 70.02% (d) 62.75%

30. A gaseous mixture of H_2 and CO_2 gases contains 66 mass % of CO_2. The vapour density of the mixture is:
(a) 6.1 (b) 5.4 (c) 2.7 (d) 10.8

31. The vapour density of a mixture containing NO_2 and N_2O_4 is 27.6. The mole fraction of N_2O_4 in the mixture is:
(a) 0.1 (b) 0.2 (c) 0.5 (d) 0.8

32. Density of an ideal gas at 2 atm and 600 K is 2 g/L.
Calculate relative density of this gas with respect to Ne(g) under similar conditions :
(Given : R 1/12 atm L/mol.K)
(a) 2.5 (b) 2 (c) 3 (d) 5

33. Average atomic mass of magnesium is 24.31 amu. This magnesium is composed of 79 mole % of ^{24}Mg and remaining 21 mole % of ^{25}Mg and ^{26}Mg. Calculate mole % of ^{26}Mg.
(a) 10 (b) 11 (c) 15 (d) 16

34. Indium (atomic mass = 114.82) has two naturally occurring isotopes, the predominant one form has isotopic mass 114.9041 and abundance of 95.72%. Which of the following isotopic mass is the most likely for the other isotope ?
(a) 112.94 (b) 115.90 (c) 113.90 (d) 114.90

35. Calculate density of a gaseous mixture which consist of 3.01 10^{24} molecules of N_2 and 32 g of O_2 gas at 3 atm pressure and 860 K temperature (Given : R 1/12 atm L/mole.K)
(a) 0.6 g/L (b) 1.2 g/L (c) 0.3 g/L (d) 12 g/L

36. A mixture of O_2 and gas "Y" (mol. mass 80) in the mole ratio a : b has a mean molecular mass 40. What would be mean molecular mass, if the gases are mixed in the ratio b : a under identical conditions ? (Assume that gases are non-reacting):
(a) 40 (b) 48 (c) 62 (d) 72

37. If water sample are taken from sea, rivers or lake, they will be found to contain hydrogen and oxygen in the approximate ratio of 1 : 8. This indicates the law of:
(a) multiple proportion (b) definite proportion
(c) reciprocal proportions (d) none of these

38. Hydrogen and oxygen combine to form H_2O_2 and H_2O containing 5.93% and 11.2% hydrogen respectively. The data illustrates :
(a) law of conservation of mass (b) law of constant proportion
(c) law of reciprocal proportion (d) law of multiple proportion

39. Which one of the following combinations illustrate law of reciprocal proportions ?
(a) N_2O_3, N_2O_4, N_2O_5 (b) NaCl, NaBr, NaI
(c) CS_2, CO_2, SO_2 (d) PH_3, P_2O_3, P_2O_5

40. Carbon and oxygen combine to form two oxides, carbon monoxide and carbon dioxide in which the ratio of the masses of carbon and oxygen is respectively 12 : 16 and 12 : 32. These figures illustrate the :
(a) law of multiple proportions (b) law of reciprocal proportions
(c) law of conservation of mass (d) law of constant proportions

For More Material Join: @JEEAdvanced_2024

41. A sample of calcium carbonate ($CaCO_3$) has the following percentage composition : Ca = 40%, C = 12%, O = 48%. If the law of constant proportions is true, then the mass of calcium in 4 g of a sample of calcium carbonate obtained from another source will be :
(a) 0.016 g (b) 0.16 g (c) 1.6 g (d) 16 g

42. The law of multiple proportion is illustrated by the two compounds :
(a) Sodium chloride and sodium bromide (b) Ordinary water and heavy water
(c) Caustic soda and caustic potash (d) Sulphur dioxide and sulphur trioxide

43. All the substances listed below are fertilizers that contribute nitrogen to the soil. Which of these is the richest source of nitrogen on a mass percentage basis ?
(a) Urea, $(NH_2)_2CO$ (b) Ammonium nitrate, NH_4NO_3
(c) Nitric oxide, NO (d) Ammonia, NH_3

44. One mole of element *X* has 0.444 times the mass of one mole of element *Y*. One atom of element *X* has 2.96 times the mass of one atom of ^{12}C. What is the atomic mass of *Y* ?
(a) 80 (b) 15.77 (c) 46.67 (d) 40.0

45. A given sample of pure compound contains 9.81 gm of Zn, 1.8 10^{23} atoms of chromium and 0.60 mole of oxygen atoms. What is the simplest formula ?
(a) $ZnCr_2O_7$ (b) $ZnCr_2O_4$ (c) $ZnCrO_4$ (d) $ZnCrO_6$

46. The formula of an acid is HXO_2. The mass of 0.0242 moles of the acid is 1.657 g. What is the atomic mass of *X* ?
(a) 35.5 (b) 28.1 (c) 128 (d) 19.0

47. What is the empirical formula of vanadium oxide, if 2.74 g of the metal oxide contains 1.53 g of metal ?
(a) V_2O_3 (b) VO (c) V_2O_5 (d) V_2O_7

48. Determine the empirical formula of Kelvar, used in making bullet proof vests, is 70.6% C, 4.2% H, 11.8% N and 13.4% O :
(a) $C_7H_5NO_2$ (b) $C_7H_5N_2O$ (c) C_7H_9NO (d) C_7H_5NO

49. The hydrated salt $Na_2CO_3\,xH_2O$ undergoes 63% loss in mass on heating and becomes anhydrous. The value of *x* is:
(a) 10 (b) 12 (c) 8 (d) 18

50. A 6.85 g sample of the hydrated $Sr(OH)_2\,xH_2O$ is dried in an oven to give 3.13 g of anhydrous $Sr(OH)_2$. What is the value of *x* ? (Atomic masses : Sr = 87.60, O = 16.0, H = 1.0)
(a) 8 (b) 12 (c) 10 (d) 6

51. What percentage of oxygen is present in the compound $CaCO_3\,3Ca_3(PO_4)_2$?
(a) 23.3% (b) 45.36% (c) 41.94% (d) 17.08%

52. Dieldrin, an insecticide, contains C, H, Cl and O. Combustion of 29.72 mg of Dieldrin gave 41.21 mg CO_2 and 5.63 mg of H_2O. In a separate analysis 25.31 mg of Dieldrin was converted into 57.13 mg AgCl. What is the empirical formula of Dieldrin ?
(a) $C_6H_4Cl_3O$ (b) C_8H_8ClO (c) $C_{12}H_8Cl_6O$ (d) $C_6H_4Cl_3O_2$

53. A gaseous compound is composed of 85.7% by mass carbon and 14.3% by mass hydrogen. Its density is 2.28 g/litre at 300 K and 1.0 atm pressure. Determine the molecular formula of the compound.
(a) C_2H_2 (b) C_2H_4 (c) C_4H_8 (d) C_4H_{10}

For More Material Join: @JEEAdvanced_2024

54. Complete combustion of 0.86 g of compound X gives 2.64 g of CO_2 and 1.26 g of H_2O. The lowest molecular mass X can have :
(a) 43 g (b) 86 g (c) 129 g (d) 172 g

55. The sulphate of a metal M contains 9.87% of M. This sulphate is isomorphous with $ZnSO_4$ $7H_2O$. The atomic mass of M is :
(a) 40.3 (b) 36.3 (c) 24.3 (d) 11.3

56. In an organic compound of molar mass 108 gm mol ^{1} C, H and N atoms are present in 9 : 1 : 3.5 by mass. Molecular formula can be :
(a) $C_6H_8N_2$ (b) $C_7H_{10}N$ (c) $C_5H_6N_3$ (d) $C_4H_{18}N_3$

57. On analysis, a certain compound was found to contain 254 g of iodine (at. mass 127) and 80 g oxygen (at. mass 16). What is the formula of the compound ?
(a) IO (b) I_2O (c) I_5O_3 (d) I_2O_5

58. An element A is tetravalent and another element B is divalent. The formula of the compound formed from these elements will be :
(a) A_2B (b) AB (c) AB_2 (d) A_2B_3

59. A compound used in making nylon, contains 43.8% oxygen. There are four oxygen atoms per molecule. What is the molecular mass of compound?
(a) 36 (b) 116 (c) 292 (d) 146

60. Suppose two elements X and Y combine to form two compounds XY_2 and X_2Y_3. 0.05 mole of XY_2 weighs 5 g while 3.011 10^{23} molecules of X_2Y_3 weigh 85 g. The atomic masses of X and Y are respectively :
(a) 20, 30 (b) 30, 40 (c) 40, 30 (d) 80, 60

61. 44 g of a sample on complete combustion gives 88 g CO_2 and 36 g of H_2O. The molecular formula of the compound may be:
(a) C_4H_6 (b) C_2H_6O (c) C_2H_4O (d) C_3H_6O

62. 40 milligram diatomic volatile substance (X_2) is converted to vapour that displaced 4.92 mL of air at 1 atm and 300 K. Atomic mass of element X is nearly:
(a) 400 (b) 240 (c) 200 (d) 100

63. Two elements X (at. mass 75) and Y (at. mass 16) combine to give a compound having 75.8% of X. The formula of the compound is:
(a) XY (b) X_2Y (c) X_2Y_2 (d) X_2Y_3

64. A sample of phosphorus that weighs 12.4 g exerts a pressure 8 atm in a 0.821 litre closed vesel at 527°C. The molecular formula of the phosphorus vapour is:
(a) P_2 (b) P_4 (c) P_6 (d) P_8

65. Manganese forms non-stoichiometric oxides having the general formula MnO_x. The value of x for the compound that analyzed 64% by mass Mn:
(a) 1.16 (b) 1.83 (c) 2 (d) 1.93

66. 1.44 gram of titanium (At. mass 48) reacted with excess of O_2 and produce x gram of non-stoichiometric compound $Ti_{1.44}O$. The value of x is:
(a) 2 (b) 1.77 (c) 1.44 (d) none of these

67. Which statement is false for the balanced equation given below ?

$$CS_2 + 3O_2 \quad 2SO_2 + CO_2$$

For More Material Join: @JEEAdvanced_2024

(a) One mole of CS_2 will produce one mole of CO_2
(b) The reaction of 16 g of oxygen produces 7.33 g of CO_2
(c) The reaction of one mole of O_2 will produce 2/3 mole of SO_2
(d) Six molecules of oxygen requires three molecules of CS_2

68. Which of the following setups is correct to calculate the mass (in g) of $KClO_3$ produced from the reaction of 0.150 moles of Cl_2?

$$3Cl_2 + 6KOH \quad 5KCl + KClO_3 + 3H_2O$$

(a) 0.150 moles Cl_2 1 mole $KClO_3$/3 moles Cl_2 122.5 g/1 mole $KClO_3$
(b) 0.150 moles Cl_2 1 mole $KClO_3$/3 moles Cl_2 1 mole $KClO_3$/122.5 g
(c) 0.150 moles Cl_2 3 moles Cl_2/1 mole $KClO_3$ 122.5 g/1 mole $KClO_3$
(d) 0.150 moles Cl_2 3 moles Cl_2/1 mole $KClO_3$ 1 mole $KClO_3$/122.5 g

69. 2.0 g of a sample contains mixture of SiO_2 and Fe_2O_3. On very strong heating, it leaves a residue weighing 1.96 g. The reaction responsible for loss of mass is given below.

$$Fe_2O_3(s) \quad Fe_3O_4(s) + O_2(g), \text{ (unbalance equation)}$$

What is the percentage by mass of SiO_2 in original sample ?
(a) 10% (b) 20% (c) 40% (d) 60%

70. What volume of air at 1 atm and 273 K containing 21% of oxygen by volume is required to completely burn sulphur (S_8) present in 200 g of sample, which contains 20% inert material which does not burn. Sulphur burns according to the reaction

$$\frac{1}{8}S_8(s) + O_2(g) \quad SO_2(g)$$

(a) 23.52 litre (b) 320 litre (c) 112 litre (d) 533.33 litre

71. For the reaction, $2Fe(NO_3)_3 + 3Na_2CO_3 \quad Fe_2(CO_3)_3 + 6NaNO_3$
initially 2.5 mole of $Fe(NO_3)_3$ and 3.6 mole of Na_2CO_3 are taken. If 6.3 mole of $NaNO_3$ is obtained then % yield of given reaction is:
(a) 50 (b) 84 (c) 87.5 (d) 100

72. How many moles of P_4 can be produced by reaction of 0.10 moles $Ca_5(PO_4)_3F$, 0.36 moles SiO_2 and 0.90 moles C according to the following reaction ?

$$4\,Ca_5(PO_4)_3F + 18\,SiO_2 + 30\,C \quad 3P_4 + 2CaF_2 + 18CaSiO_3 + 30\,CO$$

(a) 0.060 (b) 0.030 (c) 0.045 (d) 0.075

73. Some older emergency oxygen masks contains potassium superoxide, KO_2 which reacts with CO_2 and water present in exhaled air to produce oxygen according to the given equation. If a person exhales 0.667 g of CO_2 per minute, how many grams of KO_2 are consumed in 5.0 minutes?

$$4KO_2 + 2H_2O + 4CO_2 \quad 4KHCO_3 + 3O_2$$

(a) 10.7 (b) 0.0757 (c) 1.07 (d) 5.38

74. The mass of N_2F_4 produced by the reaction of 2.0 g of NH_3 and 80 g of F_2 is 3.56 g. What is the per cent yield?

$$2NH_3 + 5F_2 \quad N_2F_4 + 6HF$$

(a) 79.0 (b) 71.2 (c) 84.6 (d) None of these

For More Material Join: @JEEAdvanced_2024

75. Calculate the mass of lime (CaO) obtained by heating 200 kg of 95% pure lime stone ($CaCO_3$) :
(a) 104.4 kg (b) 105.4 kg (c) 212.8 kg (d) 106.4 kg

76. Phosphoric acid (H_3PO_4) prepared in a two step process.
(1) $P_4 + 5O_2 \rightarrow P_4O_{10}$ (2) $P_4O_{10} + 6H_2O \rightarrow 4H_3PO_4$
We allow 62 g of phosphorus to react with excess oxygen which form P_4O_{10} in 85% yield. In the step (2) reaction 90% yield of H_3PO_4 is obtained. Mass of H_3PO_4 produced is:
(a) 37.485 g (b) 149.949 g (c) 125.47 g (d) 564.48 g

77. 9 moles of "*D*" and 14 moles of *E* are allowed to react in a closed vessel according to given reactions. Calculate number of moles of *B* formed in the end of reaction, if 4 moles of *G* are present in reaction vessel. (Percentage yield of reaction is mentioned in the reaction)
Step-1 $3D + 4E \xrightarrow{80\%} 5C + A$
Step-2 $3C + 5G \xrightarrow{50\%} 6B + F$
(a) 2.4 (b) 30 (c) 4.8 (d) 1

78. The chief ore of Zn is the sulphide, ZnS. The ore is concentrated by froth floatation process and then heated in air to convert ZnS to ZnO.

$$2ZnS + 3O_2 \xrightarrow{80\%} 2ZnO + 2SO_2$$

$$ZnO + H_2SO_4 \xrightarrow{100\%} ZnSO_4 + H_2O$$

$$2ZnSO_4 + 2H_2O \xrightarrow{80\%} 2Zn + 2H_2SO_4 + O_2$$

The number of moles of ZnS required for producing 2 moles of Zn will be :
(a) 3.125 (b) 2 (c) 2.125 (d) 4

79. 0.8 mole of a mixture of CO and CO_2 requires exactly 40 gram of NaOH in solution for complete conversion of all the CO_2 into Na_2CO_3. How many moles more of NaOH would it require for conversion into Na_2CO_3, if the mixture (0.8 mole) is completely oxidised to CO_2 ?
(a) 0.2 (b) 0.6 (c) 1 (d) 1.5

80. Silver oxide (Ag_2O) decomposes at temperature 300°C yielding metallic silver and oxygen gas. A 1.60 g sample of impure silver oxide yields 0.104 g of oxygen gas. What is the per cent by mass of the silver oxide in the sample ?
(a) 5.9 (b) 47.125 (c) 94.25 (d) 88.2

81. 342 g of 20% by mass of $Ba(OH)_2$ solution (sp. gr. 0.57) is reacted with 1200 mL of 2 *M* HNO_3. If the final density of solution is same as pure water then molarity of the ion in resulting solution which decides the nature of the above solution is:
(a) 0.25 (b) 0.5 *M* (c) 0.888 *M* (d) None of these

82. 100 mL of H_2SO_4 solution having molarity 1 *M* and density 1.5 g/mL is mixed with 400 mL of water. Calculate final molarity of H_2SO_4 solution, if final density is 1.25 g/mL :
(a) 4.4 *M* (b) 0.145 *M* (c) 0.52 *M* (d) 0.227 *M*

83. What volume of HCl solution of density 1.2 g/cm^3 and containing 36.5% by mass HCl, must be allowed to react wtih zinc (Zn) in order to liberate 4.0 g of hydrogen ?
(a) 333.33 mL (b) 500 mL (c) 614.66 mL (d) None of these

84. An ideal gaseous mixture of ethane (C_2H_6) and ethene (C_2H_4) occupies 28 litre at 1 atm and 273 K. The mixture reacts completely with 128 g O_2 to produce CO_2 and H_2O. Mole fraction at C_2H_6 in the mixture is:
(a) 0.6 (b) 0.4 (c) 0.5 (d) 0.8

For More Material Join: @JEEAdvanced_2024

85. Wood's metal contains 50.0% bismuth, 25.0% lead, 12.5% tin and 12.5% cadmium by mass. What is the mole fraction of tin ?
(Atomic mass : Bi 209, Pb 207, Sn 119, Cd 112)
(a) 0.202 (b) 0.158 (c) 0.176 (d) 0.221

86. The density of a 56.0% by mass aqueous solution of 1-propanol ($CH_3CH_2CH_2OH$) is 0.8975 g/cm^3. What is the mole fraction of the 1-propanol ?
(a) 0.292 (b) 0.227 (c) 0.241 (d) 0.276

87. What is the molartiy of SO_4^{2} ion in aqueous solution that contain 34.2 ppm of $Al_2(SO_4)_3$?
(Assume complete dissociation and density of solution 1 g/mL)
(a) 3 10 ^{4} *M* (b) 2 10 ^{4} *M* (c) 10 ^{4} *M* (d) None of these

88. The relation between molarity (*M*) and molality (*m*) is given by :
(density of solution (g/mL), M_1 molecular mass of solute)
(a) $m = \dfrac{1000\ M}{1000 \quad M_1}$ (b) $m = \dfrac{1000 \quad M}{1000 \quad MM_1}$
(c) $m = \dfrac{1000\ MM}{1000 \quad MM_1}$ (d) $m = \dfrac{1000\ M}{1000 \quad MM_1}$

89. Molarity and molality of a solution of an liquid (mol. mass 50) in aqueous solution is 9 and 10 respectively. What is the density of solution?
(a) 1 g/cc (b) 0.95 g/cc (c) 1.05 g/cc (d) 1.35 g/cc

90. An aqueous solution of ethanol has density 1.025 g/mL and it is 2 *M*. What is the molality of this solution ?
(a) 1.79 (b) 2.143 (c) 1.951 (d) None of these

91. 0.2 mole of HCl and 0.2 mole of barium chloride were dissolved in water to produce a 500 mL solution. The molarity of the Cl ions is :
(a) 0.06 *M* (b) 0.09 *M* (c) 1.2 *M* (d) 0.80 *M*

92. Calculate the mass of anhydrous HCl in 10 mL of concentrated HCl (density = 1.2 g / mL) solution having 37% HCl by mass.
(a) 4.44 g (b) 4.44 mg (c) 4.44 10 ^{3} mg (d) 0.444 g

93. Calculate the molality of 1 L solution of 80% H_2SO_4 (*w/V*), given that the density of the solution is 1.80 g mL ^{1}.
(a) 8.16 (b) 8.6 (c) 1.02 (d) 10.8

94. Fluoxymesterone, $C_{20}H_{29}FO_3$, is an anabolic steroid. A 500 mL solution is prepared by dissolving 10.0 mg of the steroid in water. 1.0 mL portion of this solution is diluted to a final volume of 1.00 L. What is the resulting molarity ?
(a) 1.19 10 ^{10} (b) 1.19 10 ^{7} (c) 5.95 10 ^{8} (d) 2.38 10 ^{11}

95. The 25 mL of a 0.15 *M* solution of lead nitrate, $Pb(NO_3)_2$ reacts with all of the aluminium sulphate, $Al_2(SO_4)_3$, present in 20 mL of a solution. What is the molar concentration of the $Al_2(SO_4)_3$?

$$3Pb(NO_3)_2(aq) + Al_2(SO_4)_3(aq) \qquad 3PbSO_4(s) + 2Al(NO_3)_3(aq)$$

(a) 6.25 10 ^{2} *M* (b) 2.421 10 ^{2} *M* (c) 0.1875 *M* (d) None of these

96. Concentrated HNO_3 is 63% HNO_3 by mass and has a density of 1.4 g/mL. How many millilitres of this solution are required to prepare 250 mL of a 1.20 *M* HNO_3 solution?
(a) 18.0 (b) 21.42 (c) 20.0 (d) 14.21

For More Material Join: @JEEAdvanced_2024

97. 50 mL of 20.8% (*w/V*) $BaCl_2$ (*aq*) and 100 mL of 9.8% (*w/V*) H_2SO_4 (*aq*) solutions are mixed. Molarity of Cl ions in the resulting solution is: (At. mass of Ba = 137)
(a) 0.333 *M* (b) 0.666 *M* (c) 0.1 *M* (d) 1.33 *M*

98. 100 mL of 10% NaOH (*w/V*) is added to 100 mL of 10% HCl (*w/V*). The nature of resultant solution is:
(a) alkaline (b) strongly alkaline (c) acidic (d) neutral

99. How many millilitres of 0.1 *M* H_2SO_4 must be added to 50 mL of 0.1 *M* NaOH to give a solution that has a concentration of 0.05 *M* in H_2SO_4 ?
(a) 400 mL (b) 200 mL (c) 100 mL (d) None of these

100. 1 *M* HCl and 2 *M* HCl are mixed in volume ratio of 4 : 1. What is the final molarity of HCl solution?
(a) 1.5 (b) 1 (c) 1.2 (d) 1.8

101. Three solutions *X*, *Y*, *Z* of HCl are mixed to produce 100 mL of 0.1 *M* solution. The molarities of *X*, *Y* and *Z* are 0.07 *M*, 0.12 *M* and 0.15 *M* respectively. What respective volumes of *X*, *Y* and *Z* should be mixed ?
(a) 50 mL, 25 mL, 25 mL (b) 20 mL, 60 mL, 20 mL
(c) 40 mL, 30 mL, 30 mL (d) 55 mL, 20 mL, 25 mL

102. A bottle of an aqueous H_2O_2 solution is labelled as '28.375 *V*' H_2O_2 and the density of the solution (in g/mL) is 1.25. Choose the correct option:
(a) Molality of H_2O_2 solution is 2 (b) Molarity of H_2O_2 solution is 5
(c) Molality of H_2O_2 solution is 2.15 (d) None of these

103. The impure 6 g of NaCl is dissolved in water and then treated with excess of silver nitrate solution. The mass of precipitate of silver chloride is found to be 14 g. The % purity of NaCl solution would be:
(a) 95% (b) 85% (c) 75% (d) 65%

104. $Al_2(SO_4)_3$ solution of 1 molal concentration is present in 1 litre solution of density 2.684 g/cc. How many moles of $BaSO_4$ would be precipitated on adding excess of $BaCl_2$ in it ?
(a) 2 moles (b) 3 moles (c) 6 moles (d) 12 moles

105. A certain public water supply contains 0.10 ppb (part per billion) of chloroform ($CHCl_3$). How many molecules of $CHCl_3$ would be obtained in 0.478 mL drop of this water ?
(assumed d 1 g/mL)
(a) $4 \quad 10^{13} \quad N_A$ (b) $10^{3} \quad N_A$ (c) $4 \quad 10^{10} \quad N_A$ (d) None of these

106. Decreasing order (first having highest and then others following it) of mass of pure NaOH in each of the aqueous solution:
(i) 50 g of 40% (*w/W*) NaOH
(ii) 50 mL of 50% (*w/V*) NaOH [$d_{soln.} = 1.2\ g/mL$]
(iii) 50 g of 15 *M* NaOH [$d_{soln.} = 1\ g/mL$]
(a) i, ii, iii (b) iii, ii, i (c) ii, iii, i (d) ii, i, iii

107. What is the molar mass of diacidic organic Lewis base (*B*), if 12 g of its chloroplatinate salt ($B\,H_2PtCl_6$) on ignition produced 5 g residue of Pt?
(a) 52 (b) 58 (c) 88 (d) None of these

For More Material Join: @JEEAdvanced_2024

108. On strong heating, one gram of the silver salt of an organic dibasic acid yields 0.5934 g of silver. If the mass percentage of carbon in it 8 times the mass percentage of hydrogen and one-half the mass percentage of oxygen, determine the molecular formula of the acid.
(a) $C_4H_6O_4$ (b) $C_4H_6O_6$ (c) $C_2H_6O_2$ (d) $C_5H_{10}O_5$

109. 0.607 g of a silver salt of tribasic organic acid was quantitatively reduced to 0.37 g of pure Ag. What is the molecular mass of the acid?
(a) 207 (b) 210 (c) 531 (d) 324

110. A sample of peanut oil weighing 2 g is added to 25 mL of 0.40 *M* KOH. After saponification is complete, 8.5 mL of 0.28 *M* H_2SO_4 is needed to neutralize excess of KOH. The saponification number of peanut oil is :
(saponification number is defined as the milligrams of KOH consumed by 1 g of oil)
(a) 146.72 (b) 223.44 (c) 98.9 (d) None of these

111. 20 mL of a mixture of CO and H_2 were mixed with excess of O_2 and exploded and cooled. There was a volume contraction of 18 mL. All volume measurements corresponds to room temperature (27°C) and one atmospheric pressure. Determine the volume ratio $V_1 : V_2$ of CO and H_2 in the original mixture.
(a) 1 : 2 (b) 3 : 2 (c) 2 : 3 (d) 4 : 1

112. In the reaction,

$$2Al(s) \quad 6HCl(aq) \qquad 2Al^3 \ (aq) \quad 6Cl \ (aq) \quad 3H_2(g)$$

(a) 6 L HCl(*aq*) is consumed for every 3 L $H_2(g)$ produced
(b) 33.6 L $H_2(g)$ is produced regardless of temperature and pressure for every mole Al that reacts
(c) 67.2 L $H_2(g)$ at 1 atm and 273 K is produced for every mole Al that reacts
(d) 11.2 L $H_2(g)$ at 1 atm and 273 K is produced for every mole HCl(*aq*) consumed

113. The percentage by volume of C_3H_8 in a gaseous mixture of C_3H_8, CH_4 and CO is 20. When 100 mL of the mixture is burnt in excess of O_2, the volume of CO_2 produced is:
(a) 90 mL (b) 160 mL (c) 140 mL (d) none of these

114. 40 mL gaseous mixture of CO, CH_4 and Ne was exploded with 15 mL of oxygen. After treatment with KOH the volume reduced by 9 mL and again on treatment with alkaline pyrogallol, the volume further reduced by 1.5 mL. Percentage of CH_4 in the original mixture is:
(a) 22.5 (b) 77.5 (c) 7.5 (d) 15

115. A gaseous mixture of propane and butane of volume 3 litre on complete combustion produces 11.0 litre CO_2 under standard conditions of temperature and pressure.
The ratio of volume of butane to propane is:
(a) 1 : 2 (b) 2 : 1 (c) 3 : 2 (d) 3 : 1

116. Phosphorous has the oxidation state of +1 in:
(a) Orthophosphoric acid (b) Phosphorous acid
(c) Hypophosphorous acid (d) Metaphosphoric acid

117. The oxidation state(s) of Cl in $CaOCl_2$ (bleaching powder) is/are:
(a) +1 only (b) –1 only (c) +1 and 1 (d) none of these

118. The oxidation number of sulphur in S_8, S_2F_2, H_2S and H_2SO_4 respectively are:
(a) 0, 1, 2 and 6 (b) 2, 0, 2 and 6
(c) 0, 1, 2 and 4 (d) 2, 0, 2 and 6

For More Material Join: @JEEAdvanced_2024

119. Fe shows an oxidation state of +1 in:
(a) $Fe(CO)_5$ (b) $[Fe(H_2O)_5NO]SO_4$
(c) $Fe_4[Fe(CN)_6]_3$ (d) $FeCl_4$

120. When SO_2 is passed into an acidified potassium dichromate solution, the oxidation numbers of sulphur and chromium in the final products respectively are:
(a) +6, +6 (b) +6, +3 (c) 0, +3 (d) +2, +3

121. The oxidation number of nitrogen atoms in NH_4NO_3 are:
(a) +3, +3 (b) +3, 3 (c) 3, 5 (d) 5, 3

122. The oxidation states of S-atoms in Caro's and Marshall's acids are:
(a) 6, 6 (b) 6, 4 (c) 6, 6 (d) 4, 6

123. In which of the following the oxidation number of oxygen has been arranged in increasing order :
(a) OF_2 KO_2 BaO_2 O_3 (b) BaO_2 KO_2 O_3 OF_2
(c) BaO_2 O_3 OF_2 KO_2 (d) KO_2 OF_2 O_3 BaO_2

124. The oxidation numbers of oxygen in KO_3, Na_2O_2 respectively are :
(a) 3, 2 (b) 1, 0 (c) 0, 1 (d) –0.33, –1

125. The oxidation number of phosphorus in $Ba(H_2PO_2)_2$ is :
(a) –1 (b) +1 (c) +2 (d) +3

126. If it is known that in $Fe_{0.96}O$, Fe is present in +2 and +3 oxidation state, what is the mole fraction of Fe^2 in the compound ?
(a) 12/25 (b) 25/12 (c) 1/12 (d) 11/12

127. Which of the following sequence of compounds is according to the decreasing order of the oxidation state of nitrogen ?
(a) HNO_3, NO, NH_4Cl, N_2 (b) HNO_3, NO, N_2, NH_4Cl
(c) HNO_3, NH_4Cl, NO, N_2 (d) NO, HNO_3, NH_4Cl, N_2

128. 2 mole of N_2H_4 loses 16 mole of electron is being converted to a new compound *X*. Assuming that all of the N appears in the new compound. What is the oxidation state of 'N' in *X* ?
(a) 1 (b) 2 (c) 2 (d) 4

129. When $K_2Cr_2O_7$ is converted to K_2CrO_4, the change in the oxidation state of chromium is :
(a) 0 (b) 6 (c) 4 (d) 3

130. When a manganous salt is fused with a mixture of KNO_3 and solid NaOH, the oxidation number of Mn changes from +2 to :
(a) +4 (b) +3 (c) +6 (d) +7

131. In Fe(II)–MnO_4 titration, HNO_3, is not used because:
(a) it oxidises Mn^2 (b) it reduces MnO_4
(c) it oxidises Fe^{2+} (d) it reduces Fe^{3+} formed

132. Which species are oxidized and reduced in the reaction ?
FeC_2O_4 $KMnO_4$ Fe^3 CO_2 Mn^2
(a) Oxidised : Fe, C ; Reduced : Mn (b) Oxidised : Fe ; Reduced : Mn
(c) Reduced : Fe, Mn ; Oxidised : C (d) Reduced : C ; Oxidised : Mn, Fe

For More Material Join: @JEEAdvanced_2024

133. In which of the following reactions, H_2O_2 is acting as a reducing agent ?
(a) $SO_2 \quad H_2O_2 \quad H_2SO_4$ (b) $2KI \quad H_2O_2 \quad 2KOH \quad I_2$
(c) $PbS \quad 4H_2O_2 \quad PbSO_4 \quad 4H_2O$ (d) $Ag_2O \quad H_2O_2 \quad 2Ag \quad H_2O \quad O_2$

134. Following reaction describes the rusting of iron

$$4Fe \quad 3O_2 \quad 4Fe^3 \quad 6O^2$$

Which one of the following statement is incorrect ?
(a) This is an example of a redox reaction (b) Metallic iron is reduced to Fe^3
(c) Fe^3 is an oxidising agent (d) Metallic iron is a reducing agent

135. Which reaction does not represent auto redox or disproportionation ?
(a) $Cl_2 \quad OH \quad Cl \quad ClO_3 \quad H_2O$ (b) $2H_2O_2 \quad H_2O \quad O_2$
(c) $2Cu \quad Cu^2 \quad Cu$ (d) $(NH_4)_2Cr_2O_7 \quad N_2 \quad Cr_2O_3 \quad 4H_2O$

136. Which of the following is redox reaction ?
(a) H_2SO_4 reacts with NaOH
(b) In atmosphere, O_3 is formed from O_2 by lightning
(c) Evaporation of H_2O
(d) Oxides of nitrogen are formed form nitrogen & oxygen by lightning

137. Which of the following is a redox-reaction ?
(a) $2Na[Ag(CN)_2] \quad Zn \quad Na_2[Zn(CN)_4] \quad 2Ag$
(b) $BaO_2 \quad H_2SO_4 \quad BaSO_4 \quad H_2O_2$
(c) $N_2O_5 \quad H_2O \quad 2HNO_3$
(d) $AgNO_3 \quad KI \quad AgI \quad KNO_3$

138. For the redox reaction, $MnO_4 \quad C_2O_4^2 \quad H^+ \quad Mn^{2+} + CO_2 + H_2O$
the correct coefficients of the reactants for the balanced reaction are respectively MnO_4, C_2O_4, H^+:
(a) 2, 5, 16 (b) 16, 3, 12 (c) 15, 16, 12 (d) 2, 16, 5

139. In a chemical reaction, $K_2Cr_2O_7 + xH_2SO_4 + ySO_2 \quad K_2SO_4 + Cr_2(SO_4)_3 + zH_2O$; the value of x, y and z respectively are :
(a) $x = 1, y = 3, z = 1$ (b) $x = 4, y = 1, z = 4$
(c) $x = 3, y = 2, z = 1$ (d) $x = 2, y = 2, z = 1$

140. Balance the following equation and choose the quantity which is the sum of the coefficients of the products :

$$\ldots\ldots CS_2 + \ldots\ldots Cl_2 \quad CCl_4 + \ldots\ldots S_2Cl_2$$

(a) 5 (b) 3 (c) 6 (d) 2

141. Balance the following equation and choose the quantity which is the sum of the coefficients of reactants and products:

$$\ldots PtCl_4 + \ldots XeF_2 \quad PtF_6 + \ldots ClF + \ldots Xe$$

(a) 16 (b) 13 (c) 18 (d) 12

142. It 0.1 mole H_3PO_x is completely neutralised by 5.6 g KOH then select the true statement.
(a) $x = 3$ and given acid is dibasic
(b) $x = 4$ and given acid has no P–H linkage
(c) $x = 2$ and given acid does not form acid salt
(d) all of these

For More Material Join: @JEEAdvanced_2024

143. When potassium permanganate is titrated against ferrous ammonium sulphate in acidic medium, the equivalent mass of potassium permanganate is:

(a) $\frac{\text{molecular mass}}{3}$ (b) $\frac{\text{molecular mass}}{5}$

(c) $\frac{\text{molecular mass}}{2}$ (d) $\frac{\text{molecular mass}}{10}$

144. Equivalent mass of FeS_2 in the half reaction, $FeS_2 \quad Fe_2O_3 + SO_2$ is :

(a) $M/10$ (b) $M/11$ (c) $M/6$ (d) $M/1$

145. The equivalent mass of HCl in the given reaction is:

$K_2Cr_2O_7 + 14HCl \quad 2KCl + 2CrCl_3 + 3Cl_2 + H_2O$

(a) 16.25 (b) 36.5 (c) 73 (d) 85.1

146. Equivalent mass of H_3PO_2 when it disproportionate into PH_3 and H_3PO_3 is:

(a) M (b) $M/2$ (c) $M/4$ (d) $3M/4$

147. In the following reaction, $As_2S_3 + H^+ + NO_3^- \quad NO + H_2O + AsO_4^{3-} \quad SO_4^{2-}$

the equivalent mass of As_2S_3 is related to its molecular mass by :

(a) $M/2$ (b) $M/4$ (c) $M/24$ (d) $M/28$

148. Sulphur forms the chlorides S_2Cl_2 and SCl_2. The equivalent mass of sulphur in SCl_2 is :

(a) 8 g/mol (b) 16 g/mol (c) 64.8 g/mol (d) 32 g/mol

149. The equivalent mass of an element is 4. Its chloride has a vapour density 59.25. Then, the valency of the elements is :

(a) 4 (b) 3 (c) 2 (d) 1

150. 6 10^{-3} mole $K_2Cr_2O_7$ reacts completely with 9 10^{-3} mole X^{n+} to give XO_3 and Cr^{3+}. The value of n is :

(a) 1 (b) 2 (c) 3 (d) None of these

151. What mass of $H_2C_2O_4 \cdot 2H_2O$ (mol. mass 126) should be dissolved in water to prepare 250 mL of centinormal solution which act as a reducing agent ?

(a) 0.63 g (b) 0.1575 g (c) 0.126 g (d) 0.875 g

152. The equivalent mass of the salt, $KHC_2O_4 \cdot H_2C_2O_4 \cdot 4H_2O$ when it act as reducing agent is :

(a) $\frac{\text{Mol. mass}}{1}$ (b) $\frac{\text{Mol. mass}}{2}$ (c) $\frac{\text{Mol. mass}}{3}$ (d) $\frac{\text{Mol. mass}}{4}$

153. The equivalent mass of divalent metal is W. The molecular mass of its chloride is :

(a) W + 35.5 (b) W + 71 (c) 2W + 71 (d) 2W + 35.5

154. When BrO_3^- ion reacts with Br^- in acid medium, Br_2 is liberated. The equivalent mass of Br_2 in this reaction is:

(a) $\frac{5M}{8}$ (b) $\frac{5M}{3}$ (c) $\frac{3M}{5}$ (d) $\frac{4M}{6}$

155. If m_A gram of a metal A displaces m_B gram of another metal B from its salt solution and if the equivalent mass are E_A and E_B respectively then equivalent mass of A can be expressed as:

(a) $E_A = \frac{m_A}{m_B} E_B$ (b) $E_A = \frac{m_A m_B}{E_B}$ (c) $E_A = \frac{m_B}{m_A} E_B$ (d) $E_A = \sqrt{\frac{m_A}{m_B} E_B}$

For More Material Join: @JEEAdvanced_2024

156. Hydrazine reacts with KIO_3 in presence of HCl as

$$N_2H_4 + IO_3^- \quad 2H^+ + Cl^- \qquad ICl + N_2 + 3H_2O$$

The equivalent masses of N_2H_4 and KIO_3 respectively are:

(a) 8 and 53.5 (b) 16 and 53.5 (c) 8 and 35.6 (d) 8 and 87

157. What will be the normality of a solution obtained by mixing 0.45 *N* and 0.60 *N* NaOH in the ratio 2 : 1 by volume ?

(a) 0.4 *N* (b) 0.5 *N* (c) 1.05 *N* (d) 0.15 *N*

158. A solution containing 2.7 10^{-3} mol of A^{2+} ions required 1.6 10^{-3} mole of MnO_4^- for the oxidation of A^{2+} to AO_3^- the medium used is :

(a) neutral (b) acidic (c) strong basic (d) none of these

159. H_2O_2 is used as bleaching reagent because on dissociation it gives oxygen ($H_2O_2 \quad H_2O + \frac{1}{2}O_2$).

"Chachi 420" used H_2O_2 solution to bleach her hair and she required 2.24 L O_2 gas at 1 atm and 273 K. She has a H_2O_2 solution labelled as '5.6 V' then what volume of such solution must she required to bleach her hair ?

(a) 200 mL (b) 300 mL (c) 400 mL (d) 500 mL

160. 1.25 g of a solid dibasic acid is completely neutralised by 25 mL of 0.25 molar $Ba(OH)_2$ solution. Molecular mass of the acid is :

(a) 100 (b) 150 (c) 120 (d) 200

161. 10 mL of *N*-HCl, 20 mL of *N*/2 H_2SO_4 and 30 mL of *N*/3 HNO_3 are mixed together and volume made to one litre. The normality of H in the resulting solution is :

(a) 3*N*/100 (b) *N*/10 (c) *N*/20 (d) *N*/40

162. 0.45 g of an acid of mol. mass 90 was neutralised by 20 mL of 0.54 *N* caustic potash (KOH). The basicity of acid is :

(a) 1 (b) 2 (c) 3 (d) 4

163. A 3.4 g sample of H_2O_2 solution containing *x*% H_2O_2 by mass requires *x* mL of a $KMnO_4$ solution for complete oxidation under acidic condition. The molarity of $KMnO_4$ solution is :

(a) 1 (b) 0.5 (c) 0.4 (d) 0.2

164. What volume of $O_2(g)$ measured at 1 atm and 273 K will be formed by action of 100 mL of 0.5 *N* $KMnO_4$ on hydrogen peroxide in an acid solution ? The skeleton equation for the reaction is

$$KMnO_4 + H_2SO_4 + H_2O_2 \qquad K_2SO_4 + MnSO_4 + O_2 + H_2O$$

(a) 0.12 litre (b) 0.028 litre (c) 0.56 litre (d) 1.12 litre

165. A sample of 1.0 g of solid Fe_2O_3 of 80% purity is dissolved in a moderately concentrated HCl solution which is reduced by zinc dust. The resulting solution required 16.7 mL of a 0.1 *M* solution of the oxidant. Calculate the mole of electrons taken up by the oxidant.

(a) 2 (b) 4 (c) 6 (d) 5

166. $KMnO_4$ reacts with oxalic acid according to the equation

$$2MnO_4^- \quad 5C_2O_4^{2-} \quad 16H^+ \qquad 2Mn^{2+} + 10CO_2 + 8H_2O$$

Here, 20 mL of 0.1 *M* $KMnO_4$ is equivalent to :

(a) 120 mL of 0.25 *M* $H_2C_2O_4$ (b) 150 mL of 0.10 *M* $H_2C_2O_4$

(c) 25 mL of 0.20 *M* $H_2C_2O_4$ (d) 50 mL of 0.20 *M* $H_2C_2O_4$

For More Material Join: @JEEAdvanced_2024

167. Ratio of moles of Fe (II) oxidised by equal volumes of equimolar $KMnO_4$ and $K_2Cr_2O_7$ solutions in acidic medium will be:
(a) 5 : 3 (b) 1 : 1 (c) 1 : 2 (d) 5 : 6

168. The mass of a mixture containing HCl and H_2SO_4 is 0.1 g. On treatment with an excess of an $AgNO_3$ solution, reacted with this acid mixture gives 0.1435 g of AgCl. Mass % of the H_2SO_4 mixture is :
(a) 36.5 (b) 63.5 (c) 50 (d) None of these

169. A solution of $Na_2S_2O_3$ is standardized iodometrically against 0.167 g of $KBrO_3$. This process requires 50 mL of the $Na_2S_2O_3$ solution. What is the normality of the $Na_2S_2O_3$?
(a) 0.2 *N* (b) 0.12 *N* (c) 0.72 *N* (d) 0.02 *N*

170. 0.80 g of impure $(NH_4)_2SO_4$ was boiled with 100 mL of a 0.2 *N* NaOH solution till all the $NH_3(g)$ evolved. The remaining solution was diluted to 250 mL. 25 mL of this solution was neutralized using 5 mL of a 0.2 *N* H_2SO_4 solution. The percentage purity of the $(NH_4)_2SO_4$ sample is:
(a) 82.5 (b) 72.5 (c) 62.5 (d) 17.5

171. The NH_3 evolved due to complete conversion of N from 1.12 g sample of protein was absorbed in 45 mL of 0.4 *N* HNO_3. The excess acid required 20 mL of 0.1 *N* NaOH. The % N in the sample is:
(a) 8 (b) 16 (c) 20 (d) 25

172. Find out % of oxalate ion in a given sample of an alkali metal oxalate salt, 0.30 g of it is dissolved in 100 mL water and its required 90 mL of centimolar $KMnO_4$ solution in acidic medium :
(a) 66% (b) 55% (c) 44% (d) 6.6%

173. 320 mg of a sample of magnesium having a coating of its oxide required 20 mL of 0.1 *M* hydrochloric acid for the complete neutralisation of the latter. The composition of the sample is:
(a) 87.5% Mg and 12.5% MgO (b) 12.5% Mg and 87.5% MgO
(c) 80% Mg and 20% MgO (d) 20% Mg and 80% MgO

174. The concentration of bivalent lead ions in a sample of polluted water that also contains nitrate ions is determined by adding solid sodium sulphate (M = 142) to exactly 500 mL water. Calculate the molarity of lead ions if 0.355 g of sodium sulphate was needed for complete precipitation of lead ions as sulphate.
(a) 1.25 10^{-3} *M* (b) 2.5 10^{-3} *M* (c) 5 10^{-3} *M* (d) None of these

175. What volume of HNO_3 (sp. gravity 1.05 g mL $^{-1}$ containing 12.6 (*w/W*) of HNO_3) that reduce into NO is required to oxidise iron 1 g $FeSO_4$ $7H_2O$ in acid medium is:
(a) 70 mL (b) 0.57 mL (c) 80 mL (d) 0.65 mL

176. The total volume of 0.1 *M* $KMnO_4$ solution that are needed to oxidize 100 mg each of ferrous oxalate and ferrous sulphate in a mixture in acidic medium is:
(a) 1.096 mL (b) 1.32 mL (c) 5.48 mL (d) none of these

177. When 2.5 g of a sample of Mohr's salt reacts completely with 50 mL of $\frac{N}{10}$ $KMnO_4$ solution. The % purity of the sample of Mohr's salt is :
(a) 78.4 (b) 70 (c) 37 (d) 40

178. 4 mole of a mixture of Mohr's salt and $Fe_2(SO_4)_3$ requires 500 mL of 1 *M* $K_2Cr_2O_7$ for complete oxidation in acidic medium. The mole % of the Mohr's salt in the mixture is:
(a) 25 (b) 50 (c) 60 (d) 75

For More Material Join: @JEEAdvanced_2024

179. The equivalent mass of a metal is twice to that of oxygen. How many times is the equivalent mass of it's oxide than the equivalent mass of the metal ?
(a) 1.5 (b) 2 (c) 3 (d) 4

180. A metal oxide has the formula X_2O_3. It can be reduced by hydrogen to give free metal and water. 0.1596 g of metal oxide requires 6 mg of hydrogen for complete reduction. The atomic mass of the metal (in amu) is:
(a) 15.58 (b) 155.8 (c) 5.58 (d) 55.8

181. Calculate the mass of anhydrous oxalic acid, which can be oxidised to $CO_2(g)$ by 100 mL of an MnO_4 solution, 10 mL of which is capable of oxidising 50 mL of 1N I to I_2.
(a) 45 g (b) 22.5 g (c) 30 g (d) 12.25 g

182. A mixture of $NaHC_2O_4$ and $KHC_2O_4\,H_2C_2O_4$ required equal volumes of 0.2 N $KMnO_4$ and 0.12 N NaOH separately. What is the molar ratio of $NaHC_2O_4$ and $KHC_2O_4\,H_2C_2O_4$ in the mixture ?
(a) 6 : 1 (b) 1 : 6 (c) 1 : 3 (d) 3 : 1

183. Stannous sulphate ($SnSO_4$) and potassium permanganate are used as oxidising agents in acidic medium for oxidation of ferrous ammonium sulphate to ferric sulphate. The ratio of number of moles of stannous sulphate required per mole of ferrous ammonium sulphate to the number of moles of $KMnO_4$ required per mole of ferrous ammonium sulphate, is :
(a) 2.5 (b) 0.2 (c) 0.4 (d) 2.0

184. If a g is the mass of $NaHC_2O_4$ required to neutralize 100 mL of 0.2 M NaOH and b g that required to reduce 100 mL of 0.02 M $KMnO_4$ in acidic medium, then:
(a) $a = b$ (b) $2a = b$ (c) $a = 2b$ (d) None of these

185. 2 mole, equimolar mixture of $Na_2C_2O_4$ and $H_2C_2O_4$ required V_1 L of 0.1 M $KMnO_4$ in acidic medium for complete oxidation. The same amount of the mixture required V_2 L of 0.2 M NaOH for neutralization. The ratio of V_1 to V_2 is:
(a) 1 : 2 (b) 2 : 1 (c) 4 : 5 (d) 5 : 4

186. A mixture containing 0.05 mole of $K_2Cr_2O_7$ and 0.02 mole of $KMnO_4$ was treated with excess of KI in acidic medium. The liberated iodine required 1.0 L of $Na_2S_2O_3$ solution for titration. Concentration of $Na_2S_2O_3$ solution was:
(a) 0.40 mol L^{-1} (b) 0.20 mol L^{-1} (c) 0.25 mol L^{-1} (d) 0.30 mol L^{-1}

187. 25 mL of 2 N HCl, 50 mL of 4 N HNO_3 and x mL of 2 M H_2SO_4 are mixed together and the total volume is made up to 1 L after dilution. 50 mL of this acid mixture completely reacted with 25 mL of a 1 N Na_2CO_3 solution. The value of x is:
(a) 250 mL (b) 62.5 mL (c) 100 mL (d) None of these

188. In an iodometric estimation, the following reactions occur
$2Cu^{2+} + 4I \quad Cu_2I_2 + I_2; \quad I_2 + 2Na_2S_2O_3 \quad 2NaI + Na_2S_4O_6$
0.12 mole of $CuSO_4$ was added to excess of KI solution and the liberated iodine required 120 mL of hypo. The molarity of hypo solution was:
(a) 2 (b) 0.20 (c) 0.1 (d) 1.0

189. 1 g mixture of equal number of mole of Li_2CO_3 and other metal carbonate (M_2CO_3) required 21.6 mL of 0.5 N HCl for complete neutralisation reaction. What is the approximate atomic mass of the other metal ?
(a) 25 (b) 23 (c) 51 (d) 118

For More Material Join: @JEEAdvanced_2024

190. 32 g of a sample of $FeSO_4\ 7H_2O$ were dissolved in dilute sulphuric acid and water and its volume was made up to 1 litre. 25 mL of this solution required 20 mL of 0.02 *M* $KMnO_4$ solution for complete oxidation. Calculate the mass % of $FeSO_4\ 7H_2O$ in the sample.
(a) 34.75 (b) 69.5
(c) 89.5 (d) None of these

191. In the mixture of ($NaHCO_3 + Na_2CO_3$), volume of HCl required is *x* mL with phenolphthalein indicator and *y* mL with methyl orange indicator in the same titration. Hence, volume of HCl for complete reaction of Na_2CO_3 is:
(a) 2*x* (b) *y*
(c) *x*/2 (d) (*y* *x*)

192. 0.1 g of a solution containing Na_2CO_3 and $NaHCO_3$ requires 10 mL of 0.01 *N* HCl for neutralization using phenolphthalein as an indicator. Mass % of Na_2CO_3 in solution is:
(a) 25 (b) 32
(c) 50 (d) None of these

193. A mixture of NaOH and Na_2CO_3 required 25 mL of 0.1 *M* HCl using phenolphthalein as the indicator. However, the same amount of the mixture required 30 mL of 0.1 *M* HCl when methyl orange was used as the indicator. The molar ratio of NaOH and Na_2CO_3 in the mixture was:
(a) 2 : 1 (b) 1 : 2 (c) 4 : 1 (d) 1 : 4

194. When 100 mL solution of NaOH and Na_2CO_3 was first titrated with *N*/10 HCl in presence of HPh, 17.5 mL were used till end point is obtained. After this end point MeOH was added and 2.5 mL of same HCl were required to attain new end point. The amount of NaOH in mixture is:
(a) 0.06 g per 100 mL (b) 0.06 g per 200 mL
(c) 0.05 g per 100 mL (d) 0.012 g per 200 mL

195. 1 gram of a sample of $CaCO_3$ was strongly heated and the CO_2 liberated was absorbed in 100 mL of 0.5 *M* NaOH solution. Assuming 90% purity for the sample, how many mL of 0.5 *M* HCl would be required to react with the resulting solution to reach the end point in presence of phenolphthalein?
(a) 73 mL (b) 41 mL
(c) 82 mL (d) 100 mL

196. A sample of pure sodium carbonate 0.318 g is dissolved in water and titrated with HCl solution. A volume of 60 mL is required to reach the methyl orange end point. Calculate the molarity of the acid.
(a) 0.1 *M* (b) 0.2 *M*
(c) 0.4 *M* (d) None of these

197. 10 L of hard water required 5.6 g of lime for removing hardness. Hence temporary hardness in ppm of $CaCO_3$ is:
(a) 1000 (b) 2000
(c) 100 (d) 1

198. 1 L of pond water contains 20 mg of Ca^{2} and 12 mg of Mg^{2+} ions. What is the volume of a 2 *N* Na_2CO_3 solution required to soften 5000 L of pond water?
(a) 500 L (b) 50 L (c) 5 L (d) None of these

For More Material Join: @JEEAdvanced_2024

199. One litre of a sample of hard water contain 4.44 mg $CaCl_2$ and 1.9 mg of $MgCl_2$. What is the total hardness in terms of ppm of $CaCO_3$?

(a) 2 ppm (b) 3 ppm

(c) 4 ppm (d) 6 ppm

200. If hardness of water sample is 200 ppm, then select the incorrect statement :

(a) Mass ratio of $CaCO_3$ to H_2O is $\frac{0.02}{100}$

(b) Mole ratio of $CaCO_3$ to H_2O is $3.6 \quad 10^{5}$

(c) Mass of $CaCO_3$ present in hard water is 0.2 g/L

(d) 1 miliequivalent of $CaCO_3$ present in 1 kg of hard water

For More Material Join: @JEEAdvanced_2024



Level 2

1. A mixture of NH_4NO_3 and $(NH_4)_2HPO_4$ contain 30.40% mass per cent of nitrogen. What is the mass ratio of the two components in the mixture ?

(a) 2 : 1 (b) 1 : 2 (c) 3 : 4 (d) 4 : 1

2. What volume of 75% alcohol by weight ($d = 0.80\ g/cm^3$) must be used to prepare 150 cm^3 of 30% alcohol by mass ($d = 0.90\ g/cm^3$)?

(a) 67.5 mL (b) 56.25 mL (c) 44.44 mL (d) None of these

3. Calculate the number of millilitres of NH_3 (aq) solution ($d = 0.986\ g/mL$) contain 2.5% by mass NH_3, which will be required to precipitate iron as $Fe(OH)_3$ in a 0.8 g sample that contains 50% Fe_2O_3.

(a) 0.344 mL (b) 3.44 mL (c) 17.24 mL (d) 10.34 mL

4. In the preparation of iron from haematite (Fe_2O_3) by the reduction with carbon

$$Fe_2O_3 + C \qquad Fe + CO_2$$

how much 80% pure iron may be produced from 120 kg of 90% pure Fe_2O_3 ?

(a) 94.5 kg (b) 60.48 kg (c) 116.66 kg (d) 120 kg

5. A mineral consists of an equimolar mixture of the carbonates of two bivalent metals. One metal is present to the extent of 12.5% by mass. 2.8 g of the mineral on heating lost 1.32 g of CO_2. What is the % by mass of the other metal ?

(a) 87.5 (b) 35.71 (c) 65.11 (d) 23.21

6. 6.2 g of a sample containing Na_2CO_3, $NaHCO_3$ and non-volatile inert impurity on gentle heating loses 5% of its mass due to reaction $2NaHCO_3 \quad Na_2CO_3 \quad H_2O + CO_2$. Residue is dissolved in water and formed 100 mL solution and its 10 mL portion requires 7.5 mL of 0.2 M aqueous solution of $BaCl_2$ for complete precipitation of carbonates.

Determine mass (in gram) of Na_2CO_3 in the original sample.

(a) 1.59 (b) 1.06 (c) 0.53 (d) None of these

7. Nitric acid can be produced from NH_3 in three steps process given below

(I) $4NH_3(g) + 5O_2(g) \qquad 4NO(g) + 6H_2O(g)$

(II) $2NO(g) + O_2(g) \qquad 2NO_2(g)$


(III) $3NO_2(g) + H_2O(l) \qquad 2HNO_3(aq) + NO(g)$

percent yield of 1st, 2nd and 3rd steps are respectively 50%, 60% and 80% respectively then what volume of $NH_3(g)$ at 1 atm and 0°C required to produced 1575 g of HNO_3.

(a) 156.25 (b) 350 L (c) 3500 L (d) None of these

8. 1 M NaOH solution was slowly added into 1000 mL of 183.75 g impure H_2SO_4 solution and the following plot was obtained. The percentage purity of H_2SO_4 sample and slope of the curve respectively are:

For More Material Join: @JEEAdvanced_2024



(a) 75%, $\frac{1}{3}$ (b) 80%, $\frac{1}{2}$ (c) 80%, 1 (d) None of these

9. MnO_2 on ignition converts into Mn_3O_4. A sample of pyrolusite having 75% MnO_2, 20% inert impurities and rest water is ignited in air to constant mass. What is the percentage of Mn in the ignited sample ?

(a) 24.6% (b) 37% (c) 55.24% (d) 74.05%

10. A 1.0 g sample of a pure organic compound containing chlorine is fused with Na_2O_2 to convert chlorine to NaCl. The sample is then dissolved in water, and the chloride precipitated with $AgNO_3$, giving 1.96 g of AgCl. If the molecular mass of organic compound is 147, how many chlorine atoms does each molecule contain ?

(a) 1 (b) 2 (c) 3 (d) 4

11. A 0.60 g sample consisting of only CaC_2O_4 and MgC_2O_4 is heated at 500°C, converting the two salts of $CaCO_3$ and $MgCO_3$. The sample then weighs 0.465 g. If the sample had been heated to 900°C, where the products are CaO and MgO. What would the mixtures of oxides have weighed ?

(a) 0.12 g (b) 0.21 g (c) 0.252 g (d) 0.3 g

12. A metal M forms the sulphate $M_2(SO_4)_3$. A 0.596 gram sample of the sulphate reacts with excess $BaCl_2$ to give 1.220 g $BaSO_4$. What is the atomic mass of M ?

(Atomic mass : S = 32, Ba = 137.3)

(a) 26.9 (b) 69.7 (c) 55.8 (d) 23

13. Urea (H_2NCONH_2) is manufactured by passing $CO_2(g)$ through ammonia solution followed by crystallization. CO_2 for the above reaction is prepared by combustion of hydrocarbons. If combustion of 236 kg of a saturated hydrocarbon (C_nH_{2n+2}) produces as much CO_2 as required for production of 999.6 kg urea then molecular formula of hydrocarbon is:

(a) $C_{10}H_{22}$ (b) $C_{12}H_{26}$ (c) $C_{13}H_{28}$ (d) C_8H_{18}

14. 11.6 g of an organic compound having formula C_nH_{2n+2} is burnt in excess of $O_2(g)$ initially taken in a 22.41 litre steel vessel. Before reaction the gaseous mixture was at 273 K with pressure reading 2 atm. After complete combustion and loss of considerable amount of heat, the mixture of product and excess of O_2 had a temperature of 546 K and 4.6 atm pressure. The formula of organic compound is:

(a) C_2H_6 (b) C_3H_8 (c) C_5H_{12} (d) C_4H_{10}

For More Material Join: @JEEAdvanced_2024

15. $H_2O_2 + 2KI \xrightarrow{40\%\text{ yield}} I_2 + 2KOH$

$H_2O_2 + 2KMnO_4 + 3H_2SO_4 \xrightarrow{50\%\text{ yield}} K_2SO_4 + 2MnSO_4 + 3O_2 + 4H_2O$

150 mL of H_2O_2 sample was divided into two parts. First part was treated with KI and formed KOH required 200 mL of $M/2$ H_2SO_4 for neutralisation. Other part was treated with $KMnO_4$ yielding 6.74 litre of O_2 at 1 atm. and 273 K. Using % yield indicated find volume strength of H_2O_2 sample used.

(a) 5.04 (b) 10.08 (c) 3.36 (d) 33.6

16. SO_2Cl_2 (sulphuryl chloride) reacts with water to given a mixture of H_2SO_4 and HCl. What volume of 0.2 M $Ba(OH)_2$ is needed to completely neutralize 25 mL of 0.2 M SO_2Cl_2 solution:

(a) 25 mL (b) 50 mL (c) 100 mL (d) 200 mL

17. 5 g sample contain only Na_2CO_3 and Na_2SO_4. This sample is dissolved and the volume made up to 250 mL. 25 mL of this solution neutralizes 20 mL of 0.1 M H_2SO_4.

Calculate the % of Na_2SO_4 in the sample.

(a) 42.4 (b) 57.6 (c) 36.2 (d) none of these

18. 20 mL of 0.2 M NaOH(aq) solution is mixed with 35 mL of 0.1M NaOH(aq) solution and the resultant solution is diluted to 100 mL. 40 mL of this diluted solution reacted with 10% impure sample of oxalic acid ($H_2C_2O_4$). The mass of impure sample is:

(a) 0.15 gram (b) 0.135 gram (c) 0.59 gram (d) None of these

19. A silver coin weighing 11.34 g was dissolved in nitric acid. When sodium chloride was added to the solution all the silver (present as $AgNO_3$) was precipitated as silver chloride. The mass of the precipitated silver chloride was 14.35 g. Calculate the percentage of silver in the coin.

(a) 4.8% (b) 95.2% (c) 90% (d) 80%

20. Two elements X (at. mass 16) and Y (at. mass 14) combine to form compounds A, B and C. The ratio of different masses of Y which combine with a fixed mass of X in A, B and C is 1 : 3 : 5. If 32 parts by mass of X combines with 84 parts by mass of Y in B, then in C, 16 parts by mass of X will combine with :

(a) 14 parts by mass of Y (b) 42 parts by mass of Y

(c) 70 parts by mass of Y (d) 84 parts by mass of Y

21. The conversion of oxygen to ozone occurs to the extent of 15% only. The mass of ozone that can be prepared from 67.2 L of oxygen at 1 atm and 273 K will be :

(a) 14.4 g (b) 96 g (c) 640 g (d) 64 g

22. RH_2 (ion exchange resin) can replace Ca^{2+} ions in hard water as $RH_2 + Ca^{2+}$ $RCa + 2H^+$. If 1 L of hard water after passing through RH_2 has pH = 3 then hardness in parts per million of Ca^{2+} is:

(a) 20 (b) 10 (c) 40 (d) 100

23. 100 cm^3 of a solution of an acid (Molar mass = 98) containing 29.4 g of the acid per litre were completely neutralized by 90.0 cm^3 of aq. NaOH containing 20 g of NaOH per 500 cm^3. The basicity of the acid is:

(a) 3 (b) 2 (c) 1 (d) data insufficient

For More Material Join: @JEEAdvanced_2024

24. 20 mL of 0.1 *M* solution of compound Na_2CO_3 $NaHCO_3$ $2H_2O$ is titrated against 0.05 *M* HCl. *x* mL of HCl is used when phenolphthalein is used as an indicator and *y* mL of HCl is used when methyl orange is the indicator in two separate titrations. Hence (*y* *x*) is:

(a) 40 mL (b) 80 mL
(c) 120 mL (d) None of these

25. A sample containing $HAsO_2$ (mol. mass 108) and weighing 3.78 g is dissolved and diluted to 250 mL in a volumetric flask. A 50 mL sample (aliquot) is withdrawn with a pipet and titrated with 35 mL of 0.05 *M* solution of I_2. Calculate the percentage $HAsO_2$ in the sample:

(a) 25% (b) 20%
(c) 10% (d) none of these

26. A mixture of FeO and Fe_2O_3 is completely reacted with 100 mL of 0.25 *M* acidified $KMnO_4$ solution. The resultant solution was then treated with Zn dust which converted Fe^{3+} of the solution to Fe^{2+}. The Fe^{2+} required 1000 mL of 0.10 *M* $K_2Cr_2O_7$ solution. Find out the weight % Fe_2O_3 in the mixture.

(a) 80.85 (b) 19.15
(c) 50 (d) 89.41

27. To a 10 mL, 1 *M* aqueous solution of Br_2, excess of NaOH is added so that all Br_2 is disproportionated to Br and BrO_3. The resulting solution is free from Br , by extraction and excess of OH neutralised by acidifying the solution. The resulting solution is sufficient to react with 2 g of impure CaC_2O_4 (M = 128 g/mol) sample. The % purity of oxalate sample is:

(a) 85.3% (b) 12.5%
(c) 90% (d) 64%

28. 0.10 g of a sample containing $CuCO_3$ and some inert impurity was dissolved in dilute sulphuric acid and volume made up to 50 mL. This solution was added into 50 mL of 0.04 *M* KI solution where copper precipitates as CuI and I is oxidized into I_3. A 10 mL portion of this solution is taken for analysis, filtered and made up free I_3 and then treated with excess of acidic permanganate solution. Liberated iodine required 20 mL of 2.5 *mM* sodium thiosulphate solution to reach the end point.

Determine mass percentage of $CuCO_3$ in the original sample.

(a) 7.41 (b) 74.1
(c) 61.75 (d) None of these

29. 1 mole of equimolar mixture of ferric oxalate and ferrous oxalate requires *x* mole of $KMnO_4$ in acidic medium for complete oxidation. *x* is:

(a) 0.5 mole (b) 0.9 mole
(c) 1.2 mole (d) 4.5 mole

30. An impure sample of sodium oxalate ($Na_2C_2O_4$) weighing 0.20 g is dissolved in aqueous solution of H_2SO_4 and solution is titrated at 70°C, requiring 45 mL of 0.02 *M* $KMnO_4$ solution. The end point is overrun, and back titration in carried out with 10 mL of 0.1 *M* oxalic acid solution. Find the % purity of $Na_2C_2O_4$ in sample:

(a) 75 (b) 83.75
(c) 90.25 (d) None of these

For More Material Join: @JEEAdvanced_2024

31. 0.5 g mixture of $K_2Cr_2O_7$ and $KMnO_4$ was treated with excess of KI in acidic medium. Iodine liberated required 150 cm^3 of 0.10 *N* solution of thiosulphate solution for titration.

Find the percentage of $K_2Cr_2O_7$ in the mixture:

(a) 14.64 (b) 34.2

(c) 65.69 (d) 50

32. A 150 mL of solution of I_2 is divided into two unequal parts. I part reacts with hypo solution in acidic medium. 15 mL of 0.4 *M* hypo was consumed. II part was added with 100 mL of 0.3 *M* NaOH solution. Residual base required 10 mL of 0.3 *M* H_2SO_4 solution for complete neutralization. What was the initial concentration of I_2 ?

(a) 0.08 *M* (b) 0.1 *M*


(c) 0.2 *M* (d) None of these

33. A mixture of H_2SO_4 and $H_2C_2O_4$ (oxalic acid) and some inert impurity weighing 3.185 g was dissolved in water and the solution made up to 1 litre. 10 mL of this solution required 3 mL of 0.1 *N* NaOH for complete neutralization. In another experiment 100 mL of the same solution in hot condition required 4 mL of 0.02 *M* $KMnO_4$ solution for complete reaction. The mass % of H_2SO_4 in the mixture was:

(a) 40 (b) 50

(c) 60 (d) 80

34. During developing of an exposed camera film, one step involves in the following reaction :



(Quinol)

Which of the following best describes the role of quinol ?

(a) It acts as an acid (b) It acts as reducing agent

(c) It acts as oxidant (d) It acts as a base

35. The concentration of an oxalic acid solution is '*x*' mol litre ^{1}. 40 mL of this solution reacts with 16 mL of 0.05 M acidified $KMnO_4$. What is the pH of '*x*' *M* oxalic acid solution ? (Assume that oxalic acid dissociates completely.)

(a) 1.3 (b) 1.699

(c) 1 (d) 2

For More Material Join: @JEEAdvanced_2024

Level 3

Passage 1

Oleum is considered as a solution of SO_3 in H_2SO_4, which is obtained by passing SO_3 in solution of H_2SO_4. When 100 g sample of oleum is diluted with desired mass of H_2O then the total mass of H_2SO_4 obtained after dilution is known as % labelling in oleum.

For example, a oleum bottle labelled as '109% H_2SO_4' means the 109 g total mass of pure H_2SO_4 will be formed when 100 g of oleum is diluted by 9 g of H_2O which combines with all the free SO_3 present in oleum to form H_2SO_4 as $SO_3 + H_2O \quad H_2SO_4$.

1. What is the % of free SO_3 in an oleum that is labelled as '104.5% H_2SO_4' ?
(a) 10 (b) 20 (c) 40 (d) None of these

2. 9.0 g water is added into oleum sample labelled as "112%" H_2SO_4 then the amount of free SO_3 remaining in the solution is : (STP 1 atm and 273 K)
(a) 14.93 L at STP (b) 7.46 L at STP (c) 3.73 L at STP (d) 11.2 L at STP

3. If excess water is added into a bottle sample labelled as "112% H_2SO_4" and is reacted with 5.3 g Na_2CO_3, then find the volume of CO_2 evolved at 1 atm pressure and 300 K temperature after the completion of the reaction :
(a) 2.46 L (b) 24.6 L (c) 1.23 L (d) 12.3 L

4. 1 g of oleum sample is diluted with water. The solution required 54 mL of 0.4 *N* NaOH for complete neutralization. The % of free SO_3 in the sample is :
(a) 74 (b) 26 (c) 20 (d) None of these

Passage 2

The strength of H_2O_2 is expressed in several ways like molarity, normality, % (*w/V*), volume strength, etc. The strength of "10 V" means 1 volume of H_2O_2 on decomposition gives 10 volumes of oxygen at 1 atm and 273 K or 1 litre of H_2O_2 gives 10 litre of O_2 at 1 atm and 273 K The decomposition of H_2O_2 is shown as under :

$$H_2O_2(aq) \qquad H_2O(l) \quad \frac{1}{2}O_2(g)$$

H_2O_2 can acts as oxidising as well as reducing agent. As oxidizing agent H_2O_2 is converted into H_2O and as reducing agent H_2O_2 is converted into O_2. For both cases its *n*-factor is 2.

Normality of H_2O_2 solution = 2 Molarity of H_2O_2 solution

1. What is the molarity of "11.2 *V*" H_2O_2?
(a) 1 *M* (b) 2 *M* (c) 5.6 *M* (d) 11.2 *M*

2. What is the percentage strength (% *w/V*) of "11.2 *V*" H_2O_2?
(a) 1.7 (b) 3.4 (c) 34 (d) None of these

For More Material Join: @JEEAdvanced_2024

3. 20 mL of H_2O_2 solution is reacted with 80 mL of 0.05 *M* $KMnO_4$ in acidic medium then what is the volume strength of H_2O_2?
(a) 2.8 (b) 5.6 (c) 11.2 (d) None of these

4. 40 g $Ba(MnO_4)_2$ (mol. mass = 375) sample containing some inert impurities in acidic medium completely reacts with 125 mL of "33.6 *V*" of H_2O_2. What is the percentage purity of the sample?
(a) 28.12% (b) 70.31% (c) 85% (d) None of these



Passage 3

A water is said to be a soft water if it produces sufficient foam with the soap and water that does not produce foam with soap is known as hard water. Hardness has been classified into two types (i) Temporary hardness (ii) Permanent hardness.

Temporary hardness is due to presence of calcium and magnesium bicarbonate. It is simply removed by boiling as

$$Ca(HCO_3)_2 \quad CaCO_3 + CO_2 + H_2O$$

$$Mg(HCO_3)_2 \quad MgCO_3 + CO_2 + H_2O$$

Temporary hardness can also be removed by addition of slaked lime, $Ca(OH)_2$

$$Ca(HCO_3)_2 + Ca(OH)_2 \quad 2CaCO_3 + 2H_2O$$

Permanent hardness is due to presence of sulphates and chlorides of Ca, Mg, etc. It is removed by washing soda as

$$CaCl_2 + Na_2CO_3 \quad CaCO_3 + 2NaCl$$

$$CaSO_4 + Na_2CO_3 \quad CaCO_3 + Na_2SO_4$$

Permanent hardness also removed by ion exchange resin process as

$$2RH \quad Ca^2 \quad R_2Ca \quad 2H$$

$$2ROH \quad SO_4^2 \quad R_2SO_4 \quad 2OH$$

The degree of hardness of water is measured in terms of ppm of $CaCO_3$. 100 ppm means 100 g of $CaCO_3$ is present in 10^6 g of H_2O. If any other water sample which contain 120 ppm of $MgSO_4$, hardness in terms of $CaCO_3$ is equal to 100 ppm.

1. One litre of a sample of hard water (d 1 g/mL) contains 136 mg of $CaSO_4$ and 190 mg of $MgCl_2$. What is the total hardness of water in terms of $CaCO_3$?
(a) 100 ppm (b) 200 ppm (c) 300 ppm (d) None of these

2. What is the mass of $Ca(OH)_2$ required for 10 litre of water remove temporary hardness of 100 ppm due to $Ca(HCO_3)_2$?
(a) 1.62 g (b) 0.74 g (c) 7.4 g (d) None of these

3. A 200 g sample of hard water is passed through the column of cation exchange resin, in which H is exchanged by Ca^2. The outlet water of column required 50 mL of 0.1 *M* NaOH for complete neutralization. What is the hardness of Ca^2 ion in ppm?
(a) 250 ppm (b) 500 ppm (c) 750 ppm (d) 1000 ppm

For More Material Join: @JEEAdvanced_2024



Passage 4

Equivalent mass $\dfrac{\text{Molecular mass / Atomic mass}}{n\text{-factor}}$

n-factor is very important in redox as well as non-redox reactions. With the help of *n*-factor we can predict the molar ratio of the reactant species taking part in reactions. The reciprocal of *n*-factor's ratio of the reactants is the molar ratio of the reactants.

In general *n*-factor of acid/base is number of moles of H /OH furnished per mole of acid/base. *n*-factor of a reactant is number of moles of electrons lost or gained per mole of reactant.

Example 1 :

1. In acidic medium : $KMnO_4$ (n 5) Mn^2
2. In neutral medium : $KMnO_4$ (n 3) Mn^2
3. In basic medium : $KMnO_4$ (n 1) Mn^6

Example 2 : FeC_2O_4 Fe^{3+} $2CO_2$

Total number of moles of e lost by 1 mole of FeC_2O_4

1 1 2 3

1. *n*-factor of $Ba(MnO_4)_2$ in acidic medium is :

(a) 2 (b) 6 (c) 10 (d) None of these

2. Consider the following reaction.

$$H_3PO_2 + NaOH \quad NaH_2PO_2 + H_2O$$

What is the equivalent mass of H_3PO_2? (mol. wt. is M)

(a) M (b) $M/2$ (c) $M/3$ (d) None of these

3. For the reaction, $Fe_{0.95}O$ (molar mass = M) Fe_2O_3, what is the eq. mass of $Fe_{0.95}O$?

(a) $\dfrac{M}{0.85}$ (b) $\dfrac{M}{0.95}$ (c) $\dfrac{M}{0.8075}$ (d) None of these

4. In the reaction, xVO yFe_2O_3 FeO V_2O_5, what is the value of x and y respectively?

(a) 1, 1 (b) 2, 3 (c) 3, 2 (d) None of these



Passage 5

Consider the following series of reactions :

Cl_2 2NaOH NaCl NaClO H_2O

3NaClO 2NaCl $NaClO_3$

$4NaClO_3$ $3NaClO_4$ NaCl

1. How much Cl_2 is required to prepare 122.5 g of $NaClO_4$ by above sequencial reactions ?

(a) 284 g (b) 213 g (c) 142 g (d) 71 g

For More Material Join: @JEEAdvanced_2024

2. How many moles of NaCl will be formed by using 1 mole Cl_2 and other reagents in excess ?
(a) $\frac{1}{12}$ mole (b) 1.67 mole (c) 1.75 mole (d) 0.75 mole

3. How many moles of $NaClO_3$ obtained after the complection of reaction by taking 1 mole Cl_2 and other reagents in excess ?
(a) $\frac{1}{3}$ mole (b) Zero (c) $\frac{1}{4}$ mole (d) 1 mole

One or More Answers is/are Correct



1. 1 g atom of nitrogen represents :
(a) 6.02 10^{23} N_2 molecules (b) 22.4 litre of N_2 at 1 atm and 273 K
(c) 11.2 litre of N_2 at 1 atm and 273 K (d) 14 g of nitrogen

2. 1 g molecule of V_2O_5 contains :
(a) 5 mole of oxygen atom (b) 2 mole of V atom
(c) 1 mole of oxygen atom (d) 2.5 mole of oxygen atom

3. Select the dimensionless quantity(ies) :
(a) vapour density (b) molality (c) specific gravity (d) mass fraction

4. Which of the following concentration terms is/are affected by a change in temperature?
(a) Molarity (b) Molality (c) Normality (d) Specific gravity

5. Which of the following statements regarding the compound A_xB_y is/are correct?
(a) 1 mole of A_xB_y contains 1 mole of A and 1 mole B
(b) 1 equivalent of A_xB_y contains 1 equivalent of A and 1 equivalent of B
(c) 1 mole of A_xB_y contains x moles of A and y moles of B
(d) equivalent mass of A_xB_y equivalent mass of A + equivalent mass of B

6. 1 mole of $Ba(OH)_2$ will exactly neutralize :
(a) 0.5 mole HCl (b) 1 mole of H_2SO_4 (c) 1 mole of H_3PO_3 (d) 2 mole of H_3PO_2

7. The pair of species having different percentage (mass)of carbon is :
(a) CH_3COOH and $C_6H_{12}O_6$ (b) CH_3COOH and C_2H_5OH
(c) $HCOOCH_3$ and HCOOH (d) C_2H_5OH and CH_3OCH_3

8. 30 mL of CH_3OH (d 0.8 g/cm^3) is mixed with 60 mL of C_2H_5OH (d 0.92 g/cm^3) at 25°C to form a solution of density 0.88 g/cm^3. Select the correct option :
(a) Molarity and molality of resulting solution are 6.33 and 13.59 respectively
(b) The mole fraction of solute and molality are 0.385 and 13.59 respectively
(c) Molarity and % change in volume are 13.59 and zero respectively
(d) Mole fraction of solvent and molality are 0.615 and 13.59 respectively

9. Which of the following is/are correct for 17 g/L of H_2O_2 solution?
(a) Volume strengths is 5.6 at 273 K and 1 atm
(b) Molarity of solution is 0.5 M
(c) 1 mL of this solution gives 2.8 mL O_2 at 273 K and 2 atm
(d) The normality of solution is 2 N

For More Material Join: @JEEAdvanced_2024

10. Solutions containing 23 g HCOOH is/are :

(a) 46 g of 70% $\frac{w}{V}$ HCOOH (d_{solution} 1.40 g/mL)

(b) 50 g of 10 M HCOOH (d_{solution} 1 g/mL)

(c) 50 g of 25% $\frac{w}{w}$ HCOOH

(d) 46 g of 5 M HCOOH (d_{solution} 1 g/mL)

11. A sample of H_2O_2 solution labelled as "28 volume" has density of 265 g/L. Mark the correct option(s) representing concentration of same solution in other units :

(a) $M_{H_2O_2}$ 2.5

(b) $\%\frac{w}{V}$ 17

(c) Mole fraction of $H_2O_2 = 0.2$

(d) $m_{H_2O_2}$ 13.88

12. A mixture of 100 mL of CO, CO_2 and O_2 was sparked. When the resulting gaseous mixture was passed through KOH solution, contraction in volume was found to be 80 mL. The composition of initial mixture may be (in the same order) :

(a) 30 mL, 60 mL, 10 mL

(b) 30 mL, 50 mL, 20 mL

(c) 50 mL, 30 mL, 20 mL

(d) 20 mL, 70 mL, 10 mL

13. If 1 mole of H_3PO_4 reacts with 1 mole of $X(OH)_2$ as shown below:

H_3PO_4 $X(OH)_2$ $XHPO_4$ $2H_2O$ then :

(a) the equivalent mass of base is $\frac{\text{mol. mass}}{2}$

(b) the eq. mass of H_3PO_4 is $\frac{98}{3}$

(c) the resulting solution requires 1 mole NaOH for complete neutralization

(d) minimum 1 mole of $X(OH)_2$ is required for complete neutralization of $XHPO_4$

14. In acidic medium dichromate ion osxidizes stannous ion as:

$$x\text{Sn}^{2} \quad y\text{Cr}_2\text{O}_7^{2} \quad z\text{H} \qquad a\text{Sn}^{4} \quad b\text{Cr}^{3} \quad c\text{H}_2\text{O}$$

(a) the value of $x : y$ is 1 : 3

(b) the value of x y z is 18

(c) $a : b$ is 3 : 2

(d) the value of z c is 7

15. When a equimolar mixture of Cu_2S and CuS is titrated with $Ba(MnO_4)_2$ in acidic medium, the final products contain Cu^{2} , SO_2 and Mn^{2} . If the mol. mass of Cu_2S, CuS and $Ba(MnO_4)_2$ are M_1, M_2 and M_3 respectively then :

(a) eq. mass of Cu_2S is $\frac{M_1}{8}$

(b) eq. mass of CuS is $\frac{M_2}{6}$

(c) eq. mass of $Ba(MnO_4)_2$ is $\frac{M_3}{5}$

(d) Cu_2S and CuS both have same equivalents in mixture

For More Material Join: @JEEAdvanced_2024

16. Which is/are **INCORRECT** statement ?

(a) Equivalent mass of H_2PO_3 is 40.5.

(b) Eq. mass of H_2PO_4 may be equal to molar mass or less than molar mass because it depends on the reaction.

(c) $KMnO_4$ has maximum eq. mass in acidic medium.

(d) Oxidation state of H in MgH_2 is greater than in H_2O_2.

Match the Column



Column-I and **Column-II** contain four entries in each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

1.

Column-I	Column-II
(A) 0.5 mole of $SO_2(g)$	(P) Occupy 11.2 L at 1 atm and 273 K
(B) 1 g of $H_2(g)$	(Q) Weighs 24 g
(C) 0.5 mole of $O_3(g)$	(R) Total no. of atoms 1.5 N_A
(D) 1 g molecule of $O_2(g)$	(S) Weighs 32 g

2.

Column-I	Column-II
(A) 44 g CO_2 gas	(P) 1g molecule
(B) 35.2 g of CH_4	(Q) N_A molecule
(C) 48 g of O_3 gas	(R) 22 N_A electrons
(D) 44 g of N_2O gas	(S) 49.28 L at 1 atm and 273 K
	(T) N_A atoms of oxygen

3.

Column-I [Atomic masses (*M*)]			Column-II (% composition of heavier isotope)
Isotope-II	Isotope-II	Average	
(A) (z 1)	(z 3)	z	(P) 25% by moles
(B) (z 1)	(z 3)	(z 2)	(Q) 50% by moles
(C) z	3z	2z	(R) % by mass dependent on z
(D) (z 1)	(z 1)	z	(S) 75% by mass

For More Material Join: @JEEAdvanced_2024

4.

Column-I	Column-II (*n*-factor)
(A) When Bi_2S_3 converted into Bi^5 and S	(P) 18
(B) When $Al_2(Cr_2O_7)_3$ reduced into Cr^3 in acidic medium	(Q) 11
(C) When FeS_2 converted into Fe_2O_3 and SO_2	(R) 2
(D) When $Mn(NO_3)_2$ converted into MnO_4^2 and NO	(S) 10

5.

Column-I	Column-II
(A) Eq. mass $\frac{\text{Molecular mass}}{33}$	(P) When CrI_3 oxidises into $Cr_2O_7^2$ and IO_4
(B) Eq. mass $\frac{\text{Molecular mass}}{27}$	(Q) When $Fe(SCN)_2$ oxidises into Fe^3, SO_4^2, CO_3^2 and NO_3
(C) Eq. mass $\frac{\text{Molecular mass}}{28}$	(R) When NH_4SCN oxidizes into SO_4^2, CO_3^2 and NO_3
(D) Eq. mass $\frac{\text{Molecular mass}}{24}$	(S) When As_2S_3 oxidises into AsO_3 and SO_4^2

6.

Column-I	Column-II
(A) $\underline{P_2H_4}$ PH_3 P_4H_2	(P) E $\frac{3M}{4}$
(B) $\underline{I_2}$ I IO_3	(Q) E $\frac{3M}{5}$
(C) MnO_4 Mn^2 H_2O $\underline{Mn_3O_4}$ H	(R) E $\frac{15M}{26}$
(D) $\underline{H_3PO_2}$ PH_3 H_3PO_3	(S) E $\frac{5M}{6}$

7. A sample of raw material contain $NaNO_3$. It contains some $NaIO_3$ also. The $NaIO_3$ can be used as a source of iodine, produced in the following reactions:

IO_3 HSO_3 I SO_4^2 ...(1)

I IO_3 I_2 H_2O ...(2)

One litre of sample solution containing 396 g of $NaIO_3$ is treated with stoichiometric quantity of $NaHSO_3$. Now a substantial amount of same solution is added to reaction mixture to bring about the reaction (2).

For More Material Join: @JEEAdvanced_2024

Column-I	Column-II
(A) n-factor of IO_3 in reaction (2)	(P) 6
(B) Number of moles of HSO_3 used in reaction (1)	(Q) 1.2
(C) Moles of I_2 produced	(R) 2
(D) Equivalents of IO_3 used in reaction (2)	(S) 5

8.

Column-I	Column-II (Type of Redox Reaction)
(A) Br_2 OH BrO_3 Br	(P) Intermolecular
(B) $FeCl_2$ $KMnO_4$ HCl $MnCl_2$ $FeCl_3$	(Q) Intramolecular
(C) Ag_2O Ag $\frac{1}{2}O_2$	(R) Disproportion
(D) NH_4NO_3 N_2O H_2O	(S) Comproportion

Assertion-Reason Type Questions



Each question contains STATEMENT-1 (Assertion) and STATEMENT-2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below

(A) If both the statement are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1

(B) If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1

(C) If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

(D) If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

1. **STATEMENT-1 :** Specific gravity is dimensionless.
STATEMENT-2 : Specific gravity is density of a substance measured w.r.t. density of water at 4°C.

2. **STATEMENT-1 :** Molarity of pure water is 55.55 M at 298 K.
STATEMENT-2 : Molarity is temperature dependent.

3. **STATEMENT-1 :** Gram molecular mass of O_2 is 32.
STATEMENT-2 : Relative atomic mass of oxygen is 32 a.m.u.

4. **STATEMENT-1 :** The oxidation state of S in $H_2S_2O_8$ is 6.
STATEMENT-2 : Maximum oxidation state of S is 6 because the maximum oxidation state of an element is equal to number of its valence electrons in it.

5. **STATEMENT-1 :** 0.1 M $H_3PO_3(aq)$ solution has normality equal to 0.3 N when completely reacted with NaOH.
STATEMENT-2 : H_3PO_3 is a dibasic acid.

For More Material Join: @JEEAdvanced_2024

6. STATEMENT-1 : MnO_2 can act as an oxidizing agent as well as reducing agent.
STATEMENT-2 : Oxidation state of Mn in MnO_2 lies between highest and lowest oxidation state.

7. STATEMENT-1 : Equivalent volume of H_2 is 11.2 L at 1 atm and 273 K.
STATEMENT-2 : 1/2 mole H_2 has produced when 1 mole of H (aq) accepted 1 mole of e .


8. STATEMENT-1 : For the reaction in titration Na_2CO_3 HCl $NaCl + NaHCO_3$, the suitable indicator is phenolphthalein.
STATEMENT-2 : Phenolphthalein provide its colour in acidic medium.

9. STATEMENT-1 : $[Fe(CN)_6]^4$ Fe^3 CO_2 NO_3, the equivalent mass of reactant is 3.74.
STATEMENT-2 : Equivalent mass of reactant $\frac{\text{Mol. mass}}{61}$.

10. STATEMENT-1 : In the balanced redox reaction,
$x As_2S_3$ $y NO_3$ $4H_2O$ $a AsO_4^3$ $b NO$ $c SO_4^2$ $8H$
the n-factor of As_2S_3 and NO_3 is 28 and 3 respectively.
STATEMENT-2 : Molar ratio is reciprocal of n-factor's ratio so $x : y$ is 3 : 28.

11. STATEMENT-1 : In the given reaction, NaOH H_3PO_4 NaH_2PO_4 H_2O equivalent mass of H_3PO_4 is $M/3$
STATEMENT-2 : H_3PO_4 is tribasic acid.

12. STATEMENT-1 : In CrO_5 oxidation number of Cr is +6.
STATEMENT-2 : CrO_5 has butterfly structure in which peroxide bonds are present.



13. STATEMENT-1 : I_2 IO_3 I , is an example of a disproportionation reaction.
STATEMENT-2 : Oxidation number of I can vary from –1 to +7.

14. STATEMENT-1 : Fluorine exists only in –1 oxidation state.
STATEMENT-2 : Fluorine has $2s^2 2p^5$ configuration.

15. STATEMENT-1 : H_2SO_4 can not act as reducing agent.
STATEMENT-2 : Sulphur can not increase its oxidation number beyond +6.

Subjective Problems



1. What volume of a liquid (in L) will contain 10 mole ? If molar mass of liquid is 280 and its density is 1.4 g/mL .

2. 16 g of SO_x gas occupies 5.6 L at 1 atm and 273 K. What will be the value of x ?

3. 200 mL of 1M HCl is mixed with 300 mL of 6 M HCl and the final solution is diluted to 1000 mL. Calculate molar concentration of [H] ion.

4. $N_2(g)$ reacts with $H_2(g)$ in either of the following ways depending upon supply of $H_2(g)$:

For More Material Join: @JEEAdvanced_2024

$N_2(g)$ $H_2(g)$ $N_2H_2(l)$

$N_2(g)$ $2H_2(g)$ $N_2H_4(g)$

If 5 L $N_2(g)$ and 3 L $H_2(g)$ are taken initially (at same temperature and pressure), calculate the contraction in volume after the reaction (in L).

5. One commercial system removes SO_2 emission from smoke at 95 C by the following set of reaction :

$$SO_2(g) \quad Cl_2(g) \qquad SO_2Cl_2(g)$$
$$SO_2Cl_2(g) \quad H_2O(l) \qquad H_2SO_4 \quad HCl$$
$$H_2SO_4 \quad Ca(OH)_2 \qquad CaSO_4 \quad H_2O$$

How many grams of $CaSO_4$ may be produced from 3.78 g of SO_2 ?

6. W is the mass of iron (in g) which will be converted into Fe_3O_4 by the action of 18 g of steam on it. What is the value of $W/7$?

$$Fe \quad H_2O \qquad Fe_3O_4 \quad H_2$$

7. Calculate the total moles of atoms of each element present in 122.5 g of $KClO_3$.

8. On dissolving 2.0 g of metal in sulphuric acid, 4.51 g of the metal sulphate was formed. The specific heat of the metal is 0.057 cal g ^{1} C ^{1}. What is the valency of the metal ?

9. One gram of a metallic chloride was found to contain 0.835 g of chlorine. Its vapour density is 85.5. If its molecular formula is M_xCl_y, then what is value of $(x \quad y)$?

10. 0.7875 g of crystalline barium hydroxide is dissolved in water. For the neutralization of this solution 20 mL of $N/4$ HNO_3 is required. How many moles of water of crystallization are present in one mole of this base ? (Given : Atomic mass Ba 137, O 16, N 14, H 1)

11. 2.0 g of polybasic organic acid (Molecular mass 600) required 100 mL of a $\frac{M}{6}$ NaOH solution for complete neutralisation. Find the basicity of acid.

12. A mixture contains 1.0 mole each of NaOH, Na_2CO_3 and $NaHCO_3$. When half of mixture is titrated with HCl, it required x mole of HCl in presence of phenolphthalein. In another experiment, half of mixture required y mole of same HCl in presence of methyl orange. Find the value of $(x \quad y)$.

13. When BrO_3 ion reacts with Br ion in acidic medium, Br_2 is liberated. Calculate the ratio of molecular mass and equivalent mass of $KBrO_3$.

14. A volume of 12.5 mL of 0.05 M SeO_2 reacts with 25 mL of 0.1 M $CrSO_4$ which is oxidised to Cr^{3} . To what oxidation state was the selenium converted by the reaction ?

15. A 0.276 g impure sample of copper ore is dissolved and Cu^{2} is titrated with KI solution. I_2 liberated required 40 mL of 0.1 M $Na_2S_2O_3$ solution for titration. What is the % of impurities in the ore ?

16. A sample of 28 mL of $H_2O_2(aq)$ solution required 10 mL of 0.1 M $KMnO_4$ (aq) solution for complete reaction in acidic medium. What is the volume strength of H_2O_2 ?

17. For the redox reaction given below, what is the value of $\frac{x}{z}$?

$xNO_3 \quad y\,As_2S_3 \quad zH_2O \qquad --- AsO_4^{3} \quad ---NO \quad --- SO_4^{2} \quad ---H$.

18. On heating 0.220 g of a metallic oxide in presence of hydrogen, 0.045 g of water is formed. If the equivalent mass of the metal is E, then what is the value of $E/9$?

For More Material Join: @JEEAdvanced_2024

19. 10 g mixture of $K_2Cr_2O_7$ and $KMnO_4$ was treated with excess of KI in acidic medium. Iodine liberated required 100 cm^3 of 2.2 *N* sodium thiosulphate solution for titration. If the mass percent of $KMnO_4$ in the mixture *Z*, then what is the value of $2Z/5$?

20. In an ore the only oxidisable material is Sn^2 . This ore is titrated with a dichromate solution containing 2.5 g $K_2Cr_2O_7$ in 0.50 litre. A 0.40 g of sample of the ore required 10.0 cm^3 of the titrant to reach equivalent point. If the percentage of tin in ore is x, then what is the value of $x/5$? (K = 39.1, Cr = 52, Sn = 118.7)

For More Material Join: @JEEAdvanced_2024

Answers

Level 1

1.	(c)	**2.**	(a)	**3.**	(b)	**4.**	(d)	**5.**	(c)	**6.**	(d)	**7.**	(c)	**8.**	(c)	**9.**	(b)	**10.**	(b)
11.	(d)	**12.**	(b)	**13.**	(b)	**14.**	(a)	**15.**	(d)	**16.**	(d)	**17.**	(d)	**18.**	(d)	**19.**	(d)	**20.**	(a)
21.	(a)	**22.**	(c)	**23.**	(b)	**24.**	(b)	**25.**	(a)	**26.**	(d)	**27.**	(d)	**28.**	(c)	**29.**	(c)	**30.**	(c)
31.	(b)	**32.**	(a)	**33.**	(a)	**34.**	(a)	**35.**	(b)	**36.**	(d)	**37.**	(b)	**38.**	(d)	**39.**	(c)	**40.**	(a)
41.	(c)	**42.**	(d)	**43.**	(d)	**44.**	(a)	**45.**	(b)	**46.**	(a)	**47.**	(c)	**48.**	(d)	**49.**	(a)	**50.**	(a)
51.	(c)	**52.**	(c)	**53.**	(c)	**54.**	(a)	**55.**	(c)	**56.**	(a)	**57.**	(d)	**58.**	(c)	**59.**	(d)	**60.**	(c)
61.	(c)	**62.**	(d)	**63.**	(d)	**64.**	(b)	**65.**	(d)	**66.**	(b)	**67.**	(d)	**68.**	(a)	**69.**	(c)	**70.**	(d)
71.	(c)	**72.**	(a)	**73.**	(d)	**74.**	(d)	**75.**	(d)	**76.**	(b)	**77.**	(a)	**78.**	(a)	**79.**	(b)	**80.**	(c)
81.	(c)	**82.**	(d)	**83.**	(a)	**84.**	(b)	**85.**	(c)	**86.**	(d)	**87.**	(a)	**88.**	(d)	**89.**	(d)	**90.**	(b)
91.	(c)	**92.**	(a)	**93.**	(a)	**94.**	(c)	**95.**	(a)	**96.**	(b)	**97.**	(b)	**98.**	(c)	**99.**	(c)	**100.**	(c)
101.	(d)	**102.**	(c)	**103.**	(a)	**104.**	(c)	**105.**	(a)	**106.**	(b)	**107.**	(b)	**108.**	(b)	**109.**	(b)	**110.**	(a)
111.	(b)	**112.**	(d)	**113.**	(c)	**114.**	(d)	**115.**	(b)	**116.**	(c)	**117.**	(c)	**118.**	(a)	**119.**	(b)	**120.**	(b)
121.	(c)	**122.**	(a)	**123.**	(b)	**124.**	(d)	**125.**	(b)	**126.**	(d)	**127.**	(b)	**128.**	(c)	**129.**	(a)	**130.**	(c)
131.	(c)	**132.**	(a)	**133.**	(d)	**134.**	(b)	**135.**	(d)	**136.**	(d)	**137.**	(a)	**138.**	(a)	**139.**	(a)	**140.**	(d)
141.	(a)	**142.**	(c)	**143.**	(b)	**144.**	(b)	**145.**	(d)	**146.**	(d)	**147.**	(d)	**148.**	(b)	**149.**	(b)	**150.**	(a)
151.	(b)	**152.**	(d)	**153.**	(c)	**154.**	(c)	**155.**	(a)	**156.**	(a)	**157.**	(b)	**158.**	(b)	**159.**	(c)	**160.**	(d)
161.	(a)	**162.**	(b)	**163.**	(c)	**164.**	(c)	**165.**	(c)	**166.**	(c)	**167.**	(d)	**168.**	(b)	**169.**	(b)	**170.**	(a)
171.	(c)	**172.**	(a)	**173.**	(a)	**174.**	(c)	**175.**	(b)	**176.**	(c)	**177.**	(a)	**178.**	(d)	**179.**	(a)	**180.**	(d)
181.	(b)	**182.**	(d)	**183.**	(a)	**184.**	(d)	**185.**	(c)	**186.**	(a)	**187.**	(b)	**188.**	(d)	**189.**	(d)	**190.**	(b)
191.	(a)	**192.**	(d)	**193.**	(c)	**194.**	(a)	**195.**	(c)	**196.**	(a)	**197.**	(a)	**198.**	(c)	**199.**	(d)	**200.**	(d)

For More Material Join: @JEEAdvanced_2024



Level 2

1.	(a)	**2.**	(a)	**3.**	(d)	**4.**	(a)	**5.**	(d)	**6.**	(b)	**7.**	(c)	**8.**	(c)	**9.**	(c)	**10.**	(b)
11.	(c)	**12.**	(a)	**13.**	(b)	**14.**	(d)	**15.**	(d)	**16.**	(b)	**17.**	(b)	**18.**	(a)	**19.**	(b)	**20.**	(c)
21.	(a)	**22.**	(a)	**23.**	(a)	**24.**	(b)	**25.**	(a)	**26.**	(a)	**27.**	(d)	**28.**	(b)	**29.**	(b)	**30.**	(b)
31.	(a)	**32.**	(b)	**33.**	(a)	**34.**	(b)	**35.**	(c)										



Level 3

Passage–1	**1.**	(b)	**2.**	(c)	**3.**	(c)	**4.** (b)
Passage–2	**1.**	(a)	**2.**	(b)	**3.**	(b)	**4.** (b)
Passage–3	**1.**	(c)	**2.**	(b)	**3.**	(b)	
Passage–4	**1.**	(c)	**2.**	(a)	**3.**	(a)	**4.** (b)
Passage–5	**1.**	(a)	**2.**	(c)	**3.**	(b)	

One or More Answers is/are correct

1. (c,d) **2.** (a,b) **3.** (a,c,d) **4.** (a,c,d) **5.** (b,c,d) **6.** (b,c,d) **7.** (b,c) **8.** (b,d)

9. (a,b,c) **10.** (a,b) **11.** (a,c,d) **12.** (a,b) **13.** (a,c) **14.** (b,c,d) **15.** (a,b) **16.** (a,c,d)

Match the Column

1.	A	P, R, S;	B	P;	C	P, Q, R;	D	S
2.	A	P, Q, R;	B	R, S ;	C	P, Q;	D	P, Q, R, T
3.	A	P, R;	B	Q, R;	C	Q, S;	D	Q, R
4.	A	S;	B	P;	C	Q;	D	R
5.	A	Q;	B	P;	C	S;	D	R
6.	A	S;	B	Q;	C	R;	D	P
7.	A	S;	B	P;	C	Q;	D	R
8.	A	P, R;	B	P;	C	Q;	D	Q, S

Assertion-Reason Type Questions

1. (A) **2.** (B) **3.** (C) **4.** (A) **5.** (D) **6.** (A) **7.** (A) **8.** (C) **9.** (D) **10.** (A)

11. (D) **12.** (A) **13.** (B) **14.** (B) **15.** (A)

Subjective Problems

1.	2	**2.**	2	**3.**	2	**4.**	6	**5.**	8	**6.**	6	**7.**	5	**8.**	3	**9.**	5	**10.**	8
11.	5	**12.**	3	**13.**	5	**14.**	0	**15.**	8	**16.**	1	**17.**	7	**18.**	4	**19.**	6	**20.**	3

For More Material Join: @JEEAdvanced_2024

Hints and Solutions

Level 1

1. (c) Number of neutrons $(17 - 8) \times 12 \times 10^{25}$
$9 \times 2 \times 6 \times 10^{23} \times 100 \times 1800\, N_A$

2. (a) Number of nucleons = mass number.
Molar mass of element
$= 3.32 \times 10^{-23} \times 6.02 \times 10^{23}$
$= 20$ = mass number.
So, number of nucleons in 2 atoms = 40

3. (b) Moles of $PO_4^{3-} = \frac{9.5}{95} = 0.1$ mol
Number of electrons in 0.1 mol PO_4^{3-}
$= 50 \times 0.1 \times N_A = 5N_A$

4. (d) 126 amu HNO_3 = 2 molecules of HNO_3 = 6 atoms of O
Moles of O-atom $= \frac{6}{N_A}$

5. (c) 96 amu of S^{2-} = $3S^{2-}$ ions
Charge on $3S^{2-}$ ions = 3 × charge of 2 electrons $= 3 \times 2 \times 1.6 \times 10^{-19} = 9.6 \times 10^{-19}$ C

6. (d) Number of sodium atoms $= \frac{46}{23} \times N_A = 2N_A$
Number of calcium atoms = Number of sodium atoms $= 2N_A$
Mass of Ca atoms $= \frac{2N_A}{N_A} \times 40 = 80$ g

7. (c) 54 mL H_2O = 54 g H_2O
Moles of $H_2O = \frac{54}{18} = 3$ mole
Number of neutrons $= 8 \times 3 \times N_A = 24\, N_A$

8. (c) Moles of $Mg^{2+} = \frac{48}{24} = 2$ mole
Number of electrons $= 2 \times 10\, N_A = 20\, N_A$

9. (b) Moles of $D_2O = \frac{5}{20} = \frac{1}{4}$
Number of neutrons $= 10 \times \frac{1}{4} \times N_A = 2.5\, N_A$

10. (b) Mass of one molecule $= 300 \times 1.66 \times 10^{-24}$ g
$= 4.98 \times 10^{-22}$ g

11. (d) Moles of $C_9H_8O_4 = \frac{360 \times 10^{-3}}{180} = 2 \times 10^{-3}$
Number of O-atoms $= 2 \times 10^{-3} \times N_A \times 4$
$= 8 \times 6.02 \times 10^{20} = 4.81 \times 10^{21}$

12. (b) $M = \frac{wRT}{PV} = \frac{20 \times 0.0821 \times 273}{5.6 \times 1} = 80$

13. (b) 1 mole $(NH_4)_3PO_4$
= 12 mole H-atoms = 4 mole O-atoms
= 6 mole H-atoms = 2 mole O-atoms

14. (a) Moles of $O_3 = \frac{3 \times 8.21}{0.0821 \times 300} = 1$
Moles of O-atoms = 3

15. (d) Moles of atoms $= \frac{3.011 \times 10^{22}}{6.02 \times 10^{23}} = \frac{1}{20}$
Atomic mass $= \frac{\text{mass}}{\text{moles}} = \frac{1.15}{1/20} = 23$

16. (d) Atomic mass of an element
$x = 6.643 \times 10^{-23} \times N_A = 40$
number of moles of $x = \frac{20 \times 1000}{40} = 500$

17. (d) 3.9854×10^{-23} g of element = 1 atom
1 g of element $= \frac{1}{3.9854 \times 10^{-23}}$
$= 2.509 \times 10^{22}$

18. (d) 4 mole $C_2H_4O_2$ = 16 mole H-atoms
= 16 g hydrogen

19. (d) 12.044×10^{22} molecules of CO_2 = 0.2 mole CO_2 = 0.4 mole O-atoms = $0.4\, N_A$ O-atoms

20. (a) (I) 0.5 mole O_3 = 24 g O_3 ;
(II) 0.5 g atom of oxygen = 8 g
(III) $\frac{3.011 \times 10^{23}}{6.022 \times 10^{23}} \times 32 = 16$ g O_2
(IV) $\frac{5.6}{22.4} \times 44$ g $CO_2 = 11$ g CO_2

For More Material Join: @JEEAdvanced_2024

21. (a) Mass of a drop of water 0.0018 g

Moles of water $\frac{0.0018}{18}$ 10^{-4} mole

Number of water molecules in one drop

$10^{-4} N_A$

Number of water molecules in two drops

$2 \quad 10^{-4} \quad 6.023 \quad 10^{23} \quad 12.046 \quad 10^{19}$

22. (c) Number of protons in ${}_6C^{14} = 6$;
Number of neutrons in ${}_6C^{14} = 8$;

As per given new atomic mass of

${}_6C^{14} = 12 + 4 = 16$

(As the mass of e is negligible as compared to neutron and proton)

% increase in mass $\frac{16 \quad 14}{14}$ $100 \quad 14.28$

23. (b) Mass of NaCl $25 \quad \frac{8.775}{100} \quad 2.193$ g

Number of NaCl units $\frac{2.193}{58.5}$ N_A

$2.25 \quad 10^{22}$

24. (b) Moles of $C_{12}H_{22}O_{11}$ $\frac{25.6}{342.3} \quad 0.0747$

Number of H-atoms

$0.0747 \quad 22 \quad 6.023 \quad 10^{23} \quad 9.91 \quad 10^{23}$

25. (a) Let there are x-atoms of N

% of N $28.9 \quad \frac{x \quad 14}{194} \quad 100 \qquad x \quad 4$

26. (d) Mass of 1 molecule of water 18 amu

$18 \quad 1.66 \quad 10^{-24}$ g

Volume of 1 molecule $\frac{\text{mass}}{\text{density}}$

$29.88 \quad 10^{-24} \quad 2.988 \quad 10^{-23}$ mL

27. (d) Volume of gold foil $25 \quad 40 \quad 0.25$ mm^3

$250 \quad 10^{-3}$ cm^3

Mass of gold foil $19.32 \quad 250 \quad 10^{-3}$ g

4.83 g

No. of gold atoms $\frac{4.83}{197}$ N_A

$1.47 \quad 10^{22}$

28. (c) Let total mass be

100 g

N_2 $\quad$ O_2

x $\quad$ $(100 - x)$

$M_{\text{Avg.}} \quad \frac{\text{Total mass}}{\text{Total moles}}$

$29 \quad \frac{100}{\frac{x}{28} \quad \frac{(100 \quad x)}{32}} \qquad x \quad 72.41\ \%$

29. (c) $d \quad \frac{PM}{RT}$

$1.15 \quad \frac{\frac{740}{760} \quad M}{0.0821 \quad 300}$

$M \quad 29.09$. Let mole fraction of N_2 is x.

$29.09 \quad 28 \quad x \quad 32 \ (1 \quad x)$;

$x \quad 0.7275$

mass % $\quad$ mole % $\quad \frac{\text{mol. mass of } N_2}{\text{average mol. mass of air}}$

$= 72.75 \quad \frac{28}{29.09} \quad 70.02$

30. (c) No. of moles of CO_2 in 100 g mixture

$\frac{66}{44} \quad 1.5$

No. of moles of H_2 in 100 g mixture

$\frac{34}{2} \quad 17$

$M_{\text{average}} \quad \frac{100}{18.5} \quad 5.40$

V. D. $\quad \frac{5.4}{2} \quad 2.7$

31. (b) Let 1 mole of mixture has x mole N_2O_4

$2 \quad 27.6 \quad x \ (92) \quad (1 \quad x) \ 46; \ x \quad 0.2$

32. (a) $PM \quad dRT$

$M_{\text{Gas}} \quad \frac{2 \quad 0.0821 \quad 600}{2} \quad 49.26$

At const. P and T $\quad d \quad M$

$\frac{d_{\text{Gas}}}{d_{\text{Ne}}} \quad \frac{M_{\text{Gas}}}{M_{\text{Ne}}} \quad \frac{49.26}{20} \quad 2.463 \quad 2.5$

For More Material Join: @JEEAdvanced_2024

33. (a) Let mole % of ^{26}Mg be x

$$\frac{(21 \quad x)\,25 \quad x\,(26) \quad 79\,(24)}{100} \quad 24.31$$

$$x \quad 10\%$$

34. (a) Let atomic weight of other isotope is M

$$114.82 \quad \frac{114.9041 \quad 95.72 \quad M \quad 4.28}{100}$$

$$M \quad 112.94$$

35. (b) Moles of N_2 $\quad \frac{3.01 \quad 10^{24}}{6.02 \quad 10^{23}} \quad 5$

Moles of O_2 $\quad \frac{32}{32} \quad 1$ mol

Average molar mass $\quad \frac{5 \quad 28 \quad 32 \quad 1}{5 \quad 1}$

$$\frac{172}{6} \quad 28.67$$

$$d_{mix} \quad \frac{PM_{avg.}}{RT} \quad \frac{3 \quad 28.67}{\frac{1}{12} \quad 860} \quad 1.2\ g/L$$

36. (d) Let mole fraction of O_2 is x

$$40 \quad 32 \quad x \quad 80\,(1 \quad x)$$

or $x \quad 5/6$

$$a : b \quad x : (1 \quad x) \quad \frac{5}{6} : \frac{1}{6}$$

When ratio is changed

$$M_{mixture} \quad 32 \quad \frac{1}{6} \quad 80 \quad \frac{5}{6} \quad 72$$

41. (c) Weight of calcium $\quad 4 \quad \frac{40}{100} \quad 1.6\ g$

43. (d) Weight % of $N \quad \frac{14}{17} \quad 100 \quad 82.35$

44. (a) Molar mass of $x \quad 0.444$ molar mass of y

Molar mass of $x \quad 2.96 \quad 12$

Molar mass of $y \quad \frac{2.96 \quad 12}{0.444} \quad 80$

45. (b) Moles of Zn $\quad \frac{9.81}{65} \quad 0.15$

Moles of Cr $\quad \frac{1.8 \quad 10^{23}}{6.02 \quad 10^{23}} \quad 0.3$

Moles of O $\quad 0.6$

$Zn_{0.15}\,Cr_{0.3}\,O_{0.6}$

$ZnCr_2O_4$

46. (a) $n \quad \frac{\text{Mass}}{\text{Molar mass}}$

$$0.0242 \quad \frac{1.657}{M} \qquad M \quad 68.47$$

$$1 \quad X \quad 32 \quad 68.47$$

$$X \quad 35.5$$

47. (c) Mass of oxide $\quad$ Mass of metal $\quad$ Mass of oxygen

$$2.74 \quad 1.53 \quad W_{Oxygen} \qquad W_{Oxygen} \quad 1.21\ g$$

Moles of $V \quad \frac{1.53}{51} \quad 0.03$

Moles of $O \quad \frac{1.21}{16} \quad 0.075$

$V_{0.03}\,O_{0.075}$

$V\,O_{2.5} \qquad V_2O_5$

48. (d)

	C	H	N	O
Moles	$\frac{70.6}{12}$	$\frac{4.2}{1}$	$\frac{11.8}{14}$	$\frac{13.4}{16}$
Moles	$\frac{5.88}{0.84}$	$\frac{4.2}{0.84}$	$\frac{0.842}{0.84}$	$\frac{0.84}{0.84}$
Simple ratio	7	5	1	1

C_7H_5NO

49. (a) Loss in mass is due to loss of water $\quad 63.1$

% of $H_2O \quad 63 \quad \frac{18x}{106 \quad 18x} \quad 100 \quad x \quad 10$

50. (a) $Sr(OH)_2 \ xH_2O \qquad Sr(OH)_2 \qquad xH_2O$

$$6.85\ g \qquad 3.13\ g \qquad 3.72\ g$$

$$\% \text{ of water} \quad \frac{3.72}{6.85} \quad 100 \quad \frac{18x}{121 \quad 18x} \quad 100$$

$$x \quad 8$$

51. (c) $\%$ of $O = \frac{16 \quad 27}{(100 + 3 \quad 310)} \quad 100$

$$= 41.94\%$$

52. (c) $C_xH_yCl_zO_w \quad O_2 \qquad CO_2 \quad H_2O$

Millimoles of C $\quad \frac{41.21}{44}$

Millimoles of H $\quad \frac{5.63}{18} \quad 2$

For More Material Join: @JEEAdvanced_2024

29.72 mg of Dieldrin contains $\frac{57.13}{25.31}$ 29.72 mg of AgCl

Millimoles of Cl $\frac{57.13}{25.31}\ \frac{29.72}{143.5}$

Weight of O $29.72\ \frac{41.21}{44}\ 12\ \frac{5.63}{18}\ 2\ 1\ \frac{57.13}{25.31}\ \frac{29.72}{143.5}\ \frac{35.5}{}$

Weight of O 29.72 [11.239 0.6255 16.59]

Weight of oxygen 1.265 mg

n_C $\frac{41.21}{44}$ $\frac{41.21}{44}\ \frac{16}{1.265}$ 11.84 12

n_H $\frac{5.63}{9}$ $\frac{5.63}{9}\ \frac{16}{1.265}$ 7.9 8

n_O $\frac{1.265}{16}$ $\frac{1.265}{16}\ \frac{16}{1.265}$ 1

n_{Cl} $\frac{57.13}{25.31}\ \frac{29.72}{143.5}$ $\frac{57.13}{25.31}\ \frac{29.72}{143.5}\ \frac{16}{1.265}$ 5.95 6

53. (c) $d\ \frac{PM}{RT}\ M\ \frac{dRT}{P} = \frac{2.28\ \ 0.0821\ \ 300}{1} = 56.15$ g/mol

E.F. $\frac{85.7}{12} : \frac{14.3}{1}$ 7.14 : 14.3 = 1 : 2; E.F. is CH_2; M.F. $(CH_2)_n$

where n $\frac{56.15}{12+2} \sim 4$;

M.F. is C_4H_8

54. (a) Moles of C n_{CO_2} $\frac{2.64}{44}$ 0.06

mass of C 0.72

Mole of H 2 Moles of H_2O 2 $\frac{1.26}{18}$ 0.14 mass of H 0.14

Compound does not contains oxygen.

So EF $C_{0.06}H_{0.14}$ C_3H_7

Lowest M.M. 43

55. (c) Formula of sulphate will be $—MSO_4\ 7H_2O$

% of metal $\frac{M}{M\ \ 96\ \ 7\ \ 18}\ 100\ \ 9.87$

M 24.31

56. (a)

	C	H	N		
Moles	$\frac{9}{12}$	$\frac{1}{1}$	$\frac{3.5}{14}$	n	$\frac{108}{54}$ 2
	$\frac{3}{4}$	: 1 :	$\frac{1}{4}$		
	3	: 4 :	1		

For More Material Join: @JEEAdvanced_2024

C_3H_4N M.F. $(C_3H_4N)_n$

MF $(C_3H_4N)_2$ $C_6H_8N_2$

57. (d)

	I	O
Moles	$\frac{254}{127}$	$\frac{80}{16}$
	2	5

I_2O_5

58. (c) A^4 B^2 A_2B_4

Empirical formula AB_2

59. (d) 100 g compound contain 43.8 g oxygen so 1 g oxygen present with $\frac{100}{43.8}$ g compound

16 4 g or 4 mole oxygen present with $\frac{100}{43.8}$ 16 4 ~146 g compound

mol. mass of compound acid is 146.

60. (c) Mol. mass of XY_2 $\frac{5}{0.05}$ 100

Mol. mass of X_2Y_3

$\frac{85}{3.011 \quad 10^{23}}$ N_A 170

Let molar mass of X and Y are a and b respectively

$a \quad 2b \quad 100$

$2a \quad 3b \quad 170;$

$a \quad 40$;

$b \quad 30$

61. (c) $CO_2 = \frac{88}{44} = 2$ mole of CO_2 2 mole of C

$H_2O = \frac{36}{18} = 2$ mole of H_2O 4 mole of H

Mass of C + Mass of H + Mass of O = 44

24 4 x 44; x 16

mole of O = 1 and molecular formula is C_2H_4O.

62. (d) M.M. of X_2

$\frac{WRT}{PV}$ $\frac{40 \quad 10^{3} \quad 0.0821 \quad 300}{1 \quad 4.92 \quad 10^{3}}$ 200

Atomic mass of X 100.

63. (d)

	X	Y
	$\frac{75.8}{75}$	$\frac{24.2}{16}$
	1	3/2
	2	3

X_2Y_3

64. (b) M $\frac{WRT}{PV}$ $\frac{12.4 \quad 0.0821 \quad 100}{8 \quad 0.821}$ 124 P_4

65. (d) % Mn $\frac{55}{55 \quad 16x}$ 100 64

x 1.93

66. (b) Ti + O_2 $Ti_{1.44}O$

$\frac{1.44}{48}$ mole $\frac{x}{48\,(1.44) \quad 16}$ mole

$\frac{1.44}{48}$ $\frac{1.44x}{48\,(1.44) \quad 16}$

x 1.77 g

67. (d) 3 molecules of O_2 1 molecules of CS_2

6 molecules of O_2 2 molecules of CS_2

68. (a) Factor label method.

69. (c) $3Fe_2O_3(s)$ $2Fe_3O_4 + \frac{1}{2}O_2$

480 g Fe_2O_3 provide 16 g O_2. For loss of 0.04 g O_2 0.04 $\frac{480}{16} = 1.2$ g Fe_2O_3

% by mass of $SiO_2 = \frac{0.8}{2.0}$ 100 40%

70. (d) Mass of S_8 in sample 160 g;

Moles of S_8 $\frac{160}{32 \quad 8}$ 0.625

Number of moles of O_2 required

= 0.625 8

Volume of O_2 required = 22.4 5

Vol. of air required 22.4 5 $\frac{100}{21}$

= 533.33 L

For More Material Join: @JEEAdvanced_2024

71. (c) $2Fe(NO_3)_3 + 3Na_2CO_3 \quad Fe_2(CO_3)_3 + 6NaNO_3$

mole 2.5 3.6

mole/stoichiometric coefficient 1.25 1.2

Limiting reagent is Na_2CO_3 so moles of $NaNO_3$ formed 3.6 2 7.2

% yield $\frac{6.3}{7.2}$ 100 87.5

72. (a) $\underset{0.1}{4Ca_5(PO_4)_3F} \quad \overset{\text{L.R.}}{\underset{0.36}{18SiO_2}} \quad \underset{0.9}{30C} \quad 3P_4 \quad 2CaF_2 \quad 18CaSiO_3 \quad 30CO$

18 mole SiO_2 gives 3 mole P_4

0.36 mole SiO_2 will give $\frac{3}{18}$ 0.36 0.06 mole

73. (d) CO_2 exhaled in 5 minutes 0.667 5 3.33 g

moles of CO_2 0.0756 mole

moles of KO_2 consumed 0.0756 mole

mass of KO_2 consumed 5.38 g

74. (d) L.R.

$2NH_3 \quad 5F_2 \quad N_2F_4 \quad 6HF$

2 17 g NH_3 gives 66 g N_2F_4

2*g* will give $\frac{66}{34}$ 2 3.88*g* N_2F_4

% yield $\frac{3.56}{3.88}$ 100 91.75%

75. (d) Mass of pure $CaCO_3$ 200 $\frac{95}{100}$ 190 kg

$CaCO_3 \quad CaO \quad CO_2$

100 g 56

1 g $\frac{56}{100}$

190 kg $\frac{56}{100}$ 190 kg 106.4 kg

76. (b) Mass of H_3PO_4 produced = $\frac{62}{4 \quad 31}$ 0.85 0.9 4 98 149.94 g

77. (a) L.R.

$3D \quad 4E \quad \overset{80\%}{\quad} \quad 5C \quad A$

9 mole 14 mole $\frac{5}{3}$ 9 0.8 12 mole

$3C \quad 5G \quad \overset{50\%}{\quad} \quad 6B \quad F$

12 mole 4 mole

L.R. is G

Moles of *B* formed $\frac{6}{5}$ 4 0.5 2.4

78. (a) Apply PO AC for Zn

1 moles of ZnS 1 mol Zn

Moles of ZnS $\frac{2}{0.8 \quad 0.8 \quad 1}$ 3.125 mole

For More Material Join: @JEEAdvanced_2024

79. (b) $CO_2 + 2NaOH \quad Na_2CO_3 + H_2O$

n_{NaOH} 1; CO_2 present in mixture 0.5 and CO present 0.3 mole

Moles of CO_2 obtained from CO 0.3, extra moles of NaOH required 0.3 2 0.6 mole

80. (c) Let mass of Ag_2O x g

$Ag_2O \quad 2Ag \quad \frac{1}{2}O_2$

232 g $\frac{1}{2}$ 32 16 g O_2

x g $\frac{16}{232}$ x 0.104 x 1.508 g

% of Ag_2O $\frac{1.508}{1.6}$ 100 94.25%

81. (c) Number of moles of $Ba(OH)_2 = \frac{20 \quad 342}{100 \quad 171} = 0.4$ mole

or moles of OH 0.4 2

Number of moles of HNO_3 1.2 2 = 2.4 mole

Hence, the final solution is acidic due to presence of excess H

$[H^+] \quad \frac{(2.4 \quad 0.8) \quad 1000}{1200 + \frac{342}{0.57}} \quad 0.888\ M$

82. (d) Total moles of $H_2SO_4 = 0.1$ mole

Total volume $\frac{150 \quad 400}{1.25} \quad \frac{550}{1.25} \quad 440$

$M \quad \frac{0.1}{440} \quad 1000 \quad \frac{1}{4.4} \quad 0.227\ M$

83. (a) $Zn + 2HCl \quad ZnCl_2 + H_2$;

moles of H_2 evolved 2

moles of HCl required 4

$\frac{V \quad 1.2 \quad 0.365}{36.5} \quad 4$; V 333.33 mL

84. (b) $C_2H_6 + 3.5O_2 \quad 2CO_2 + 3H_2O$;

$C_2H_4 + 3O_2 \quad 2CO_2 + 2H_2O$

Let volume of ethane is x litre, $22.4 \quad 4 = 3.5x \quad 3\ (28 \quad x)$

x 11.2 litre

at constant T and P, $V \quad n$;

Mole fraction of C_2H_6 in mixture $\frac{11.2}{28}$ 0.4

85. (c) $X_{Sn} \quad \frac{12.5 / 119}{\frac{50}{209} \quad \frac{25}{207} \quad \frac{12.5}{119} \quad \frac{125}{112}} \quad 0.176$

86. (d) Mass of solution 100 g

Mass of solute 56 g

moles of solute $\frac{56}{60}$ 0.93

Mass of solvent 44 g

For More Material Join: @JEEAdvanced_2024

moles of solvent $\frac{44}{18}$ 2.44

X_{Solute} $\frac{0.93}{0.93 \quad 2.44}$ $\frac{0.93}{3.37}$ 0.276

87. (a) $\frac{\text{mass of } Al_2(SO_4)_3}{\text{mass of water}}$ 10^6 34.2

1 litre solution contains 1000 g of water

In 1 litre solution, mass of $Al_2(SO_4)_3$ $\frac{34.2 \quad 1000}{10^6}$ 34.2 mg

molarity of $Al_2(SO_4)_3 = \frac{34.2 \quad 10^{\ 3}}{342} M = 10^{\ 4} M$

$Al_2(SO_4)_3(aq)$ $2Al^{3+}(aq) + 3SO_4^{2}(aq)$

$10^{\ 4} M$ 2 $10^{\ 4} M$ 3 $10^{\ 4} M$

$[SO_4^{2}] = 3 \quad 10^{\ 4} M$

88. (d) M $\frac{n_{\text{Solute}}}{\text{Vol of solution (in mL)}}$ 1000 ...(1)

m $\frac{n_{\text{Solute}}}{\text{Weight of solvent (in g)}}$ 1000 ...(2)

m $\frac{n_{\text{Solute}}}{\text{Weight of soution (in g)} - \text{wt of solute (in g)}}$ 1000

m $\frac{M \quad \text{vol of solution (in mL)} \quad 1000}{1000 \quad (\text{wt of solution (in g)} - \text{wt of solute (in g)})}$

m $\frac{M \quad 1000}{\text{Density of solution (g / mL)} \quad 1000 \quad \frac{\text{wt of solute} \quad M_{\text{Solute}} \quad 1000}{M_{\text{Solute}} \quad \text{volume of solution (in mL)}}}$

m $\frac{M \quad 1000}{1000 \quad M.M_{\text{Solute}}}$

89. (d) m $\frac{1000\,M}{1000\,d \quad M \quad M_{\text{Solute}}}$

d $\frac{1000 \quad 9}{10}$ 9 50 $\frac{1}{1000}$ 1.35 g/mL

90. (b) 1000 mL solution contain 2 mole of ethanol or 1000 1.025 g solution contain 2 mole of ethanol

Mass of solvent 1000 1.025 2 46

m $\frac{2}{1000 \quad 1.025 \quad 2 \quad 46}$ 1000

m $\frac{2}{933}$ 1000 2.143

91. (c) Total Cl 0.2 0.4 mole (from $BaCl_2$) 0.6 mole

M_{Cl} $\frac{0.6}{500}$ 1000 1.2

92. (a) Mass of solution 10 1.2 12 g

Mass of solute 12 $\frac{37}{100}$ 4.44 g

93. (a) The sample of H_2SO_4 is 80% by volume , $W_{H_2SO_4}$ 800 g in 1000 mL

Volume of solution 1000 mL

For More Material Join: @JEEAdvanced_2024

Mass of solution = 1000.0 1.80 = 1800 g
Mass of water 1800 800 1000 g
Molality $\frac{800}{98}\frac{1000}{1000}$ 8.16 mol kg ^{1}

94. (c) Initial molarity (M_1) $\frac{10}{336}\frac{10^{3}}{500}$ 1000
0.0595 10 ^{3}

M_1V_1 M_2V_2

M_2 $\frac{M_1V_1}{V_2}$ $\frac{0.0595\ 10^{3}\ 1}{1000}$

0.0595 10 ^{6} 5.95 10 ^{8}

95. (a) Molar mass of $Pb(NO_3)_2$ 25 0.15
3.75 m. moles
Molar mass of $Al_2(SO_4)_3$ $\frac{1}{3}$ 3.75 M 20

M 0.0625 6.25 10 ^{2}M

96. (b) m-moles of HNO_3 required
250 1.2 300
100 g solution contains 63 g HNO_3 or
$\frac{100}{1.4}$ mL solution contain 1 mole HNO_3
molarity of HNO_3 solution
$\frac{1000}{100}$ 1.40 14

14 V 300 or V 21.42 mL

97. (b) 50 mL $BaCl_2$ (aq) solution contain 10.4 g $BaCl_2$

n_{BaCl_2} $\frac{10.4}{137+71}$ 0.05

moles of H_2SO_4 in 100 mL H_2SO_4 (aq) solution $\frac{9.8}{98}$ 0.1

$BaCl_2 + H_2SO_4$ $BaSO_4 + 2HCl$;
moles of HCl formed 0.1
[Cl] $\frac{0.1\ \ 1000}{50\ \ 100}$ 0.666 M

98. (c) No. of moles of HCl are more than NaOH, so final solution becomes acidic.

99. (c) Let volume is x mL, x 0.1 2 50 0.1
(x 50) 0.05 2
x 100 mL

100. (c) [HCl] $\frac{1\ \ 4V\ \ 2\ \ V}{5V}$ 1.2 M

101. (d) M $\frac{55\ \ 0.07\ \ 20\ \ 0.12\ \ 25\ \ 0.15}{55\ \ 20\ \ 25}$ 0.1

102. (c) Molarity $\frac{28}{11.2}$ 2.5 M
1000 mL of solution contains 2.5 mole H_2O_2
molality, m $\frac{2.5}{1250\ \ 85}$ 1000 2.15 m

103. (a) The reaction that takes place is
$NaCl + AgNO_3$ $AgCl + NaNO_3$
$\because$ 143.5 g of AgCl is produced from 58.5 g NaCl
14 g of AgCl will be produced from
$\frac{58.5\ \ 14}{143.5}$ 5.70 g NaCl
This is the amount of NaCl in common salt;
% purity $\frac{5.70}{6}$ 100 95%

104. (c) Let no. of moles of $Al_2(SO_4)_3$ in solution is x

$\frac{x}{2684\ \ x\ \ 342}$ 1000 1

x 2
$2Al_2(SO_4)_3$ 6 moles SO_4^{2}
6 moles of $BaSO_4$

105. (a) 0.478 mL 0.478 g of water; 10^{9} g water contain 0.10 g $CHCl_3$
0.478 g water contain
$\frac{0.1}{10^{9}}$ 0.478 g $CHCl_3$

n_{CHCl_3} $\frac{0.1}{10^{9}}$ $\frac{0.478}{119.5}$

No. of molecules $\frac{0.1}{10^{9}}$ $\frac{0.478}{119.5}$ N_A

4 10 ^{13} N_A

106. (b) (i) Mass of NaOH 50 $\frac{40}{100}$ 20 g

(ii) $\frac{w}{v}$ NaOH 50 g NaOH in 100 mL
So, in 50 mL, mass of NaOH will be 25 g
(iii) 50 g of 15 M NaOH (d 1 g/mL)
15 M NaOH 15 moles NaOH in 1000 mL solution
Mass of solution 1000 1 1000 g
Mass of NaOH 15 40 600 g
In 1000 g solution mass of NaOH 600 g
So, in 50 g solution, mass of NaOH
$\frac{600}{1000}$ 50 30 g

Order (iii) > (ii) > (i)

For More Material Join: @JEEAdvanced_2024

107. (b) $B\ H_2PtCl_6$ Pt; $\frac{12}{M_B} \quad \frac{410}{195}$

moles of Pt; Molecular mass of base 58

108. (b) $\underset{\text{Dibasic acid}}{H_2A} \quad \underset{\text{Silver salt}}{Ag_2A} \quad Ag$

Apply PoAC for Ag

$2\ n_{Salt} \quad 1\ n_{Ag}$

$2\ \frac{1}{216\ A} \quad \frac{0.5934}{108} \quad A \quad 148$

i.e., Molar mass of acid (H_2A) 150,

Let % of C is x

C H O wt. % of H $\frac{x}{8}$

Mass $x \quad \frac{x}{8} \quad 2x$ wt % of O 2 % of C $2x$

Moles $\frac{x}{12} \quad \frac{x}{8} \quad \frac{2x}{16}$

$\frac{x}{3} \quad \frac{x}{2} \quad \frac{x}{2}$ $C_2H_3O_3$ E.F.

M.F. $(C_2H_3O_3)_n \quad n \quad \frac{\text{M.M.}}{\text{EFM}} \quad \frac{150}{75} \quad 2$

M.F. $C_4H_6O_6$

109. (b) Moles of $Ag_3A = \frac{\text{moles of Ag}}{3} \quad \frac{0.607}{M}$

$\frac{0.37}{108} \quad \frac{1}{3} \quad M \quad 531$

mol. mass of H_3A= mol. mass of Ag_3A 3 At. mass of Ag + 3 At. mass of H 210

110. (a) Equivalents of KOH used by oil

[25 0.40 8.5 0.28 2] 10^{-3}

Moles of KOH used 5.24 10^{-3}

Mass of KOH used in milligrams

5.24 10^{-3} 56 1000 223.44

Saponification number $\frac{223.44}{2}$

146.72

111. (b) $\underset{x}{CO} \quad \underset{x/2}{\frac{1}{2}O_2} \quad \underset{x}{CO_2}$ V.C. *VR VP*

$\underset{(20\ x)}{H_2} \quad \underset{\frac{20\ x}{2}}{\frac{1}{2}O_2} \quad H_2O(l)$

18 (20 10) x

x 12 mL CO

H_2 8 mL

$V_{CO} : V_{H_2}$

12 : 8

3 : 2

112. (d) For 6 mole HCl, 3 mole $H_2(g)$ is produced

So, For 1 mole HCl, $\frac{1}{2}$ mole H_2 will be produced

$\frac{1}{2}$ 22.4 L at STP 11.2 L H_2 gas

113. (c) 100 mL gaseous mixture contain 20 mL C_3H_8

So, volume of CH_4 and CO

(100 20) 80 mL

$C_3H_8 + 5O_2 \quad 3CO_2 + 4H_2O$

$CH_4 + 2O_2 \quad CO_2 + 2H_2O;$

$CO + \frac{1}{2}O_2 \quad CO_2$

80 mL (CH_4 and CO) will produce 80 mL CO_2; C_3H_8 will produce 3 20

60 mL CO_2

Total CO_2 produced 80 60 140 mL

114. (d) Let the volume of CO, CH_4 and Ne be x, y & z respectively

$CO + \frac{1}{2}O_2 \quad CO_2;$

$x \quad x/2 \quad x$

$\underset{y}{CH_4} + \underset{2y}{2O_2} \quad \underset{y}{CO_2} + 2H_2O\ (l)$

Remaining volume of O_2 15 $\frac{x}{2}$ 2y 1.5

Volume after reaction :

$x \quad y \quad 9$ ...(i)

$x \quad y \quad z \quad 40$ ...(ii)

by Eq. (i) and (ii)

Volume of CH_4 = 6 mL;

% of $CH_4 = \frac{6}{40} \quad 100 \quad 15$

115. (b) Suppose, the volume of propane V L

$\underset{V}{C_3H_8(g)} + \underset{5V}{5O_2(g)} \quad \underset{3V}{3CO_2(g)} + 4H_2O(l)$

$C_4H_{10}(g) + \frac{13}{2}O_2(g) \quad 4CO_2(g) + 5H_2O(l)$

$(3\ V) \quad \frac{13}{2}(3\ V) \quad 4(3\ V)$

$\because$ Total volume of CO_2 produced = 10 L;

$\because$ $3V \quad 4(3\ V) \quad 11; \quad V \quad 1$

Volume of butane (3 1) 2 L

For More Material Join: @JEEAdvanced_2024

Thus, the ratio of volume of butane to propane 2 : 1

116. (c) Hypophosphorous Acid —H_3PO_2

117. (c) $Ca^{2}\begin{cases}{}^{2(1)}OCl\\ Cl^{(1)}\end{cases}$

118. (a) $\overset{0}{S_8}$, $\overset{1}{S_2}F_2$, $H_2\overset{2}{S}$, $H_2\overset{6}{S}O_4$

119. (b) Let oxidation state is x;

$(x \quad 5 \quad 0 \quad 1) \quad 2 \quad 0 \quad x \quad 1$

120. (b) The redox reaction is :

$Cr_2O_7^{2} \quad 3SO_2 + 2H^{+} \qquad 2Cr^{3+} + 3SO_4^{2} + H_2O$

121. (c) $\overset{3}{N}H_4\overset{5}{N}O_3$

122. (a) Caro's acid H_2SO_5

= Peroxomono-sulphuric acid

Marshal's acid $H_2S_2O_8$

= Peroxodi-sulphuric acid

The oxidation state in both the acids is +6 each because oxidation state cannot be greater than the number of valence electrons.

123. (b) $\underline{\overset{1}{Ba}O_2}$ $\underline{\overset{1/2}{K}O_2}$ $\underline{\overset{0}{O}_3}$ $\underline{\overset{2}{O}F_2}$

124. (d) $\underline{\overset{1/3}{K}O_3}$, $Na_2\underline{\overset{1}{O}_2}$

125. (b) $Ba(H_2PO_2)_2 \qquad Ba^{2} \quad \overset{1}{H_2}PO_2$

126. (d) $Fe_{0.96}O \rightarrow \begin{cases}Fe^{2+} & Fe^{3+}\\ x & (0.96-x)\end{cases}$

Charge balance $2x \quad 3(0.96 \quad x) \quad 2 \quad 0$,

$x \quad \frac{11}{12}$

127. (b) $\overset{5}{H}N\overset{2}{O}_3$ $\overset{0}{N}O$ $\overset{3}{N}_2$ $\overset{3}{N}H_4Cl$

128. (c) 1 mole N_2H_4 loses 8 mole e ; 1 mole N loses 4 mole of e

New oxidation state of N is $2 \quad 4 \quad 2$

129. (a) $\overset{6}{Cr}_2O_7^{2}$ $\overset{6}{Cr}O_4^{2}$

130. (c) $Mn^{2} \xrightarrow{(KNO_3 \quad NaOH)} \overset{6}{Mn}O_4^{2}$

131. (c) $Fe^{2} \quad MnO_4 \quad H \qquad Fe^{3} \quad Mn^{2} \quad H_2O$

132. (a) Both Fe and C are oxidised and MnO_4 is reduced.

133. (d) H_2O_2 it self be oxidised to O_2.

134. (b) Metallic iron is oxidised to Fe^{3} .

137. (a) $2Na[\underline{\overset{1}{Ag}}(CN)_2] \quad \underline{\overset{0}{Zn}}$

$Na_2[\underline{\overset{2}{Zn}}(CN)_4] \quad 2\underline{\overset{0}{Ag}}$

138. (a) The balanced redox reaction is

$2MnO_4 \quad 5C_2O_4^{2} \quad 16H^{+} \qquad 2Mn^{2+} + 10CO_2 + 8H_2O$

139. (a) Balancing the equation, we have

$K_2Cr_2O_7 + H_2SO_4 + 3SO_2 \qquad K_2SO_4 + Cr_2(SO_4)_3 + H_2O$

140. (d) $CS_2 \quad 3Cl_2 \qquad CCl_4 \quad S_2Cl_2$

141. (a) Balanced reaction is

$PtCl_4 + 5XeF_2 \qquad PtF_6 + 4ClF + 5Xe$

Sum of stoichiometric coefficients

$1 \quad 5 \quad 1 \quad 4 \quad 5 \quad 16$

142. (c) If $x \quad 4$

$H_3PO_4 \qquad 3KOH \qquad Na_3PO_4 \quad 3H_2O$

0.1 mole 0.3 mole

If $x \quad 3$

$H_3PO_3 \qquad 2KOH \qquad Na_2PO_3 \quad 2H_2O$

0.1 mole 0.2 mole

If $x \quad 2$

$H_3PO_2 \qquad KOH \qquad NaPO_2 \quad H_2O$

0.1 mole 0.1 mole

So, if $x \quad 2{,}5.6$ g KOH will be required.

143. (b) In acidic medium $MnO_4 \qquad Mn^{2}$

$n_f \quad 5, \quad E \quad \frac{M}{5}$

144. (b) $n \quad |1 \quad 5 \quad 2|$

145. (d) 14 mole HCl loses 6 mole e ;

1 mole HCl loses $\frac{6}{14}$ mole e

eq. mass of HCl $\frac{M}{\frac{6}{14}}$

$\frac{36.5 \quad 14}{6} \quad 85.1$

146. (d) n-factor $\frac{n_1 \quad n_2}{n_1 \quad n_2} \quad \frac{4 \quad 2}{4 \quad 2} \quad \frac{4}{3}$

Eq. mass $\frac{\text{mol mass}}{n\text{-factor}} \quad \frac{M \quad 3}{4}$

147. (d) n-factor $|2 \quad 2 \quad 8 \quad 3| \quad 28$

148. (b) In SCl_2

71 g chlorine combines with 32 g S

For More Material Join: @JEEAdvanced_2024

So, 35.5 g chlorine will combine with 16 g

So, Eq. mass of S 16 g

As the mass which combines with 35.5 g chlorine is equivalent mass.

149. (b) E_{Element} 4, Formula of chloride $M\,Cl_x$

Eq. mass of chloride E_{Element} E_{Chlorine}

$\frac{M_{\text{Mass}}}{n_f}$ E_{Element} E_{Chlorine}

$\frac{59.25 \quad 2}{x}$ 4 35.5

x 3

150. (a) $K_2Cr_2O_7$ + X^{n+} $X^{5+}O_3$ Cr^{3+}

6 10^{-3} 6 $(5 \quad n)$ 9 10^{-3} n 1

151. (b) Normality $\frac{1}{100}$ (Centinormal)

Number of eq. Normality Volume (in L)

Number of eq. $\frac{1}{100}$ $\frac{250}{1000}$ $\frac{1}{400}$


$\frac{\text{Mass}}{\text{Molar mass}}$ n_f $\frac{1}{400}$

As R.A. $C_2O_4^{2-}$ CO_2

$\frac{w}{126}$ 2 $\frac{1}{400}$

w 0.1575 g

152. (d) As R.A.



n_f 1 4 4

E $\frac{M}{4}$

153. (c) E_{Chloride} E_{Element} E_{Chlorine}

E_{Chloride} w 35.5

Metal is divalent, n_f for chloride 2

E $\frac{M}{2}$ M $2(w \quad 35.5)$ $2w$ 71

154. (c) $2BrO_3^- + 12H^+ + 10Br^- \rightarrow 6Br_2 + 6H_2O$

10 mole e^- are required for formation of 6 moles of Br_2

n-factor of $Br_2 = \frac{10}{6} \quad \frac{5}{3}$

eq. wt. $\frac{\text{mol. wt.}}{n}$ $\frac{m}{5/3}$ $\frac{3M}{5}$

155. (a) Eq. of A Eq. of B

$\frac{m_A}{E_A}$ $\frac{m_B}{E_B}$

156. (a) O.N. of N in N_2H_4 is –2 which changes to 0 in N_2. Hence, equivalent mass of

$N_2H_4 = \frac{\text{molar mass}}{2 \quad 2}$

O.N. of iodine changes from +5 in IO_3^- to 1 in ICl. Hence, equivalent mass of

IO_3^- $\frac{\text{molar mass}}{4}$

157. (b) N_R $\frac{N_1V_1 \quad N_2V_2}{V_1 \quad V_2}$ $\frac{2 \quad 0.45 \quad 0.6 \quad 1}{3}$ 0.5

158. (b) Eq. of A^{2-} Eq. of MnO_4^-

2.7 10^{-3} 3 1.6 10^{-3} n_f

n_f 5 Acidic medium.

159. (c) Moles of O_2 gas obtained $\frac{2.24}{22.4}$ 0.1 mole

H_2O_2 H_2O $\frac{1}{2}O_2$ Moles of H_2O_2 0.2

1 mole $\frac{1}{2}$ mole M V(L) 0.2

2 mole 1 mole $\frac{5.6}{11.2}$ V(L) 0.2

2 mole 1 mole V 0.4 L 400 mL

0.2 mole 0.1 mole

160. (d) Eq. of acid Eq. of base

1000 $\frac{1.25}{M}$ 2 25 0.25 2

M 200

161. (a) $N_{(H^+)}$ $\frac{\left|10 \quad 1 \quad 20 \quad \frac{1}{2} \quad 30 \quad \frac{1}{3}\right|}{1000}$ $\frac{3N}{100}$

162. (b) Eq. of acid Eq. of base

$\frac{0.45}{90}$ n_f $\frac{20 \quad 0.54}{1000}$ n_f 2

163. (c) Eq. mass of $H_2O_2 = \frac{34}{2} = 17$

Eq. of $H_2O_2 = \frac{3.4 \quad x}{100 \quad 17}$

= Eq. of $KMnO_4$ x N 10^{-3} N 2

Molarity of $KMnO_4 = \frac{2}{5} = 0.4$

164. (c) 50 m-eq. $KMnO_4$ = 10 m-mole $KMnO_4$

or 25 m-mole of O_2 = 25 22.4 10^{-3}

0.56 L

For More Material Join: @JEEAdvanced_2024

165. (c) Zn will reduce Fe^3 to Fe^2

$$Fe_2O_3 + Zn \xrightarrow{+3} Fe^{2+} + Zn^{2+}$$

n_f 2 POAC for Fe

Fe^2 O. A. Fe^3

2 $\frac{0.8}{160}$ 1 n_{Fe^2} n_{Fe^2} 0.01

eq. of Fe^2 Eq. of O.A.

Moles of Fe^2 1 16.7 0.1 n_f

0.01 1 $\frac{16.7 \quad 0.1}{1000}$ n_f

n_f 6

166. (c) Eq. of $KMnO_4$ 20 0.1 5 10

Eq. of oxalic acid 25 0.2 2 10

167. (d) Fe^2 MnO_4 Fe^3 Mn^2

n_f 1 n_f 5

Moles of Fe^2 1 by $KMnO_4$

moles of MnO_4 5 ...(1)

Fe^2 $Cr_2O_7^2$ Fe^3 Cr^3

n_f 1 n_f 6

Moles of Fe^2 1 by $K_2Cr_2O_7$

moles of MnO_4 6 ...(2)

(1) (2)

$\frac{n_1}{n_2}$ $\frac{5}{6}$

168. (b) Moles of AgCl moles of $AgNO_3$

= moles of HCl $\frac{0.1435}{143.5}$ 10^{3}

mass of HCl 0.0365

mass of H_2SO_4 = 0.0635

mass % of H_2SO_4 $\frac{.0635 \quad 100}{0.1}$ 63.5

169. (b) Eq. mass of $KBrO_3 = \frac{1}{6}$ of its mol. mass

$\frac{1}{6}$ 167

$N_{Na_2S_2O_3}$ $\frac{0.167}{167}$ 6 $\frac{1}{0.05}$ 0.12 *N*

170. (a) Total m-eq. of NaOH taken 20

m-eq. of H_2SO_4 = m-eq. of NaOH reacted

$\frac{5 \quad 0.2}{25}$ 250 10

m-eq. of NaOH reacted 20 10 10

$(NH_4)_2SO_4 + 2NaOH$

$Na_2SO_4 + 2NH_3 + 2H_2O$

m-moles of $(NH_4)_2SO_4$ reacted = 5

mass of $(NH_4)_2SO_4$ 5 10^{3} 132 0.66

Percentage of $(NH_4)_2SO_4$ in sample

$= \frac{0.66}{0.80}$ 100 = 82.5

171. (c) milli-equivalent of NH_3 reacted with HNO_3

45 0.4 20 0.1 = 16

$\frac{W}{17}$ 1000 16; W_{NH_3} 0.272 g;

mass of N 0.272 $\frac{14}{17}$ 0.224

% N in the sample $\frac{0.224}{1.12}$ 100 20%

172. (a) 1000 $\frac{W}{24 \quad 64}$ 2 90 $\frac{1}{100}$ 5;

W 0.198 g

% of oxalate ion in a given sample

$\frac{0.198}{0.3}$ 100 66

173. (a) m-moles of HCl 20 0.1 = 2

m-moles of MgO reacted with HCl $\frac{2}{2}$ 1

mass of MgO present 1 40 mg

% of MgO $\frac{40}{320}$ 100 12.5%

174. (c) $Pb^{2+} + SO_4^2$ (from Na_2SO_4) $PbSO_4$

moles of Na_2SO_4 required = moles of Pb^{2+} ions

$\frac{0.355}{142}$ 0.0025

$[Pb^{2+}]$ $\frac{0.0025}{0.50}$ 5 10^{3} *M*

175. (b) Let volume of HNO_3 is V then

V 1.05 $\frac{12.6}{100}$ $\frac{3}{63} = \frac{1}{278}$ 1

V 0.57 mL

176. (c) m-eq. of $KMnO_4$ m-eq. of FeC_2O_4 + m-eq. of $FeSO_4$

(0.1 5) V $\frac{\frac{100}{144}}{3}$ $\frac{\frac{100}{152}}{1}$

V 5.48 mL

177. (a) meq. $FeSO_4$ $(NH_4)_2SO_4$ $6H_2O$

meq of $KMnO_4$

(n 1)

For More Material Join: @JEEAdvanced_2024

$\frac{W}{392}$ 1 1000 0.1 50; W 1.96 g

Hence, % purity of Mohr's salt

$\frac{1.96}{2.5}$ 100 78.4%

178. (d) $Cr_2O_7^{2-}$ $6Fe^{2+}$ (n 1) (Mohr's salt) $+ 14H^+$ $2Cr^{3+} + 6Fe^{3+} + 7H_2O$

Equivalents of Fe^{2+} = moles of Mohr's salt = equivalents of $K_2Cr_2O_7$

500 10^{-3} 6 1 3.0

Hence, mole per cent of Mohr's salt

$\frac{3}{4}$ 100 75

179. (a) Eq. mass of metal 2 8 16

Eq. mass of oxide E_{Element} E_{Oxygen}

16 8 24

$\frac{E_{\text{Oxide}}}{E_{\text{Metal}}}$ $\frac{24}{16}$ $\frac{3}{2}$

180. (d) X_2O_3 $2X^{3+} + 3O^{2-}$

i.e., the valency of metal in oxide is 3

$\because$ 6 mg 6 10^{-3} g of H_2 reacts with X_2O_3

1 g or 1 equivalent of H_2 reacts with

$X_2O_3 = 0.1596$ g

Equivalents of $X_2O_3 = \frac{0.1596}{6 \quad 10^{-3}} = 26.6$ g;

Equivalent mass of X_2O_3 = 26.6

= Equivalent mass of X + Equivalent mass of O E 8

E 8 26.6; E 18.6

$\because$ At. mass = Eq. mass Valency

18.6 3 = 55.8

181. (b) Let normality of $KMnO_4$ solution is N

N 10 50 1 N 5

m-eq. of MnO_4^- m-eq. of oxalic acid

N_1V_1 $\frac{W}{E}$ 1000

5 100 $\frac{W}{(90/2)}$ 1000 W 22.5 g

182. (d) Let the moles of $NaHC_2O_4$ in the mixture are 'a' and that of KHC_2O_4 $H_2C_2O_4$ are 'b'. The volumes of each component consumed are V litre.

In redox reaction :

when $KMnO_4$ is used : 0.2 V 2a 4b. ..(1)

In acid-base neutralization :

when NaOH is used: 0.12 V a 3b ...(2)

from (1) and (2) a 0.06 V

b 0.02 V or $a : b$ 3 : 1

183. (a) $\underset{n_f\ 2}{Sn^{2+}}$ SO_4 $\underset{n_f\ 1}{Fe^{2+}}$ $SO_4(NH_4)_2SO_4$ Sn^{4+} Fe^{3+}

Eq. of $SnSO_4$ eq. of FAS

Moles 2 Moles 1

Moles of $SnSO_4$ required for 1 mole FAS $\frac{1}{2}$

$\underset{n_f\ 5}{KMnO_4}$ $\underset{n_f\ 1}{FeSO_4(NH_4)SO_4}$ Fe^{3+} Mn^{2+}

Eq. of $KMnO_4$ Eq. of FAS

Moles 5 Moles 1

Moles of $KMnO_4$ req. $\frac{1}{5}$

Ratio $\frac{1/2}{1/5}$ $\frac{5}{2}$ 2.5

184. (d) eq. mass of $NaHC_2O_4$ as an acid 112

eq. of $NaHC_2O_4$ = eq. of NaOH

100 10^{-3} 0.2

0.02

a 0.02 112 = 2.24 g

eq. mass of $NaHC_2O_4$ as reductant

($C_2O_4^{2-}$ $2CO_2 + 2e^-$) $\frac{112}{2}$ 56

eq. of $NaHC_2O_4$ as reductant = eq. of $KMnO_4$ = 100 10^{-2} 0.02 5 = 0.01

b 0.01 56 = 0.56 g

Hence, $a : b$ $\frac{2.24}{0.56}$ 4 : 1 or a 4b

185. (c) eq. of $Na_2C_2O_4$ + eq. of $H_2C_2O_4$ = eq. of $KMnO_4$ V 0.1 5

2 2 0.5V; V = 8L

eq. of $H_2C_2O_4$ = eq. of NaOH

1 2 0.2 V_2; V_2 10 L

Hence, $V_1 : V_2$ 8 : 10 4 : 5

186. (a) eq. mass of $K_2C_2O_7$ = $\frac{\text{molar mass}}{6}$

eq. mass of $KMnO_4$ = $\frac{\text{molar mass}}{5}$

eq. of $Na_2S_2O_3$ = eq. of I_2 liberated = eq. of $KMnO_4$ + eq. of $K_2Cr_2O_7$

or N 1 0.02 5 0.05 6

N 0.4 or M 0.4

For More Material Join: @JEEAdvanced_2024

187. (b) m-eq. of acids in 50 its mL m-eq. of Na_2CO_3

$$25 \quad 2 \quad 50 \quad 4 \quad 2 \quad 2 \quad x \quad \frac{1000}{50} \quad 25 \quad 1$$

$x \quad 62.5$ mL

188. (d) Moles of pure $CuSO_4 = 0.12 =$ moles of $Na_2S_2O_3$ reacted with I_2

Moles of $Na_2S_2O_3 = 120 \quad 10^{-3} \quad M$

$$0.12$$

$$M \quad 1.0$$

189. (d) Let x g of Li_2CO_3 and $(1 \quad x)$ g of M_2CO_3 present in given mixture

2 Total moles of carbonates moles of HCl and

$$2 \quad \frac{x}{74} \quad \frac{1 \quad x}{2M \quad 60} \quad 21.6 \quad 0.5 \quad 10^{-3}$$

$$x \quad 0.20$$

$\because$

$$\frac{x}{74} \quad \frac{1 \quad x}{2M \quad 60}$$

$$M \quad 118$$

190. (b) m-eq. of $FeSO_4\ 7H_2O$ in 1 litre

$$20 \quad 0.02 \quad 5 \quad \frac{1000}{25} \quad 80$$

$$\frac{W}{278} \quad 1 \quad 1000 \quad 80 \quad W \quad 22.24$$

mass % of $FeSO_4\ 7H_2O$ in given sample

$$\frac{22.24}{32} \quad 100 \quad 69.5$$

191. (a) Since, phenolphthalein indicates conversion of Na_2CO_3 into $NaHCO_3$ only hence, x mL, of HCl will be further required to convert $NaHCO_3$ to H_2CO_3. So, total volume of HCl required to convert Na_2CO_3 into $H_2CO_3 \quad x \quad x \quad 2x$ mL

192. (d) In presence of phenolphthalein

$(Na_2CO_3 \quad NaHCO_3)$

50% No reaction

$\frac{1}{2}$ eq. of Na_2CO_3 eq. of HCl

$$\frac{1}{2} \quad \frac{x}{106} \quad 2 \quad \frac{10 \quad 0.01}{1000}$$

$$x \quad 0.0106$$

$$\% \text{ of } Na_2CO_3 \quad \frac{0.0106}{0.1} \quad 100 \quad 10.6$$

193. (c) Phenolphthalein gives the end point corresponding to the reactions :

$$NaOH + HCl \quad NaCl + H_2O$$

$$Na_2CO_3 + HCl \quad NaHCO_3 + NaCl$$

m-moles of NaOH + m-moles of Na_2CO_3

m-moles of HCl 2.5

Methyl orange gives the end point corresponding to the reactions :

$$NaOH + HCl \quad NaCl + H_2O$$

$$Na_2CO_3 + 2HCl \quad 2NaCl + CO_2 + H_2O$$

m-moles of NaOH

+ m moles of Na_2CO_3 2

= m-moles of HCl 30 0.1 3

m-moles of $Na_2CO_3 = 0.5$

m-moles of NaOH = 2.5 0.5 2

Ratio of m-moles of NaOH and Na_2CO_3

$$= \frac{2}{0.5} = 4 : 1$$

194. (a) $\underset{xg}{NaOH} \quad \underset{yg}{Na_2CO_3}$

In presence of HPh

Eq. of NaOH $\frac{1}{2}$ eq. of Na_2CO_3

Eq. of HCl

$$\frac{x}{40} \quad 1 \quad \frac{1}{2} \quad \frac{y}{106} \quad 2 \quad \frac{17.5}{1000} \quad \frac{1}{10} \qquad \ldots(1)$$

After this MeOH is added

$\frac{1}{2}$ eq. of Na_2CO_3 eq. of HCl

$$\frac{1}{2} \quad \frac{y}{106} \quad 2 \quad 2.5 \quad \frac{1}{10} \quad \frac{1}{1000} \qquad \ldots(2)$$

Placing the value of Eqn. ...(1)

$$\frac{x}{40} \quad \frac{2.5}{10000} \quad \frac{17.5}{10000}$$

$$\frac{x}{40} \quad 15 \quad 10^{-4} \qquad x \quad 15 \quad 40 \quad 10^{-4}$$

$x \quad 0.06$ g

195. (c) $CaCO_3 \qquad CaO \quad CO_2$

Mass of $CaCO_3 \quad 0.9$ g

Moles $\quad 9 \quad 10^{-3}$

$CO_2 \qquad 2NaOH \qquad Na_2CO_3 \quad H_2O$

$9 \quad 10^{-3} \qquad 50 \quad 10^{-3}$ moles

0

$$- \qquad 32 \quad 10^{-3} \qquad 9 \quad 10^{-3}$$

moles moles

$Na_2CO_3 \quad HCl \qquad NaHCO_3 \quad NaCl$

Eq. of HCl Eq. of NaOH Eq. of Na_2CO_3

For More Material Join: @JEEAdvanced_2024

$0.5 \quad V \quad 1 \quad 32 \quad 10^{-3} \quad 1 \quad 9 \quad 10^{-3} \quad 1$

$V \quad 82\ \text{mL}$

196. (a) $Na_2CO_3 + 2HCl \quad 2NaCl + H_2O + CO_2$

At the equivalent point m-mole of HCl

$2 \quad$ m-mole of Na_2CO_3

$$60 \quad M \quad 2 \quad \frac{0.318}{106} \quad 1000$$

$M \quad 0.1$

197. (a) $Ca(HCO_3)_2 \quad CaO \quad CaCO_3 \quad H_2O$

$162\ \text{g} \quad — \quad 56\ \text{g}$

$16.2\ \text{g} \quad — \quad 5.6\ \text{g}$

$162\ \text{g}\ Ca(HCO_3)_2 \quad 100\ \text{g}\ CaCO_3$

$16.2\ \text{g}\ Ca(HCO_3)_2 \quad 10\ \text{g}\ CaCO_3$

$$\text{Hardness} \quad \frac{\text{Mass of } CaCO_3}{\text{Mass of water}} \quad 10^6$$

$$\frac{10}{10^4} \quad 10^6 \quad 1000\ \text{ppm}$$

198. (c) Milli-equivalents of Ca^{2+}

$+$ milli-equivalents of Mg^{2+}

$=$ milli-equivalents of Na_2CO_3

$$\frac{20}{20} \quad \frac{12}{12} \quad V \quad 2$$

$V \quad 1$ mL for 1 L of tap water

For 5000 L of pond water

$V \quad 5000$ mL or 5 L

199. (d) 4.44 mg $CaCl_2$ is equivalent to 4 mg $CaCO_3$

1.9 mg $MgCl_2$ is equivalent to 2 mg $CaCO_3$

Total mass of $CaCO_3 = 6$ mg

$$\text{Hardness in ppm} \quad \frac{6 \quad 10^{-3}}{10^{-3}} \quad 10^6 \quad 6\ \text{ppm}$$

200. (d) $$\text{Hardness of water sample} \quad \frac{200\ \text{g of } CaCO_3}{10^6\ \text{g of } H_2O}$$

10^6 g of $H_2O \quad 200$ g of $CaCO_3$

10^3 g of $H_2O \quad \dfrac{200 \quad 10^3}{10^6}$ g of $CaCO_3$

Millimoles of $CaCO_3 \quad 2$

Milliequivalents of $CaCO_3 \quad 2 \quad 2 \quad 4$

Level 2

1. (a) Let wt. of NH_4NO_3 and $(NH_4)_2HPO_4$ are x and y gram respectively

$$\frac{\dfrac{x}{80} \quad 2 \quad 14 \quad \dfrac{y}{132} \quad 2 \quad 14}{x \quad y} \quad 100 \quad 30.4$$

$x : y \quad 2 : 1$

2. (a) Let V mL of alcohol be required

$\because$ mass of alcohol is same in both solutions

$$\frac{75}{100} \quad 0.8 \quad V \quad \frac{30}{100} \quad 0.9 \quad 150$$

$V \quad 67.5$ mL

3. (d) The precipitation reaction is

$Fe^{3+} + 3NH_3 + 3H_2O \quad Fe(OH)_3 + 3NH_4^+$

$$\text{mole of } Fe_2O_3 \text{ in sample} \quad \frac{0.80 \quad 0.5}{160}$$

$2.5 \quad 10^{-3}$

mole of $Fe^{3+} \quad 2 \quad 2.5 \quad 10^{-3} \quad 5 \quad 10^{-3}$

$$M_{NH_3} \quad \frac{0.986\ \text{g/mL} \quad 1000\ \text{mL/litre} \quad 0.025}{17}$$

$M_{NH_3} \quad 1.45$

3 moles of $Fe^{3+} =$ moles of NH_3

$$1.45 \quad V\ (\text{in L})$$

$$V \quad \frac{3 \quad 5 \quad 10^{-3}}{1.45} \quad 10.34 \quad 10^{-3}\ \text{L or } 10.34\ \text{mL}$$

4. (a) Balanced reaction is

$2F_2O_3 + 3C \quad 4Fe + 3CO_2$

Number of moles of Fe_2O_3

$$= \frac{120 \quad 1000}{2 \quad 56 \quad 48} \quad \frac{90}{100}$$

Mass of 80% pure iron produced

$$\frac{120 \quad 1000 \quad 0.9}{2 \quad 56 + 48} \quad \frac{2 \quad 56}{0.8}$$

94500 gram or 94.5 kg

5. (d) Let two equimolar carbonates are ACO_3 and BCO_3

mass of metal $A = 2.8 \quad 0.125 = 0.35$ g

$ACO_3 + BCO_3 \quad AO + BO + 2CO_2$

Total two moles CO_3^{2-} provide 2 moles of CO_2 or 60 g CO_3^{2-} provide 44 g of CO_2

Total mass of CO_3^{2-} required to produce 1.32 g of $CO_2 \quad \dfrac{60}{44} \quad 1.32 \quad 1.8$ g

mass of metal $(A + B) = 2.8 \quad 1.8 = 1$ g

mass of $B = 1 \quad 0.35 = 0.65$ g

$$\%\ \text{of } B \quad \frac{0.65}{2.80} \quad 100 \quad 23.21$$

6. (b) Loss in mass due to heating $\quad 6.2 \quad \dfrac{5}{100}$

$= 0.31$ g

$2NaHCO_3 \quad Na_2CO_3 + CO_2 + H_2O$

$2 \quad 84$ g $\quad 44 \quad 18$ g loss

1 mole $NaHCO_3 \quad 31$ g loss due to heating

For More Material Join: @JEEAdvanced_2024

0.31 g loss from 0.01 mole of $NaHCO_3$

Moles of Na_2CO_3 produced $\frac{0.01}{2}$ 0.005

Total moles of carbonate reacted with $BaCl_2$

7.5 0.2 $\frac{100}{10}$ 10^{-3} 0.015

Moles of carbonates in original sample

0.015 0.005 0.01

Mass of Na_2CO_3 in original sample

0.01 106 = 1.06 gram

7. (c) Moles of NO_2 required

$\frac{1575}{63}$ $\frac{3}{2}$ $\frac{1}{0.8}$ 46.875

moles of NO required $\frac{46.875}{0.60}$

moles of NH_3 required $\frac{46.875}{0.60}$ $\frac{1}{0.50}$

156.25

Volume of NH_3 at STP required

156.25 22.4 = 3500 L

8. (c) Moles of NaOH consumed when it reacts with H^+ = 3 1

moles of H_2SO_4 present in sample = 1.5

mass of H_2SO_4 in sample = 1.5 98 147 g

% purity $\frac{147}{183.75}$ 100 80

9. (c) Let mass of sample is x g; mass of MnO_2 is 0.75 x g

moles of MnO_2 $\frac{0.75x}{87}$

mass of Mn $\frac{0.75x}{87}$ 55 0.474x

$3MnO_2$ $Mn_3O_4 + O_2$

mass of Mn_3O_4 produced

$\frac{0.75x}{87}$ $\frac{1}{3}$ 229 0.658x g

Total mass of residue 0.658x 0.2 x

(mass of impurities)

% of Mn in residue $\frac{0.474x}{0.858x}$ 100

55.24

10. (b) Moles of AgCl = moles of chloride

$\frac{1.96}{143.5}$ 0.0136

Moles of organic compound $\frac{1}{147}$

6.8 10^{-3}

Chlorine atoms in each molecule of organic compound $\frac{0.0136}{6.8 \quad 10^{-3}}$ 2

11. (c) Let x g is mass of CaC_2O_4 and (0.6 x) g mass of MgC_2O_4

CaC_2O_4 $CaCO_3 + CO$

MgC_2O_4 $MgCO_3 + CO$

mass of $CaCO_3$ produced $\frac{x}{128}$ 100

mass of $MgCO_3$ produced $\frac{(0.6 \quad x)}{112}$ 84

$\frac{x}{128}$ 100 $\frac{(0.6 \quad x)}{112}$ 84 0.465

x 0.48 g

Due to further heating

$CaCO_3$ $CaO + CO_2$

– $\frac{x}{128}$

$MgCO_3$ $MgO + CO_2$

– $\frac{0.6 \quad x}{112}$

mass of CaO and MgO produced

$\frac{0.48}{128}$ 56 $\frac{0.12}{112}$ 40 0.252 g

12. (a) Moles of $BaSO_4 = \frac{1.22}{233.3}$

moles of $M_2(SO_4)_3 = \frac{1.22}{233.3}$ $\frac{1}{3}$

1.743 10^{-3}

wt. of $M_2(SO_4)_3$ = 0.596

1.743 10^{-3} (2M 96 3) 0.596

M 26.9

13. (b) $2NH_3 + CO_2$ $NH_2—CO—NH_2 + H_2O$

moles of urea $\frac{999.6 \quad 10^3}{60}$ 16.66 10^3

moles of CO_2 = 16.66 10^3

$C_nH_{2n \quad 2}$ O_2 $nCO_2 + H_2O$

$\frac{16.66 \quad 10^3}{n}$ [12n (2n 2)] 236 10^3

n 12

For More Material Join: @JEEAdvanced_2024

molecular formula of hydrocarbon

$C_{12}H_{26}$

14. (d) $C_nH_{2n\ \ 2}$ $\frac{3n\ \ 1}{2}$ O_2 nCO_2

$(n\ \ 1)\,H_2O$

Let initial pressure of $C_nH_{2n\ \ 2}$ is P then increase in pressure

$P\ \ (2n\ \ 1)\ \ 1\ \ \frac{3n\ \ 1}{2}$

$\frac{n\ \ 1}{2}\ \ P$

546 K and 4.6 atm 273 K and 2.3 atm;
Increase in pressure 0.3 atm

$P\ \ \frac{nRT}{V}\ \ \frac{11.6}{M}\ \ \frac{0.0821\ \ 273}{22.41}$

$\frac{n\ \ 1}{2}\ \ \frac{11.6}{(14n\ \ 2)}\ \ 0.3$

$\frac{n\ \ 1}{14n\ \ 2}\ \ \frac{0.6}{11.6}\ \ n\ \ 4$

Compound is C_4H_{10}

15. (d) Moles of $H_2SO_4 = 0.1$; mole of KOH 0.2
moles of H_2O_2 used in first reaction

$\frac{0.2}{2}\ \ \frac{1}{0.4}\ \ 0.25$

moles of O_2 produce $\frac{6.74}{22.4} = 0.3$

moles of H_2O_2 used in second reaction

$\frac{0.3}{3\ \ 0.5}\ \ 0.2$

Total mole of H_2O_2 consumed 0.45
Molarity of $H_2O_2 = \frac{0.45}{0.15} = 3\ M$

Volume strength 11.2 3 = 33.6 volumes

16. (b) $SO_2Cl_2 + 2H_2O$ $H_2SO_4 + 2HCl$
m-moles of H_2SO_4 produced 5; m-moles of HCl produced 10
m-moles of $Ba(OH)_2$ required

5 (for H_2SO_4) + $\frac{10}{2}$ (for HCl) 10

$M\ \ V\ \ 10, 0.2\ \ V\ \ 10, V\ \ 50$ mL

17. (b) Only Na_2CO_3 reacts with H_2SO_4;

$Na_2CO_3 + H_2SO_4$ $Na_2SO_4 + H_2CO_3$

m-moles of Na_2CO_3 = m-moles of H_2SO_4

20 0.1 2

m-moles of Na_2CO_3 in 250 mL solution

$\frac{250}{25}\ \ 2\ \ 20$

wt. of Na_2CO_3 20 106 $10^{\ 3}$ 2.12 g

% of Na_2CO_3 $\frac{2.12}{5}$ 100 42.4

% of Na_2SO_4 100 42.4 57.6

18. (a) M_{NaOH} resultant

$\frac{20\ \ 0.2\ \ 35\ \ 0.1}{100}\ \ 0.075\ M$

Milli-equivalent of NaOH = milli-equivalent of $H_2C_2O_4$
Let mass of impure sample is x gram

40 0.075 $\frac{x\ \ 0.90}{90}$ 2 1000

x 0.15 gram

19. (b) Ag HNO_3 $AgNO_3$
$AgNO_3$ NaCl $NaNO_3$ AgCl
POAC on Ag
Mole of Ag in coin mole of Ag in AgCl

$a\ \ \frac{14.35}{143.5}\ \ 0.1$ mole

Mass of Ag in coin 0.1 108 10.8 g
% silver in coin $\frac{10.8}{11.34}$ 100% 95.2%

20. (c) Mass ratio of Y in compound A, B and C is 1 : 3 : 5, in compound B, 32 parts of X combine with 84 parts of Y
16 parts of X combine with 42 parts of Y
For Y 3 parts in compound B, then in compound C 5 parts
42 parts in compound B, then in the compound C $\frac{5}{3}$ 42 70 parts

21. (a) Mole of O_2 $\frac{67.2}{22.4}$ 3 mole

$3\,O_2$ $2\,O_3$

Mole of Ozone formed $\frac{2}{3}\ \ \frac{15}{100}\ \ 3$

0.3 mole

Mass of Ozone formed 0.3 48 g 14.4 g

22. (a) $[H^+] = 10^{\ 3}\ M$

So $[Ca^{2+}] = \frac{1}{2}\ \ 10^{\ 3}\ M$

Mass of $Ca^{2+} = \frac{10^{\ 3}}{2}\ \ 40$

For More Material Join: @JEEAdvanced_2024

So, mass of Ca^{2+} ions in 10^6 mL hard water

$$\frac{\frac{1}{2} \quad 10^{-3} \quad 40 \quad 10^6}{10^3} \quad 20$$

23. (a) m-eq. of acid = m-eq. of base

$$N_1V_1 \quad N_2V_2$$

$$\frac{29.4}{\frac{98}{n}} \quad 100 \quad 90 \quad \frac{20}{40} \quad \frac{1000}{500}$$

$$n \quad 3$$

24. (b) When phenolphthalein is used as an indicator : $0.05 \; x \quad 20 \quad 0.1 \quad 1; x \quad 40$ mL

When methyl orange is used as an indicator:

$$0.05 \; y \quad 20 \quad 0.1 \quad 3$$

$$y \quad 120 \text{ mL}$$

$$y \quad x \quad 80 \text{ mL}$$

25. (a) $I_2 + AsO_2 \quad 2H_2O \qquad HAsO_4^{2-} \quad 2I^- \qquad 3H^+$

m-eq. of $HAsO_2$ (in 50 mL) m-eq. of I_2

$$= 35 \quad 0.05 \quad 2 \quad 3.5$$

m-eq. of $HAsO_2$ in 250 mL $= 3.5 \quad \frac{250}{50}$

$$17.5$$

mass of $HAsO_2$ in sample

$$\frac{17.5}{2} \quad (108) \quad 10^{-3} \quad 0.945 \text{ g}$$

% of $HAsO_2$ in the sample $\frac{0.945}{3.78} \quad 100$

$$25\%$$

26. (a) m-eq. of FeO m-eq. of $KMnO_4$

$$0.25 \quad 5 \quad 100$$

m-mole of FeO ($n \quad 1$) $\frac{0.25 \quad 100 \quad 5}{1}$

$$125$$

total m-eq. or m-mole. of Fe^{2+}

$$1000 \quad 0.1 \quad 6 \quad 600$$

(from FeO and Fe_2O_3 after reaction with Zn dust)

m-mole of Fe^{2+} from $Fe_2O_3 = 600 \quad 125$

$$475$$

m-mole of $Fe_2O_3 = \frac{475}{2}$

Mass of FeO $\frac{125 \quad (56 \quad 16)}{1000}$ g $= 9$ g

Mass of $Fe_2O_3 = \frac{475}{2} \quad \frac{160}{1000} = 38$ g

$$\% \; Fe_2O_3 = \frac{38}{38 + 9} \quad 100 = 80.85\%$$

27. (d) $3Br_2 + 6OH^- \qquad 5Br^- \quad BrO_3^- \quad 3H_2O$ ...(1)

$6H^+ + BrO_3^- \quad 3C_2O_4^{2-} \qquad Br^- \quad 6CO_2 + 3H_2O$ ...(2)

10 m-mole of Br_2 produce (10/3) m-mole BrO_3^- and m-moles of

CaC_2O_4 required $= \frac{10}{3} \quad 3 \quad 10$

Mass of $CaC_2O_4 \quad 10 \quad 10^{-3} \quad 128$

% purity $\frac{10 \quad 10^{-3} \quad 128}{2} \quad 100 \quad 64$

28. (b) m-moles of KI $= 50 \quad 0.04 \quad 2$

m-equivalent of $Na_2S_2O_3$ reacted with 10 mL solution $20 \quad 2.5 \quad 10^{-3} \quad 50 \quad 10^{-3}$

m-equivalent of I left unreacted in 100 mL solution $50 \quad 10^{-3} \quad \frac{100}{10} \quad 0.5$

m-moles of I which reacts with $CuCO_3 \quad 2 \quad 0.5 \quad 1.5$

$\because \quad 2Cu^{2+} + 5I^- \qquad 2CuI + I_3^-$

m-moles of $Cu^{2+} = \frac{2}{5} \quad 1.5 \quad 0.6$

weight of $CuCO_3 = 0.6 \quad 10^{-3} \quad 123.5$

$$0.0741$$

weight % of $CuCO_3 = \frac{0.0741}{0.1} \quad 100$

$$74.1$$

29. (b) $\underset{(n \quad 6)}{Fe_2(C_2O_4)_3} + \underset{(n \quad 5)}{MnO_4^-} \quad H^+ \qquad 2Fe^{3+} + 6CO_2 + H_2O$

$FeC_2O_4 + MnO_4^- \quad H^+ \qquad Fe^{3+}$

$(n \quad 3) \quad (n \quad 5) \qquad +2CO_2 + H_2O$

Total equivalents of $(Fe_2(C_2O_4)_3 + FeC_2O_4)$

equivalents of $KMnO_4$

$$0.5 \quad 6 + 0.5 \quad 3 = x \quad 5$$

moles of $KMnO_4 = x \quad 0.9$

30. (b) m-eq. of $Na_2C_2O_4$ = m-eq. of $KMnO_4$ reacted

total m-eq. of $KMnO_4$– excess m-eq. of $KMnO_4$ reacted with $H_2C_2O_4$

$$45 \quad 0.02 \quad 5 \quad 10 \quad 0.1 \quad 2 = 2.5$$

$$1000 \quad \frac{W}{134} \quad 2 \quad 2.5$$

$$W_{Na_2C_2O_4} \quad 0.1675 \text{ g}$$

For More Material Join: @JEEAdvanced_2024

% purity of $Na_2C_2O_4$ in sample

$$\frac{0.1675}{0.2} \quad 100 \quad 83.75$$

31. (a) Reactions of $K_2Cr_2O_7$ and $KMnO_4$ with KI may be given as :

$K_2Cr_2O_7 + 7H_2SO_4 + 6KI \quad 4K_2SO_4 + Cr_2(SO_4)_3 + 7H_2O + 3I_2$

$2KMnO_4 + 8H_2SO_4 + 10KI \quad 6K_2SO_4 + 2MnSO_4 + 5I_2$

Thus, equivalent mass of $K_2Cr_2O_7$

$$= \frac{294}{6} = 49$$

equivalent mass of $KMnO_4 = \frac{158}{5} = 31.6$

$\because$ m-eq. of $K_2Cr_2O_7$ + m-eq. of $KMnO_4$ = m-eq. of I_2 = m-eq. of hypo.

Let the mass of $K_2Cr_2O_7$ x g;

mass of $KMnO_4 = (0.5 \quad x)$ g

$$\frac{x}{49} \quad \frac{(0.5 \quad x)}{31.6} \quad 150 \quad 0.1 \quad 10^{-3}$$

or $x \quad 0.0732$

$$\% \text{ of } K_2Cr_2O_7 \quad \frac{0.0732}{0.5} \quad 100$$

14.64

32. (b) $I_2 + 2Na_2S_2O_3 \quad 2NaI + Na_2S_4O_6$ ...(i)

m-moles of $Na_2S_2O_3$ consumed

15 0.4 = 6 m-mole

m-moles of I_2 consumed 3 m-mole

$3I_2 + 6NaOH \quad 5NaI + NaIO_3 + 3H_2O$ ...(ii)

m-mol of I_2 reacted with NaOH are

$$\frac{30 \quad 2 \quad 3}{2} \quad 12 \text{ m-mole}$$

Total m-mole of I_2 consumed in reaction (i) and (ii) 3 12 15 m-mole

$$\text{Molarity of } I_2 = \frac{15}{150} = 0.1\,M$$

33. (a) Let x and y are milli-moles H_2SO_4 and $H_2C_2O_4$ present in the given mixture reacts with base

m-eq. of acids m-eq. of base;

$$(x \quad y) \quad 2 \quad 3 \quad 0.1 \quad \frac{1000}{10} \quad \text{or} \quad x \quad y \quad 15$$

In second experiment only $KMnO_4$ reacted with $H_2C_2O_4$ in which Mn^{7+} converted into Mn^{2+} and $C_2O_4^{2-}$ converted into CO_2

milli-equivalent of $H_2C_2O_4$ = milli-equivalent of $KMnO_4$

$$y \quad 2 \quad 4 \quad 0.02 \quad 5 \quad \frac{1000}{100}$$

$y \quad 2$

$x \quad 13$, mass of $H_2SO_4 \quad 13 \quad 10^{-3} \quad 98$

1.274 g

wt. % of H_2SO_4 in sample $\frac{1.274}{3.185} \quad 100$

40%

35. (c) $KMnO_4 \quad C_2O_4^{2-} \quad H \quad Mn^{2} \quad CO_2$

$n_f \quad 5 \qquad n_f \quad 2$

16 mL 40 mL

0.05 M XM

m. eq. of $C_2O_4^{2-}$ m.eq. of $KMnO_4$

$2(40)X \quad 5 \quad 16 \quad 0.05$

$X \quad 0.05M$

$[H^+] \quad 2X \quad 0.1\,M$

pH 1

Level 3

Passage-1

1. (b) $H_2O + SO_3 \quad H_2SO_4$;

18 g water combines with 80 g SO_3

4.5 g of H_2O combines with 20 g of SO_3.

100 g of oleum contains 20 g of SO_3

or 20% free SO_3.

2. (c) Initial moles of free SO_3 present in oleum

$$\frac{12}{18} \quad \frac{2}{3} \text{ moles}$$

moles of water that can combines with SO_3

moles of free SO_3 combined with water

$$\frac{9}{18} \quad \frac{1}{2} \text{ mole}$$

moles of free SO_3 left $\frac{2}{3} \quad \frac{1}{2} = \frac{1}{6}$ mole

volume of free SO_3 at STP $\frac{1}{6} \quad 22.4$

= 3.73 L

3. (c) $Na_2CO_3 + H_2SO_4 \quad Na_2SO_4 + H_2O + CO_2$

moles of CO_2 formed = moles of Na_2CO_3 reacted $\frac{5.3}{106} = 0.05$

For More Material Join: @JEEAdvanced_2024

volume of CO_2 formed at 1 atm pressure and 300 K 0.05 24.63 1.23 L

4. (b) eq. of H_2SO_4 + eq. of SO_3 eq. of NaOH

$$\frac{x}{98} \quad 2 \quad \frac{(1 \quad x) \quad 2}{80} \quad 54 \quad 0.4 \quad 10^{3}$$

$$x \quad 0.74$$

$$\% \text{ of free } SO_3 \quad \frac{1 \quad 0.74}{1} \quad 100 = 26\%$$

Passage-2

1. (a) 1 litre of $H_2O_2(aq)$ provide 11.2 litre of O_2 at 1 atm and 273 K

moles of O_2 $\frac{11.2}{22.4} = 0.5$

$n_{H_2O_2}$ required 0.5 2

$$M_{H_2O_2} \quad \frac{n_{H_2O_2}}{V_{solution}} = 1 \quad M$$

2. (b) Strength in percentage means how many g H_2O_2 present per 100 mL

$\because$ M 1 and mol. mass of H_2O_2 34

34 g H_2O_2 present per litre of solution or 3.4 g H_2O_2 present per 100 mL of solution.

3. (b) m-eq. of H_2O_2 m-eq. of $KMnO_4$

20 N 0.05 5 80 N 1

$$N \quad \frac{\text{volume strength of } H_2O_2}{5.6}$$

volume strength of $H_2O_2 = 5.6$

4. (b) m-eq. of $Ba(MnO_4)_2$ m-eq. of H_2O_2

$$\because M \quad \frac{33.6}{11.2} \quad 3$$

$$\frac{w}{375} \quad 10 \quad 1000 \quad 3 \quad 125 \quad 2; \quad w \quad 28.125$$

$$\% \text{ purity} \quad \frac{w}{40} \quad 100$$

$$\frac{28.125}{40} \quad 100 = 70.31$$

Passage-3

1. (c) 136 mg of $CaSO_4$ 100 mg of $CaCO_3$;
190 mg of $MgCl_2$ 2 100 mg of $CaCO_3$
1 litre or 1000 g water contain 300 mg of $CaCO_3$
10^6 g water contain

$$\frac{300 \quad 10^{3}}{1000} \quad 10^6 = 300 \text{ ppm}$$

2. (b) 10^6 g water contain 100 g $CaCO_3$
10 L or 10^4 g water contain

$$\frac{100}{10^6} \quad 10^4 = 1 \text{ g of } CaCO_3$$

100 g $CaCO_3$ 162 g of $Ca(HCO_3)_2$
or 1 g $CaCO_3$ 1.62 g of $Ca(HCO_3)_2$
or 0.01 mole

$Ca(HCO_3)_2 + Ca(OH)_2$ $2CaCO_3 + 2H_2O$

moles of $Ca(OH)_2$ required = 0.01
or mass of $Ca(OH)_2$ required = 0.74 g

3. (b) m-moles of H present in outlet water
= m-moles of OH 50 0.1 = 5
m-moles of Ca^2 present in hard water $\frac{5}{2}$
= 2.5; (1 Ca^2 replaced by 2H)
no. of mg of Ca^2 ion 2.5 40 = 100
200 g sample hard water contain 100 mg of Ca^2

10^6 g sample hard water contain

$$\frac{100}{200} \quad 10^6 \quad 10^{3} = 500 \text{ ppm}$$

Passage-4

1. (c) n-factor 5 2 = 10

2. (a) H_3PO_2 is a monobasic acid n-factor 1

3. (a) n-factor 3 $\frac{2}{0.95}$ 0.95 = 0.85

$$E \quad \frac{M}{0.85}$$

4. (b) n-factor of VO 3; Fe_2O_3 1 2 = 2;
x and y are 2 and 3

Passage-5

1. (a) Cl_2 2NaOH NaCl NaClO H_2O
4 mole 4 mole
3NaClO 2NaCl $NaClO_3$
$\frac{3}{1}$ $\frac{4}{3}$ 4 mole $\frac{4}{3}$ mole
$4NaClO_3$ $3NaClO_4$ NaCl
$\frac{4}{3}$ mole 1 mole

For More Material Join: @JEEAdvanced_2024

Moles of $NaClO_4$ $\frac{122.5}{122.5}$ 1 mole

4 mol Cl_2 284 g Cl_2 will be required.

2. (c) Cl_2 2NaOH NaCl NaClO H_2O

1 mole 1 mole 1 mole

3NaClO 2NaCl $NaClO_3$

1 mole $\frac{2}{3}$ 0.67 mole

$4NaClO_3$ $3NaClO_4$ NaCl

$\frac{1}{3}$ mole $\frac{1}{4}$ $\frac{1}{3}$ $\frac{1}{12}$ mole 0.083 mole

Total NaCl 1 0.67 0.083 1.75 mole

3. (b) As $NaClO_3$ is intermediate it will be consumed in the complete reaction.

One or More Answers is/are Correct

1. (c, d)

1g-atom of nitrogen 1 mol *N* atoms

$\frac{1}{2}$ mol N_2 gas

$\frac{1}{2}$ mol N_2 gas 11.2 L N_2 at STP 14 g N_2

2. (a, b)

1g molecule V_2O_5 1 mole V_2O_5 2 mole V atom 5 mole O atom

5. (b, c, d)

xA *yB* A_xB_y

As per equivalent concept

Number of equivalent of *A* No. of equivalent of *B* No. of equivalent of A_xB_y.

6. (b, c, d)

Equivalent of $Ba(OH)_2$ 1 2 2

Acids having 2 equivalents will be exactly neutralised by $Ba(OH)_2$.

7. (b, c)

Species having same empirical formula (EF) will have same percentage of C.

	EF
CH_3COOH	CH_2O
$C_6H_{12}O_6$	CH_2O
C_2H_5OH	C_2H_6O
$HCOOCH_3$	CH_2O
HCOOH	CH_2O_2
CH_3OCH_3	C_2H_6O

8. (b, d)

$\because$ both are liquid CH_3OH is solute (less amount)

Mass of CH_3OH 30 0.8 24 g,

Mass of C_2H_5OH 60 0.92 55.2 g

Mass of solution 24 55.2 79.2 g

Volume of solution $\frac{79.2}{0.88}$ 90 mL

Molarity $\frac{nCH_3OH}{V(L)}$ $\frac{24/32}{90}$ 1000

8.33 mol L^{-1}

Molality $\frac{n_{Solute}}{w_{Solvent}(kg)}$ $\frac{24/32}{55.2}$ 1000

13.59

Mole fraction of solute $\frac{\frac{24}{32}}{\frac{24}{32}\ \frac{55.2}{46}}$ 0.385

Mole fraction of solvent 1 0.385 0.615

9. (a, b, c)

17 g/L H_2O_2 $\frac{17}{34}$ mol L^{-1} H_2O_2

Molarity of H_2O_2 $\frac{1}{2}$

M $\frac{V}{11.2}$ V $\frac{1}{2}$ 11.2 5.6

5.6 volume H_2O_2 means 1 mL will give 5.6 mL at 273 K and 1 atm

P_1V_1 P_2V_2

1 5.6 2 V_2

V_2 2.8 mL

10. (a, b)

(a) 46g of 70% $\frac{W}{V}$ HCOOH ($d_{Solution}$ 1.4 g/mL)

70% $\frac{W}{V}$ HCOOH 70 g HCOOH in 100 mL solution.

Mass of solution 1.4 100 140 g

So, in 140 g solution, mass of HCOOH 70 g

in 46 g, mass of HCOOH $\frac{70}{140}$ 46 23 g

(b) 10 *M* HCOOH 10 mole HCOOH in 1000 mL solution mass of solution 1000 g

Mass of HCOOH 10 46 460 g

For More Material Join: @JEEAdvanced_2024

So in 50 g solution mass of

$$\text{HCOOH} \quad \frac{460}{1000} \quad 50 \quad 23\,\text{g}$$

(c) $25\%\ \frac{W}{W}$ HCOOH 25 g HCOOH in 100 g solution.

So in 50 g solution, mass of HCOOH 12.5 g

11. (a,c,d) V_{strength} 56;

$$M \quad \frac{28}{11.2} \quad 2.5$$

1 L contain 2.5 moles of H_2O_2

or 2.5 34 = 85 g H_2O_2

Mass of 1 litre solution 265 g

($\because$ d 265 g/L)

w_{H_2O} 180 g or moles of H_2O 10

$$x_{H_2O_2} \quad \frac{2.5}{2.5 \quad 10} \quad 0.2$$

$$\%\ \frac{w}{v} \quad \frac{2.5 \quad 34}{1000} \quad 100 \quad 8.5$$

$$m \quad \frac{2.5}{180} \quad 1000 = 13.88$$

12. (a, b)

$$CO \quad \frac{1}{2}O_2 \qquad CO_2$$

(A) O_2 is L.R.

So, Volume of CO_2 produced 20 mL

Total volume of CO_2 20 60 80 mL

(B) CO is L.R.

Volume of CO_2 produced 30 mL

Total volume of CO_2 30 50 80 mL

(C) O_2 is L.R.

Volume of CO_2 produced 40 mL

Total volume of CO_2 30 40 70 mL

13. (a, c)

$$\underset{n\text{-factor} = 2}{H_3PO_4} \qquad \underset{n\text{-factor} = 2}{X(OH)_2} \qquad XHPO_4 \quad 2H_2O$$

(A) Equivalent weight of base $\frac{\text{Mol. wt}}{2}$

(B) Equivalent weight of H_3PO_4 $\frac{98}{2}$

(C) $\underset{n\text{-factor} = 1}{XHPO_4}$ $\underset{n\text{-factor} = 1}{\text{NaOH}}$

Moles of NaOH 1 moles of $XHPO_4$ 1

Moles of NaOH 1 1 1

(D)

$$2XHPO_4 \quad X(OH)_2 \qquad X_3(PO_4)_2 \quad 2H_2O$$

Moles 1 $\frac{1}{2}$

Moles of $X(OH)_2$ required $\frac{1}{2}$

14. (b, c, d)

$$x\text{Sn}^{2} \quad y\text{Cr}_2\text{O}_7^{2} \quad z\text{H}$$

$$a\text{Sn}^{4} \quad b\text{Cr}^{3} \quad c\text{H}_2\text{O}$$

Upon balancing the redox-reaction

$$3\text{Sn}^{2} \quad \text{Cr}_2\text{O}_7^{2} \quad 14\,\text{H}$$

$$3\text{Sn}^{4} \quad 2\text{Cr}^{3} \quad 7\text{H}_2\text{O}$$

15. (a, b)

Let n_{Cu_2S} a; n_{CuS} a

$$\underset{n\text{-factor} = 8}{Cu_2S} \quad \underset{6}{CuS} \quad \underset{10}{Ba(MnO_4)_2} \quad H \qquad Cu^{2} \quad SO_2 \quad Mn^{2}$$

16. (a, c, d)

Equivalent weight $\frac{\text{Molecular weight}}{n\text{-factor}}$

(A) n-factor of H_2PO_3 as acid 1

as base 1

In both case, eq. wt. of H_2PO_3 $\frac{81}{1}$ 81

(B) n-factor of H_2PO_4 as acid 1 or 2

as base 1

(C) In acidic medium

Eq. wt. of $KMnO_4$ $\frac{\text{Molecular weight}}{5}$

In basic medium

Eq. wt. of $KMnO_4$ $\frac{\text{Molecular weight}}{1}$

In neutral/slightly basic medium

Eq. wt. of $KMnO_4$ $\frac{\text{Molecular weight}}{3}$

(D) In MgH_2, O.S. of H 1

In H_2O_2, O.S. of H 1

Match the Column

1. (A) 0.5 mole of $SO_2(g)$ 32 g

weight 1.5 N_A atoms

11.2 L at 1 atm and 273 K

(B) 1 g of $H_2(g)$ 0.5 mole

H_2 11.2 L at 1 atm and 273 K

N_A atoms

For More Material Join: @JEEAdvanced_2024

(C) 0.5 moles of O_3 $1.5N_A$ atoms 11.2 L at 1 atm and 273 K 24 g weight

(D) 1 g molecules of O_2 1 mole of O_2 $2N_A$ atom 22.4 L at 1 atm and 273 K 32 g weight

2. (A) Moles of $CO_2(g)$ $\frac{44}{44}$ 1

1 mole $CO_2(g)$ 1 g-molecule $CO_2(g)$

(B) Moles of $CH_4(g)$ $\frac{35.2}{16}$ 2.2 moles

V 2.2 22.4 49.28 L at 1 atm and 273 K

(C) Moles of $O_3(g)$ $\frac{48}{48}$ 1

(D) Moles of $N_2O(g)$ $\frac{44}{44}$ 1

Number of O atoms N_A 1 N_A

3. Use % by moles $\frac{M_{avg} \quad M_1}{M_2 \quad M_1}$ 100

% by mass = % by moles $\frac{M_2}{M_{avg}}$

4. (A) $\overset{3}{Bi}_2\overset{2}{S}_3$ Bi^5 $\overset{0}{S}$

n-factor (2 2) (2 3) 10

(B) $Al_2(\overset{6}{Cr}_2O_7)_3$ Cr^3

n-factor (3 6) 18

(C) $\overset{2}{Fe}\overset{1}{S}_2$ $\overset{3}{Fe}_2O_3$ $\overset{4}{S}O_2$

n-factor (1 1) (5 2) 11

(D) $\overset{2}{Mn}(\overset{5}{N}O_3)_2$ $\overset{6}{Mn}O_4^2$ $\overset{2}{N}O$

n-factor |(4 1) (3 2)| 2

5. (P) $\overset{3}{Cr}\overset{1}{I}_3$ $\overset{6}{Cr}_2O_7^2$ $\overset{7}{I}O_4$

n-factor (3 1) (8 3) 27

(Q) $\overset{2}{Fe}(\overset{2}{S}\overset{4}{C}\overset{3}{N})_2$ Fe^3 $\overset{6}{S}O_4^2$ $\overset{4}{C}O_3^2$ $\overset{5}{N}O_3$

n-factor (1 1) (8 2) (8 2) 33

(R) $\overset{3}{N}H_4\overset{2}{S}\overset{4}{C}\overset{3}{N}$ $\overset{6}{S}O_4^2$ $\overset{4}{C}O_3^2$ $\overset{5}{N}O_3$

n-factor (8 2) (8 1) 24

(S) $\overset{3}{A}S_2\overset{2}{S}_3$ $\overset{5}{A}sO_3$ $\overset{6}{S}O_4^2$

n-factor (2 2) (8 3) 28

7. Moles of $NaIO_3$ $\frac{396}{198}$ 2

(1) $\overset{5}{I}O_3$ $3\overset{4}{H}SO_3$ $\overset{1}{I}$ $3\overset{6}{S}O_4^2$ 3H

n-factor $\frac{6}{1}$ $\frac{6}{3}$ $\frac{6}{1}$ $\frac{6}{3}$ $\frac{6}{3}$

(2) 5I IO_3 $\overset{6H}{}$ $3I_2$ $3H_2O$

n-factor $\frac{5}{5}$ 1, $\frac{5}{1}$ $\frac{5}{3}$ $\frac{5}{3}$

(A) n-factor of IO_3 in reaction (2) 5

(B) Moles of HSO_3 used in (1) reaction 3 moles of IO_3 3 2 6

(C) Moles of I_2 produced moles of I produced $\frac{3}{5}$ 2 $\frac{3}{5}$ 1.2

(D) Equivalents of IO_3 used in reaction (2) Equivalents of I

2 1 2

8. (A) $\overset{0}{Br}_2$ OH $\overset{5}{Br}O_3$ $\overset{1}{Br}$

Interamolecular, Disproportionation Redox Reaction

(B) $\overset{2}{Fe}Cl_2$ $K\overset{7}{Mn}O_4$ HCl $\overset{2}{Mn}Cl_2$ $\overset{3}{Fe}Cl_3$

Interamolecular Redox Reaction

(C) $\overset{1}{Ag}_2\overset{2}{O}$ $\overset{0}{Ag}$ $\frac{1}{2}\overset{0}{O}_2$

Interamolecular Redox

(D) $\overset{3}{N}H_4$ $\overset{5}{N}O_3$ $\overset{1}{N}_2O$ H_2O

Interamolecular comproportionation Redox Reaction

Assertion-Reason Type Question

5. N M n-factor and n-factor of H_3PO_3 2 (max).

9. $[\overset{2}{Fe}(\overset{2}{C}\overset{3}{N})_6]^4$ $\overset{3}{Fe}^3$ $\overset{4}{C}O_2$ $\overset{5}{N}O_3$

n-factor (1 1) (2 6) (8 6) 61

10. $\overset{3}{As}_2\overset{2}{S}_3$ $\overset{5}{N}O_3$ $\overset{5}{As}O_4^3$ $\overset{2}{N}O$ $\overset{6}{S}O_4^2$

n-factor (2 2) (8 3) n f (3 1) 28 3

11. NaOH H_3PO_4 NaH_2PO_4 H_2O

n-factor 1 1

Eq. wt of H_3PO_4 $\frac{M}{1}$

13. $\overset{0}{I}_2$ $\overset{5}{I}O_3$ $\overset{1}{I}$

For More Material Join: @JEEAdvanced_2024

Subjective Problems

1. Mass of liquid in 10 moles 280 10 2800 g

Volume $\frac{2800}{1.4}$ 2000 mL 2L

2. (M) $\frac{16}{5.6}$ (22.4)

M 64

64 32 $16x$

x 2

3. [H] $\frac{(200\ \ 1)\ \ (300\ \ 6)}{1000}$

[H] 2

4. $N_2(g)$ $H_2(g)$ $N_2H_2(l)$

5L 3L

2L

V (8) (2) 6

5. n_{SO_2} $\frac{3.78}{64}$ 0.06; Let moles of $CaSO_4$ produced x

POAC on *S*

SO_2 $CaSO_4$

$(n_S)_{SO_2}$ $(n_S)_{CaSO_4}$

0.06 1 x 1 x 0.06

Mass of $CaSO_4$ 0.06 136 8.16

6. Moles of H_2O 1

3Fe $4H_2O$ Fe_3O_4 $4H_2$

Consumed moles of Fe $\frac{3}{4}$

Mass of Fe $\frac{3}{4}$ 56 42

7. n_{KClO_3} $\frac{122.5}{122.5}$ 1

Moles of atoms 1 1 3 5

8. Let metal sulfate is $M_2(SO_4)_x$

x valency of metal

Specific heat molar mass 6.4

Molar mass $\frac{6.4}{0.057}$ 112.2 g/mol

POAC on metal

$$\frac{2}{112.2}\ \ \frac{4.51}{2(112.2)\ \ (96)(x)}\ \ 2$$

224.4 $96x$ 506.02

x 3

9. M_xCl_y MCl_y ($\because$ valency of Cl 1)

Molar mass of M_x Cl_y 85.5 2

171 g/mol

$(M)(x)$ $(35.5)(y)$ 171 ...(1)

$\frac{\text{Mass of chlorine}}{\text{Mass of metalic chloride}}$ $\frac{0.835}{1}$

$\frac{35.5\ y}{M(x)\ \ (35.5)(y)}$

On solving y 4 and x 1

x y 5

10. $Ba(OH)_2$ $xH_2O(aq)$. $HNO_3(aq)$

$Ba(NO_3)_2$ $H_2O(l)$

Eq. of $Ba(OH)_2$ xH_2O eq. of HNO_3

$\frac{0.7875}{(171\ \ 18x)}$ 2 $\frac{1}{4}$ $\frac{20}{1000}$

x 8

11. n-factor of acid Basicity (n)

Eq. of acid Eq. of base

$\frac{2}{600}$ n $\frac{1}{6}$ $\frac{100}{1000}$ 1

n 5

12. In HPh Eq. of NaOH Eq. of Na_2CO_3

Eq. of HCl

$(0.5)(1)$ $(0.5)(1)$ $(x)(1)$

x 1

In MeOH Eq. of NaOH + Eq. of Na_2CO_3

Eq. of $NaHCO_3$ Eq. of HCl

$(0.5)(1)$ $(0.5)(2)$ $(0.5)(1)$ $(y)(1)$

y 2

x y 3

13. $\overset{5}{BrO_3}$ $\overset{1}{Br}$ H $\overset{0}{Br_2}(l)$

n-factor 5

E_{KBrO_3} $\frac{M_{KBrO_3}}{5}$

14. $\overset{4}{SeO_2}$ $\overset{2}{CrSO_4}$ Se^x Cr^3

Eq. of SeO_2 Eq. of $CrSO_4$

$(12.5)(0.05)(4\ \ x)$ $(25)(0.1)(1)$

$(4\ \ x)$ 4

x 0

15. $\underset{n\text{-factor} = 1}{Cu^2}$ KI $\underset{n\text{-factor} = 2}{I_2}$ Cu

$\underset{n\text{-factor} = 2}{I_2}$ $\underset{n\text{-factor} = 1}{Na_2S_2O_3}$ NaI $Na_2S_4O_6$

For More Material Join: @JEEAdvanced_2024

Eq. of Cu^{2} Eq. of I_2 produced Eq. of I_2 reacted Eq. of $Na_2S_2O_3$.

$(x)(1) \quad \frac{40}{1000} \quad (0.1)(1)$

$x \quad 4 \quad 10^{3}$ mol

Mass of pure Cu $\quad 4 \quad 10^{3} \quad 63.5 \quad 0.254$ g

% Purity $\quad \frac{0.254}{0.276} \quad 100 \quad 92\%$

% impurity $\quad 8\%$

16. Eq. of H_2O_2 Eq. of $KMnO_4$

$(25 \quad M \quad 2) \quad (10 \quad 0.1 \quad 5)$

$M \quad 0.089$

Volume strength $\quad M \quad 11.2 \quad 1$

17. Balanced redox reaction

$28NO_3 \quad 3As_2S_3 \quad 4H_2O$

$6AsO_4^{3} \quad 28NO \quad 9SO_4^{2} \quad 8H$

18. $\underset{0.220g}{M_2O_x}$ (Metallic oxide) $\quad H_2$

$M(s) \quad \underset{0.045g}{H_2O(l)}$

POAC on O

$\frac{0.220}{2M \quad 16x} \quad x \quad \frac{0.045}{18} \quad 1$

$3.96x \quad 0.09\,M \quad 0.72x$

$E \quad \frac{M}{x} \quad 36$

$\frac{E}{9} \quad 4$

19. Let mass of $K_2Cr_2O_7 \quad a$ g

Mass of $KMnO_4 \quad (10 \quad a)$ g

Eq. of $K_2Cr_2O_7$ Eq. of $KMnO_4$

Eq. of $Na_2S_2O_3$

$\frac{9}{294} \quad 6 \quad \frac{10 \quad a}{158} \quad 5 \quad (2.2)(0.1)$

$a \quad 8.5$ g

Mass % of $KMnO_4 \quad Z \quad 15$

20. $a \quad 5 \quad 10^{4}$

% of Sn^{2} in sample

$(x) \quad \frac{5 \quad 10^{4} \quad 11.8}{0.4} \quad 100$

$x \sim 15$

$\frac{x}{5} \quad \frac{15}{5} \quad 3$

For More Material Join: @JEEAdvanced_2024


2

ATOMIC STRUCTURE

Atomic Number of an Element

Total number of protons present in the nucleus = Total number of electrons present in the atom

Mass number of an element = Number of protons + Number of neutrons.



e.g., $^{23}_{11}Na$, $^{35}_{17}Cl$ and so on.

Terms associated with elements

- Isotopes : Atoms having same number of protons but different mass number.
- Isobars : Elements having same mass number.
- Isotones : Elements having same number of neutrons (A Z).
- Isoelectronic : Species/elements having same number of electrons.
- Isosters : Species having same number of atoms and electrons
- Isodiaphers : Elements having same number of $|N$ $Z|$ or $|A$ $2Z|$
- Paramagnetic : Species having non-zero unpaired electron.
- Diamagnetic : Species having zero unpaired electron.

Rutherford's Model

- Electrons, protons & neutrons are the most important fundamental particles of atoms of all elements (Except hydrogen)
- $^{A}_{Z}X$, Mass number (A) Atomic number (Z) number of neutrons (n)
- R_N $R_0(A)^{1/3}$, R_0 1.33 10 ^{13}cm A mass number, R_N Radius of nucleus

For More Material Join: @JEEAdvanced_2024

❖ $\frac{1}{2}m$ v^2 $K\frac{q_1 \quad q_2}{r}$; r distance of closest approach , v Velocity of a -particle

m mass of -particle

q_1 charge on -particle


q_2 charge on metal foil.

Size of the Nucleus

The volume of the nucleus is very small and is only a minute fraction of the total volume of the atom. Nucleus has a diameter of the order of 10 ^{12} to 10 ^{13} cm and the atom has a diameter of the order of 10 ^{8} cm.

Thus, diameter (size) of the atom is 1,00,000 times the diameter of the nucleus.

Electromagnetic Spectrum



Light

❖ Photon is considered as massless bundle of energy.

❖ Energy of light E mc^2, where m mass of light particle, c speed of light

❖ E_{Photon} h $hc/$ hc^{-} $\frac{1240\,\text{eV.nm}}{(\text{nm})}$

where h Planck constant, wavelength of photon, $\bar{\nu}$ wave number.

❖ Quantum efficiency or Quantum Yield $\frac{\text{no.of molecules reacting}}{\text{no.of quanta absorbed}}$

Bohr's Model

❖ Electrostatic force $\frac{Kq_1q_2}{r^2}$ where K $\frac{1}{4 \quad _0}$ 9 10^9N - m^2/C^2

For More Material Join: @JEEAdvanced_2024

- Potential energy due to electrostatic force $\frac{Kq_1q_2}{r}$

  q_1 charge of electron, q_2 charge of nucleus

- Potential due to a charge (Q) particle at a distance (r) $\frac{KQ}{r}$
- Bohr quantization rule $mvr \quad n \frac{h}{2} \quad n.\hbar$
- According to Newtons' second law, in a uniform circular motion resultant of all the forces towards centre must be equal to $\frac{mv^2}{r}$.
- $\frac{Kq_1q_2}{r^2} \quad \frac{mv^2}{r}$,

  where q_1 charge of electron, q_2 charge of nucleus, m mass of electron,
  r radius of Bohr's orbit
- Total energy of electron in n^{th} Bohr orbit

  $E_n \quad \frac{E_1}{n^2}Z^2 \quad \frac{2\ ^2me^4K^2}{n^2h^2}Z^2; \qquad E_1 \quad \frac{2\ ^2me^4K^2Z^2}{h^2}$

  $E_n \quad 13.6 \quad \frac{z^2}{n^2}$eV/atom, where Z atomic number of single electron atoms/ion,

  n principal quantum number of shell, E_1 total energy of electron in I Bohr orbit.
- Radius of nth Bohr orbit, $r_n \quad \frac{h^2}{4\ ^2e^2mK} \quad \frac{n^2}{Z} \quad 0.529 \quad \frac{n^2}{Z}$Å $\quad r_1 \quad \frac{n^2}{Z}$ Å,

  where r_1 radius of I Bohr orbit.
- Velocity of electron in n^{th} Bohr orbit, $v_n \quad \frac{2\ e^2K}{h} \quad \frac{Z}{n} \quad 2.18 \quad 10^6 \quad \frac{Z}{n}$m/s $\quad v_1 \quad \frac{Z}{n}$ m/s

  where v_1 velocity of electron in I Bohr orbit.
- Revolutions per sec $\quad v/2\ r \quad 0.657 \quad 10^{16} \quad \frac{Z^2}{n^3}$
- Time for one revolution $\quad 2\ r/v \quad 1.52 \quad 10^{\ 16} \quad \frac{n^3}{Z^2}$
- Energy difference between n_1 and n_2 energy level.


  $E \quad E_{n_2} \quad E_{n_1} \quad 13.6Z^2 \quad \frac{1}{n_1^2} \quad \frac{1}{n_2^2}$ eV/atom $\quad$ IE $\quad \frac{1}{n_1^2} \quad \frac{1}{n_2^2}$

  where IE = ionization energy of single electron species.
- Ionization energy $\quad E \quad E_{G.S.} \quad E_{G.S.}$; $\quad E_{G.S.}$ Energy of electron in ground state
- Total energy of electron in terms of kinetic energy (KE) and potential energy (PE)

  $E_n \quad KE \quad PE \quad KE \quad \frac{PE}{2}$

For More Material Join: @JEEAdvanced_2024

Spectral Lines

- Rydberg's Equation $\frac{1}{} = R_H \frac{1}{n_1^2} \frac{1}{n_2^2} Z^2$; R_H 109700 cm ^{1} Rydberg constant
- For First line of a series n_2 n_1 1
- Limiting spectral line (series limit) means n_2
- H line means n_2 n_1 1; also known as line of longest shortest , least E
- Similarly H line means n_2 n_1 2
- When electrons de-excite from higher energy level (n) to ground state in atomic sample, then number of spectral lines observed in the spectrum $\frac{n(n \quad 1)}{2}$
- When electrons de-excite from higher energy level (n_2) to lower energy level (n_1) in atomic sample, then number of spectral line observed in the spectrum $\frac{(n_2 \quad n_1)(n_2 \quad n_1 \quad 1)}{2}$
- When electron de-excites from higher energy level (n_2) to lower energy level (n_1) in isolated atom, then number of spectral line observed in the spectrum $(n_2 \quad n_1)$
- 

Photoelectric Effect

- When radiation with certain minimum frequency ($_0$ called threshold frequency), strikes the surface of a metal, electrons (called photoelectrons) are ejected from the surface.
- Kinetic energy of photoelectron $h \quad w \quad h \quad h_0$
  where w work function
  $_0$ Threshold frequency
- If $_0$, then photoelectric effect takes place.
- Accelerating potential eV $KE \quad \frac{1}{2} m v^2$

For More Material Join: @JEEAdvanced_2024

de-Broglie Hypothesis

- All material particles possess wave character as well as particle character.
- h/mv h/p
- The circumference of the nth orbit is equal to n times of wavelength of moving electron $i.e.$,
  $2\ r_n$ n
  Number of waves n principal quantum number
- Wavelength of electron ( ) $\sqrt{\frac{150}{V(\text{volts})}}$ Å
- Wave nature of electron was confirmed by Davisson and Germer experiment.

Heisenberg Uncertainty Principle

- According to this principle, "it is impossible to measure simultaneously the position and momentum of a microscopic particle with absolute accuracy".
  If one of them is measured with greater accuracy, the other becomes less accurate.
- x p $h/4$ or (x)(v) $\frac{h}{4\ m}$ or (x)( ) $\frac{^2}{4}$
  where x Uncertainty in position, p Uncertainty in momentum.
  v Uncertainty in velocity, Uncertainty in wavelength.
  m mass of microscopic particle, Wavelength of microscopic particle
- Heisenberg replaced the concept of orbit by that of orbital.

Schrodinger Equation

- Schrodinger equation is central equation of wave mechanics according to following equation.

$$\frac{^2}{x^2}\ \frac{^2}{y^2}\ \frac{^2}{z^2}\ \frac{8\ ^2m}{h^2}(E\ V)\ \ 0$$

  Wave function $f(x, y, z)$
  E Total energy of particle
  V Potential energy of particle.
- A Schrodinger equation leads to infinite solutions.
- Most of the solution are not realistic (or acceptable). Only few solution can be accepted.
- Each solution - (x, y, z) correspond to a definite energy state which depends on quantum number n, l & m.
  By proper mathematical manipulation the main equation is broken in two parts and solved separately.
  (i) Radial part contain only 'r', depends on quantum numbers n & l.
  (ii) Angular part contain and , depends on quantum numbers l & m.
  Each value of contain all the information about that particular quantum state.
- **Atomic Orbital :** This is a three dimensional space around the nucleus within which the probability of finding the electron is maximum.
- **Degenerate Orbital :** Orbitals with same value of n and l of same subshell are degenerate orbitals, which have same energy.
  For example, $2p_x, 2p_y, 2p_z$ etc.

For More Material Join: @JEEAdvanced_2024

- **Radial Probability Density** $4\ r^2R^2(r)$

  It is the probability of finding electron in the region between r and $r\quad dr$
- **Radial Node :** It is zero electron density region. $R^2(r)\quad 0$ or $R(r)\quad 0$
- **Nodal Point :** It is a point $(r\quad 0)$ where electron density is zero.
- **Nodal Planes :** It is plane by which two lobes are separated and electron density is zero here.

Quantum Number

There are four of quantum number which are required in order to explain e in atom.

- **Principal quantum number (n) :** It determines the size of an orbital. Each value of n represents a shell of orbital. Possible values of $n\quad 1, 2, 3, 4,$.....
- **Azimuthal quantum number (l) :** It determines shape of an orbital. Each value of l represents a subshell of an orbital. Possible values of $l\quad 0,1, 2,$ ..... $(n\quad 1)$
- **Magnetic quantum number (m) :** It decides orientation of orbital in space.

  Possible values of $m\quad l,\ l\quad 1,$ ..... $0, 1, 2, l$


  Total $(2l\quad 1)$ values are possible form for each value of l.
- **Spin quantum number (s) :** It is intrinsic property of an electron. The electron has two spin states. Possible values of $s\quad \frac{1}{2},\ \frac{1}{2}$

Shell, sub-shells and orbitals

Shell (n)	Sub-shells (l)	Orbitals (m)
1	0	0
2	0	0
	1	$1, 0$
3	0	0
	1	$1, 0$
	2	$2,\quad 1, 0$

For More Material Join: @JEEAdvanced_2024

Various Curves



Important Points on Quantum Number

- Orbital angular momentum $\frac{h}{2}\sqrt{l(l+1)}$
- Spin angular momentum $\frac{h}{2}\sqrt{S(S+1)}$
- Spin Magnetic moment ( ) $\sqrt{n(n+2)}$ B.M. ; n number of unpaired electron
- Maximum number of electrons in a shell $2n^2$
- Maximum number of electrons in a subshell $2(2l+1)$
- Maximum number of electrons in an orbital 2
- Total number of orbitals in a subshell $2l+1$
- Number of subshells in a shell n
- Number of orbitals in a shell n^2
- Radial Nodes $(n-l-1)$
- Angular nodes l
- Total nodes $(n-1)$
- Azimuthal quantum number 0 1 2 3 4
  Name of sub-shell s p d f g

For More Material Join: @JEEAdvanced_2024

Pauli's Exclusion Principle

No two electrons in an atom can have the same set of all the four quantum numbers, *i.e.*, an orbital cannot have more than 2 electrons because three quantum numbers (principal, azimuthal and magnetic) at the most may be same but the fourth will be different, *i.e.*, spins will be in opposite directions.

Aufbau Principle

Electrons are filled in various orbitals in order of their increasing energies. An orbital of lowest energy is filled first. The sequence of orbitals in order of their increasing energy is :

$$1s, 2s, 2p, 3s, 3p, 4s, 3d, 4p, 5s, 4d, 5p, 6s, 4f, 5d, 6p, 7s, 5f, 6d, \ldots\ldots$$

The energy of the orbitals is governed by $(n \quad l)$ rule.

Hund's Rule

No electron pairing takes place in the orbitals in a sub energy shell until each orbital is occupied by an electron with parallel spin. Exactly half filled and fully filled orbitals make the atoms more stable, *i.e.*, $p^3, p^6, d^5, d^{10} f^7$ and f^{14} configurations are more stable.

For More Material Join: @JEEAdvanced_2024



Level 1

1. Which of the following pair is isodiaphers?
(a) $^{14}_{6}C$ and $^{23}_{11}Na$ (b) $^{24}_{12}Mg$ and $^{23}_{11}Na$
(c) $^{4}_{2}He$ and $^{16}_{8}O$ (d) $^{12}_{6}C$ and $^{15}_{7}N$

2. Which of the following does not characterise X-rays?
(a) The radiation can ionise the gas
(b) It causes fluorescence effect on ZnS
(c) It is deflected by electric and magnetic fields
(d) Its wavelength is shorter than ultraviolet rays

3. The ratio of specific charge of a proton and an -particle is :
(a) 2 : 1 (b) 1 : 2 (c) 1 : 4 (d) 1 : 1

4. The increasing order for the values of e/m (charge/mass) is :
(a) $e, p, n,$ (b) $n, p, e,$ (c) $n, p,$ $, e$ (d) $n,$ $, p, e$

5. The mass to charge ratio (m/e) for a cation is 1.5 10 ^{8} kg/C. What is the mass of this cation?
(a) 2.4 10 ^{19} g (b) 2.4 10 ^{27} g (c) 2.4 10 ^{24} g (d) None of these

6. Rutherford's experiment on scattering of alpha particles showed for the first time that atom has :
(a) electrons (b) protons (c) nucleus (d) neutrons

7. -particles are represented by :
(a) lithium atoms (b) helium nuclei (c) hydrogen nuclei (d) None of these

8. In Bohr's stationary orbits :
(a) electrons do not move
(b) electrons move emitting radiations
(c) energy of the electron remains constant
(d) angular momentum of the electron is $h/2$

9. On the basis of Bohr's model, the radius of the 3rd orbit is :
(a) equal to the radius of first orbit (b) three times the radius of first orbit
(c) five times the radius of first orbit (d) nine times the radius of first orbit

10. The correct expression derived for the energy of an electron in the nth energy level is for H-atom :
(a) $E_n \quad \frac{2\ ^2me^4K^2}{n^2h^2}$ (b) $E_n \quad \frac{^2me^4K^2}{2n^2h^2}$ (c) $E_n \quad \frac{2\ ^2me^2K^2}{n^2h^2}$ (d) $E_n \quad \frac{2\ ^2me^4K^2}{n^2h^2}$

11. Ionization energy for hydrogen atom in ergs, Joules and eV respectively is :
(a) 21.8 10 ^{12},218 10 ^{20},13.6 (b) 13.6 218 10 ^{20},21.8 10 ^{13}
(c) 21.8 10 ^{20},13.6,21.8 10 ^{13} (d) 21.8 10 ^{13},13.6,21.8 10 ^{20}

12. For any H like system, the ratio of velocities of electron in I, II & III orbit *i.e.*, $V_1 : V_2 : V_3$ will be:
(a) 1 : 2 : 3 (b) 1 : 1/2 : 1/3 (c) 3 : 2 : 1 (d) 1 : 1 : 1

For More Material Join: @JEEAdvanced_2024

13. The volume of nucleus is about :
(a) 10^{-4} times to that of an atom (b) 10^{-15} times to that of an atom
(c) 10^{-5} times to that of an atom (d) 10^{-10} times to that of an atom

14. An electron in an atom jumps in such a way that its kinetic energy changes from x to $\frac{x}{4}$. The change in potential energy will be :
(a) $\frac{3}{2}x$ (b) $\frac{3}{8}x$ (c) $\frac{3}{4}x$ (d) $\frac{3}{4}x$

15. The potential energy of an electron in the hydrogen atom is 6.8 eV. Indicate in which excited state, the electron is present?
(a) first (b) second (c) third (d) fourth

16. What is the potential energy of an electron present in N-shell of the Be^{3} ion?
(a) 3.4 eV (b) 6.8 eV (c) 13.6 eV (d) 27.2 eV

17. The kinetic and potential energy (in eV) of electron present in third Bohr's orbit of hydrogen atom are respectively :
(a) 1.51, 3.02 (b) 1.51, 3.02 (c) 3.02, 1.51 (d) 1.51, 1.51

18. The distance between 4th and 3rd Bohr orbits of He is :
(a) $2.645 \quad 10^{-10}$ m (b) $1.322 \quad 10^{-10}$ m (c) $1.851 \quad 10^{-10}$ m (d) None of these


19. What atomic number of an element "X" would have to become so that the 4th orbit around X would fit inside the I Bohr orbit of H atom ?
(a) 3 (b) 4 (c) 16 (d) 25


20. The ratio of velocity of the electron in the third and fifth orbit of Li^{2} would be :
(a) 3 : 5 (b) 5 : 3 (c) 25 : 9 (d) 9 : 25


21. If radius of second stationary orbit (in Bohr's atom) is R then radius of third orbit will be :
(a) $R/3$ (b) $9R$ (c) $R/9$ (d) $2.25\,R$


22. Which Bohr's orbit of Be^{3} has the same orbit radius as that of the ground state of hydrogen atom?
(a) 3 (b) 2 (c) 4 (d) 5

23. Select the incorrect graph for velocity of e in an orbit *vs.* Z, $\frac{1}{n}$ and n :

(a) 

(b) 

(c) 

(d) 

24. What is the frequency of revolution of electron present in 2nd Bohr's orbit of H-atom?
(a) $1.016 \quad 10^{16}\ s^{-1}$ (b) $4.065 \quad 10^{16}\ s^{-1}$ (c) $1.626 \quad 10^{15}\ s^{-1}$ (d) $8.2 \quad 10^{14}\ s^{-1}$

25. According to Bohr's atomic theory, which of the following relation is correct ?
(a) Potential energy of electron $\frac{Z^2}{n^2}$
(b) The product of velocity of electron and principal quantum number (n) Z^2
(c) Frequency of revolution of electron in an orbit $\frac{Z^2}{n^3}$
(d) Coulombic force of attraction on the electron $\frac{Z^2}{n^2}$

For More Material Join: @JEEAdvanced_2024

26. Number of waves produced by an electron in one complete revolution in n^{th} orbit is :
(a) n (b) n^2 (c) $(n \quad 1)$ (d) $(2n \quad 1)$

27. Which of the following statement does not form part of Bohr's model of the hydrogen atom ?
(a) Energy of the electrons in the orbit is quantized
(b) The electron in the orbit which is nearest to the nucleus has the lowest energy
(c) Electrons revolve in different orbits around the nucleus
(d) The position and velocity of the electrons in the orbit cannot be determined simultaneously

28. What is the separation energy (in eV) for electron present in the first excited state of Be^{3} ?
(a) 13.6 eV (b) 27.2 eV (c) 40.8 eV (d) 54.5 eV

29. If in Bohr's model, for unielectronic atom, time period of revolution is represented as $T_{n, Z}$ where n represents shell no. and Z represents atomic number then the value of $T_{1, 2} : T_{2, 1}$ will be :
(a) 8 : 1 (b) 1 : 8 (c) 1 : 1 (d) 1 : 32

30. Which of the following is discreted in Bohr's theory?
(a) Potential energy (b) Kinetic energy
(c) Velocity (d) Angular momentum


31. What is the ratio of time periods (T_1 / T_2) in second orbit of hydrogen atom to third orbit of He ion?
(a) 8/27 (b) 32/27 (c) 27/32 (d) None of these


32. The mass of an electron is m, charge is e and it is accelerated from rest through a potential difference of V volts. The velocity acquired by electron will be :
(a) $\sqrt{\frac{V}{m}}$ (b) $\sqrt{\frac{eV}{m}}$ (c) $\sqrt{\frac{2eV}{m}}$ (d) zero


33. If the ionization energy of He is 19.6 10^{18} J per atom then the energy of Be^{3} ion in the second stationary state is :
(a) 4.9 10^{18} J (b) 44.1 10^{18} J (c) 11.025 10^{18} J (d) None of these


34. The energy of the second Bohr orbit in the hydrogen atom is 3.41 eV. The energy of the second Bohr orbit of He ion would be :
(a) 0.85 eV (b) –13.64 eV (c) –1.70 eV (d) –6.82 eV

35. The energy of an electron moving in n^{th} Bohr's orbit of an element is given by $E_n \quad \frac{-13.6}{n^2} Z^2$ eV/atom (Z atomic number). The graph of E *vs.* Z^2 (keeping "n" constant) will be :

(a) 

(b) 

(c) 

(d) 

36. If $_0$ be the permittivity of vacuum and r be the radius of orbit of H -atom in which electron is revolving, then velocity of electron is given by :

For More Material Join: @JEEAdvanced_2024

(a) $v = \frac{e}{\sqrt{4 \quad _0 rm}}$ (b) $v = e\sqrt{4 \quad _0 rm}$ (c) $v = \frac{4 \quad _0 rm}{e}$ (d) $v = \frac{4 \quad _0 rm}{e^2}$

37. Which of the following statement(s) is/are consistent with the Bohr theory of the atom (and no others)?

(1) An electron can remain in a particular orbit as long as it continuously absorbs radiation of a definite frequency.

(2) The lowest energy orbits are those closest to the nucleus.

(3) All electrons can jump from the *K* shell to the *M* shell by emitting radiation of a definite frequency.

(a) 1, 2, 3 (b) 2 only (c) 3 only (d) 1, 2

38. The ionization potential for the electron in the ground state of the hydrogen atom is 13.6 eV atom ^{1}. What would be the ionization potential for the electron in the first excited state of Li^{2} ?

(a) 3.4 eV (b) 10.2 eV (c) 30.6 eV (d) 6.8 eV


39. What is the energy content per photon (J) for light of frequency 4.2 10^{14} Hz?


(a) 2.8 10^{21} (b) 2.5 10^{19} (c) 2.8 10^{19} (d) 2.5 10^{18}


40. Wavelength for high energy EMR transition in H-atom is 91 nm. What energy is needed for this transition?


(a) 1.36 eV (b) 1240 eV (c) 13 eV (d) 13.6 eV

41. Which graph shows how the energy *E* of a photon of light is related to its wavelengths ( ) ?

(a) 

(b) 

(c) 

(d) 

42. Assume that 2 10^{17} J of light energy is needed by the interior of the human eye to see an object. How many photons of yellow light with 595.2 nm are needed to generate this minimum energy?

(a) 6 (b) 30 (c) 45 (d) 60

43. Line spectra is characteristic of :

(a) molecules (b) atoms (c) radicals (d) none of these

44. The spectrum produced from an element is :

(a) atomic spectrum (b) line spectrum

(c) absorption spectrum (d) any one of the above

For More Material Join: @JEEAdvanced_2024

45. Electronic transition in He ion takes from n_2 to n_1 shell such that :

$2n_2 \quad 3n_1 \quad 18$

$2n_2 \quad 3n_1 \quad 6$

What will be the total number of photons emitted when electrons transit to n_1 shell ?

(a) 21 (b) 15 (c) 20 (d) 10

46. Which of the following expressions represents the spectrum of Balmer series (If n is the principal quantum number of higher energy level) in Hydrogen atom ?

(a) $\bar{v} \quad \frac{R(n \quad 1)(n \quad 1)}{n^2}$ (b) $\bar{v} \quad \frac{R(n \quad 2)(n \quad 2)}{4n^2}$

(c) $\bar{v} \quad \frac{R(n \quad 2)(n \quad 2)}{n^2}$ (d) $\bar{v} \quad \frac{R(n \quad 1)(n \quad 1)}{4n^2}$

47. Multiple or fine structure of spectral lines is due to :

(a) presence of main energy levels (b) presence of sub-levels

(c) presence of electronic configuration (d) is not a characteristics of the atom

48. With increasing principal quantum number, the energy difference between adjacent energy levels in H-atom:

(a) decreases

(b) increases

(c) remains constant

(d) decreases for low value of Z and increases for higher value of Z.

49. Find the value of wave number (¯) in terms of Rydberg's constant, when transition of electron takes place between two levels of He ion whose sum is 4 and difference is 2.

(a) $\frac{8R}{9}$ (b) $\frac{32R}{9}$ (c) $\frac{3R}{4}$ (d) none of these

50. What is the wavelength in nm of the spectral line associated with a transition from $n \quad 3$ to $n \quad 2$ for the Li^{2} ion?

(a) 219 (b) 656 (c) 73.0 (d) 486

51. What is the energy (kJ/mol) associated with the de-excitation of an electron from $n \quad 6$ to $n \quad 2$ in He ion?

(a) $1.36 \quad 10^{6}$ (b) $1.36 \quad 10^{3}$ (c) $1.16 \quad 10^{3}$ (d) $1.78 \quad 10^{3}$

52. What is the shortest wavelength line in the Paschen series of Li^{2} ion?

(a) $\frac{R}{9}$ (b) $\frac{9}{R}$ (c) $\frac{1}{R}$ (d) $\frac{9R}{4}$

53. What is the maximum wavelength line in the Lyman series of He ion?

(a) $3R$ (b) $\frac{1}{3R}$ (c) $\frac{4}{4R}$ (d) None of these

54. Which of the following electron transitions in a hydrogen atom will require the largest amount of energy?

(a) from $n \quad 1$ to $n \quad 2$ (b) from $n \quad 2$ to $n \quad 4$

(c) from $n \quad 5$ to $n \quad 1$ (d) from $n \quad 3$ to $n \quad 5$

55. Which electronic transition in a hydrogen atom, starting from the orbit $n \quad 7$, will produce infrared light of wavelength 2170 nm? (Given : $R_H \quad 1.09677 \quad 10^{7} \text{ m}^{1}$)

For More Material Join: @JEEAdvanced_2024

(a) n 7 to n 6 (b) n 7 to n 5
(c) n 7 to n 4 (d) n 7 to n 3

56. Hydrogen atoms in the ground state are excited by monochromatic radiation of wavelength Å. The resulting spectrum consists of maximum 15 different lines. What is the wavelength ? (R_H 109737 cm ^{1})

(a) 937.3 Å (b) 1025 Å (c) 1236 Å (d) None of these

57. Electromagnetic radiation (photon) with highest wavelength results when an electron in the hydrogen atom falls from n 6 to :

(a) n 1 (b) n 2 (c) n 3 (d) n 5

58. When an electron jumps from L to K shell -

(a) energy is absorbed
(b) energy is released
(c) energy is neither absorbed nor released
(d) energy is sometimes absorbed and some times released

59. How do the energy gaps between successive electron energy levels in an atom vary from low to high n values?

(a) All energy gaps are the same
(b) The energy gap decreases as n increases
(c) The energy gap increases as n increases
(d) The energy gap changes unpredictably as n increases

60. The H-spectrum shows:

(a) Heisenberg's uncertainty principle (b) diffraction
(c) polarization (d) presence of quantized energy level

61. Splitting of spectral lines under the influence of magnetic field is called

(a) Zeeman effect (b) Stark effect
(c) photoelectric effect (d) None of these

62. The colour of sky is due to

(a) absorption of light by atmospheric gases (b) transmission of light
(c) wavelength of scattered light (d) all of the above


63. In photoelectric effect, the kinetic energy of photoelectrons increases linearly with the


(a) wavelength of incident light (b) frequency of incident light
(c) velocity of incident light (d) atomic mass of an element

64. Slope of V_0 *vs* curve is (where V_0 Stopping potential, subjected frequency)


(a) e (b) $\frac{h}{e}$ (c) (d) h


65. According to Einstein's photoelectric equation, the graph between kinetic energy of photoelectrons ejected and the frequency of the incident radiation is :

(a) 

(b) 

For More Material Join: @JEEAdvanced_2024

(c) 

(d) 

66. The photoelectric emission from a surface starts only when the light incident upon the surface has certain minimum :
(a) intensity (b) wavelength (c) frequency (d) velocity

67. If $_0$ and be the threshold wavelength and the wavelength of incident light, the speed of photo-electrons ejected from the metal surface is :
(a) $\sqrt{\frac{2h}{m}(\ _0 \ )}$ (b) $\sqrt{\frac{2hc}{m}(\ _0 \ )}$ (c) $\sqrt{\frac{2hc}{m} \ \frac{ \ _0}{ \ _0}}$ (d) $\sqrt{\frac{2h}{m} \ \frac{1}{ \ _0} \ \frac{1}{ \ }}$

68. A light source of wavelength illuminates a metal and ejects photo-electrons with $(K.E.)_{max}$ 1 eV
Another light source of wavelength $\frac{ }{3}$, ejects photo-electrons from same metal with $(K.E.)_{max}$ 4 eV
Find the value of work function?
(a) 1 eV (b) 2 eV (c) 0.5 eV (d) None of these

69. Electromagnetic radiation having 310 Å is subjected to a metal sheet having work function 12.8 eV. What will be the velocity of photo-electrons having maximum kinetic energy.
(a) 0, no emission will occur (b) 4.352 10^6 m/s
(c) 3.09 10^6 m/s (d) 8.72 10^6 m/s

70. The ratio of slopes of K_{max} *vs.* ν and V_0 *vs.* curves in the photoelectric effect gives (ν frequency, K_{max} maximum kinetic energy, V_0 stopping potential) :
(a) charge of electron
(b) Planck's constant
(c) work function
(d) the ratio of Planck's constant and electronic charge

71. Radiation corresponding to the transition n 4 to n 2 in hydrogen atoms falls on a certain metal (work function 2.5 eV). The maximum kinetic energy of the photo-electrons will be :
(a) 0.55 eV (b) 2.55 eV (c) 4.45 eV (d) None of these

72. Select the incorrect statement.
(a) K.E. of photo-electron does not depend upon the wavelength of incident radiation
(b) Photoelectric current depends on intensity of incident radiation and not on frequency
(c) Stopping potential depends on frequency of radiation and not on intensity
(d) None of these


73. Which is the de-Broglie equation?
(a) h p (b) h p^{1} (c) h p^{1} (d) h p

For More Material Join: @JEEAdvanced_2024

74. Which of the following has the largest de Broglie wavelength (all have equal velocity)?
(a) CO_2 molecule (b) NH_3 molecule (c) Electron (d) Proton


75. The de-Broglie wavelength associated with a particle of mass 10^{-6} kg with a velocity of $10\ m\,s^{-1}$ is :
(a) $6.63 \quad 10^{-22}$ m (b) $6.63 \quad 10^{-29}$ m (c) $6.63 \quad 10^{-31}$ m (d) $6.63 \quad 10^{-34}$ m


76. For two particles A and B, curves are plotted $\sqrt{V}$ against de-Broglie wavelengths, where V is the potential on the particles. Which of the following relation is correct about the mass of particles?





(a) $m_A \quad m_B$ (b) $m_A \quad m_B$ (c) $m_A \quad m_B$ (d) $m_A \quad m_B$

77. Which of following graphs correctly represents the variation of particle momentum with de-Broglie wavelength ?

(a)


(b)


(c)


(d)


78. An excited state of H atom emits a photon of wavelength and returns in the ground state. The principal quantum number of excited state is given by :
(a) $\sqrt{R(R \quad 1)}$ (b) $\sqrt{\dfrac{R}{(R \quad 1)}}$ (c) $\sqrt{R(R \quad 1)}$ (d) $\sqrt{\dfrac{(R \quad 1)}{R}}$

79. A dye absorbs a photon of wavelength and re-emits the same energy into two photons of wavelengths $_1$ and $_2$ respectively. The wavelength is related with $_1$ and $_2$ as :
(a) $\dfrac{1 \quad 2}{1 \quad 2}$ (b) $\dfrac{1 \quad 2}{1 \quad 2}$ (c) $\dfrac{^2_1 \ ^2_2}{1 \quad 2}$ (d) $\dfrac{1 \quad 2}{(1 \quad 2)^2}$

80. Be^{3} and a proton are accelerated by the same potential. Their de-Broglie wavelengths have the ratio (assume mass of proton mass of neutron) :
(a) 1 : 2 (b) 1 : 4 (c) 1 : 1 (d) $1:3\sqrt{3}$

For More Material Join: @JEEAdvanced_2024

81. The de-Broglie wavelength of an electron accelerated by an electric potential of V volts is given by :

(a) $\frac{1.23}{\sqrt{m}}$ (b) $\frac{1.23}{\sqrt{h}}\ m$ (c) $\frac{1.23}{\sqrt{V}}$ nm (d) $\frac{1.23}{V}$

82. An electron travels with a velocity of x ms ^{1}. For a proton to have the same de-Broglie wavelength, the velocity will be approximately :

(a) $\frac{1840}{x}$ (b) $\frac{x}{1840}$ (c) $1840\ x$ (d) x

83. The momentum (in kg-m/s) of photon having 6 MeV energy is :

(a) 3.2 10 ^{21} (b) 2.0 (c) 1.6 10 ^{21} (d) none of these

84. The number of photons of light having wave number 'x' in 10 J of energy source is :

(a) $10hcx$ (b) $\frac{hc}{10x}$ (c) $\frac{10}{hcx}$ (d) none of these

85. Which of the following relates to photons both as wave motion and as a stream of particles?

(a) Interference (b) E mc^2 (c) Diffraction (d) E h

86. If a_0 be the radius of first Bohr's orbit of H-atom, the de-Broglie's wavelength of an electron revolving in the second Bohr's orbit will be :

(a) 6 a_0 (b) 4 a_0 (c) 2 a_0 (d) None of these

87. Energy required to ionise 2 mole of gaseous He ion present in its ground state is :

(a) 54.4 eV (b) 108.8 N_A eV (c) 54.4 N_A eV (d) 108.8 eV

88. Which of the following is the most correct expression for Heisenberg's uncertainty principle ?

(a) $x.\ p$ $\frac{h}{4}$ (b) $x.\ p$ $\frac{h}{4}$ (c) $x.\ p$ $\frac{h}{4}$ (d) $x.\ v$ $\frac{h}{4}$

89. The Heisenberg uncertainty principle can be applied to :

(a) a cricket ball (b) a football (c) a jet aeroplane (d) an electron

90. The mass of a particle is 10 ^{10} g and its radius is 2 10 ^{4} cm. If its velocity is 10 ^{6} cm sec ^{1} with 0.0001% uncertainty in measurement, the uncertainty in its position is :

(a) 5.2 10 ^{8} m (b) 5.2 10 ^{7} m (c) 5.2 10 ^{6} m (d) 5.2 10 ^{9}

91. If an electron is travelling at 200 m/s within 1 m/s uncertainty, what is the theoretical uncertainty in its position in m (micrometer)?

(a) 14.5 (b) 29 (c) 58 (d) 114

92. The wave character of moving electron was experimentally verified by :

(a) de-Broglie (b) A. Einstein (c) Germer (d) Schrodinger

93. "The exact path of electron in 2p-orbital cannot be determined." The above statement is based upon :

(a) Hund's Rule (b) Bohr's Rule
(c) uncertainty principle (d) aufbau principle

94. Which series of subshells is arranged in the order of increasing energy for multi-electron atoms?

(a) $6s, 4f, 5d, 6p$ (b) $4f, 6s, 5d, 6p$ (c) $5d, 4f, 6s, 6p$ (d) $4f, 5d, 6s, 6p$

95. The correct Schrodinger's wave equation for a electron with total energy E and potential energy V is given by :

For More Material Join: @JEEAdvanced_2024

(a) $\frac{}{x^2}$ $\frac{}{y^2}$ $\frac{}{z^2}$ $\frac{8\ ^2}{mh^2}(E\quad V)\quad 0$ (b) $\frac{}{x^2}$ $\frac{}{y^2}$ $\frac{}{z^2}$ $\frac{8\ m}{h^2}(E\quad V)\quad 0$

(c) $\frac{}{x^2}$ $\frac{}{y^2}$ $\frac{}{z^2}$ $\frac{8\ ^2 m}{h^2}(E\quad V)\quad 0$ (d) None of the above

96. Wave mechanical model of the atom depends upon :
(a) de-Broglie concept of dual nature of electron
(b) Heisenberg uncertainty principle
(c) Schrodinger uncertainty principle
(d) All of these

97. In Schrodinger wave mechanical model, $^2(r, ,)$ represents :
(a) amplitude of electron wave
(b) probability density of electron
(c) total probability of finding electron around nucleus
(d) orbit

98. Radial amplitude of electron wave can be represented by :
(a) $R(r)$ (b) $R^2(r)$ (c) $4\ r^2$ (d) $4\ r^2R^2(r)$

99. Arrange the following orbitals of H-atom in the increasing order of their energy.
$3p_x, 2s, 4d_{xy}, 3s, 4p_z, 3p_y, 4s$
(a) $2s\quad 3s\quad 3p_x\quad 3p_y\quad 4s\quad 4p_z\quad 4d_{xy}$
(b) $2s\quad 3s\quad 3p_x\quad 3p_y\quad 4s\quad 4p_z\quad 4d_{xy}$
(c) $2s\quad 3s\quad 3p_x\quad 3p_y\quad 4s\quad 4p_z\quad 4d_{xy}$
(d) $2s\quad 3s\quad 3p_x\quad 3p_y\quad 4s\quad 4p_z\quad 4d_{xy}$

100. In a hydrogen atom, which orbital is higher in energy than a $3s$-orbital ?
(a) $2s$ (b) $3p$ (c) $3d$ (d) $4s$

101. The radii of maximum probability for $3s, 3p$ and $3d$ electrons are in the order :
(a) $(r_{max})_{3d}\quad (r_{max})_{3p}\quad (r_{max})_{3s}$ (b) $(r_{max})_{3d}\quad (r_{max})_{3s}\quad (r_{max})_{3p}$
(c) $(r_{max})_{3s}\quad (r_{max})_{3p}\quad (r_{max})_{3d}$ (d) None of these

102. The correct order of closeness of $3s, 3p, 3d$ orbitals of nucleus is :
(a) $3d\quad 3p\quad 3s$ (b) $3s\quad 3p\quad 3d$
(c) $3s\quad 3d\quad 3p$ (d) $3d\quad 3s\quad 3p$

103. The correct order of total number of node of atomic orbitals is:
(a) $4f\quad 6s\quad 5d$ (b) $6s\quad 5d\quad 4f$
(c) $4f\quad 5d\quad 6s$ (d) $5d\quad 4f\quad 6s$

104. If the subsidiary quantum number of a subenergy level is 4, then no. of degenerate orbitals are :
(a) 3 (b) 5 (c) 9 (d) 11

105. Which two orbitals are located along the axis, and not between the axis?
(a) d_{xy}, d_{z^2} (b) d_{xy}, p_z (c) d_{yz}, p_x (d) $p_z, d_{x^2\ y^2}$

106. In a set of degenerate orbitals, the electrons distribute themselves to retain similar spins as far as possible. This statement is attributed to :
(a) Pauli's exclusion principle
(b) aufbau principles

For More Material Join: @JEEAdvanced_2024

(c) Hund's Rule
(d) Slater rule

107. Which of the following rules could explain the presence of three unpaired electrons in N-atom?
(a) Hund's rule (b) aufbau's principles
(c) Heisenberg's uncertainty principle (d) Pauli's exclusion principle

108. Pauli's exclusion principle states that:
(a) nucleus of an atom contains no negative charge
(b) electrons move in circular orbits around the nucleus
(c) electrons occupy orbitals of lowest energy
(d) All the four quantum numbers of two electrons in an atom cannot be equal.

109. For which of the following sets of quantum numbers, an electron will have the highest energy?

	n	l	m	s		n	l	m	s
(a)	3	2	1	–1/2	(b)	4	3	–1	+1/2
(c)	4	1	–1	+1/2	(d)	5	0	0	–1/2

110. Which of the following statements concerning the four quantum numbers is false ?
(a) n gives idea of the size of an orbital
(b) l gives the shape of an orbital
(c) m gives the energy of the electron in the orbital
(d) s gives the direction of spin of the electron in an orbital

111. Maximum number of electrons in a subshell is given by
(a) $(2l \quad 1)$ (b) $2(2l \quad 1)$ (c) $(2l \quad 1)^2$ (d) $2(2l \quad 1)^2$


112. In any subshell, the maximum number of electrons having same value of spin quantum number is :
(a) $\sqrt{l(l \quad 1)}$ (b) $l \quad 2$ (c) $2l \quad 1$ (d) $4l \quad 2$

113. The orbital angular momentum of 3p electron is :
(a) $\sqrt{3}\,h$ (b) $\sqrt{6}\,h$ (c) zero (d) $\sqrt{2}\,\frac{h}{2}$

114. The atomic orbitals are progressively filled in order of increasing energy. The principle is called as:
(a) Hund's rule (b) aufbau principle
(c) exclusion principle (d) de-Broglie rule

115. The orbital diagram in which both the Pauli's exclusion principle and Hund's rule are violated, is :
(a) ↑↓ | ↓ | ↓ | ↓ (b) ↑↓ | ↑↓ | ↑↓ | ↑ (c) ↑↓ | ↑↓ | ↑↓ | (d) ↑↓ | ↑↑ | ↑↓ |


116. Which of the following elements is represented by the electronic configuration ?



(a) Nitrogen (b) Fluorine (c) Oxygen (d) Neon

For More Material Join: @JEEAdvanced_2024

117. The ratio of magnetic moments of Fe (III) and Co (II) is :
(a) $\sqrt{5} : \sqrt{7}$ (b) $\sqrt{35} : \sqrt{15}$ (c) 7 : 3 (d) $\sqrt{24} : \sqrt{15}$

118. If the electronic structure of oxygen atom is written as $1s^2, 2s^2$ 

it would violate
(a) Hund's rule (b) Pauli's exclusion principle
(c) both Hund's and Pauli's principles (d) None of these

119. A compound of vanadium has a magnetic moment ( ) of 1.73 BM. If the vanadium ion in the compound is present as V^x , then, the value of x is?
(a) 1 (b) 2 (c) 3 (d) 4

120. d^6 configuration will result in total spin of:
(a) $\frac{3}{2}$ (b) $\frac{1}{2}$ (c) 2 (d) 1

121. The probability of finding electrons in d_{xy} orbital is :
(a) along *X*- and *Y*-axis (b) along *X*- and *Z*-axis
(c) along *Y*- and *Z*-axis (d) at an angle of 45° with *X*-axis

122. Select correct statement :
(a) The lower the value of (n l) for an orbital, the higher is its energy.
(b) If two orbitals have the same value of (n l), the orbital with higher value of n will have lower energy.
(c) The energy of an electron in a multi-electron atom depends on quantum number n only
(d) The energy of an electron in hydrogen atom depends on quantum number n only

123. Read the following statements and choose the correct option.
(I) If the radius of the first Bohr orbit of hydrogen atom is r, then radius of 2nd orbit of Li^2 would be $4r$
(II) For *s*-orbital electron, the orbital angular momentum is zero
(a) only I is correct (b) only II is correct (c) both are correct (d) both are incorrect

124. The quantum numbers of four electrons ($e1$ to $e4$) are given below :

	n	l	m	s		n	l	m	s
$e1$	3	0	0	+1/2	$e2$	4	0	0	1/2
$e3$	3	2	2	–1/2	$e4$	3	1	–1	1/2

The correct order of decreasing energy of these electrons is :
(a) $e4$ $e3$ $e2$ $e1$ (b) $e2$ $e3$ $e4$ $e1$ (c) $e3$ $e2$ $e4$ $e1$ (d) $e1$ $e4$ $e2$ $e3$

125. The energy of an electron of $2p_y$ orbital is :
(a) greater than $2p_x$ orbital (b) less than $2p_z$ orbital
(c) equal to 2*s* orbital (d) same as that of $2p_x$ and $2p_z$ orbital

126. The number of unpaired valence electrons in an atom of phosphorus is :
(a) 0 (b) 2 (c) 3 (d) 4

127. Which quantum number defines the orientation of orbital in the space around the nucleus?
(a) principal quantum number (n) (b) angular momentum quantum number
(c) magnetic quantum number (m_l) (d) spin quantum number (m_s)

128. What is the maximum number of electrons in a subshell that can have the quantum numbers n 3 and l 2 ?
(a) 2 (b) 5 (c) 6 (d) 10

For More Material Join: @JEEAdvanced_2024

129. Which of the following statements about an electron with m_l 2 is incorrect?
(a) The electron could be in the third shell (b) The electron is in a non-spherical orbital
(c) The electron may have m_s $\frac{1}{2}$ (d) The electron is not in a *d*-orbital

130. Which of the following set of quantum numbers is impossible for an electron?
(a) n 1, l 0, m_l 0, m_s $\frac{1}{2}$ (b) n 9, l 7, m_l 6, m_s $\frac{1}{2}$
(c) n 2, l 1, m_l 0, m_s $\frac{1}{2}$ (d) n 3, l 2, m_l 3, m_s $\frac{1}{2}$

131. In a 3*d* subshell, all the five orbitals are degenerate. What does it mean?
(a) All the orbitals have the same orientation.
(b) All the orbitals have the same shape.
(c) All the orbitals have the same energy.
(d) All the orbitals are unoccupied.

132. Which of the following subshell can accommodate as many as 10 electrons?
(a) 2*d* (b) 3*d* (c) $3d_{xy}$ (d) $3d_{z^2}$

133. Which of the following statements is correct for an electron having azimuthal quantum number l 2 ?
(a) The electron may be in the lowest energy shell.
(b) The electron is in a spherical orbital.
(c) The electron must have spin m_s $\frac{1}{2}$
(d) The electron may have a magnetic quantum number 1

134. Which of the following statements is incorrect?
(a) The concepts of "penetration" and "shielding" are important in deciding the energetic ordering of orbitals in multi-electron atoms
(b) A wave-function can have positive and negative values
(c) "Radial nodes" can appear in radial probability distribution functions
(d) The shape of an orbital is given by the principal quantum number.

135. For an $4p_y$ orbital, there are
nodal plane ....and azimuthal quantum number *l*....
(a) 1,0 (b) 0,1 (c) 1,1 (d) 2,1

136. Which of the following statement is correct?
(a) Number of angular nodes n l 1
(b) Number of radial nodes l
(c) Total number of nodes n 1
(d) All of these

137. Give the correct order of initials **T** (True) or **F** (False) for following statements.
(I) If electron has zero quantum magnetic number, then it must be present in *s*-orbital.
(II) In $\boxed{\uparrow\downarrow}$ $\boxed{\uparrow\downarrow}$ $\boxed{\ }$ $\boxed{\uparrow\downarrow}$ orbital diagram, Pauli's exclusion principle is violated.
(III) Bohr's model can explain spectrum of the hydrogen atom.

For More Material Join: @JEEAdvanced_2024

(IV) A *d*-orbital can accommodate maximum 10 electrons only.

(a) TTFF (b) FFTF
(c) TFTT (d) FFTT

138. The orbital diagram in which both the Pauli's exclusion principle and Hund's rule are violated is :

(a) $\boxed{\uparrow\downarrow}\ \boxed{\uparrow\uparrow}\ \boxed{\uparrow}\ \boxed{\ }$ (b) $\boxed{\uparrow\downarrow}\ \boxed{\uparrow\downarrow}\ \boxed{\ }\ \boxed{\uparrow\downarrow}$

(c) $\boxed{\uparrow\uparrow}\ \boxed{\uparrow}\ \boxed{\uparrow}\ \boxed{\uparrow}$ (d) $\boxed{\uparrow\downarrow}\ \boxed{\uparrow\downarrow}\ \boxed{\uparrow\downarrow}\ \boxed{\uparrow}$


139. It is not possible to explain the Pauli's exclusion principle with the help of this atom.


(a) B (b) Be (c) C (d) H


140. The subshell that arises after *f* subshell is called *g* subshell.
What is the total number of orbitals in the shell in which the *g* subshell first occur?


(a) 9 (b) 16 (c) 25 (d) 36

141. The variation of radial probability density $R^2(r)$ as a function of distance *r* of the electron from the nucleus for 3*p*-orbital :

(a) 

(b) 

(c) 

(d) 

142. In iron atom, how many electrons have $n \quad 3$ and $l \quad 2$?

(a) 2 (b) 4 (c) 6 (d) 8

143. For similar orbitals having different values of *n* :

(a) the most probable distance increases with increase in *n*
(b) the most probable distance decreases with increase in *n*
(c) the most probable distance remains constant with increase in *n*
(d) none of these

144. If *n* and *l* are principal and azimuthal quantum numbers respectively, then the expression for calculating the total number of electrons in any energy level is :

(a) $\sum_{l\ 0}^{l\ n} 2(2l \quad 1)$ (b) $\sum_{l\ 1}^{l\ n} 2(2l \quad 1)$ (c) $\sum_{l\ 0}^{l\ n} (2l \quad 1)$ (d) $\sum_{l\ 0}^{l\ n\ 1} 2(2l \quad 1)$

145. Maximum number of nodes are present in :

(a) 5*s* (b) 5*p*
(c) 5*d* (d) All have same number of nodes

146. The possible correct set of quantum numbers for the unpaired electron of Cl atom is :

(a) $2, 0, 0, \ \frac{1}{2}$ (b) $2, 1, \ \ 1, \ \frac{1}{2}$ (c) $3, 1, 1, \ \frac{1}{2}$ (d) $3, 0, \ \frac{1}{2}$

147. The aufbau principle implies that a new electron will enter an orbital for which :


(a) *n* has a lower value (b) *l* has a lower value
(c) $(n \quad l)$ value is maximum (d) $(n \quad l)$ value is minimum

148. The orbital diagram in which aufbau principle is violated, is :

(a) $\boxed{\uparrow\downarrow}\ \boxed{\uparrow\uparrow}\ \boxed{\uparrow}\ \boxed{\uparrow}$ (b) $\boxed{\uparrow}\ \boxed{\uparrow\downarrow}\ \boxed{\uparrow}\ \boxed{\uparrow}$ (c) $\boxed{\uparrow\downarrow}\ \boxed{\uparrow}\ \boxed{\uparrow}\ \boxed{\uparrow}$ (d) $\boxed{\uparrow\downarrow}\ \boxed{\uparrow\downarrow}\ \boxed{\uparrow\downarrow}\ \boxed{\uparrow}$


For More Material Join: @JEEAdvanced_2024

149. Consider the following six electronic configurations (remaining inner orbitals are completely filled) and mark the incorrect option.


I. 

II. 

III. 

IV. 

V. 

VI. 

(a) Stability order : II > I > IV > III
(b) Order of spin multiplicity : IV > III = I > II
(c) V does not violate all the three rules of electronic configuration
(d) If VI represents *A* then *A* when kept near a magnet, acts as paramagnetic substance.

150. Which of the following set of quantum numbers shows orbital of highest energy?

(a) $n \quad 4, l \quad 0, m \quad 0, s \quad \frac{1}{2}$
(b) $n \quad 2, l \quad 0, m \quad 0, s \quad \frac{1}{2}$
(c) $n \quad 3, l \quad 1, m \quad 1, s \quad \frac{1}{2}$
(d) $n \quad 3, l \quad 2, m \quad 1, s \quad \frac{1}{2}$

151. A subshell $n \quad 5, l \quad 3$ can accommodate :

(a) 10 electrons (b) 14 electrons (c) 18 electrons (d) None of these

152. In H-atom energy of electron is determined by :

(a) only *n* (b) *n, l*
(c) *n, l, m* (d) all the four quantum numbers.

153. How many electron(s) in an atom can have $n \quad 3, l \quad 2$?

(a) 1 (b) 2 (c) 5 (d) 10

154. How many electrons in an atom can have $n \quad 4, l \quad 2, m \quad 2$ and $s \quad \frac{1}{2}$?

(a) 1 (b) 2 (c) 5 (d) 10

155. Not considering the electron spin, the degeneracy of first excited state ($n \quad 2$) of H-atom is:

(a) 2 (b) 3 (c) 4 (d) 8

156. Not considering the electron spin, the degeneracy of second excited state of H-atom is 9, while the degeneracy of the first excited state of H is:

(a) 1 (b) 2 (c) 3 (d) 4

157. Which orbital has only positive values of wave function at all distances from the nucleus :

(a) 1*s* (b) 2*s* (c) 2*p* (d) 3*d*

158. Four electrons in an atom have the sets of quantum numbers as given below. Which electron in at the highest energy level?

(a) $n \quad 4, l \quad 0, m_l \quad 0, m_s \quad 1/2$
(b) $n \quad 3, l \quad 0, m_l \quad 0, m_s \quad 1/2$
(c) $n \quad 3, l \quad 2, m_l \quad 0, m_s \quad 1/2$
(d) $n \quad 4, l \quad 1, m_l \quad 1, m_s \quad 1/2$

159. The set of quantum numbers, $n \quad 3, l \quad 2, m_l \quad 0$

(a) describes an electron in a 2*s* orbital
(b) is not allowed
(c) describes an electron in a 3*p* orbital
(d) describes one of the five orbitals of same energy

For More Material Join: @JEEAdvanced_2024

160. The set of quantum numbers, $n \quad 2, l \quad 2, m_l \quad 0$:
(a) describes an electron in a $2s$ orbital
(b) describes one of the five orbitals of a similar type
(c) describes an electron in a $2p$ orbital
(d) is not allowed

161. Consider the argon atom. For how many electrons does this atom have $m_l \quad 1$?
(a) 1 (b) 6 (c) 4 (d) 2

162. An orbital is occupied by an electron with the quantum numbers $n \quad 4, l \quad 1$. How many orbitals of this type are found in a multi-electron atom?
(a) $4p, 3$ (b) $4s, 1$ (c) $4d, 5$ (d) $4p, 6$

163. Which of the following sets of quantum numbers describes the electron which is removed most easily from a potassium atom in its ground state?
(a) $n \quad 3, l \quad 1, m_l \quad 1, m_s \quad \frac{1}{2}$ (b) $n \quad 2, l \quad 1, m_l \quad 0, m_s \quad \frac{1}{2}$
(c) $n \quad 4, l \quad 0, m_l \quad 1, m_s \quad \frac{1}{2}$ (d) $n \quad 4, l \quad 0, m_l \quad 0, m_s \quad \frac{1}{2}$

164. The subshell that arises after f is called the g subshell. How many electrons may occupy the g subshell?
(a) 9 (b) 7 (c) 5 (d) 18

165. Which of the following electron configurations is correct for iron, (atomic number 26)?
(a) $[Kr]\,4s^1 3d^6$ (b) $[Kr]\,4s^1 3d^7$ (c) $[Ar]\,4s^2 3d^6$ (d) $[Kr]\,4s^2 3d^6$

166. Which of the following electron configurations is correct for copper, (atomic number 29)?
(a) $[Ar]\,3d^{10} 4s^1$ (b) $[Kr]\,3d^9 4s^1$ (c) $[Ar]\,3d^9 4s^2$ (d) $[Kr]\,3d^{10} 4s^1$

167. The electronic configurations of $_{24}Cr$ and $_{29}Cu$ are abnormal
(a) due to extra stability of exactly half filled and exactly fully filled sub shells
(b) because they belong to d-block
(c) both the above
(d) None of the above

168. Which of the following representation of excited states of atoms is impossible?
(a) $1s^1 2s^1$ (b) $[Ne]\,3s^2 3p^3 4s^1$ (c) $[Ne]\,3s^2 3p^6 4s^1 3d^6$
(d) $1s^2 2s^2 2p^7 3s^2$

169. Among the following representations of excited states of atoms which is impossible?
(a) $1s^1 2s^1$ (b) $[Ne]\,3s^2 3p^3 4s^1$
(c) $1s^2 2s^2 2p^4 3s^2$ (d) $[Ne]\,3s^2 3p^6 4s^3 3d^2$

170. Among the following series of transition metal ions, the one where all metal ions have same $3d$ electronic configuration is:
(a) Ti^2 , V^3 , Cr^4 , Mn^5 (b) Ti^3 , V^2 , Cr^3 , Mn^4
(c) Ti , V^4 , Cr^6 , Mn^7 (d) Ti^4 , V^3 , Cr^2 , Mn^3


171. Which of the following has the maximum number of unpaired electrons?
(a) Mn (b) Ti (c) V (d) Al

For More Material Join: @JEEAdvanced_2024

172. Which of the following orbitals has two spherical nodes?

(a) 2*s* (b) 4*s* (c) 3*d* (d) 6*f*

173. Wave function of an orbital is plotted against the distance from nucleus. The graphical representation is of:



(a) 1*s* (b) 2*s* (c) 3*s* (d) 2*p*

174. The Schrodinger wave equation for hydrogen atom is

$$\psi_{2s} = \frac{1}{4\sqrt{2}} \left(\frac{1}{a_0}\right)^{3/2} \left(2 - \frac{r}{a_0}\right) e^{-r/a_0}$$

where a_0 is Bohr's radius. If the radial node in 2*s* be at r_0, then r_0 would be equal to :

(a) $\frac{a_0}{2}$ (b) $2a_0$ (c) $\sqrt{2}\, a_0$ (d) $\frac{a_0}{\sqrt{2}}$

175. The Schrodinger wave equation for hydrogen atom is

$$\psi_{(\text{radial})} = \frac{1}{16\sqrt{4}} \left(\frac{Z}{a_0}\right)^{3/2} [(\quad 1)(\quad ^2 \quad 8 \quad 12)] e^{\quad /2}$$

where a_0 and Z are the constant in which answer can be expressed and $\frac{2Zr}{a_0}$

minimum and maximum position of radial nodes from nucleus are .....respectively.

(a) $\frac{a_0}{Z}, \frac{3a_0}{Z}$ (b) $\frac{a_0}{2Z}, \frac{a_0}{Z}$ (c) $\frac{a_0}{2Z}, \frac{3a_0}{Z}$ (d) $\frac{a_0}{2Z}, \frac{4a_0}{Z}$

For More Material Join: @JEEAdvanced_2024



Level 2

1. Potential energy of electron present in He is :

(a) $\frac{e^2}{2\ {}_0 r}$ (b) $\frac{3e^2}{4\ {}_0 r}$ (c) $\frac{2e^2}{4\ {}_0 r}$ (d) $\frac{e^2}{4\ {}_0 r^2}$

2. A single electron in an ion has ionization energy equal to 217.6 eV. What is the total number of neutrons present in one ion of it?

(a) 2 (b) 4 (c) 5 (d) 9

3. For a hypothetical hydrogen like atom, the potential energy of the system is given by $U(r)$ $\frac{Ke^2}{r^3}$, where r is the distance between the two particles. If Bohr's model of quantization of angular momentum is applicable then velocity of particle is given by :

(a) v $\frac{n^2h^3}{Ke^2 8\ {}^3m^2}$ (b) v $\frac{n^3h^3}{8Ke^2\ {}^3m^2}$ (c) v $\frac{n^3h^3}{24Ke^2\ {}^3m^2}$ (d) v $\frac{n^2h^3}{24Ke^2\ {}^3m^2}$

4. A small particle of mass m moves in such a way that P.E. $\frac{1}{2}mkr^2$, where k is a constant and r is the distance of the particle from origin. Assuming Bohr's model of quantization of angular momentum and circular orbit, r is directly proportional to :

(a) n^2 (b) n (c) $\sqrt{n}$ (d) none of these

5. A beam of specific kind of particles of velocity 2.1 10^7 m/s is scattered by a gold (Z 79) nuclei. Find out specific charge (charge/mass) of this particle if the distance of closest approach is 2.5 10^{14} m.

(a) 4.84 10^7 C/kg (b) 4.84 10^7 C/kg (c) 2.42 10^7 C/kg (d) 3 10^{12} C/kg

6. What is the angular velocity ( ) of an electron occupying second orbit of Li^2 ion?

(a) $\frac{8\ {}^3me^4}{h^3}K^2$ (b) $\frac{8\ {}^3me^4}{9h^3}K^2$ (c) $\frac{64}{9}\ \frac{{}^3me^4}{h^3}K^2$ (d) $\frac{9\ {}^3me^4}{h^3}K^2$

7. The ratio of the radius difference between 4th and 3rd orbit of H-atom and that of Li^2 ion is :

(a) 1 : 1 (b) 3 : 1 (c) 3 : 4 (d) 9 : 1

8. The velocity of an electron in excited state of H-atom is 1.093 10^6 m/s. What is the circumference of this orbit?

(a) 3.32 10^{10} m (b) 6.64 10^{10} m (c) 13.30 10^{10} m (d) 13.28 10^8 m

9. The angular momentum of an electron in a Bohr's orbit of He is 3.1652 10^{34} kg-m^2/sec. What is the wave number in terms of Rydberg constant (R) of the spectral line emitted when an electron falls from this level to the first excited state. [Use h 6.626 10^{34} J s]

(a) $3R$ (b) $\frac{5R}{9}$ (c) $\frac{3R}{4}$ (d) $\frac{8R}{9}$

10. If radiation corresponding to second line of "Balmer series" of Li^2 ion, knocked out electron from first excited state of H-atom, then kinetic energy of ejected electron would be:

(a) 2.55 eV (b) 4.25 eV (c) 11.25 eV (d) 19.55 eV

For More Material Join: @JEEAdvanced_2024

11. When an electron makes a transition from (n 1) state to nth state, the frequency of emitted radiations is related to n according to (n 1) :

(a) ν $\frac{2cRZ^2}{n^3}$ (b) ν $\frac{cRZ^2}{n^4}$

(c) ν $\frac{cRZ^2}{n^2}$ (d) ν $\frac{2cRZ^2}{n^2}$

12. In a collection of H-atoms, all the electrons jump from n 5 to ground level finally (directly or indirectly), without emitting any line in Balmer series. The number of possible different radiations is :

(a) 10 (b) 8 (c) 7 (d) 6

13. An electron is allowed to move freely in a closed cubic box of length of side 10 cm. The uncertainty in its velocity will be :

(a) 3.35 10^{4} m sec ^{1} (b) 5.8 10^{4} m sec ^{1}

(c) 4 10^{5} m sec ^{1} (d) 4 10^{6} m sec ^{1}

14. An element undergoes a reaction as shown :

X $2e$ X^2 , energy released 30.87 eV/atom. If the energy released, is used to dissociate 4 g of H_2 molecules, equally into H and H*, where H* is excited state of H atoms where the electron travels in orbit whose circumference equal to four times its de Broglie's wavelength. Determine the least moles of X that would be required :

Given : I.E. of H 13.6 eV/atom, bond energy of H_2 4.526 eV/molecule.

(a) 1 (b) 2 (c) 3 (d) 4

15. If the energy of H-atom in the ground state is E, the velocity of photo-electron emitted when a photon having energy E_p strikes a stationary Li^{2} ion in ground state, is given by :

(a) v $\sqrt{\frac{2(E_p \quad E)}{m}}$ (b) v $\sqrt{\frac{2(E_p \quad 9E)}{m}}$

(c) v $\sqrt{\frac{2(E_p \quad 9E)}{m}}$ (d) v $\sqrt{\frac{2(E_p \quad 3E)}{m}}$

16. At which temperature will the translational kinetic energy of H-atom equal to that for H-atom of first line Lyman transition? (Given N_A 6 10^{23})

(a) 780 K (b) 1.32 10^{5} K

(c) 7.84 10^{4} K (d) 1000 K

17. For a 3s-orbital, value of is given by following relation :

$$(3s) \quad \frac{1}{9\sqrt{3}} \quad \frac{1}{a_0}^{3/2} \quad (6 \quad 6 \quad ^2) e^{\ /2}; \quad \text{where} \quad \frac{2rZ}{3a_0}$$

What is the maximum radial distance of node from nucleus ?

(a) $\frac{(3 \quad \sqrt{3})\, a_0}{Z}$ (b) $\frac{a_0}{Z}$

(c) $\frac{3}{2}\frac{(3 \quad \sqrt{3})\, a_0}{Z}$ (d) $\frac{2a_0}{Z}$

For More Material Join: @JEEAdvanced_2024

18. Monochromatic radiation of specific wavelength is incident on H-atoms in ground state. H-atoms absorb energy and emit subsequently radiations of six different wavelength. Find wavelength of incident radiations:

(a) 9.75 nm (b) 50 nm

(c) 85.8 nm (d) 97.25 nm

19. The energy of a I, II and III energy levels of a certain atom are E, $\frac{4E}{3}$ and $2E$ respectively. A photon of wavelength is emitted during a transition from III to I. What will be the wavelength of emission for transition II to I?

(a) $\frac{}{2}$ (b) (c) 2 (d) 3

20. Calculate the minimum and maximum number of electrons which may have magnetic quantum number, m 1 and spin quantum number, s $\frac{1}{2}$ in chromium (Cr) :

(a) 0, 1 (b) 1, 2 (c) 4, 6 (d) 2, 3

21. An electron in a hydrogen atom in its ground state absorbs 1.5 times as much energy as the minimum required for it to escape from the atom. What is the velocity of the emitted electron?

(a) 1.54 10^6 m/s (b) 1.54 10^8 m/s

(c) 1.54 10^3 m/s (d) 1.54 10^4 m/s

22. In a measurement of quantum efficiency of photosynthesis in green plants, it was found that 10 quanta of red light of wavelength 6850 Å were needed to release one molecule of O_2. The average energy storage in this process for 1 mole O_2 evolved is 112 kcal.

What is the energy conversion efficiency in this experiment?

Given : 1 cal 4.18 J; N_A 6 10^{23}; h 6.63 10^{-34} J.s

(a) 23.5 (b) 26.9

(c) 66.34 (d) 73.1

23. A hydrogen like species (atomic number Z) is present in a higher excited state of quantum number n. This excited atom can make a transition to the first excited state by successive emission of two photons of energies 10.20 eV and 17.0 eV respectively. Alternatively, the atom from the same excited state can make a transition to the second excited state by successive emission of two photons of energy 4.25 eV and 5.95 eV respectively. Determine the value of Z.

(a) 1 (b) 2 (c) 3 (d) 4

24. H-atom is exposed to electromagnetic radiation of 1025.6 Å and excited atom gives out induced radiations. What is the minimum wavelength of these induced radiations?

(a) 102.6 nm (b) 12.09 nm

(c) 121.6 nm (d) 810.8 nm

25. If the lowest energy X-rays have 3.055 10^{-8} m, estimate the minimum difference in energy between two Bohr's orbits such that an electronic transition would correspond to the emission of an X-ray. Assuming that the electrons in other shells exert no influence, at what Z (minimum) would a transition from the second energy level to the first result in the emission of an X-ray?

(a) 1 (b) 2

(c) 3 (d) 4


For More Material Join: @JEEAdvanced_2024

26. An -particle having kinetic energy 5 MeV falls on a Cu-foil. The shortest distance from the nucleus of Cu to which -particle reaches is (Atomic no. of Cu 29, K 9 $10^9 Nm^2/C^2$)

(a) 2.35 10^{13}m (b) 1.67 10^{14}m

(c) 5.98 10^{15}m (d) None of these

27. In the graph between $\sqrt{\nu}$ and Z for the Mosley's equation $\sqrt{\nu}$ $a(Z$ $b)$, the intercept OX is –1 on $\sqrt{\nu}$ axis.



What is the frequency when atomic number (Z) is 51 ?

(a) 50 s^{1} (b) 100 s^{1} (c) 2500 s^{1} (d) None of these

28. Balmer gave an equation for wavelength of visible region of H-spectrum as $\frac{Kn^2}{n^2 \quad 4}$.

Where n principal quantum number of energy level, K constant in terms of R (Rydberg constant).

The value of K in terms of R is:

(a) R (b) $\frac{R}{2}$ (c) $\frac{4}{R}$ (d) $\frac{5}{R}$

29. The energy of separation of an electron in a hydrogen like atom in excited state is 3.4 eV. The de-Broglie wave length (in Å) associated with the electron is:

(Given radius of first orbit of H-atom is 0.53Å)

(a) 3.33 (b) 6.66 (c) 13.31 (d) None of these

30. If I excitation energy for the H-like (hypothetical) sample is 24 eV, then binding energy in III excited state is:

(a) 2 eV (b) 3 eV (c) 4 eV (d) 5 eV

For More Material Join: @JEEAdvanced_2024



Level 3



Passage 1

Werner Heisenberg considered the limits of how precisely we can measure the properties of an electron or other microscopic particle. He determined that there is a fundamental limit to how closely we can measure both position and momentum. The more accurately we measure the momentum of a particle, the less accurately we can determine its position. The converse is also true. This is summed up in what we now call the Heisenberg uncertainty principle.

The equation is $x \quad (mv) \quad \frac{h}{4}$

The uncertainty in the position or in the momentum of a macroscopic object like a baseball is too small to observe. However, the mass of microscopic object such as an electron is small enough for the uncertainty to be relatively large and significant.

1. If the uncertainties in position and momentum are equal, the uncertainty in the velocity is :

(a) $\sqrt{h}$ (b) $\sqrt{\frac{h}{2}}$ (c) $\frac{1}{2m}\sqrt{h}$ (d) none of these

2. If the uncertainty in velocity and position is same, then the uncertainty in momentum will be :

(a) $\sqrt{\frac{hm}{4}}$ (b) $m\sqrt{\frac{h}{4}}$ (c) $\sqrt{\frac{h}{4\ m}}$ (d) $\frac{1}{m}\sqrt{\frac{h}{4}}$

3. What would be the minimum uncertainty in de-Broglie wavelength of a moving electron accelerated by potential difference of 6 volt and whose uncertainty in position is $\frac{7}{22}$ nm?

(a) 6.25 Å (b) 6 Å (c) 0.625 Å (d) 0.3125 Å



Passage 2

One of the fundamental laws of physics is that matter is most stable with the lowest possible energy. Thus, the electron in a hydrogen atom usually moves in the $n \quad 1$ orbit, the orbit in which it has the lowest energy. When the electron is in this lowest energy orbit, the atom is said to be in its ground electronic state. If the atom receives energy from an outside source, it is possible for the electron to move to an orbit with a higher n value, in which case the atoms is in an excited state with a higher energy.

The law of conservation of energy says that we cannot create or destroy energy. Thus, if a certain amount of external energy is required to excite an electron from one energy level to another, then that same amount of energy will be liberated when the electron returns to its initial state.

For More Material Join: @JEEAdvanced_2024

Lyman series is observed when the electron returns to the lowest orbit while Balmer series is formed when the electron returns to second orbit. Similarly, Paschen, Brackett and Pfund series are formed when electrons returns to the third, fourth and fifth orbits from higher energy orbits respectively.

When electrons return from n_2 to n_1 state, the number of lines in the spectrum will equal to

$$\frac{(n_2 \quad n_1)(n_2 \quad n_1 \quad 1)}{2}$$

If the electron comes back from energy level having energy E_2 to energy level having energy E_1, then the difference may be expressed in terms of energy of photon as :

$$E_2 \quad E_1 \quad E, \quad E \quad \frac{hc}{}$$

Since, h and c are constants, E corresponds to definite energy. Thus, each transition from one energy level to another will produce a radiation of definite wavelength. This is actually observed as a line in the spectrum of hydrogen atom.

Wave number of a spectral line is given by the formula

$$- \quad R \quad \frac{1}{n_1^2} \quad \frac{1}{n_2^2} .$$

where R is a Rydberg's constant ($R \quad 1.1 \quad 10^7$ m ^{1})

1. If the wavelength of series limit of Lyman series for He ion is x Å, then what will be the wavelength of series limit of Balmer series for Li^2 ion?

(a) $\frac{9x}{4}$ Å (b) $\frac{16x}{9}$ Å (c) $\frac{5x}{4}$ Å (d) $\frac{4x}{7}$ Å

2. The emission spectra is observed by the consequence of transition of electrons from higher energy state to ground state of He ion. Six different photons are observed during the emission spectra, then what will be the minimum wavelength during the transition?

(a) $\frac{4}{27R_H}$ (b) $\frac{4}{15R_H}$ (c) $\frac{15}{16R_H}$ (d) $\frac{16}{15R_H}$

3. What transition in the hydrogen spectrum would have the same wavelength as Balmer transition, $n \quad 4$ to $n \quad 2$ in the He spectrum?

(a) $n \quad 3$ to $n \quad 1$ (b) $n \quad 3$ to $n \quad 2$ (c) $n \quad 4$ to $n \quad 1$ (d) $n \quad 2$ to $n \quad 1$

4. An electron in H-atom in M-shell on de-excitation to ground state gives maximum ..... spectrum lines.

(a) 10 (b) 6 (c) 2 (d) 1

For More Material Join: @JEEAdvanced_2024



Passage 3

If hydrogen atoms (in the ground state) are passed through an homogeneous magnetic field, the beam is split into two parts. This interaction with the magnetic field shows that the atoms must have magnetic moment. However, the moment cannot be due to the orbital angular momentum since l 0. Hence one must assume existence of intrinsic angular momentum, which as the experiment shows, has only two permitted orientations.

Spin of the electron produce angular momentum equal to S $\sqrt{s(s\ 1)}\,\frac{h}{2}$ where S $\frac{1}{2}$.

Total spin of an atom $\frac{n}{2}$ or $\frac{n}{2}$

where n is the number of unpaired electrons.

The substance which contain species with unpaired electrons in their orbitals behave as paramagnetic substances. The paramagnetism is expressed in terms of magnetic moment.

The magnetic moment of an atom

$$_s\sqrt{s(s\ 1)}\,\frac{eh}{2\ mc}\quad\sqrt{\frac{n}{2}\ \frac{n}{2}\ 1}\,\frac{eh}{2\ mc}\qquad s\ \frac{n}{2}$$

$$_s\quad\sqrt{n(n\ 2)}\ \text{B.M.}$$

n number of unpaired electrons

1. B.M. (Bohr magneton) $\frac{eh}{4\ mc}$

If magnetic moment is zero the substances is diamagnetic.

1. Which of the following ion has lowest magnetic moment ?
(a) Fe^{2} (b) Mn^{2} (c) Cr^{3} (d) V^{3}

2. If an ion of $_{25}Mn$ has a magnetic moment of 3.873 B.M. Then oxidation state of Mn in ion is :
(a) 2 (b) 3 (c) 4 (d) 5



Passage 4

Ozone in the upper atmosphere absorbs ultraviolet radiation which induces the following chemical reaction

$$O_3(g)\qquad O_2(g)\quad O(g)$$

O_2produced in the above photochemical dissociation undergoes further dissociation into one normal oxygen atom (O) and more energetic oxygen atom O*.

$$O_2(g)\qquad O\quad O^{*}$$

(O*) has 1 eV more energy than (O) and normal dissociation energy of O_2 is 480 kJ mol ^{1}.

[1 eV/ Photon 96 kJ mol ^{1}]

For More Material Join: @JEEAdvanced_2024


1. What is the maximum wavelength effective for the photochemical dissociation of O_2 molecule
(a) 2440 Å (b) 2066.67 Å (c) 1000 Å (d) 155 Å

2. If dissociation of O_3 into O_2 and O requires 400kJ mol ^{1} and O_2 produced in this reaction is further dissociated to O and O* then the total energy required to for the dissociation of O_3 into O and O* is:
(a) 1168kJ/mol (b) 976kJ/mol (c) 880kJ/mol (d) None of these




Passage 5

The existence of negatively charged particle in an atom was shown by J.J. Thomson as a result of the studies of the passage of electricity through gases at extremely low pressures known as discharge tube experiments. When a high voltage of the order of 10,000 volts or more was impressed across the electrodes, some sort of invisible rays moved from the negative electrode to the positive electrode these rays are called as cathode rays.



Cathode rays experiment



Anode rays experiment

Cathode rays travel in straight path in absence of electrical and magnetic field. Cathode rays consist of material part and charged particles? Cathode rays produce **X-rays** and light is emitted when they strike on ZnS screen. Cathode rays penetrate through thin sheets of aluminium and other metals. They affect the photographic plate and passes heating effect when they strike on metal foil. The ratio of charge to mass *i.e.* charge/mass is same for all the cathode rays irrespective of the gas used in the tube.

The existence of positively charged particles in an atom was shown by E. Goldstein. He repeated the same discharge tube experiments by using a perforated cathode. It was observed that when a high potential difference was applied between the electrodes, not only cathode rays were produced but also a new type of rays were produced simultaneously from anode moving towards cathode and passed through the holes or canal of the cathode. These termed as canal rays or anode rays.

These rays travel in straight lines and consist of positively charged particles. These rays have kinetic energy and produces heating effect also. The *e/m* ratio of these rays is smaller than that of electrons. Unlike cathode rays, their *e/m* value is is dependent upon the nature of the gas taken in the tube. These rays produce flashes of light on ZnS screen and can pass throughs thin metal foils. They can produce physical and chemical changes and are capable to produce ionisation in gases.

For More Material Join: @JEEAdvanced_2024

1. For cathode rays the value of e/m:
(a) is independent of the nature of the cathode and the gas filled in the discharge tube
(b) is constant
(c) is -1.7588×10^8 coulombs/g
(d) all of the above are correct

2. Which is not true with respect to cathode rays ?
(a) A stream of electrons (b) Charged particles
(c) Move with same speed as that of light (d) Can be deflected by the electric field

3. Select the incorrect statement.
(a) Cathode rays has charge only and no mass
(b) Anode rays are deflected by electrical and magnetic field
(c) Canal rays is named for beam of positive charged particle
(d) Anode rays particles carrying positive charge

One or More Answers is/are Correct



1. Select the correct statement(s) :
(a) The phenomena of diffraction of light can only be explained by assuming that light behaves as waves
(b) de-Broglie postulate the dual character existed with matter
(c) In his atomic model Bohr considered electron as a particle
(d) Wave nature of electrons was proved when diffraction rings were observed photographically when a stream of protons was passed through a metal foil

2. The angular momentum of electron can have the value(s) :
(a) $0.5\frac{h}{}$ (b) $\frac{h}{}$ (c) $\frac{h}{0.5}$ (d) $2.5\frac{h}{2}$

3. Select incorrect statement(s) :
(a) Only three quantum numbers n, l and m are needed to define an orbital
(b) Four quantum numbers are needed for complete description of an electron
(c) Two quantum numbers n and l are needed to identify subshell and shape of orbital
(d) Splitting of spectrum lines in presence of electric field is known as Zeeman effect

4. Select the correct statement(s):
(a) An electron near the nucleus is attracted by the nucleus and has a low potential energy
(b) According to Bohr's theory, an electron continuously radiate energy if it stays in one orbit
(c) Bohr's model could not explain the spectra of multielectron atoms
(d) Bohr's model was the first atomic model based on quantisation of energy

5. Choose the correct statement(s) :
(a) The shape of an atomic orbital depends upon azimuthal quantum number
(b) The orientation of an atomic orbital depends upon the magnetic quantum number
(c) The energy of an electron in an atomic orbital of multi-electron atom depends upon principal quantum number only
(d) The number of degenerate atomic orbitals of one type depends upon the value of azimuthal quantum number

For More Material Join: @JEEAdvanced_2024

6. For radial probability curves, which of the following is/are correct?
(a) The number of maxima in 2*s* orbital are two
(b) The number of spherical or radial nodes is equal to $n \quad l \quad 1$
(c) The number of angular nodes are '*l*'
(d) $3d_z^2$ has 3 angular nodes

7. Select the correct statement(s):
(a) Lower value of quantum number *l* indicates that there is a higher probability of finding the 3*s* electron close to the nucleus than those of 3*p* and 3*d* electrons
(b) Energy of 3*s* orbital is less than for the 3*p* and 3*d* orbitals
(c) At the node, the value of the radial function changes from positive to negative
(d) The radial function depends upon the quantum numbers *n* and *l*

8. Choose the incorrect statement(s):
(a) For a particular orbital in hydrogen atom, the wave function may have negative value
(b) Radial probability distribution function may have zero value but can never have negative value
(c) $3d_{x^2 \quad y^2}$ orbital has two angular nodes and one radial node
(d) *yz* and *xz* planes are nodal planes for d_{xy} orbital

9. Select the correct statement(s) :
(a) Heisenberg's principle is applicable to stationary electron
(b) Pauli's exclusion principle is not applicable to photons
(c) For an electron the product of velocity and principal quantum number will be independent to principal quantum number
(d) Quantum numbers *l* and *m* determine the value of angular wave function

10. Choose the correct statements among the following :
(a) A node is a point in space where the wave-function has zero amplitude
(b) The number of maxima (peaks) in radial probability distribution function is $(n \quad l)$
(c) Radial probability density is $4 \quad r^2 R^2_{n,\,l}(r)$
(d) ^{2} represents probability of finding electron

11. Select the correct statement(s) regarding $3p_y$ orbital :
(a) Total number of nodes are 2
(b) Number of maxima in the curve $4 \quad r^2 R^2$ vs *r* are two
(c) Quantum number *n, l* and *m* for an orbital may be 3, 1, –1 respectively
(d) The magnetic quantum number may have a positive value

12. Select the correct statement(s) :
(a) In wave mechanical model, the energy of electron in the orbital remains constant
(b) d_{xy} orbital is lies in *yz* plane
(c) Nodal planes are *yz* and *xy* in $d_{x^2 \quad y^2}$ orbital
(d) Rest mass of photon is zero and increases with its velocity

13. Hydrogen has :
(a) half filled subshell
(b) half filled shell
(c) one electron in valence shell
(d) half filled orbital

For More Material Join: @JEEAdvanced_2024

14. Select incorrect statement(s) .
(a) If the value of l 0, the electron distribution is spherical
(b) The shape of the orbital is given by magnetic quantum number
(c) Angular momentum of $1s, 2s, 3s$ orbit electrons are equal
(d) In an atom, all the electrons travel with the same velocity


15. The radial distribution functions $[P(r)]$ is used to determine the most probable radius, which is used to find the electron in a given orbital. $\frac{dP(r)}{dr}$ for $1s$-orbital of hydrogen like atom having atomic number Z, is $\frac{dP}{dr}$ $\frac{4Z^3}{a_0^3}$ $2r$ $\frac{2Zr^2}{a_0}$ e^{-2Zr/a_0} :

Then which of the following statements is/are correct?
(a) At the point of maximum value of radial distribution function $\frac{dP(r)}{dr}$ 0 ; one antinode is present
(b) Most probable radius of Li^{2} is $\frac{a_0}{3}$ pm
(c) Most probable radius of He is $\frac{a_0}{2}$ pm
(d) Most probable radius of hydrogen atom is a_0 pm

16. Select the correct statement(s) :
(a) An orbital with l 0 is symmetrical about the nucleus
(b) An orbital with l 1 is spherically symmetrical about the nucleus
(c) $3d_{z^2}$ is spherically symmetrical about the z-axis
(d) All are correct


17. Select the correct statement(s) :
(a) Radial function $[R(r)]$ is a part of wave function which depends upon quantum number n only
(b) Angular function depends only on the direction, and is independent to the distance from the nucleus
(c) $^{2}(r, ,)$ is the probability density of finding the electron at a particular point in space
(d) Radial distribution function $(4 r^2R^2)$ gives the probability of the electron being present at a distance r from the nucleus


18. Which is/are correct graph?





19. Select the correct curve(s) :
If v Velocity of electron in Bohr's orbit
r Radius of electron in Bohr's orbit
P.E. Potential energy of electron in Bohr's orbit
K.E. Kinetic energy of electron in Bohr's orbit

For More Material Join: @JEEAdvanced_2024

(a) 

(b) 

(c) 

(d) 

20. Select the correct set (s) of quantum numbers
(a) n 3, l 0, m_l 1 (b) n 3, l 3, m_l 2
(c) n 3, l 2, m_l 2 (d) n 3, l 1, m_l 0

21. Which is/are correct statement?
(a) Number of subshell present in M-shell 3
(b) Number of orbitals present in N-shell 16
(c) Cu (z = 29) is paramagnetic
(d) Zeeman effect explains splitting of spectral lines in magnetic field.

22. In a sample of H-atoms electrons are de-excited from 4th excited state to ground state. Which is/are correct statement?
(a) No line observed in P-fund series.
(b) Total ten lines observed in spectrum.
(c) 4 line in UV-region and 3 line in visible region observed.
(d) One line observed in Brackett series.

Match the Column

Column-I and Column-II contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

1.

Column-I	Column-II
(A) Electron	(P) Negative charge
(B) Proton	(Q) Positive charge
(C) Neutron	(R) $1.6 \quad 10^{19}$ C
(D) Positron	(S) Chargeless

2.

Column-I	Column-II
(A) Thomson model of atom	(P) Electrons are present in extra nuclear region
(B) Rutherford model of atom	(Q) Electron in the atom is described as wave
(C) Bohr model of atom	(R) Positive charge is accumulated in the nucleus
(D) Schrodinger model of hydrogen atom	(S) Uniform sphere of positive charge with embedded electrons

For More Material Join: @JEEAdvanced_2024

3.

Column-I	Column-II
(A) Atomic theory of matter	(P) Rutherford scattering experiment
(B) Quantization of charge	(Q) Muliken's oil drop experiment
(C) Quantization of electronic energy level	(R) Atomic spectra
(D) Size of nucleus	(S) Law of multiple proportions

4.

Column-I	Column-II
(A) $\frac{\text{K.E.}}{\text{P.E.}}$	(P) 2
(B) P.E. 2 K.E.	(Q) $\frac{1}{2}$
(C) $\frac{\text{P.E.}}{\text{T.E.}}$	(R) –1
(D) $\frac{\text{K.E.}}{\text{T.E.}}$	(S) 0

5.

Column-I	Column-II
(A) Lyman series	(P) Visible region
(B) Humphery series	(Q) Ultraviolet region
(C) Paschen series	(R) Infrared region
(D) Balmer series	(S) Far infrared region

6. In case of hydrogen spectrum wave number is given by

$$- \quad R_H \quad \frac{1}{n_1^2} \quad \frac{1}{n_2^2} \quad \text{where} \quad n_1 \quad n_2$$

Column-I	Column-II
(A) Lyman series	(P) n_2 2
(B) Balmer series	(Q) n_2 3
(C) Pfund series	(R) n_2 6
(D) Brackett series	(S) n_2 5

7.

Column-I (Shell)	Column-II (Value of *l*)
(A) 2nd	(P) 1
(B) 3rd	(Q) 2
(C) 4th	(R) 3
(D) 1st	(S) 0

For More Material Join: @JEEAdvanced_2024

8. In Bohr's model, for aunielectronic atom, following symbols are used

$r_{n,Z}$ Radius of nth orbit with atomic number Z;

$U_{n,Z}$ Potential energy of electron ; $K_{n,Z}$ Kinetic energy of electron ;

$V_{n,Z}$ Velocity of electron ; $T_{n,Z}$ Time period of revolution

Column-I	Column-II
(A) $U_{1,2} : K_{1,1}$	(P) 1 : 8
(B) $r_{2,1} : r_{1,2}$	(Q) –8 : 1
(C) $V_{1,3} : V_{3,1}$	(R) 9 : 1
(D) $T_{1,2} : T_{2,2}$	(S) 8 : 1

9.

Column-I	Column-II
(A) The radial node of 5*s* atomic orbital is	(P) 1
(B) The angular node of $3d_{yz}$ atomic orbital is	(Q) 4
(C) The sum of angular node and radial node of $4d_{xy}$ atomic orbital	(R) 2
(D) The angular node of 3*p* atomic orbital is	(S) 3

10.

Column-I	Column-II
(A) The *d*-orbital which has two angular nodes	(P) $3d_{x^2\ y^2}$
(B) The *d*-orbital with two nodal surfaces form cones	(Q) $3d_{z^2}$
(C) The orbital without angular node	(R) $4f$
(D) The orbital which has three angular nodes	(S) $3s$

11.

Column-I	Column-II
(A) Orbital angular momentum of an electron	(P) $\sqrt{s(s\ \ 1)}\ \frac{h}{2}$
(B) Angular momentum of an electron in an orbit	(Q) $\sqrt{n(n\ \ 2)}$
(C) Spin angular momentum of an electron	(R) $\frac{nh}{2}$
(D) Magnetic moment of atom	(S) $\sqrt{l(l\ \ 1)}\ \frac{h}{2}$

For More Material Join: @JEEAdvanced_2024



12.

Column-I	Column-II
(A) Number of orbitals in the nth shell	(P) $2(2l \quad 1)$
(B) Maximum number of electrons in a subshell	(Q) n
(C) Number of subshells in n^{th} shell	(R) $2l \quad 1$
(D) Number of orbitals in a subshell	(S) n^2


13.

Column-I	Column-II
(A) 2s	(P) $n \quad 4, l \quad 2, m \quad 0$
(B) $2p_z$	(Q) $n \quad 4, l \quad 2, m \quad \quad 2$ or 2
(C) $4d_{x^2 \quad y^2}$	(R) $n \quad 2, l \quad 1, m \quad 0$
(D) $4d_{z^2}$	(S) $n \quad 2, l \quad 0, m \quad 0$

14.

Column-I	Column-II
(A) 	(P) $4s$
(B) 	(Q) $5p_y$
(C) Angular probability depends upon and	(R) $3s$
(D) Atleast one angular node is present	(S) $6d_{xy}$

15.

Column-I	Column-II
(A) $3s$	(P) 


For More Material Join: @JEEAdvanced_2024

(B) $3p$


(C) $3d$

(D) $2p$


(Q)



(R)



(S)



Assertion-Reason Type Questions

Each question contains STATEMENT-1 (Assertion) and STATEMENT-2 (Reason).
Examine the statements carefully and mark the correct answer according to the instructions given below :

(A) If both the statement are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1

(B) If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1

(C) If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

(D) If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

1. STATEMENT-1 : The angular momentum of d-orbitals is $\sqrt{6}\,\frac{h}{2}$.

STATEMENT-2 : Angular momentum of electron in orbit is mvr $\frac{nh}{2}$.

2. STATEMENT-1 : Angular momentum of the electron in the orbit which has four subshell is $\frac{2h}{}$.

STATEMENT-2 : Angular momentum of electron is quantized.

3. STATEMENT-1 : Line emission spectra useful in the study of atomic structure.

STATEMENT-2 : Each element has a unique line emission spectrum.

4. STATEMENT-1 : Emitted radiation will fall in visible range when an electron jump from n 4 to n 2 in H-atom.

STATEMENT-2 : Balmer series radiations belong to visible range for hydrogen atom only.

For More Material Join: @JEEAdvanced_2024

5. STATEMENT-1 : Half-filled and fully-filled degenerate orbitals are more stable.
STATEMENT-2 : Extra stability is due to the symmetrical distribution of electrons and exchange energy.

6. STATEMENT-1 : The ground state configuration of Cr is $3d^5 4s^1$.
STATEMENT-2 : A set of half-filled orbitals containing one electrons each with their spin parallel provides extra stability.

7. STATEMENT-1 : The ground state electronic configuration of nitrogen is



STATEMENT-2 : Electrons are filled in orbitals as per aufbau principle, Hund's rule of maximum spin multiplicity and Pauli's principle.

8. STATEMENT-1 : An orbital cannot have more than two electrons and they must have opposite spins.
STATEMENT-2 : No two electrons in an atom can have same set of all the four quantum numbers as per Pauli's exclusion principle.

9. STATEMENT-1 : Orbital having xz plane as node may be $3d_{xy}$.
STATEMENT-2 : $3d_{xy}$ has zero radial node.

10. STATEMENT-1 : The kinetic energy of photo-electrons increases with increase in frequency of incident light where $v \quad v_0$.
STATEMENT-2 : Whenever intensity of light is increased the number of photo-electron ejected always increases.

11. STATEMENT-1 : Cu^{2} is a coloured ion.
STATEMENT-2 : Every ion with unpaired electron is coloured.

12. STATEMENT-1 : For $n \quad 3$, l may be 0,1 and 2 and m may be 0; 0 1 ; 0 1 and 2
STATEMENT-2 : For each value of n, there are 0 to $(n \quad 1)$ possible values of l; and for each value of l, there are 0 to l values of m.

Subjective Problems



1. Given $r_{n\ 1} \quad r_{n\ 1} \quad 2r_n$, where r_n, $r_{n\ 1}$, $r_{n\ 1}$ are Bohr radius for hydrogen atom in n^{th}, $(n \quad 1)^{th}$ and $(n \quad 1)^{th}$ shell respectively. Calculate the value of n.

2. The energy of separation of an electron is 30.6 eV moving in an orbit of Li^{2}. Find out the number of waves made by the electron in one complete revolution in the orbit.

3. Calculate the number of waves made by a Bohr electron in one complete revolution in n^{th} orbit of H-atom, if ratio of de-Broglie wavelength associated with electron moving in n^{th} orbit and 2nd orbit is 1.5.

4. A certain dye absorbs lights of 400 nm and then fluorescence light of wavelength 500 nm. Assuming that under given condition 40% of the absorbed energy is re-emitted as fluorescence, calculate the ratio of quanta absorbed to number of quanta emitted out.

5. A photon of energy 4.5 eV strikes on a metal surface of work function 3.0 eV. If uncertainty in position is $\frac{25}{4}$ Å, find the uncertainty in measurement of de-Broglie wavelength (in Å).

For More Material Join: @JEEAdvanced_2024

6. Find out the difference in number of angular nodes and number of radial nodes in the orbital to which last electron of chromium present.

7. What is the total number of radial and angular nodes present in 5*f* orbital ?

8. Infrared lamps are used in restaurants to keep the food warm. The infrared radiation is strongly absorbed by water, raising its temperature and that of the food. If the wavelength of infrared radiation is assumed to be 1500 nm, and the number of quanta of infrared radiation produced per second by an infrared lamp (that consumes energy at the rate of 100 W and is 12% efficient only) is $(x \quad 10^{19})$, then the value of x is :

(Given : $h \quad 6.625 \quad 10^{\ 34}$ J-s)

9. When an electron makes transition from $(n \quad 1)$ state to n state the wavelength of emitted radiations is related to $n(n \quad 1)$ according to $\quad n^{x}$.

What is the value of x ?

10. For 3*s* orbital of hydrogen atom, the normalised wave function is

$$_{3s} \quad \frac{1}{81\sqrt{3}} \quad \frac{1}{a_0}^{3/2} \quad 27 \quad \frac{18r}{a_0} \quad \frac{2r^2}{a_0^2} \quad e^{\frac{r}{3a_0}}$$

If distance between the radial nodes is d, calculate the value of $\frac{d}{1.73\, a_0}$.

11. Find the separation between two electrons (in Å) in vacuum, if electrostatic potential energy between these electrons is $7.67 \quad 10^{\ 19}$ J.

[Given: $e \quad 1.6 \quad 10^{\ 19}$ C ; $_0 \quad 8.85 \quad 10^{\ 12}$ J $^{\ 1}$ C^{2}m $^{\ 1}$, $\quad 3.14$]

12. An - particle moving with velocity $\frac{1}{30}$ th times of velocity of light. If uncertainty in position is $\frac{3.31}{}$ pm, then minimum uncertainty in kinetic energy is $y \quad 10^{\ 16}$ J. Calculate the value of y.

13. In a sample of excited hydrogen atoms electrons make transition from $n \quad 2$ to $n \quad 1$. Emitted quanta strike on a metal of work function ( ) 4.2eV.

Calculate the wavelength (in Å) associated with ejected electrons having maximum kinetic energy.

14. For 1s orbital of Hydrogen atom radial wave function is given as:

$$R(r) \quad \frac{1}{\sqrt{}} \quad \frac{1}{a_0}^{3/2} \quad e^{\ r/a_0} \qquad \text{(where } a_0 \quad 0.529\text{Å)}$$

The ratio of radial probability density of finding electron at $r \quad a_0$ to the radial probability density of finding electron at the nucleus is given as $(x.e^{\ y})$. Calculate the value of $(x \quad y)$.

15. Calculate the value of A.

$A \quad \frac{E_{1,2}}{2E_{2,1}}$ where $E_{n,Z}$ Energy of electron in n^{th} orbit, Z= atomic number of hydrogen like species.

For More Material Join: @JEEAdvanced_2024

Answers

Level 1

1.	(c)	**2.**	(c)	**3.**	(a)	**4.**	(d)	**5.**	(c)	**6.**	(c)	**7.**	(b)	**8.**	(c)	**9.**	(d)	**10.**	(d)
11.	(a)	**12.**	(b)	**13.**	(b)	**14.**	(a)	**15.**	(a)	**16.**	(d)	**17.**	(b)	**18.**	(c)	**19.**	(c)	**20.**	(b)
21.	(d)	**22.**	(b)	**23.**	(d)	**24.**	(d)	**25.**	(c)	**26.**	(a)	**27.**	(d)	**28.**	(d)	**29.**	(d)	**30.**	(d)
31.	(b)	**32.**	(c)	**33.**	(d)	**34.**	(b)	**35.**	(b)	**36.**	(a)	**37.**	(b)	**38.**	(c)	**39.**	(c)	**40.**	(d)
41.	(d)	**42.**	(d)	**43.**	(b)	**44.**	(d)	**45.**	(d)	**46.**	(b)	**47.**	(b)	**48.**	(a)	**49.**	(b)	**50.**	(c)
51.	(c)	**52.**	(c)	**53.**	(b)	**54.**	(a)	**55.**	(c)	**56.**	(a)	**57.**	(d)	**58.**	(b)	**59.**	(b)	**60.**	(d)
61.	(a)	**62.**	(c)	**63.**	(b)	**64.**	(b)	**65.**	(c)	**66.**	(c)	**67.**	(c)	**68.**	(c)	**69.**	(c)	**70.**	(a)
71.	(d)	**72.**	(a)	**73.**	(a)	**74.**	(c)	**75.**	(b)	**76.**	(b)	**77.**	(d)	**78.**	(b)	**79.**	(b)	**80.**	(d)
81.	(c)	**82.**	(b)	**83.**	(a)	**84.**	(c)	**85.**	(d)	**86.**	(b)	**87.**	(b)	**88.**	(b)	**89.**	(d)	**90.**	(a)
91.	(c)	**92.**	(c)	**93.**	(c)	**94.**	(a)	**95.**	(c)	**96.**	(d)	**97.**	(b)	**98.**	(a)	**99.**	(a)	**100.**	(d)
101.	(c)	**102.**	(b)	**103.**	(b)	**104.**	(c)	**105.**	(d)	**106.**	(c)	**107.**	(a)	**108.**	(d)	**109.**	(b)	**110.**	(c)
111.	(b)	**112.**	(c)	**113.**	(d)	**114.**	(b)	**115.**	(d)	**116.**	(d)	**117.**	(b)	**118.**	(a)	**119.**	(d)	**120.**	(c)
121.	(d)	**122.**	(d)	**123.**	(b)	**124.**	(c)	**125.**	(d)	**126.**	(c)	**127.**	(c)	**128.**	(d)	**129.**	(d)	**130.**	(d)
131.	(c)	**132.**	(b)	**133.**	(d)	**134.**	(d)	**135.**	(c)	**136.**	(c)	**137.**	(b)	**138.**	(a)	**139.**	(d)	**140.**	(c)
141.	(c)	**142.**	(c)	**143.**	(a)	**144.**	(d)	**145.**	(d)	**146.**	(c)	**147.**	(d)	**148.**	(b)	**149.**	(a)	**150.**	(d)
151.	(b)	**152.**	(a)	**153.**	(d)	**154.**	(a)	**155.**	(c)	**156.**	(a)	**157.**	(a)	**158.**	(d)	**159.**	(d)	**160.**	(d)
161.	(c)	**162.**	(a)	**163.**	(d)	**164.**	(d)	**165.**	(c)	**166.**	(a)	**167.**	(a)	**168.**	(d)	**169.**	(d)	**170.**	(a)
171.	(a)	**172.**	(d)	**173.**	(c)	**174.**	(b)	**175.**	(c)										

Level 2

1.	(c)	**2.**	(c)	**3.**	(c)	**4.**	(c)	**5.**	(a)	**6.**	(d)	**7.**	(b)	**8.**	(c)	**9.**	(b)	**10.**	(d)
11.	(a)	**12.**	(d)	**13.**	(a)	**14.**	(b)	**15.**	(c)	**16.**	(c)	**17.**	(c)	**18.**	(d)	**19.**	(d)	**20.**	(d)
21.	(a)	**22.**	(b)	**23.**	(c)	**24.**	(a)	**25.**	(b)	**26.**	(b)	**27.**	(c)	**28.**	(c)	**29.**	(b)	**30.**	(a)

For More Material Join: @JEEAdvanced_2024

Level 3

Passage–1	**1.** (c)	**2.** (a)	**3.** (c)		
Passage–2	**1.** (b)	**2.** (b)	**3.** (d)	**4.** (c)
Passage–3	**1.** (d)	**2.** (c)		
Passage–4	**1.** (b)	**2.** (b)		
Passage–5	**1.** (d)	**2.** (c)	**3.** (a)	

One or More Answers is/are Correct

1. (a,b,c)	**2.** (a,b,c)	**3.** (d)	**4.** (a,c,d)	**5.** (a,b,d)	**6.** (a,b,c)
7. (a,b,c,d)	**8.** (c)	**9.** (b,c,d)	**10.** (a,b,d)	**11.** (a,b,c,d)	**12.** (a,d)
13. (a,b,c,d)	**14.** (b,c,d)	**15.** (a,b,c,d)	**16.** (a,c)	**17.** (b,c,d)	**18.** (b,c)
19. (a,b,c,d)	**20.** (c,d)	**21.** (a,b,d)	**22.** (a,b,c,d)		

Match the Column

1.	A	P, R;	B	Q, R;	C	S;	D	Q, R
2.	A	S;	B	P, R;	C	P, R;	D	P, Q, R
3.	A	S;	B	Q;	C	R;	D	P
4.	A	Q;	B	S;	C	P;	D	R
5.	A	Q;	B	S;	C	R;	D	P
6.	A	P, Q, R, S;	B	Q, R, S;	C	R;	D	R, S
7.	A	P, S;	B	P, Q, S;	C	P, Q, R, S;	D	S
8.	A	Q;	B	S;	C	R;	D	P
9.	A	Q;	B	R;	C	S;	D	P
10.	A	P; Q	B	Q;	C	S;	D	R
11.	A	S;	B	R;	C	P;	D	Q
12.	A	S;	B	P;	C	Q;	D	R
13.	A	S;	B	R;	C	Q;	D	P
14.	A	P;	B	P, Q, S;	C	Q, S;	D	Q, S
15.	A	Q;	B	R;	C	S;	D	P

For More Material Join: @JEEAdvanced_2024

Assertion-Reason Type Questions

1. (B) **2.** (B) **3.** (A) **4.** (A) **5.** (A) **6.** (A) **7.** (A) **8.** (A) **9.** (B) **10.** (C)

11. (C) **12.** (A)

Subjective Problems

1.	2	**2.**	2	**3.**	3	**4.**	2	**5.**	4	**6.**	2	**7.**	4	**8.**	9	**9.**	3	**10.**	3
11.	3	**12.**	5	**13.**	5	**14.**	3	**15.**	8										

For More Material Join: @JEEAdvanced_2024

Hints and Solutions

Level 1

1. (c) Isodiaphers have same value of $(n - p)$ or $(A - 2Z)$.

2. (c) Travel perpendicular to both electric and magnetic fields.

3. (a) $\frac{(e/m)_p}{(e/m)} = \frac{e/m_p}{2e/4m_p} = \frac{2}{1}$

4. (d) Charge/mass for $n = 0$, for $\frac{2}{4}$,

for $p = \frac{1}{1}$, for $e = \frac{1}{1/1837}$

5. (c) $\frac{m}{e} = 1.5 \times 10^{8} \times 10^{3} = \frac{m}{1.6 \times 10^{-19}}$

$m = 2.4 \times 10^{-24}$ g

9. (d) $r = \frac{n^2}{Z}$; $\frac{r_1}{r_2} = \frac{(1)^2}{(3)^2}$

$r_2 = 9r_1$

12. (b) $v = \frac{Z}{n}$

13. (b) $r_{atom} = 10^{-10}$m

$r_{nucleus} = 10^{-5}$m

$\frac{V_{Atom}}{V_{Nucleus}} = \frac{4/3 \times (10^{-10})^3}{4/3 \times (10^{-15})^3} = 10^{15}$

14. (a) Change in P.E. $= \frac{2x}{4} \times (2x) = \frac{3}{2}x$

15. (a) $E_n = \frac{1}{2}$ P.E. $= \frac{6.8}{2} = 3.4$ eV

$\because E_n = \frac{13.6}{n^2} = 3.4$

$n = 2$ or first excited state

16. (d) Energy of N-shell $= \frac{13.6 \times (4)^2}{(4)^2} = 13.6$ eV

P.E. $= 2 \times E = 2 \times 13.6 = -27.2$ eV

17. (b) Total energy of third shell $= \frac{13.6}{3^2}$

$= 1.51$ eV

K.E. = Total energy $= 1.51$ eV

P.E. $= 2 \times$ T.E. $= -3.02$ eV

18. (c) $r = 0.529 \frac{n^2}{Z}$ Å;

$r_4 - r_3 = 0.529 \left(\frac{16}{2} - \frac{9}{2}\right)$ Å $= 1.851 \times 10^{-10}$ m

19. (c) $r_1 = 0.529$ Å; $r_{4(X)} = r_1 \frac{n^2}{Z}$;

$r_{4(X)} = \frac{0.529 \times (4)^2}{Z}$; $Z = 16$

20. (b) $\frac{v_1}{v_2} = \frac{n_2}{n_1} = \frac{5}{3}$

21. (d) $\frac{r_1}{r_2} = \left(\frac{n_1}{n_2}\right)^2 = \frac{4}{9}$

22. (b) r_1 of H-atom $= 0.529$ Å

r_n (H like atom) $= \frac{n^2}{Z} r_1$ (H-atom)

r_n of $Be^{3-} = \frac{n^2}{Z} r_1$ (H-atom)

$= 0.529$ Å ($Z = 4$ for Be^{3+})

$\frac{n^2}{Z} = 0.529 = 0.529 \times n^2 \times Z$

$n^2 = 4 \Rightarrow n = 2$

23. (d) $v = 2.188 \times 10^6 \times Z/n$

24. (d) Frequency of revolution $= \frac{\text{velocity in second orbit } (V_2)}{2\pi r_2}$

$= \frac{1.082 \times 10^6 \text{ ms}^{-1}}{2\pi \times (2.12 \times 10^{-10})\text{ m}} = 8.2 \times 10^{14} \text{ s}^{-1}$

25. (c) $v = \frac{Ze^2}{n}$; $r = \frac{n^2}{Z}$;

frequency of revolution $= \frac{v_n}{2\pi r_n}$;

Coulombic force of attraction $= \frac{Ze^2}{(4\pi\varepsilon_0) r^2}$

26. (a) Number of wave in one complete revolution = shell number.

For More Material Join: @JEEAdvanced_2024

28. (d) For Be^{3} E E_2 $13.6\,\frac{z^2}{n^2}$

$13.6\,\frac{4^2}{2^2}$ 54.4 eV

29. (d) T $\frac{n^3}{Z^2}$; $\frac{T_{1,2}}{T_{2,1}}$ $\frac{1}{4}$ $\frac{1}{8}$ $\frac{1}{32}$

30. (d) mvr $\frac{nh}{2}$

31. (b) T $\frac{n^3}{Z^2}$; $\frac{T_1}{T_2}$ $\frac{n_1^3}{Z_1^2}$ $\frac{Z_2^2}{n_2^3}$

$\frac{Z^3}{1}$ $\frac{2^2}{3^3}$ $\frac{32}{27}$

32. (c) When an electron of charge e and mass m is accelerated with a potential difference of V volts K.E. eV

$\frac{1}{2}mv^2$ eV or v^2 $\frac{2\,eV}{m}$

v $\sqrt{\frac{2\,eV}{m}}$

33. (d) E_H in first orbit $\frac{19.6$ $10^{-18}}{4}$ J

$E_{Be^{3+}}$ in second orbit

$\frac{19.6$ $10^{-18}}{4}$ $\frac{16}{4}$

19.6 10^{-18} J

34. (b) $\frac{E_1}{E_2}$ $\frac{Z_1}{Z_2}$ $\frac{1}{2}$

E_2 $4E_1$ 13.64 eV

35. (b) $\because$ E_n $\frac{Z^2}{n^2}$ E_n Z^2

36. (a) $\because$ $\frac{mv^2}{r}$ $\frac{1}{4$ $_0}$ $\frac{Ze$ $e}{r^2}$

v^2 $\frac{e^2}{4$ $_0rm}$ (For hydrogen Z 1)

v^2 $\frac{e^2}{4$ $_0rm}$ v $\frac{e}{\sqrt{4$ $_0rm}}$

38. (c) E $13.6\,Z^2$ $\frac{1}{n_1^2}$ $\frac{1}{n_2^2}$ eV atom $^{-1}$

For the ionization of Li^{2} (Z 3) from first excited state, n_1 2 and n_2 .

Hence, IP E 13.6 3^2 $\frac{1}{2^2}$ $\frac{1}{}$ ^{2}

30.6 eV

39. (c) E $h\nu$ 6.62 10^{-34} 4.2 10^{14}

2.8 10^{-19} J

40. (d) E $\frac{1240}{}$ nm $\frac{1240}{91}$ 13.6 eV

41. (d) E $\frac{hc}{}$

42. (d) E $n\,\frac{hc}{}$

$\frac{2$ $10^{17}}{1.6$ $10^{-19}}$ eV n $\frac{1240}{595.2}$ $\frac{\text{eV.nm}}{\text{nm}}$

n 60

45. (d) n_1 2

n_2 6

Total number of photon emitted

$\frac{(6$ $2)(6$ 2 $1)}{2}$ 10

46. (b) n_1 2

n_2 n

$\bar{\nu}$ $R(1)^2$ $\frac{1}{(2)^2}$ $\frac{1}{n^2}$

R $\frac{n^2$ $2^2}{4n^2}$

49. (b) n_1 n_2 4; n_2 n_1 2; n_1 1, n_2 3

$-$ $R(2)^2$ $\frac{1}{1^2}$ $\frac{1}{3^2}$ $\frac{32R}{9}$

50. (c) $\frac{91.2}{9\,\frac{1}{(2)^2}$ $\frac{1}{(3)^2}}$ $91.2\,\frac{36}{9$ $5}$ 73 nm

51. (c) E 13.6 $(2)^2$ $\frac{1}{4}$ $\frac{1}{36}$ 96.43

1.16 10^3 kJ/mol

52. (c) $\frac{1}{}$ RZ^2 $\frac{1}{n_1^2}$ $\frac{1}{n_2^2}$ R 3^2 $\frac{1}{3^2}$ $\frac{1}{}$ ^{2}

R or $\frac{1}{R}$

53. (b) $\frac{1}{}$ RZ^2 $\frac{1}{n_1^2}$ $\frac{1}{n_2^2}$ R 2^2 $\frac{1}{1^2}$ $\frac{1}{2^2}$

$3R$; $\frac{1}{3R}$

54. (a) For the transition n 1 to n 2, the energy change, E is positive, *i.e.,* energy is absorbed. For the transition n 5 to n 1, E is negative, *i.e.,* energy is released.

For More Material Join: @JEEAdvanced_2024

55. (c) For hydrogen atom

$$\frac{1}{} \quad R_H \quad \frac{1}{n_1^2} \quad \frac{1}{n_2^2} ;$$

$$2170 \text{ nm} \quad 2170 \quad 10^{-9} \text{ m};$$

$$R_H \quad 1.09677 \quad 10^7 \text{ m}^{-1}$$

$$\frac{10^9}{2170} \quad 1.09677 \quad 10^7 \quad \frac{1}{n_1^2} \quad \frac{1}{7^2} ; \quad n_1 \quad 4$$

So, electronic transition from $n \quad 7$ to $n \quad 4$ will produce infrared radiation of wavelength 2170 nm.

56. (a) Total number of spectral lines given by

$$\frac{1}{2}[n \quad 1] \quad n \quad 15; \qquad n \quad 6$$

Thus, electron are excited upto 6th energy level from ground state. Therefore,

$$\frac{1}{} \quad R_H \quad \frac{1}{1^2} \quad \frac{1}{n^2} \qquad 109737 \quad \frac{35}{36};$$

$$9.373 \quad 10^{-6} \text{ cm} \quad 937.3 \text{ Å}$$

57. (d) $E \quad \frac{1}{}$ for $_{max}$, E_{min}

58. (b) $K\ L\ M\ N$ ..... energy increases.

59. (b) $E \quad 13.6Z^2 \quad \frac{1}{n^2} \quad \frac{1}{(n \quad 1)^2}$

63. (b) $h\nu \quad h\nu_0 \quad KE$

64. (b) $h\nu \quad h\nu_0 \quad eV_0$

$$V_0 \quad \frac{h\nu}{e} \quad \frac{h\nu_0}{e}$$

65. (c) $h\nu \quad h\nu_0 \quad KE$

67. (c) As per Einstein's equation of photoelectric effect $h \quad h_0$ K.E.

$$\frac{1}{2}mv^2 \quad h \quad h_0 \quad \frac{hc}{} \quad \frac{hc}{0}$$

$$v^2 \quad \frac{2hc}{m} \quad \frac{1}{} \quad \frac{1}{0} ;$$

$$v \quad \frac{2hc}{m} \quad \frac{1}{} \quad \frac{1}{0}^{1/2}$$

$$\frac{2hc}{m} \quad \frac{0}{0}^{1/2}$$

68. (c) $\frac{hc}{} \quad 1 \qquad \ldots(1)$

$$3 \quad \frac{hc}{} \quad 4 \qquad \ldots(2)$$

from, eq. (1) and (2) 0.5 eV

69. (c) $\frac{1}{2}mv^2 \quad \frac{1240 \text{ eV nm}}{31 \text{ nm}} \quad 40 \quad 12.8 \text{ eV}$

$$= 27.2 \text{ eV}$$

$$\frac{1}{2} \quad 9.1 \quad 10^{-31} \quad v^2 \quad 27.2 \quad 1.6 \quad 10^{-19}$$

$$v \quad 3.09 \quad 10^6 \text{ m/s}$$

70. (a) $h \quad h_0 \quad eV_0$; $eV_0 \quad h \quad h_0$ or $V_0 \quad \frac{h}{e} \quad \frac{h}{e} \quad _0$; slope$_1$ $\frac{h}{e}$

Similarly, $h \quad h_0 \quad K_{max}$ or $K_{max} \quad h \quad h_0$;

slope$_2$ h, $\frac{\text{slope}_2}{\text{slope}_1} \quad \frac{h}{h/e} \quad e$

71. (d) $E_n \quad \frac{13.6}{n^2}$ eV; $E_2 \quad \frac{13.6}{2^2}$;

$E_4 \quad \frac{13.6}{4^2}$ eV/atom

$E \quad E_4 \quad E_2 \quad 2.55$ eV

Absorbed energy work function of metal K.E.

$2.55 \quad 2.5 \quad$ K.E.; K.E. 0.05 eV

74. (c) $\frac{1}{m}$

75. (b) $\frac{h}{mv} \quad \frac{6.63 \quad 10^{-34}}{10^{-6} \quad 10} \quad 6.63 \quad 10^{-29}$ m

76. (b) $\frac{h}{\sqrt{2qVm}}$

77. (d) $\frac{h}{p}$

78. (b) $\because \quad \frac{1}{} \quad R \quad \frac{1}{n_1^2} \quad \frac{1}{n_2^2} ; \quad n_1 \quad 1, n_2 \quad ?;$

$$\frac{1}{} \quad R \quad \frac{1}{1} \quad \frac{1}{n_2^2} \qquad n_2^2 \quad \frac{R}{R \quad 1}$$

$$n_2 \quad \sqrt{\frac{R}{R \quad 1}}$$

79. (b) $E \quad E_1 \quad E_2$; $\frac{hc}{} \quad \frac{hc}{1} \quad \frac{hc}{2}$

$$\frac{hc}{} \quad hc \quad \frac{_2 \quad _1}{_1 \quad _2} ; \qquad \frac{_1 \quad _2}{_1 \quad _2}$$

80. (d) $_p \quad \frac{h}{\sqrt{2 \ eV \ m_p}}$

$$_{Be^3} \quad \frac{h}{\sqrt{2 \quad 3 \ eV \ m_{Be^{3+}}}} \quad \frac{h}{\sqrt{2 \quad 3 \ eV \quad 9m_p}}$$

Hence, $\frac{_{Be^3}}{_p} \quad \sqrt{\frac{2 \ eV \ m_p}{2 \quad 3 \ eV \quad 9m_p}} \quad \frac{1}{3\sqrt{3}}$

For More Material Join: @JEEAdvanced_2024

81. (c) $\frac{h}{\sqrt{2\,eV\,m}}$ $\frac{1.23}{\sqrt{V}}$ nm

82. (b)

$\frac{h}{m_e x}$ $\frac{h}{m_p V}$ $\frac{h}{1840\,m_e V}$ $[m_p$ $1840\,m_e]$

Hence, V $\frac{x}{1840}$

83. (a) E $(mC)\,C$ or P $\frac{E}{C}$

$$\frac{6\ \ 10^{6}\ \ 1.6\ \ 10^{-19}}{3\ \ 10^{8}}$$

$3.2\ \ 10^{-21}$ kg-m/s

84. (c) E $\frac{nhc}{}$ nhc^{-} $\because$ $-$ $\frac{1}{}$

10 nhc^{-} or n $\frac{10}{hc^{-}}$ $\frac{10}{hcx}$

86. (b) $\because$ mvr_n $\frac{nh}{2}$ and p $\frac{h}{}$

$2\ r$ n

$\because$ r $4a_0$

2 $4a_0$ 2

$4a_0$

or $\frac{h}{}$ r_2 $\frac{h}{}$ r_2,

r_2 $4a_0$;

$4a_0$

87. (b) I.E. 13.6 $\frac{Z^2}{n^2}$ eV 13.6 4 54.4 eV

for 2 mole 54.4 2 N_A eV $108.8\,N_A$ eV

90. (a) m 10^{-10} g 10^{-13} kg;

v $\frac{0.0001}{100}$ 10^{-6} 10^{2} 10^{-14} m sec^{-1}

x p $\frac{h}{4}$

x $\frac{h}{4\ \ p}$ $\frac{h}{4\ \ m\ \ v}$;

$$x\ \ \frac{6.62\ \ 10^{-34}}{4\ \ 3.14\ \ 10^{-13}\ \ 10^{-14}}$$

x $\frac{6.62}{12.56}$ $\frac{10^{-34}}{10^{-27}}$; x 5.2 10^{-8} m

91. (c) x $\frac{h}{4\ \ p}$ $\frac{h}{4\ \ m\ \ v}$

$$\frac{6.63\ \ 10^{-14}}{4\ \ 3.14\ \ 9.1\ \ 10^{-31}\ \ 1}$$

($\because$ v 1 m / s)

x 58 m

94. (a) Use $(n$ $l)$ rule.

101. (c) Radii of maximum probability of electron is smaller for orbital having higher value of 'l' for given same value of n.

103 (b) Total no. of nodes $(n$ $1)$

104. (c) l 4;

number of degenerate orbitals $2l$ 1 9

109. (b) $4f$ has the highest energy among $3d$, $4f$, $4p$, $5s$ orbitals.

110. (c) m orientation of orbital.

112. (c) Maximum number of electrons with same spin is equal to maximum number of orbitals, *i.e.*, $(2l$ $1)$.

113. (d) Orbital angular momentum $\sqrt{l(l\ \ 1)}\,\frac{h}{2}$; l 1 for p-orbital.

116. (d) [Ne] $1s^2 2s^2, 2p^6$

117. (b) Fe(III)—[Ar] $3d^5$; unpaired electrons 5;

magnetic moment $\sqrt{5(5\ \ 2)}$ BM


Co(II)—[Ar] $3d^7$; unpaired electrons 3;

magnetic moment $\sqrt{3\ \ (3\ \ 2)}$ BM

119. (d) Given $\sqrt{n(n\ \ 2)}$ 1.73 BM


(where n is number of unpaired electrons)

n 1; $_{23}$V $1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 3d^3$




$3d^3$ $4s^2$

for n 1 it must release 4 electrons, first two from $4s$-orbital and then next two electrons from $3d$-orbital. So, x 4.

120. (c) 

$|s|$ $\frac{\text{No. of unpaired } e}{2}$

121. (d) 

For More Material Join: @JEEAdvanced_2024

123. (b) I $r \quad 0.529\frac{n^2}{Z}$ Å

II $_l \quad \sqrt{l(l \quad 1)}\frac{h}{2}$

$l \quad 0$

124. (c)

	$n \quad l$
e_1	3
e_2	4
e_3	5
e_4	4

$e_3 \quad e_2 \quad e_4 \quad e_1$

125. (d) $E_{2p_x} \quad E_{2p_y} \quad E_{2p_z}$

126. (c) $P(33) \quad [Ar]\,4s^2 4p^3$

$n \quad 3$

128. (d) 3*d*-subshell contains maximum 10 electrons.

130. (d) $|m| \quad l$

131. (c) Degenerate orbitals having equal energy.

140. (c) For *f* subshell $l \quad 3$; *g* subshell $l \quad 4$

for principal shell, $l \quad 4, n \quad 5$

total no. of orbital in shell $\quad n^2 \quad 5^2 \quad 25$

142. (c) Fe(26) $[Ar]\,3d^6 4s^2$

143. (a) $r \quad n^2$

145. (d) Total nodes $\quad n \quad 1$

146. (c) Unpaired electron of Cl atom is $3p^1$ for which $n \quad 3$,

$l \quad 1, m \quad 1, 0$ or $+1$ and $s \quad \frac{1}{2}$

147. (d) Cl(17) $3s^2, 3p^5$ $\boxed{\quad}\boxed{\uparrow}\boxed{\quad}$

$n \quad 3, l \quad 1, m \quad 1, s \quad \frac{1}{2}$

149. (a) *A* : excitation possible only in *d*-orbitals

B : Spin multiplicity $\quad 2|S| \quad 1; \ |S|$ =total spin

C : *V* not violated Hund's rule

D : *A* is paramagnetic due to unpaired *e*

A, B, C are correct.

150. (d) Orbitals are $4s, 2s, 3p$ and $3d$. Out of these $3d$ has highest energy.

151. (b) $5f$ subshell

154. (a) Pauli's exclusion principle

155. (c) For H-atom, energy of *s*-orbital energy of $2p$-orbital so degeneracy is 1 3 4

156. (a) For multi electron species (H) first excited state means *e* will transfer from $1s$ to $2s$ subshell & $2s$ subshell has only one orbital so degeneracy is 1.

158. (d) $(n \quad l)$ rule

160. (d) $n \quad l$

161. (c) $1s^2 2s^2 \ \boxed{2p^6} \ 3s^2 \ \boxed{3p^6}$

$\boxed{\quad}\boxed{\uparrow\downarrow}\boxed{\quad} \quad \boxed{\quad}\boxed{\uparrow\downarrow}\boxed{\quad}$

$0 \quad +1 \quad -1 \quad 0 \quad +1 \quad -1$

163. (d) $4s$-orbital

164. (d) *g*-subhsell $l \quad 4, 2(2l \quad 1) \quad 18e$

171. (a) Mn(25) $[Ar]3d^5, 4s^2 \quad n \quad 5$

Ti(22) $[Ar]3d^2, 4s^2 \quad n \quad 2$

V(23) $[Ar]3d^3, 4s^2 \quad n \quad 3$

Al(13) $[Ne]3s^2, 3p^1 \quad n \quad 1$

172. (d) No. of spherical nodes $\quad n \quad l \quad 1$;

for *s, p, d, f* values of $l \quad 0, 1, 2, 3$ respectively.

173. (c) Radial node $(n \quad l \quad 1) \quad 2$

$l \quad 0, \ n \quad 3$

174. (b) For radial node $\quad _{2s} \quad 0, 2 \quad \frac{r}{a_0} \quad 0$

$\boxed{r \quad 2a_0}$

175. (c) Probability of finding *e* is zero implies that

$^2 \quad 0$ or $\quad 0$

$(\quad 1) \quad 0 \quad 1$

or $\quad r_1 \quad \frac{a_0}{2z}$ or $(\ ^2 \quad 8 \quad 12) \quad 0$

and $\quad (\quad 6)(\quad 2) \quad 0$

$6, \ r \quad \frac{3a_0}{z}$

and $\quad 2, \ r \quad \frac{a_0}{z}$

$r_2 \quad \frac{3a_0}{z}$

For More Material Join: @JEEAdvanced_2024

Level 2

1. (c) P.E. $\frac{1}{4\ _0}\frac{(\ Ze)(\ e)}{r}$

$\frac{1}{4\ _0}\frac{(\ 2e)(\ e)}{r}$ $\frac{e^2}{2\ _0 r}$

2. (c) Ionization energy :

$217.6 = -13.6\ \frac{Z^2}{1^2}$; Z 4 m

So, it is $^{9}_{4}Be^{3}$; no. of neutrons 9 4 5

3. (c) $\frac{d[U(r)]}{dr}$ $\frac{3Ke^2}{r^4}$ Magnitude of the force

$\frac{3Ke^2}{r^4}$ $\frac{mv^2}{r}$

and we know mvr $\frac{nh}{2}$ or r $\frac{nh}{2\ m\ v}$,

$3Ke^2$ $\frac{8\ ^3m^3v^3}{n^3h^3}$ mv^2, v $\frac{n^3h^3}{24Ke^2\ ^3m^2}$

4. (c) $\frac{1}{2}$ P.E. = K.E

$\frac{1}{2}$ $\frac{1}{2}mkr^2$ $\frac{1}{2}mv^2$, mvr $\frac{nh}{2}$,

v^2 $\frac{n^2h^2}{4\ ^2m^2r^2}$; r^4 $\frac{n^2h^2}{2\ ^2m^2k}$

or r $\sqrt{n}$

5. (a) $\frac{1}{2}mv^2$ $\frac{k(q_1)q_2}{r}$ $\frac{q_2}{m}$ $\frac{r.v^2}{2k.q_1.Z}$

$\frac{q_2}{m}$ $\frac{2.5\ \ 10^{-14}\ \ (2.1\ \ 10^7)^2}{2\ \ 9\ \ 10^9\ \ 79\ \ 1.6\ \ 10^{-19}}$

$4.84\ \ 10^7$ coulomb/kg

6. (d) v r where r_n $\frac{n^2h^2}{4\ ^2me^2Z\ K}$

and v_n $\frac{2\ Ze^2\ K}{n\ h}$

$\frac{2\ Ze^2\ K}{n\ h}$ $\frac{n^2h^2}{4\ ^2me^4Z\ K}$;

$\frac{8\ ^3me^4\ Z^2\ K^2}{n^3\ h^3}$

$\frac{9\ ^3me^4\ K^2}{h^3}$ ($\because n$ 2 and Z 3)

7. (b) r_n $\frac{v^2}{Z}$; for H, r_4 r_3 $0.529(16\ \ 9)$

$0.529\ \ 7$ Å

r_4 r_3 for Li^{2} $0.529\ \frac{16}{3} - \frac{9}{3}$

$0.529\ \ \frac{7}{3}$ so ratio $\frac{7}{7/3}$ 3 : 1

8. (c) v_n $2.186\ \ 10^6\ \frac{Z}{n}$

$1.093\ \ 10^6$ $2.186\ \ 10^6\ \ \frac{1}{n}$; n 2

from Bohr theory we know 2 r n

2 , where $\frac{h}{mv}$

or r $0.529\ \frac{n^2}{Z}$ 0.529 4 Å

Circumference of the orbit

$2\ \ \frac{22}{7}\ \ 0.529\ \ 4\ \ 10^{-10}$

$13.30\ \ 10^{-10}$ m

9. (b) Angular momentum $\frac{nh}{2}$

$3.1652\ \ 10^{-34}$ $\frac{n\ \ 6.626\ \ 10^{-34}}{2}$;

n 3

$-$ $R\ Z^2\ \frac{1}{n_1^2}\ \frac{1}{n_2^2}$;

$-$ $R\ 2^2\ \frac{1}{2^2}\ \frac{1}{3^2}$ $\frac{5R}{9}$

10. (d) Energy of photon corresponding to second line of Balmer series for Li^{2+} ion

$= (13.6)\ \ (3)^2\ \ \frac{1}{2^2}\ \ \frac{1}{4^2}$

$= 13.6\ \ \frac{27}{16}$

Energy needed to eject electron from n 2 level in H-atom;

$= 13.6\ \ 1^2\ \ \frac{1}{2^2}\ \ \frac{1}{\ \ 2}$ $\frac{13.6}{4}$

K.E. of ejected electron

$13.6\ \ \frac{9\ \ 3}{16} - \frac{13.6}{4}$ $13.6\ \ \frac{27\ \ 4}{16}$

19.55 eV

11. (a) $\frac{1}{\ }$ $RZ^2\ \frac{1}{n_1^2}\ \frac{1}{n_2^2}$, where n_1 n, n_2 n 1

$\frac{1}{\ }$ $RZ^2\ \frac{1}{n^2}\ \ \frac{1}{(n\ \ 1)^2}$

$\frac{1}{\ }$ $\frac{2n\ \ 1}{n^2(n\ \ 1)^2}\ RZ^2$

Since, n 1;

Therefore, $2n$ 1 $2n$


For More Material Join: @JEEAdvanced_2024

and $(n \quad 1)^2 \quad n^2$

$$\frac{1}{} \quad RZ^2 \quad \frac{2n}{n^2 \; n^2}$$

$$\frac{}{c} \quad \frac{2RZ^2}{n^3} \quad \text{or} \quad \nu \quad \frac{2cRZ^2}{n^3}$$

12. (d)



Total radiations are = 6

13. (a) If a is side of cube, then $x \quad a\sqrt{3}$

$$x \quad 10\sqrt{3} \text{ cm} = 10\sqrt{3} \quad 10^{-2}\text{m}$$

$$x \quad p \quad \frac{h}{4}; \quad x \; m \quad \nu \quad \frac{h}{4}$$

$$\nu \quad \frac{h}{4 \; m \quad x}$$

$$\frac{6.63 \quad 10^{\;34}}{4 \quad 3.14 \quad 9.1 \quad 10^{\;31} \quad 10 \quad \sqrt{3} \quad 10^{\;2}}$$

$$\nu \quad 3.34 \quad 10^{-4} \text{ ms}^{-1}$$

14. (b) $2 \; r \quad 4 \; ; \; n \quad 4$

Total energy required total energy released

0

$$2 \quad 4.526 \quad N_A \quad 2 \quad 13.6 \quad N_A \quad 2 \quad 13.6$$

$$1 \quad \frac{1}{16} \quad N_A \quad 30.87 \quad x \quad N_A \quad 0$$

$x \quad 2$ moles of X required 2

15. (c) Work function for $Li^{2+} = 9E$.

$$E_p \quad w \quad \frac{1}{2}mv^2; \quad E_p \quad 9E \quad \frac{1}{2}mv^2$$

$$\nu \quad \sqrt{\frac{2(E_p \quad 9E)}{m}}$$

16. (c) $E \quad hc^{-} \quad 1.63 \quad 10^{-18}$ J

where $^{-} \quad R(1)^2 \quad \frac{1}{1^2} \quad \frac{1}{2^2} \quad \frac{3}{4}R$

Translational K.E. of H-atom $\frac{3}{2} \quad \frac{R}{N_A} \quad T$

$$\frac{3}{2} \quad \frac{8.314}{6 \quad 10^{23}} \quad T \quad 1.63 \quad 10^{\;18}$$

$$T \quad 7.84 \quad 10^4 \text{ K}$$

17. (c) Radial node occurs where probability of finding e is zero.

$$^2 \quad 0 \quad \text{or} \quad 0$$

$$6 \quad 6 \quad ^2 \quad 0; \quad 3 \quad \sqrt{3}$$

For max. distance $r \quad \frac{3}{2}\frac{(3 \quad \sqrt{3})a_0}{Z}$

18. (d) $\frac{n(n \quad 1)}{2} \quad 6; n \quad 4,$

$$n \quad 4 \quad E_4 \quad 0.85 \text{ eV}$$

$$n \quad 1 \quad E_1 \quad 13.6 \text{ eV}$$

$$E \quad 12.75 \text{ eV}$$

$$12.75 \text{ eV} = \frac{1240 \text{ eV-nm}}{}$$

$$97.25 \text{ nm}$$

19. (d) For II to I transition

$$E \quad \frac{4E}{3} \quad E \quad \frac{hc}{_{\text{II}} \quad _{\text{I}}}; \quad \frac{E}{3} \quad \frac{hc}{_{\text{II}} \quad _{\text{I}}}$$

For III to I transition

$$E \quad 2E \quad E \quad \frac{hc}{} \quad \text{or} \quad E \quad \frac{hc}{}$$

$$\frac{hc}{3} \quad \frac{hc}{_{\text{II}} \quad _{\text{I}}} \quad _{\text{II}} \quad _{\text{I}} \quad 3$$

20. (d)



Out of 6 electrons in $2p$ and $3p$ must have one electron with $m \quad 1$ and $s \quad \frac{1}{2}$ but in $3d$-subshell an orbital having $m \quad 1$ may have spin quantum no. $\frac{1}{2}$ or $\frac{1}{2}$.

Therefore, minimum and maximum possible values are 2 and 3 respectively.

21. (a) Energy absorbed $13.6 \quad 1.5 = 20.4$ eV out of this 6.8 eV is converted to K.E.

$$6.8 \text{ eV} \quad 6.8 \quad 1.6 \quad 10^{-19} \text{ J};$$

$$6.8 \quad 1.6 \quad 10^{-19} = \text{K.E.} \quad \frac{1}{2} \quad mv^2$$

$$\nu \quad \sqrt{\frac{2KE}{m}} \quad \sqrt{\frac{2 \quad 1.088 \quad 10^{\;18}}{9.1 \quad 10^{\;31}}}$$

$$1.54 \quad 10^6 \text{ m/s}$$

22. (b) $E \quad \frac{hc}{} \quad 2.9 \quad 10^{-19}$ J

For More Material Join: @JEEAdvanced_2024

Total energy of 10 quanta
10 2.9 10 ^{19} 29 10 ^{19} J
Energy stored for process
$\frac{112 \quad 4.18 \quad 10^3}{6 \quad 10^{23}}$ 7.80 10 ^{19} J

% efficiency $\frac{7.8 \quad 10^{-19}}{29 \quad 10^{19}}$ 100 26.9%

23. (c) Total energy emitted by photo-electron
$10.2 + 17 = 27.20$ eV
Since, E_1 Photon of energy emitted through the transition
n n to n 2 $\frac{hc}{1}$ 27.20 eV

We have $\frac{1}{1}$ R_H Z^2 $\frac{1}{2^2}$ $\frac{1}{n^2}$

or $\frac{hc}{1}$ $(hc)R_H$ Z^2 $\frac{1}{2^2}$ $\frac{1}{n^2}$

27.20 $(hc)\,R_H$ Z^2 $\frac{1}{4}$ $\frac{1}{n^2}$ ...(1)


Similarly, total energy liberated during transition of electron from n n to n 3 is
E_2 $\frac{hc}{2}$ $(4.25 + 5.95) = 10.20$ eV

10.20 $(hc)\,R_H$ Z^2 $\frac{1}{9}$ $\frac{1}{n^2}$ ...(2)

Dividing Eq. (1) by (2), we get n 6 and putting n 6 in Eq. (1) or (2), we get, Z 3.

24. (a) E $\frac{hc}{}$ $\frac{1240 \text{ eV-nm}}{1025.6 \quad 10^{10} \quad 10^9}$

E 12.09 eV



In three different radiations, minimum wavelength for 3 1 transition
3 1 $\frac{hc}{E}$ $\frac{1240 \text{ eV-nm}}{12.09 \text{ eV}}$ 102.6 nm

25. (b) E $\frac{hc}{}$ $\frac{(6.63 \quad 10^{-34} \text{J-s})\,(3.00 \quad 10^8 \text{ m/s})}{3.055 \quad 10^{-8} \text{ m}}$

6.52 10^{-18} J

E_H $\frac{3}{4}(2.176 \quad 10^{-18}$ J)

1.63 10^{-18} J; E $E_H(Z^2)$

Z^2 $\frac{E}{E_H}$ $\frac{(6.52 \quad 10^{-18})}{(1.63 \quad 10^{-18})}$ 4;

Z 2 (helium)

26. (b) K.E. $\frac{K.Ze.2e}{r}$

r $\frac{9 \quad 10^9 \quad 29 \quad 2 \quad (1.6 \quad 10^{-19})^2}{5 \quad 1.6 \quad 10^{-19} \quad 10^6}$

r $\frac{9 \quad 10^9 \quad 29 \quad 2 \quad 1.6 \quad 10^{-19}}{5 \quad 10^6}$

1.67 10^{-14}m

27. (c) $\sqrt{v}$ aZ ab
ab 1, a tan 45 1
$\sqrt{v}$ 51 1 50
v 50^2 2500 s^{-1}

28. (c) $\frac{1}{}$ R $\frac{1}{2^2}$ $\frac{1}{n^2}$ R $\frac{n^2 \quad 4}{4n^2}$

$\frac{4}{R}$ $\frac{n^2}{n^2 \quad 4}$ ...(1)

Given: k $\frac{n^2}{n^2 \quad 4}$ ...(2)

Comparing equation (1) and (2) we have
K $\frac{4}{R}$

29. (b) 2 r_n n 2 $0.53\frac{n^2}{z}$ n

2 0.53 $\frac{n}{z}$ ...(1)

E_{sep} 3.4 $13.6\frac{z^2}{n^2}$ $\frac{n}{z}$ 2

2p 0.53 2 6.66 Å

30. (a) $\frac{E}{E_{4th}}$ $\frac{24}{E_{4th}}$ $\frac{\frac{1}{1} \quad \frac{1}{4}}{\frac{1}{16}}$ $\frac{3/4}{1/16}$

E_{4th} 2 eV.

Level 3

Passage-1

1. (c) x p $\frac{h}{4}$ p^2 $\frac{h}{4}$

m^2 v^2 $\frac{h}{4}$ v $\frac{1}{2m}\sqrt{h}$

For More Material Join: @JEEAdvanced_2024

2. (a) $x \sqrt{\frac{h}{4 m}}$; $x p \frac{h}{4}$

$\sqrt{\frac{h}{4 m}} p \frac{h}{4}$, $p \sqrt{\frac{mh}{4}}$

3. (c) D.B. $\sqrt{\frac{150}{6}}$ Å = 5 Å

and $x p \frac{h}{4}$; $p \frac{h}{}$ or $p \frac{h}{2}$

$x \frac{h}{2} \frac{h}{4}$

$\frac{1}{} \frac{10^{-9}}{2} \frac{1}{4} \frac{2.5}{4} 10^{-10}$

0.625 Å

Passage-2

1. (b) $\because \frac{1}{} R_H Z^2 1 \frac{1}{n^2}$ for Lyman's series

For He ion series limit n ;

$\frac{1}{1} R_H 2^2 1 \frac{1}{2} \frac{1}{1} R_H 4$

Similarly, Balmer series limit for Li^{2} ion

$\frac{1}{2} R_H 3^2 9R_H 9 \frac{1}{4} {}_2 \frac{4x}{9}$

2. (b) $6 (n_2 n_1) \frac{(n_2 n_1 1)}{2}$; $n_2 4$, $n_1 1$

$\frac{1}{} R_H 2^2 1 \frac{1}{16}$

$R \frac{15}{4}$; $\frac{4}{15R_H}$

3. (d) $\frac{1}{} R_{H Z^2} \frac{1}{n_1^2} \frac{1}{n_2^2}$

$\frac{1}{\text{He}} R_H 2^2 \frac{1}{2^2} \frac{1}{4^2}$

$R_H \frac{1}{1^2} \frac{1}{2^2} \frac{1}{H}$

for n_2 2 to n_1 1

Passage-3

1. (d) Magnetic moment $\sqrt{n(n 2)}$

(A) Fe^{2} $3d^6 4s^0$ $n 4$

(B) Mn^{2} $3d^5 4s^0$ $n 5$

(C) Cr^{3} $3d^3 4s^0$ $n 3$

(D) V^{3} $3d^2 4s^0$ $n 2$

Smaller n, smaller

2. (c) $\sqrt{n(n 2)}$ 3.873

$n 3$

Mn $3d^5 4s^2$

Mn^{4} $3d^3 4s^0$

Passage-4

1. (b) Total energy required for dissociation of O_2 molecule and then assigning (O*) 1eV more energy than (O) (480+96) 576 kJ/mol

Maximum wavelength $E \frac{hc}{}$;

maximum $\frac{hc}{E_{min}}$; $_{max}$ (Å) $\frac{12400}{E_{min}(eV)}$

$\frac{12400}{6}$ 2066.67Å

2. (b) $O_3(g)$ $O_2(g)$ O ....(i)

$O_2(g)$ O O* ...(ii)

Energy required for (i) reaction is 400 kJ/mol

Normal dissociation of O_2 required 480 kJ/mol (given)

each (O*) has 1 eV more energy than (0) and given 1 eV/photon 96 kJ/mol

So total energy required for the dissociation of O_3 into O and O* is 400 480 96 976 kJ/mol

One or More Answers is/are Correct

2. (a, b, c)

$mvr n \frac{h}{2}$

n integer value 1,2,3,4, …

15. (a,b,c,d) At the point of maximum value of RDF

$\frac{dP}{dr} 0$

$2r \frac{2Zr^2}{a_0} 0$; $r \frac{a_0}{Z}$

where Z 3 for Li^{2} and Z 2 for the He ;

Z 1 for hydrogen.

18. (b, c)

$h W KE$

Kinetic energy of electrons does not depend on intensity of light $\frac{h}{\sqrt{2KE m}}$

For More Material Join: @JEEAdvanced_2024

19. (a, b, c, d)

(a) $v \quad \frac{Z}{n}$ (b) $r \quad \frac{n^2}{Z}$

(c) P.E. $\frac{Z^2}{n^2}$ (d) K.E. $\frac{Z^2}{n^2}$

20. (c, d)

Given n, l 0 to n 1

$m_l \quad l$.....0..... l

21. (a, b, d)

(a) No. of subshell n, n 3

(b) No. of orbital $2n^2$, n 4

(c) cu $3d^{10}4s^0$ diamagnetic

22. (a, b, c, d)

5 1 5 2 5 3 $\underbrace{5 \quad 4}_{\text{Brackett}}$

4 1 4 2 $\underbrace{4 \quad 3}_{\text{Paschen}}$

3 1 $\underbrace{3 \quad 2}_{\text{Balmer}}$

$\underbrace{2 \quad 1}_{\text{Lyman}}$

Match the Column

13. (a) s-orbital $\because r$ 0, 0 and 3 radial nodes $4s$

(b) 3 radial nodes (s, p, d) $4s, 5p_x\ 6d_{xy}$

(c) Angular probability is dependent of and for $5p_y, 6d_{xy}$

(d) At least one angular node $5p_x(1)$; $6d_{xy}(2)$

Subjective Problems

1. $r_{n\ 1} \quad r_{n\ 1} \quad 2\,r_n$

$0.529\frac{(n\ 1)^2}{Z} \quad 0.529\frac{(n\ 1)^2}{Z} \quad 2 \quad 0.529\frac{n^2}{Z}$

$n^2 \quad 2n \quad 1 \quad [n^2 \quad 2n \quad 1] \quad 2n^2$

$4n \quad 2n^2$

$n \quad 2$

2. $E \quad 30.6 \quad 13.6\frac{(3)^2}{(n)^2}$

$n \quad 2$ no. of wave in one complete evolution in the orbit.

3. n

$\underline{\quad}_{2\text{nd}}$ orbit $1.5 \quad \frac{n}{2}$

$n \quad 3$ no. of wave in one complete revolution in the orbit.

4. $E_{\text{abs}} \quad \frac{40}{100} \quad E_{\text{Emitted.}}$

$n_{\text{ab}} \quad \frac{hc}{400} \quad \frac{40}{100} \quad n_{\text{em}} \quad \frac{hc}{500}$

$\frac{n_{\text{ab}}}{n_{\text{em}}} \quad \frac{400}{500} \quad \frac{100}{40} \quad 2$

5. $E_{\text{inc}} \quad KE$

4.5 3.0 KE

$KE \quad 1.5$ eV

$\sqrt{\frac{150}{1.5}} \quad 10$ Å

$P \quad x \quad \frac{}{4}$

$\frac{h}{2} \quad x \quad \frac{h}{4}$

$\frac{}{(10 \quad 10^{\ 10})^2} \quad \frac{25}{4} \quad 10^{\ 10} \quad \frac{1}{4}$

4 Å

6. Cr $3d^5 4s^1$

No. of angular nodes 2

No. of radial nodes 3 2 1 0

Angular node radial node 2

7. Total node Radial node angular node

$n \quad 1 \quad 1 \quad 1 \quad n \quad 1$

$5f \quad 5 \quad 1 \quad 4$

8. $P \quad \frac{n\frac{hc}{}}{t}$

$\frac{100 \quad 12}{100} \quad (x \quad 10^{19}) \quad \frac{6.63 \quad 10^{\ 34} \quad 3 \quad 10^8}{1500 \quad 10^{\ 9} \quad 1}$

$x \quad 9$

9. $\frac{1}{} \quad RZ^2 \quad \frac{1}{n^2} \quad \frac{1}{(n \quad 1)^2}$

$\frac{1}{} \quad RZ^2\frac{(n \quad 1)^2 \quad n^2}{n^2(n \quad 1)^2}$

$\frac{1}{} \quad RZ^2\frac{(n \quad 1 \quad n)(n \quad 1 \quad n)}{n^2(n \quad 1)^2}$

$\frac{1}{} \quad RZ^2\frac{(2n \quad 1)}{n^2(n \quad 1)^2}$

n^3

$x \quad 3$

For More Material Join: @JEEAdvanced_2024

10. For radial node

$$\psi_{3s} = 0 = \frac{1}{81\sqrt{3}}\left(\frac{1}{a_0}\right)^{3/2}\left(27 - \frac{18r}{a_0} + \frac{2r^2}{a_0^2}\right)e^{-r/3a_0}$$

$$27 - \frac{18r}{a_0} + \frac{2r^2}{a_0^2} = 0$$

$$r = \frac{(18a_0) \pm \sqrt{(-18a_0)^2 - 4 \cdot 2 \cdot 27a_0^2}}{2 \cdot 2}$$

$$= \frac{18a_0 - 10.39\,a_0}{4}$$

distance between to radial node (d)

$$\frac{d}{1.73a_0} = \frac{2}{4} \cdot \frac{10.39a_0}{1.73a_0}$$

$$d = 3$$

11. $7.67 \times 10^{-19} = \frac{(1.6 \times 10^{-19})^2}{4 \times 3.14 \times 8.85 \times 10^{-12} \times r}$

$r = 3.00 \times 10^{-10}\text{ m} = 3\text{Å}$

12. $d(KE) = mv\,dv = mv\,\frac{h}{4\,m\,\Delta x}$

$$= \frac{3 \times 10}{3} \cdot \frac{6.62 \times 10^{-34}}{4 \times \frac{3.31}{10^{12}}}$$

$= 5 \times 10^{-16}\text{J}$

13. $E_{in} = 10.2\text{ eV}$

$= 4.2\text{ eV}$

$KE_{max} = 10.2 - 4.2 = 6\text{eV}$

$e = \sqrt{\frac{150}{6}}\text{Å} = 5\text{Å}$

14. $\frac{\text{Radial probability density at } r = a_0}{\text{Radial probability density at } r = 0} = \frac{R^2(a_0)}{R^2(0)}$

For 1s orbital: $R_{(r)} = \frac{1}{\sqrt{\pi}}\left(\frac{1}{a_0}\right)^{3/2} e^{-\frac{r}{a_0}}$

$$\frac{R^2(a_0)}{R^2(0)} = \frac{(1/\pi a_0^3)e^{-2r/a_0}}{(1/\pi a_0^3)e^0} = e^{-2}$$

15. $A = \frac{E_{1,2}}{2E_{2,1}} = \frac{-13.6 \times 2^2 \times 2^2}{2 \times 1^2 \times (-13.6) \times 1^2} = \frac{16}{2} = 8$

For More Material Join: @JEEAdvanced_2024



3

GASEOUS STATE

Gaseous State

This state is characterized by sensitivity of volume change with change of pressure and temperature. It is due to large distance between molecules as compared to their own dimensions. There exists weak vander Waal's forces. (but no force in ideal state).

Ideal Gas: A gas with no intermolecular attractions & having very negligible volume occupied by molecules when compared with volume of gas is termed as ideal gas. An ideal gas is a hypothetical gas and none of the gases known is ideal. These gases are called real gases and they may behave like an ideal gas under specific conditions.

Parameter associated with the ideal gas : P, V, T, n

where

P represents pressure exerted by the gas molecules on the walls of the container assuming negligible intermolecular attractions,

V represents free volume available for motion (equal to the volume of the container),

T represents absolute temperature, n represents no. of moles.

❖ **Conversion factors**

Pressure : atm= 1.013 10^5 Pa 760 mm of Hg 760 torr 1.013 bar

Volume : 1 L 1 dm^3 10 $^3m^3$ 1000 mL 1000 cm^3

Temperature : T_k T_C° 273 $\frac{5}{9}T_F$ 255.22

Instruments for Pressure Measurement

❖ **Barometer:** The instrument used for the measurement of atmospheric pressure is called a barometer.

P $\frac{A \quad h \quad d \quad g}{A}$ d g h

where d density of fluid

h vertical height

g acceleration due to gravity

For More Material Join: @JEEAdvanced_2024

❖ **Manometer:** The instrument used for the measurement of the pressure of a gas is called a manometer. manometer are of two types

(i) Open end (ii) Closed end

Experimental Gas Laws

I. **Boyle's law** $V \quad \frac{1}{P}$ (T & n constant) $P_1V_1 \quad P_2V_2$

II. **Charle's law** $V \quad T$ (P & n constant) $\frac{V_1}{T_1} \quad \frac{V_2}{T_2}$

III. **Gay Lussac's law** $P \quad T$ (V & n constant) $\frac{P_1}{T_1} \quad \frac{P_2}{T_2}$

IV. **Avogadro's law** $V \quad n$ (T & P constant) $\frac{V_1}{n_1} \quad \frac{V_2}{n_2}$

Combined Gas law : $\frac{P_1V_1}{T_1} \quad \frac{P_2V_2}{T_2}$

Equation of State: $PV \quad nRT$

$PV \quad \frac{w}{M}RT$ R is Universal Gas constant,

R 0.0821 atm. L/K. mol 8.314 joule/K. mol 2cal/ K.mol

density of gas (d) $\frac{PM}{RT}$

❖ **Dalton's law of partial pressure:**

$P_{total} \quad P_A \quad P_B \quad .....$

P_A, P_B are partial pressures; P_A mole fraction$_A$ Total pressure

and % of gas in mixture $\frac{\text{Partial pressure}}{\text{Total pressure}}$ 100

$P_{\text{wet gas}} \quad P_{\text{dry gas}} \quad P_{H_2O \text{ vapour}}$ (aqueous tension)

❖ **Amagat's law:** At constant temperature and pressure, the total volume of a mixture of gases is equal to the sum of the partial volumes of the constituent gases.

❖ **Relative humidity:**

% Relative humidity $\frac{\text{Partial pressure of } H_2O(g)}{\text{Vapour pressure of } H_2O(l)}$ 100

❖ **Graham's law of Diffusion or Effusion:**

$r \quad \frac{1}{\sqrt{d}}$ or $r \quad \frac{1}{\sqrt{M}} \quad r \quad \frac{P}{\sqrt{M}}$ [For gases effusing at different pressures]

r is rate of diffusion of any gas, d is density at same temperature.

$\frac{r_1}{r_2} \quad \sqrt{\frac{d_2}{d_1}} \quad \sqrt{\frac{M_2}{M_1}}$; $\frac{V_1 / t_1}{V_2 / t_2} \quad \sqrt{\frac{M_2}{M_1}} \quad \frac{\text{Pressure drop I}}{\text{Pressure drop II}}$

$r \quad \frac{\text{moles diffused}}{\text{time taken}} \quad \frac{\text{distance travelled in a narrow tube}}{\text{time taken}}$

For More Material Join: @JEEAdvanced_2024

❖ **Isotopic Separation Factor:**

After x steps of separation, isotopic separation factor (f) $\frac{(n_1/n_2)_{\text{final}}}{(n_1/n_2)_{\text{initial}}} = \sqrt{\frac{M_2}{M_1}}^{x}$

or $x = \frac{2\log f}{\log \frac{M_2}{M_1}}$

Kinetic Theory of Gases

$PV = \frac{1}{3}mNu^2 = \frac{1}{3}Mu^2$ (For 1 mole)

Types of speeds:

❖ Mean square speed, $u^2 = \frac{u_1^2 + u_2^2 + \ldots + u_N^2}{N}$

❖ Root mean square speed, $u = \sqrt{\frac{3RT}{M}} = \sqrt{\frac{3PV}{nM}} = \sqrt{\frac{3P}{d}}$

❖ Average speed $\frac{u_1 + u_2 + \ldots + u_N}{N} = \sqrt{\frac{8RT}{M}} = \bar{u}$

❖ Most probable speed $\sqrt{\frac{2RT}{M}} = u_{\text{mps}}$

Relationship between three types of speeds

❖ Most probable: average : r.m.s. $\sqrt{2} : \sqrt{\frac{8}{\pi}} : \sqrt{3}$ = 1:1.13:1.22

❖ $u_{\text{rms}} > u_{\text{av}} > u_{\text{mp}}$

❖ Average kinetic energy of a single molecule $= \frac{3}{2}\frac{R}{N}T = \frac{3}{2}kT$

where, k = Boltzman constant = 1.3806×10^{-16} erg deg^{-1}


❖ Total kinetic energy for **one mole** of a gas $= \frac{3}{2}RT$.

❖ Kinetic energy of ***n* moles** of a gas $= n\frac{3}{2}RT$

Maxwells speed distribution Law:

$$dN_u = 4N\left(\frac{M}{2\pi RT}\right)^{3/2} \exp(-Mu^2/2RT)\, u^2 du$$

$$= 4N\left(\frac{m}{2\pi kT}\right)^{3/2} \exp(-mu^2/2kT)\, u^2 du$$




For More Material Join: @JEEAdvanced_2024

Real Gases

Real gases: Gases which do not obey perfect gas laws over a wide range of pressure and temperature are called real or non-ideal gases. These deviations are due to the interactions of molecules with each other and volume occupied by gas molecules. The repulsive forces between molecules assist expansion and attractive forces assist compression.

❖ **Deviation from ideal behaviour**

The curve for the real gas has a tendency to coincide with that of an ideal gas at low pressures when the volume is large. At higher pressures, however deviations are observed.



❖ **van der Waals equation of state**

$$P \quad \frac{an^2}{V^2} \quad (V \quad nb) \quad nRT$$

a, b are van der Waals constants; different for each gas

unit of a atm L^2 mol^{-2} ; S.I. unit Pa m^6 mol^{-2}


unit of b L mol^{-1} ; S.I. unit m^3 mol^{-1}

greater the value of 'a' more easily the gas is liquefiable;

greater the value of 'b' greater the molecular size,

❖ **Compressibility factor**

$$Z \quad \frac{PV}{nRT} \quad \frac{\text{observed molar volume}}{\text{ideal molar volume}}$$



❖ **Interpretation of deviation using van der Waals equation**

(i) At low pressure $Z \quad \frac{PV}{RT} \quad 1 \quad \frac{a}{VRT}$

(ii) At high pressure $Z \quad \frac{PV}{RT} \quad 1 \quad \frac{Pb}{RT}$

(iii) At extremely low pressure $Z \quad \frac{PV}{RT} \quad 1$; $PV \quad nRT$

❖ **Virial equation of state for 1 mole of gas**

$$Z \quad \frac{PV}{RT} \quad 1 \quad B\frac{1}{V} \quad C\frac{1}{V^2} \quad D\frac{1}{V^3} \quad \ldots.$$

B second virial coefficient $b \quad \frac{a}{RT}$ gas dependent

C third virial coefficient b^2

For More Material Join: @JEEAdvanced_2024

❖ **Boyle temperature (T_B):** The temperature at which a real gas obeys Boyle's law (*i.e.*, behaves as an ideal gas) in low pressure range.

$$T_B \quad \frac{a}{bR}$$

❖ **Inversion temperature:** The temperature at which a real gas show no Joule-Thomson effect is called its inversion temperature (T_i)

$$T_i \quad 2T_B \quad \frac{2a}{bR}$$

In Joule-Thomoson's effect, all gases except H_2 and He show cooling effect while H_2 and He show heating effect.

❖ **Critical constants**

(i) Critical temperature (T_c) : T_c is the maximum temperature at which a gas can be liquefied, *i.e.*, the temperature above which a liquid can not exist.


$$T_c \quad \frac{8a}{27Rb}$$

(ii) Critical pressure (P_c) : P_c is the minimum pressure required to cause liquefaction at the temperature T_c.

$$P_c \quad \frac{a}{27b^2}$$

(iii) Critical volume (V_c) : V_c is the volume occupied by one mole of gas at critical temperature and critical pressure.

$$V_c \quad 3b$$



(iv) Compressibility factor under critical state of a gas

$$Z_c \quad \frac{P_c V_c}{RT_c}$$

Now substituting the values of P_c, V_c and T_c we get

$$Z_c \quad \frac{3}{8}$$

❖ **The law of corresponding states**

$$p_r \quad \frac{p}{p_c}, \qquad T_r \quad \frac{T}{T_c} \quad \text{and } V_r \quad \frac{V_m}{V_c}$$

$$(p_r \quad 3 / V_r^2)(3V_r \quad 1) \quad 8T_r$$


$$Z \quad \frac{pV_m}{RT} \quad \frac{(p_r p_c)(V_r V_c)}{R(T_r T_c)} \quad \frac{p_c V_c}{T_c} \quad \frac{p_r V_r}{T_r} \quad \frac{3P_r V_r}{8T_r}$$


For More Material Join: @JEEAdvanced_2024


Level 1


1. Which one of the following statements is not correct about the three states of matter *i.e.,* solid, liquid and gaseous ?
(a) Molecules of a solid possess least energy whereas those of a gas possess highest energy
(b) The density of solids is highest whereas that of gases is lowest
(c) Gases and liquids possess definite volumes
(d) Molecules of a solids possess vibratory motion

2. Which of the following curve does not represent Boyle's law?

(a) 


(b) 


(c) 


(d) 


3. A certain sample of gas has a volume of 0.2 litre measured at 1 atm pressure and 0°C. At the same pressure but at 273°C, its volume will be:
(a) 0.4 litre (b) 0.8 litre (c) 27.8 litres (d) 55.6 litres

4. Among the following curves, which is not according to Charle's law ?

(a) 

(b) 


(c) 


(d) 


5. Initial temperature of an ideal gas is 75°C. At what temperature, the sample of neon gas would be heated to double its pressure, if the initial volume of gas is reduced by 15%?
(a) 319°C (b) 592°C (c) 128°C (d) 60°C


6. Which is correct curve for Charle's law, when the curve is plotted at 0.821 atm pressure for 10 mole ideal gas?

For More Material Join: @JEEAdvanced_2024

(a) 


(b) 


(c) 


(d) 


7. At constant volume, for a fixed number of moles of a gas, the pressure of the gas increases with increase in temperature due to:
(a) increase in the average molecular speed
(b) decrease in rate of collision amongst molecules
(c) increase in molecular attraction
(d) decrease in mean free path

8. Which in not correct curve for Gay-lusacc's law ?

(a) 


(b) 


(c) 

(d) 


9. Three flasks of equal volumes contain CH_4, CO_2 and Cl_2 gases respectively. They will contain equal number of molecules if :
(a) the mass of all the gases is same
(b) the mass of all the gas is same but temperature is different
(c) temperature and pressure of all the flasks are same
(d) temperature, pressure and masses same in the flasks


10. Which is **incorrect** curve for Boyle's law ?

(a) 

(b) 

For More Material Join: @JEEAdvanced_2024


(c) 


(d) 


11. "Equal volumes of all gases at the same temperature and pressure contain equal number of particles." This statement is a direct consequence of :
(a) Avogadro's law (b) Charle's law
(c) ideal gas equation (d) law of partial pressure


12. A 2.24L cylinder of oxygen at 1 atm and 273 K is found to develop a leakage. When the leakage was plugged the pressure dropped to 570 mm of Hg. The number of moles of gas that escaped will be :
(a) 0.025 (b) 0.050 (c) 0.075 (d) 0.09

13. Which of the following curve is correct for an ideal gas ?

(a) 

(b) 

(c) 


(d) 


14. In the equation of state of an ideal gas $PV = nRT$, the value of the universal gas constant is not correct :
(a) 8.314 JK ^{1}mol ^{1} (b) 0.0821 atm L mol ^{1}K ^{1}
(c) 0.8314 bar L mol ^{1}K ^{1} (d) 2 cal mol ^{1}K ^{1}


15. At 0°C and one atm pressure, a gas occupies 100 cc. If the pressure is increased to one and a half-time and temperature is increased by one-third of absolute temperature, then final volume of the gas will be:
(a) 80 cc (b) 88.9 cc (c) 66.7 cc (d) 100 cc


16. 10 g of a gas at 1 atm and 273 K occupies 5 litres. The temperature at which the volume becomes double for the same mass of gas at the same pressure is:
(a) 273 K (b) –273°C (c) 273°C (d) 546°C

17. Which of the following curve does not represent Gay-lusacc's law?

(a) 

(b) 

(c) 

(d) 


For More Material Join: @JEEAdvanced_2024

18. If densities of two gases are in the ratio 1 : 2 and their temperatures are in the ratio 2 : 1, then the ratio of their respective molar mass at certain pressure is:
(a) 1 : 1 (b) 1 : 2 (c) 2 : 1 (d) 4 : 1

19. Two separate bulbs contain ideal gases *A* and *B*. The density of gas *A* is twice that of gas *B*. The molecular mass of *A* is half that of gas *B*. The two gases are at the same temperature. The ratio of the pressure of *A* to that of gas *B* is :
(a) 2 (b) 1/2 (c) 4 (d) 1/4

20. Volume of the air that will be expelled from a vessel of 300 cm^3 when it is heated from 27°C to 37°C at the constant pressure will be:
(a) 310 cm^3 (b) 290 cm^3 (c) 10 cm^3 (d) 37 cm^3

21. *V versus T* curves at constant pressure P_1 and P_2 for an ideal gas are shown in fig. Which is correct?



(a) P_1 P_2 (b) P_1 P_2
(c) P_1 P_2 (d) All of these

22. Two flasks *A* and *B* of 500 mL each are respectively filled with O_2 and SO_2 at 300 K and 1 atm. pressure. The flasks will contain:
(a) the same number of atoms
(b) the same number of molecules
(c) more number of moles of molecules in flask *A* as compared to flask *B*
(d) the same amount of gases

23. 2.8 g of a gas at 1atm and 273K occupies a volume of 2.24 litres. The gas can not be:
(a) O_2 (b) CO (c) N_2 (d) C_2H_4

24. Five grams each of the following gases at 87°C and 750 mm pressure are taken. Which of them will have the least volume ?
(a) HF (b) HCl (c) HBr (d) HI

25. At what pressure a quantity of gas will occupy a volume of 60 mL, if it occupies a volume of 100 mL at a pressure of 720 mm (while temperature is constant) :
(a) 700 mm (b) 800 mm (c) 100 mm (d) 1200 mm

26. At 1 atm and 273 K the density of gas, whose molecular weight is 45, is:
(a) 44.8 g/L (b) 11.4 g/L (c) 2 g/L (d) 3 g/L

27. A small bubble rises from the bottom of a lake, where the temperature and pressure are 8 C and 6.0 atm, to the water's surface, where the temperature is 25°C and pressure is 1.0 atm. Calculate the final volume of the bubble if its initial volume was 2 mL.
(a) 14 mL (b) 12.72 mL (c) 11.31 mL (d) 15 mL

28. Argon is an inert gas used in light bulbs to retard the vaporization of the filament. A certain light-bulb containing argon at 1.25 atm and 18°C is heated to 85°C at constant volume. Calculate its final pressure.
(a) 1.53 atm (b) 1.25 atm (c) 1.35 atm (d) 2 atm

29. Calculate the volume of O_2 at 1 atm and 273 K required for the complete combustion of 2.64 L of acetylene (C_2H_2) at 1 atm and 273 K. $2C_2H_2(g) + 5O_2(g)$ $4CO_2(g) + 2H_2O(l)$
(a) 3.6 L (b) 1.056 L (c) 6.6 L (d) 10 L


For More Material Join: @JEEAdvanced_2024


30. The density of O_2 (*g*) is maximum at :
(a) STP (b) 273 K and 2 atm
(c) 546 K and 1 atm (d) 546 K and 2 atm


31. At 27°C a sample of ammonia gas exerts a pressure of 5.3 atm. What is the pressure when the volume of the gas is reduced to one-tenth of the original value at the same temperature?
(a) 0.53 atm (b) 5.3 atm (c) 53 atm (d) None of these


32. A certain amount of gas at 25°C and at a pressure of 0.80 atm is kept in a glass vessel. Suppose that the vessel can withstand a pressure of 2.0 atm. How high can you raise the temperature of the gas without bursting the vessel?
(a) 745°C (b) 472°C (c) 500°C (d) None of these

33. Which one of these graphs for an ideal gas, the arrow indication is incorrectly marked ?

(a) 

(b) 

(c) 

(d) 

34. The pressure of sodium vapour in a 1.0 L container is 10 torr at 1000°C. How many atoms are in the container?
(a) 9.7 10^{17} (b) 7.6 10^{19}
(c) 4.2 10^{17} (d) 9.7 10^{19}

35. An ideal gaseous mixture of ethane (C_2H_6) and ethene (C_2H_4) occupies 28 litre at 1 atm and 273 K. The mixture reacts completely with 128 gm O_2 to produce CO_2 and H_2O. Mole fraction at C_2H_4 in the mixture is :
(a) 0.6 (b) 0.4 (c) 0.5 (d) 0.8

36. A certain hydrate has the formula $MgSO_4 . xH_2O$. A quantity of 54.2 g of the compound is heated in an oven to drive off the water. If the steam generated exerts a pressure of 24.8 atm in a 2.0 L container at 120°C, calculate *x*.
(a) 2 (b) 5 (c) 6 (d) 7

37. Air entering the lungs ends up in tiny sacs called alveoli. From the alveoli, the oxygen diffuses into the blood. The average radius of the alveoli is 0.0050 cm and the air inside contains 14 per cent oxygen. Assuming that the pressure in the alveoli is 1.0 atm and the temperature is 37°C, calculate the number of oxygen molecules in one of the alveoli.
(a) 6 10^{13} (b) 10^{24} (c) 1.7 10^{22} (d) 1.7 10^{12}

For More Material Join: @JEEAdvanced_2024

38. Nitrogen forms several gaseous oxides. One of them has a density of 1.33 g/L measured at 764 mmHg and 150°C. Write the formula of the compound.
(a) NO (b) N_2O (c) NO_2 (d) N_2O_5

39. Starting out on a trip into the mountains, you inflate the tires on your automobile to a recommended pressure of 3.21 10^5 Pa on a day when the temperature is 5.0 C. You drive to the beach, where the temperature is 28.0°C. Assume that the volume of the tire has increased by 3%. What is the final pressure in the tyres?
(a) 350 Pa (b) 3500 Pa (c) 3.5 10^5 Pa (d) None of these

40. A compressed cylinder of gas contains 1.50 10^3 g of N_2 gas at a pressure of 2.0 10^7 Pa and a temperature of 17.1°C. What volume of gas has been released into the atmosphere if the final pressure in the cylinder is 1.80 10^5 Pa? Assume ideal behaviour and that the gas temperature is unchanged.
(a) 1264 L (b) 126 L (c) 12600 L (d) 45 L

41. A high-altitude balloon is filled with 1.41 10^4 L of hydrogen at a temperature of 21°C and a pressure of 745 torr. What is the volume of the balloon at a height of 20 km, where the temperature is – 48°C and the pressure is 63.1 torr?
(a) 1.274 10^5 L (b) 1.66 10^5 L (c) 1.66 10^4 L (d) None of these

42. The atmospheric pressure on Mars is 0.61 kPa. What is the pressure in mm Hg?
(a) 0.63 (b) 4.6 (c) 6.3 (d) 3.2

43. The density of liquid gallium at 30°C is 6.095 g/mL. Because of its wide liquid range (30 to 2400°C), gallium is used as a barometer fluid at high temperature. What height (in cm) of gallium will be observed on a day when the mercury barometer reads 740 torr?
(The density of mercury is 13.6 g/mL.).
(a) 322 (b) 285 (c) 165 (d) 210

44. A weather balloon is inflated with helium. The balloon has a volume of 100 m^3 and it must be inflated to a pressure of 0.10 atm. If 50 L gas cylinders of helium at a pressure of 100 atm are used, how many cylinders are needed? Assume that the temperature is constant.
(a) 2 (b) 3 (c) 4 (d) 1

45. A balloon contains 14.0 L of air at 760 torr. What will be the volume of the balloon when it is taken to a depth of 10 ft. in a swimming pool? Assume that the temperature of the air and water are equal. (density : Hg = 13.6 g/mL.)
(a) 11.0 (b) 11.3 (c) 10 (d) 10.8

46. A 0.50 L container is occupied by nitrogen at a pressure of 800 torr and a temperature of 0°C. The container can only withstand a pressure of 3.0 atm. What is the highest temperature (°C) to which the container may be heated?
(a) 505 (b) 450 (c) 625 (d) 560


47. Equal volumes of oxygen gas and a second gas weigh 1.00 and 2.375 grams respectively under the same experimental conditions. Which of the following is the unknown gas?
(a) NO (b) SO_2 (c) CS_2 (d) CO


48. A high altitude balloon contains 6.81 g of helium in 1.16 10^4 L at – 23°C. Assuming ideal gas behaviour, how many grams of helium would have to be added to increase the pressure to 4.0 10 ^{3} atm?
(a) 1.27 g (b) 1.58 g (c) 2.68 g (d) 2.23 g


For More Material Join: @JEEAdvanced_2024


49. A 4.40 g piece of solid CO_2 (dry ice) is allowed to sublime in a balloon. The final volume of the balloon is 1.00 L at 300 K. What is the pressure (atm) of the gas?
(a) 0.122 (b) 2.46 (c) 122 (d) 24.6

50. For a closed (not rigid) container containing n 10 moles of an ideal gas fitted with movable, frictionless, weightless piston operating such that pressure of gas remains constant at 0.821 atm. Which graph represents correct variation of log V *vs* log T where V is in litre and T in kelvin.

(a) 

(b) 

(c) 

(d) 


51. The intercept on y-axis and slope of curve plotted between P/T *vs.* T

(For an ideal gas having 10 moles in a closed rigid container of volume 8.21 L. (P Pressure in atm and T Temp. in K, $\log_{10} 2$ 0.30)) are respectively :
(a) 0.01, 0 (b) 0.1, 0 (c) 0.1, 1 (d) 10, 1

52. A He atom at 300 K is released from the surface of the earth to travel upwards. Assuming that constant temperature and g 9.81ms ^{2}, how high will it be before coming to the rest?
(a) 9.53 m (b) 95.3 m (c) 953 m (d) 9.53 10^4 m

53. The density of gas A is twice that to B at the same temperature. The molecular weight of gas B is twice that of A. The ratio of pressure of gas A and B will be :
(a) 1 : 6 (b) 1 : 1 (c) 4 : 1 (d) 1 : 4

54. Two inflated ballons I and II (thin skin) having volume 600 mL and 1500 mL at 300 K are taken as shown in diagram. If maximum volume of inner and outer balloons are 800 mL and 1800 mL respectively then find the balloon which will burst first on gradual heating.
(a) inner balloon
(b) outer balloon
(c) both simultaneously
(d) unpredictable




For More Material Join: @JEEAdvanced_2024

55. An open flask containing air is heated from 300 K to 500 K. What percentage of air will be escaped to the atmosphere, if pressure is keeping constant?
(a) 80 (b) 40 (c) 60 (d) 20


56. The value of universal gas constant R depends on :
(a) temperature of gas (b) volume of gas
(c) number of moles of gas (d) units of volume and pressure

57. A manometer attached to a flask contains with ammonia gas have no difference in mercury level initially as shown in diagram. After sparking into the flask, ammonia is partially dissociated as $2NH_3\,(g)$ $N_2\,(g) + 3H_2\,(g)$ now it have difference of 6 cm in mercury level in two columns, what is partial pressure of $H_2\,(g)$ at equilibrium?



(a) 9 cm Hg (b) 18 cm Hg
(c) 27 cm Hg (d) None of these

58. An ideal gas is collected by downward displacement of water. Select the correct expression for P_{gas} according to the diagram [d_{Hg} 13.6 g/cm^3] :




(a) P_{gas} P_{atm} aq. Tension $\frac{h}{13.6}$ (b) P_{gas} P_{atm} hdg

(c) P_{gas} P_{atm} aq. Tension $\frac{h}{13.6}$ (d) none of these

59. A bubble of gas released at the bottom of a lake increases to four times its original volume when it reaches the surface. Assuming that atmospheric pressure is equivalent to the pressure exerted by a column of water 10 m high, what is the depth of the lake?
(a) 20 m (b) 10 m (c) 30 m (d) 40 m

60. Calculate the number of moles of gas present in the container of volume 10 L at 300 K, if the manometer containing glycerine shows 5 m difference in level as shown diagram.
(Given : $d_{glycerin}$ = 2.72 g/mL; $d_{mercury}$ = 13.6 g/mL)

For More Material Join: @JEEAdvanced_2024



(a) 0.94 mole (b) 0.49 mole (c) 0.64 mole (d) none of these

61. A rigid vessel of volume 0.50 m^3 containing H_2 at 20.5°C and a pressure of 611 10^3 Pa is connected to a second rigid vessel of volume 0.75 m^3 containing Ar at 31.2°C at a pressure of 433 10^3 Pa. A valve separating the two vessels is opened and both are cooled to a temperature of 14.5°C. What is the final pressure in the vessels?

(a) 2 10^5 (b) 3.22 10^5 Pa

(c) 4840 Pa (d) 4.84 10^5 Pa

62. Two glass bulbs *A* and *B* at same temperature are connected by a very small tube having a stop-cock. Bulb *A* has a volume of 100 cm^3 and contained the gas while bulb *B* is empty. On opening the stop-cock, the pressure fell down to 20%. The volume of the bulb *B* is :

(a) 100 cm^3 (b) 200 cm^3

(c) 250 cm^3 (d) 400 cm^3

63. A mixture of C_2H_2 and C_3H_8 occupied a certain volume at 80 mm Hg. The mixture was completely burnt to CO_2 and $H_2O(l)$. When the pressure of CO_2 was found to be 230 mm Hg at the same temperature and volume, the mole fraction of C_3H_8 in the mixture is :

(a) 0.125 (b) 0.875

(c) 0.6 (d) 0.8

64. The total pressure of a mixture of oxygen and hydrogen is 1.0 atm. The mixture is ignited and the water is removed. The remaining gas is pure hydrogen and exerts a pressure of 0.40 atm when measured at the same values of *T* and *V* as the original mixture. What was the composition of the original mixture in mole fraction ?

(a) x_{O_2} 0.2; x_{H_2} 0.8 (b) x_{O_2} 0.4; x_{H_2} 0.6

(c) x_{O_2} 0.6; x_{H_2} 0.4 (d) x_{O_2} 0.8; x_{H_2} 0.2

65. Two closed vessel *A* and *B* of equal volume of 8.21 L are connected by a narrow tube of negligible volume with open valve. The left hand side container is found to contain 3 mole CO_2 and 2 mole of He at 400 K. What is the partial pressure of He in vessel *B* at 400 K?

(a) 2.4 atm (b) 8 atm (c) 12 atm (d) None of these

66. At STP, a container has 1 mole of He, 2 mole Ne, 3 mole O_2 and 4 mole N_2. Without changing total pressure if 2 mole of O_2 is removed. The partial pressure of O_2 will be decreased by :

(a) 26% (b) 40%

(c) 58.33% (d) 66.66%

For More Material Join: @JEEAdvanced_2024

67. 821 mL N_2 (g) was collected over liquid water at 300 K and 1 atm. If vapour pressure of H_2O is 30 torr then moles of N_2 (g) in moist gas mixture is :
(a) 0.39 (b) 0.032
(c) 0.96 (d) 0.0013

68. Let p and p_s be the partial pressure of H_2O (g) and vapour pressure of H_2O (l) respectively. Then the % relative humidity is given by:
(a) $\frac{p_s \quad p}{p_s}$ 100 (b) $\frac{p}{p_s}$ 100 (c) $\frac{p_s}{p}$ 100 (d) ($p \quad p_s$) 100

69. The vapour pressure of water at 80°C is 355 mm of Hg. 1 L vessel contains O_2 at 80°C, which is saturated with water and the total pressure being 760 mm of Hg. The contents of the vessel were pumped into 0.3 L vessel at the same temperature. What is the partial pressure of O_2 ?
(a) 1350 Hg (b) 2178.3 Hg (c) 121.5 Hg (d) 355 Hg

70. Which of the following gaseous mixture does not follow Dalton's law of partial pressure?
(a) SO_2 and Cl_2 (b) CO_2 and N_2 (c) CO and CO_2 (d) CO and N_2

71. Equal masses of methane and oxygen are mixed in an empty container at 25°C. The fraction of the total pressure exerted by oxygen is:
(a) $\frac{2}{3}$ (b) $\frac{1}{3} \quad \frac{273}{298}$ (c) $\frac{1}{3}$ (d) $\frac{1}{2}$

72. A box of 1 L capacity is divided into two equal compartments by a thin partition which are filled with 2g H_2 and 16 g CH_4 respectively. The pressure in each compartment is recorded as P atm. The total pressure when partition is removed will be:
(a) P (b) $2P$ (c) $P/2$ (d) $P/4$

73. If $10^{\ 4}$ dm^3 of water is introduced into a 1.0 dm^3 flask at 300 K, how many moles of water are in the vapour phase when equilibrium is established?
(Given : Vapour pressure of H_2O at 300 K is 3170 Pa; R 8.314 $JK^{\ 1}mol^{\ 1}$)
(a) 1.27 $10^{\ 3}$ mole (b) 5.56 $10^{\ 3}$ mole
(c) 1.53 $10^{\ 2}$ mole (d) 4.46 $10^{\ 2}$ mole

74. At room temperature Dalton's law of partial pressure is not applicable to :
(a) H_2 and N_2 mixture (b) H_2 and Cl_2 mixture
(c) H_2 and CO_2 mixture (d) none of these

75. 56 g of nitrogen and 96 g of oxygen are mixed isothermaly and at a total pressure of 10 atm. The partial pressures of oxygen and nitrogen (in atm) are respectively :
(a) 4, 6 (b) 5, 5 (c) 2, 8 (d) 6, 4

76. The closed containers of the same capacity and at the same temperature are filled with 44 g of H_2 in one and 44 g of CO_2 in the other. If the pressure of carbon dioxide in the second container is 1 atm, then pressure of hydrogen in the first container would be :
(a) 1 atm (b) 10 atm (c) 22 atm (d) 44 atm

77. A jar contains a gas and a few drops of water. The pressure in the jar is 830 mm of Hg. The temperature of the jar is reduced by 1%. The vapour pressure of water at two temperatures are 30 and 25 mm of Hg. Calculate the new pressure in jar.
(a) 792 mm of Hg (b) 817 mm of Hg
(c) 800 mm of Hg (d) 840 mm of Hg

For More Material Join: @JEEAdvanced_2024

78. O_2 and SO_2 gases are filled in ratio of 1 : 3 by mass in a closed container of 3 L at temperature of 27°C. The partial pressure of O_2 is 0.60 atm, the concentration of SO_2 would be
(a) 0.36 mol/L (b) 0.036 mol/L (c) 3.6 mol/L (d) 36 mol/L

79. A gaseous mixture contains three gases *A, B* and *C* with a total number of moles of 10 and total pressure of 10 atm. The partial pressure of *A* and *B* are 3 atm and 1 atm respectively. If *C* has molecular weight of 2 g/mol. then, the weight of *C* present in the mixture will be :
(a) 8 g (b) 12 g (c) 3 g (d) 6 g


80. A rigid container contains 5 mole H_2 gas at some pressure and temperature. The gas has been allowed to escape by simple process from the container due to which pressure of the gas becomes half of its initial pressure and temperature become (2/3) rd of its initial. The mass of gas remaining is :
(a) 7.5 g (b) 1.5 g (c) 2.5 g (d) 3.5 g

81. Pressure of 1 g ideal gas *X* at 300 K is 2 atm. When 2 g of another gas *Y* is introduced in the same vessel at same temperature, the pressure become 3 atm. The correct relationship between molar mass of *X* and *Y* is :
(a) M_Y $2\,M_X$ (b) M_Y $4\,M_X$
(c) M_X $4\,M_Y$ (d) None of these

82. Dry ice is solid carbon dioxide. A 0.050 g sample of dry ice is placed in an evacuated 4.6 L vessel at 30°C. Calculate the pressure inside the vessel after all the dry ice has been converted to CO_2 gas.
(a) 6.14 atm (b) 0.614 atm (c) 0.0614 atm (d) 6.14 10^{-3} atm

83. A mixture of helium and neon gases is collected over water at 28.0°C and 745 mmHg. If the partial pressure of helium is 368 mmHg, what is the partial pressure of neon?
(Vapour pressure of water at 28 C = 28.3 mmHg)
(a) 348.7 mmHg (b) 377 mmHg (c) 384.7 mmHg (d) none of these

84. Consider the following apparatus. Calculate the partial pressure of helium after the opening valve. The temperature remains constant at 16°C.



(a) 0.164 atm (b) 1.64 atm (c) 0.328 atm (d) 1 atm

85. Oxygen gas generated by the decomposition of potassium chlorate is collected over water. The volume of oxygen collected at 24°C and atmospheric pressure of 760 mmHg is 128 mL. Calculate the mass of oxygen gas obtained. The vapour pressure of the water vapour at 24°C is 22.4 mm Hg.
(a) 1.36 g (b) 1.52 g (c) 0.163 g (d) 1.63 g

For More Material Join: @JEEAdvanced_2024

86. The quantity $\frac{PV}{k_B T}$ represents the (k_B : Boltzmann constant)

(a) number of particles of the gas (b) mass of the gas
(c) number of moles of the gas (d) translation energy of the gas

87. Which of the following statements about kinetic energy (K.E.) is true?
(a) All objects moving with the same velocity have the same K.E.
(b) The K.E. of a body will quadruple if its velocity doubles
(c) As the velocity of a body increases, its K.E. decreases
(d) The K.E. of a body is independent of its mass

88. The Ne atom has 10 times the mass of H_2 molecule. Which of the following statements is true?
I. At 25°C both of them have the same kinetic energy.
II. Ten moles of H_2 would have the same volume as 1 mole of Ne at same temp. and pressure.
III. One mole of Ne exerts the same pressure as one mole of H_2 at STP.
IV. A H_2 molecule travels 10 times faster than Ne atom at same temperature.
V. At STP, one litre of Ne has 10 times the density of 1 litre of H_2.
(a) II, IV, V (b) I, III, V (c) I, II, III (d) I, II

89. Which of the following is NOT a postulate of the kinetic molecular theory of gases?
(a) The gas molecules possess a volume that is negligibly small compared to the container
(b) The pressure and volume of a gas are inversely related
(c) Gases consist of discrete particles that are in constant chaotic motion
(d) The average kinetic energy of the molecules is directly proportional to the absolute temperature

90. Which one of the following relationships when graphed does not give a straight line for helium gas?
I. K.E. and T at constant pressure and volume
II. P *v/s* V at constant temperature for a constant mass
III. V *v/s* $1/T$ at constant pressure for a constant mass
(a) II (b) II and III (c) III (d) I

91. Consider three one-litre flasks labeled *A*, *B* and *C* filled with the gases NO, NO_2, and N_2O, respectively, each at 1 atm and 273 K. In which flask do the molecules have the highest average kinetic energy?
(a) Flask *C* (b) All are the same (c) Flask *A* (d) None

92. Which of the following statements is false?
(a) The product of pressure and volume of fixed amount of a gas is independent of temperature.
(b) Molecules of different gases have the same kinetic energy at a given temperature.
(c) The gas equation is not valid at high pressure and low temperature.
(d) The gas constant per molecule is known as Boltzmann's constant.

93. Which is not correct in terms of kinetic theory of gases?
(a) Gaseous particles are considered as point mass.
(b) The gaseous molecules are in random motion.
(c) When gaseous molecules collide, they lose energy.
(d) When the gas is heated , the molecules moves faster.

For More Material Join: @JEEAdvanced_2024

94. Two flasks *A* and *B* have equal volumes. *A* is maintained at 300 K and *B* at 600 K. *A* contains H_2 gas, *B* has an equal mass of CO_2 gas. Find the ratio of total K.E. of gases in flask *A* to that of *B*.

(a) 1 : 2 (b) 11 : 1
(c) 33 : 2 (d) 55 : 7

95. Kinetic energy and pressure of a gas of unit volume are related as :

(a) $P \quad \frac{2}{3}E$ (b) $P \quad \frac{3}{2}E$ (c) $P \quad \frac{E}{2}$ (d) $P \quad 2E$

96. Two flask *A* and *B* of equal volumes maintained at temperature 300 K and 700 K contain equal mass of He(*g*) and $N_2(g)$ respectively. What is the ratio of translational kinetic energy of gas in flask *A* to that of flask *B*?

(a) 1 : 3 (b) 3 : 1 (c) 3 : 49 (d) None of these

97. Which of the following change is observed occurs when a substance *X* is converted from liquid to vapour phase at the standard boiling point?

I. Potential energy of the system decreases

II. The distance between molecules increases

III. The average kinetic energy of the molecules in both phases are equal

(a) I only (b) II only
(c) III only (d) II and III only

98. A mixture of Ne and Ar kept in a closed vessel at 250 K has a total K.E. = 3 kJ. The total mass of Ne and Ar is 30 g. Find mass % of Ne in gaseous mixture at 250 K.

(a) 61.63 (b) 38.37 (c) 50% (d) 28.3

99. In two vessels of 1 litre each at the same temperature 1g of H_2 and 1g of CH_4 are taken. For these gases:

(a) V_{rms} values will be same (b) Kinetic energy per mol will be same
(c) Total kinetic energy will same (d) Pressure will be same

100. Four particles have speed 2,3,4 and 5 cm/s respectively. Their rms speed is:

(a) 3.5 cm/s (b) (27/2) cm/s (c) $\sqrt{54}$ cm/s (d) $(\sqrt{54}/2)$ cm/s

101. A gaseous mixture contains 4 molecules with a velocity of 6 cm sec ^{1}, 5 molecules with a velocity of 2 cm sec ^{1} and 10 molecules with a velocity of 3 cm sec ^{1}. What is the rms speed of the gas:

(a) 2.5 cm sec ^{1} (b) 1.9 cm sec ^{1} (c) 3.6 cm sec ^{1} (d) 4.6 cm sec ^{1}

102. The ratio between the root mean square velocity of H_2 at 50 K and that of O_2 at 800 K is:

(a) 4 (b) 2 (c) 1 (d) 1/4

103. If C_1, C_2, C_3 represent the speeds of n_1, n_2, n_3 molecules respectively, then the root mean square speed will be:

(a) $\sqrt{\frac{n_1C_1^2 \quad n_2C_2^2 \quad n_3C_3^2}{n_1 \quad n_2 \quad n_3}}$ (b) $\sqrt{\frac{(n_1 \quad n_2 \quad n_3 \quad)^2}{n_1C_1^2 \quad n_2C_2^2 \quad n_3C_3^2}}$


(c) $\sqrt{\frac{(n_1C_1)}{n_1} \quad \frac{(n_2C_2)}{n_2} \quad \frac{(n_3C_3)}{n_3}}$ (d) $\sqrt{\frac{(n_1C_1 \quad n_2C_2 \quad n_2C_3 \quad)^2}{n_1 \quad n_2 \quad n_3}}$

104. The root mean square speed of hydrogen is $\sqrt{5}$ times than that of nitrogen. If *T* is the temperature of the gas, then :

For More Material Join: @JEEAdvanced_2024

(a) T_{H_2} T_{N_2} (b) T_{H_2} T_{N_2}
(c) T_{H_2} T_{N_2} (d) T_{H_2} $\sqrt{7}\,T_{N_2}$

105. At a definite temperature (T), the distribution of speeds is given by the curve. In the curve points A, B and C indicates the speeds corresponding to :



(a) most probable, average and root mean square speeds
(b) average, root mean square and most probable speeds
(c) root mean square, average and most probable speeds
(d) most probable, root mean square and average speeds

106. The most probable speed of 8 g of H_2 is 200 ms ^{1}. Average kinetic energy (neglect rotational and vibrational energy) of H_2 gas is :
(a) 480 J (b) 240 J (c) 120 J (d) none of these

107. At what temperature will average speed of the molecules of the second member of the series C_nH_{2n} be the same to that of Cl_2 at 627°C?
(a) 259.4 K (b) 400 K
(c) 532.4 K (d) None of these

108. If u_{rms} of a gas is 30 $R^{1/2}$ ms ^{1} at 27°C then the molar mass of gas is :
(a) 0.02 kg/mol (b) 0.001 kg/mol
(c) 0.003 kg/mol (d) 1 kg/mol

109. 6 10^{22} gas molecules each of mass 10 ^{24} kg are taken in a vessel of 10 litre. What is the pressure exerted by gas molecules? The root mean square speed of gas molecules is 100 m/s.
(a) 20 Pa (b) 2 10^{4} Pa (c) 2 10^{5} Pa (d) 2 10^{7} Pa

110. At what temperature will most probable speed of the molecules of the second member of homologous series $C_nH_{2n\ 2}$ be the same as that of SO_2 at 527°C.
(a) 500°C (b) 727°C (c) 227°C (d) None of these

111. The root mean square speed of 8 g of He is 300 ms ^{1}. Total kinetic energy of He gas is :
(a) 120 J (b) 240 J (c) 360 J (d) None of these

112. If $T_{1,}$ T_2 and T_3 are the temperatures at which the u_{rms}, $u_{average}$, u_{mp} of oxygen gas are all equal to 1500 m/s then the correct statement is :
(a) T_1 T_2 T_3 (b) T_1 T_2 T_3
(c) T_1 T_2 T_3 (d) None of these

113. The density of a gas filled in electric lamp is 0.75 kg/m ^{3}. After the lamp has been switched on, the pressure in it increases from 4 10^{4} Pa to 9 10^{4} Pa. What is increases in u_{rms} ?
(a) 100 (b) 200 (c) 300 (d) None of these

For More Material Join: @JEEAdvanced_2024

114. The root mean square speed of an ideal gas at constant pressure varies with density (d) as :
(a) d^2 (b) d (c) $\sqrt{d}$ (d) $1/\sqrt{d}$

115. The ratio among most probable speed, mean speed and root mean square speed is given by
(a) 1 : 2 : 3 (b) $1 : \sqrt{2} : \sqrt{3}$
(c) $\sqrt{2} : \sqrt{3} : \sqrt{8/}$ (d) $\sqrt{2} : \sqrt{8/} : \sqrt{3}$

116. The average speed at temperature T°C of $CH_4(g)$ is $\sqrt{\frac{28}{88}}$ 10^3 ms ^{1}. What is the value of T ?
(a) 240.55°C (b) – 32.45°C
(c) 3000°C (d) – 24.055°C

117. At what temperature most probable speed of SO_2 molecules have the same value as root mean square speed of O_2 molecules at 300 K?
(a) 150 K (b) 600 K
(c) 750 K (d) 900 K

118. The rms speed of N_2 gas molecules is u. If the temperature is doubled and the nitrogen molecules dissociate into nitrogen atoms, the rms speed becomes:
(a) $u/2$ (b) $2u$ (C) $4u$ (D) $14u$

119. The rate of diffusion of a gas is proportional to:
(a) $\frac{P}{\sqrt{M}}$ (b) $\frac{P}{M}$ (c) $\sqrt{\frac{P}{M}}$ (d) $\frac{\sqrt{P}}{M}$

120. At constant volume and temperature conditions, the rates of diffusion r_A and r_B of gases A and B having densities $_A$ and $_B$ are related by the expression :
(a) $r_A \quad r_B.(\ _A/\ _B)^2$ (b) $r_A \quad r_B(\ _B/\ _A)^{1/2}$
(c) $r_A \quad (r_B.\ _B/\ _A)^{1/2}$ (d) $r_A \quad r_B(\ _A/\ _B)^{1/2}$

121. What is the ratio of diffusion rate of oxygen to hydrogen?
(a) 1 : 4 (b) 4 : 1 (c) 1 : 8 (d) 8 : 1

122. The molecular weight of a gas which diffuses through a porous plug at $1/6^{th}$ of the speed of hydrogen under identical conditions is:
(a) 27 (b) 72
(c) 36 (d) 48

123. X mL of H_2 gas effuses through a hole in a container in 5 seconds. The time taken for the effusion of the same volume of the gas specified below under identical condition is:
(a) 10 sec : He (b) 20 sec : O_2
(c) 25 sec : CO (d) 55 sec : CO_2

124. At identical temperature and pressure, the rate of diffusion of hydrogen gas is $3\sqrt{3}$ times that of a hydrocarbon having molecular formula $C_nH_{2n\ 2}$. What is the value of n?
(a) 1 (b) 4 (c) 3 (d) 8

125. Calculate relative rate of effusion of O_2 to CH_4 from a container containing O_2 and CH_4 in 3 : 2 mass ratio.
(a) $\frac{3\sqrt{2}}{4}$ (b) $\frac{3}{4\sqrt{2}}$ (c) $\frac{3}{\sqrt{2}}$ (d) none of these

For More Material Join: @JEEAdvanced_2024

126. Calculate relative rates of effusion of SO_2 to CH_4, in the mixture obtained by effusing out a mixture with initial molar ratio $\frac{n_{SO_2}}{n_{CH_4}}$ $\frac{8}{1}$ for three effusing steps.

(a) 2 : 1 (b) 1 : 4
(c) 1 : 2 (d) none of these

127. A gaseous mixture containing He, CH_4 and SO_2 was allowed to effuse through a fine hole. If mixture contain He, CH_4 and SO_2 in 1 : 2 : 3 mole ratio, calculate the molar ratio of gases effusing out initially.

(a) 2 : 2 : 3 (b) 6 : 6 : 1 (c) $\sqrt{2} : \sqrt{2} : 3$ (d) 4 : 4 : 3

128. 80 mL of O_2 takes 2 minute to pass through the hole. What volume of SO_2 will pass through the hole in 3 minute?

(a) $\frac{120}{\sqrt{2}}$ mL (b) 120 $\sqrt{2}$ mL (c) $\frac{12}{\sqrt{2}}$ mL (d) None of these

129. When two cotton plugs, one moistened with ammonia and the other with hydrochloric acid, are simulataneously inserted into opposite ends of a glass tube 87.0 cm long, a white ring of NH_4Cl forms where gaseous NH_3 and gaseous HCl first come into contact.

$$NH_3(g) + HCl(g) \qquad NH_4Cl(s)$$

At what distance from the ammonia-moistened plug does this occur?

(a) 51.7 cm from NH_3 end (b) 51.7 cm from HCl end
(c) 43.5 at mid point (d) None of these

130. Dimethyl ether decomposes as

$$CH_3OCH_3(g) \qquad CH_4(g) + CO(g) + H_2(g)$$

When CH_3OCH_3 decomposes to 20% extent under certain conditions, what is the ratio of diffusion of pure CH_3OCH_3 with methane?

(a) 0.59 : 1 (b) 1.18 : 1 (c) 2.36 : 1 (d) 1.77 : 1

131. 4 gm of sulphur dioxide gas diffuses from a container in 8 min. Mass of helium gas diffusing from the same container over the same time interval is :

(a) 0.5 g (b) 1 g (c) 2 g (d) None of these

132. Under identical conditions of pressure and temperature, 4 L of gaseous mixture (H_2 and CH_4) effuses through a hole in 5 min whereas 4 L of a gas *X* of molecular mass 36 takes to 10 min to effuse through the same hole. The mole ratio of $H_2 : CH_4$ in the mixture is :

(a) 1 : 2 (b) 2 : 1 (c) 2 : 3 (d) 1 : 1

133. A balloon weighing 50 kg is filled with 685 kg of helium at 1 atm pressure and 25°C. What will be its pay load if it displaced 5108 kg of air?

(a) 4373 kg (b) 4423 kg
(c) 5793 kg (d) none of these

134. According to the law of equipartition of energy, the energy associated with each degree of freedom is :

(a) $\frac{1}{3} K_B T$ (b) $\frac{1}{2} K_B T$
(c) $K_B T$ (d) $\frac{3}{2} K_B T$

For More Material Join: @JEEAdvanced_2024

135. Calculate (ratio of C_p and C_v) for triatomic linear gas at high temperature. Assume that the contribution of vibrational degree of freedom is 75% :
(a) 1.222 (b) 1.121
(c) 1.18 (d) 1.33

136. If one mole each of a monoatomic and diatomic gases are mixed at low temperature then C_p/C_v ratio for the mixture is :
(a) 1.40 (b) 1.428
(c) 1.5 (d) 1.33

137. If one mole of a mono-atomic gas (5/3) is mixed with one mole of a diatomic gas (7/5), the value of for the mixture is :
(a) 1.4 (b) 1.5
(c) 1.53 (d) 3.07

138. Select the correct statement :
(a) Internal energy of a real gas at a given temperature increases as the volume increases
(b) Internal energy of an ideal gas at given temperature increase as the volume increases
(c) Internal energy of an ideal gas molecules is not a function of temperature
(d) The internal energy of a real gas at a constant temperature is independent of change in volume

139. Which gas shows real behaviour?
(a) 16 g O_2 at 1 atm & 273 K occupies 11.2 L
(b) 1 g H_2 in 0.5 L flask exerts pressure of 24.63 atm at 300 K
(c) 1 mole NH_3 at 300 K and 1 atm occupies volume 22.4 L
(d) 5.6 L of CO_2 at 1 atm & 273 K is equal to 11 g

140. Consider a real gas placed in a container. If the intermolecular attractions are supposed to disappear suddenly which of the following would happen?
(a) Pressure decreases (b) Pressure increases
(c) Pressure remains unchanged (d) Gas collapses

141. The pressure of a real gas is less than the pressure of an ideal gas because of :
(a) increase in number of collisions (b) finite size of molecule
(c) increase in KE of molecules (d) intermolecular forces of attraction

142. A gas behaves like an ideal gas at :
(a) high pressure and low temperature (b) low pressure and high temperature
(c) high pressure and high temperature (d) low pressure and low temperature

143. If temperature and volume are same, the pressure of a gas obeying van der Waal's equation is :
(a) less than that of an ideal gas (b) more than that of an ideal gas
(c) same as that of an ideal gas (d) none of these

144. 1 mole of each of X_1, X_2, X_3 with van der Waal's constants a (in atm L^3 mol^{-2}) 1.0, 3.8, 2.1 respectively is kept separately in three different vessels of equal volume at identical temperature. Their pressures are observed to $P_1, P_2,$ and P_3 respectively. On the basis of this data alone, select the correct option (neglect the effect of '*b*'):
(a) P_1 P_2 P_3 (b) P_2 P_1 P_3
(c) P_2 P_3 P_1 (d) P_1 P_2 P_3

For More Material Join: @JEEAdvanced_2024

145. A gas obeys the equation of state $P(V \quad b) \quad RT$ (The parameter b is a constant). The slope for an isochore will be :
(a) negative (b) zero
(c) $R/(V \quad b)$ (d) R/P

146. van der Waals constant b of helium is 24 mL mol ^{1}. Find molecular diameter of helium.
(a) $1.335 \quad 10^{10}$ cm (b) $1.335 \quad 10^{8}$ cm
(c) $2.67 \quad 10^{8}$ cm (d) $4.34 \quad 10^{8}$ cm

147. Which of the following can be most readily liquefied?
(Given: value of 'a' for NH_3 4.17 atm .L^3mol ^{2}, CO_2 3.59 atm .L^3mol ^{2}, SO_2 6.71 atm .L^3mol ^{2}, Cl_2 6.49 atm .L^3mol ^{2})
(a) NH_3 (b) Cl_2
(c) SO_2 (d) CO_2

148. For which of the following gases should the correction for the molecular volume be largest : CO, CO_2, NH_3 or SF_6 ?
(a) CO (b) CO_2
(c) NH_3 (d) SF_6

149. Under which of the following sets of conditions is a real gas expected to deviate from ideal behaviour?
(I) High pressure, small volume
(II) High temperature, low pressure
(III) Low temperature, high pressure
(a) only I (b) only II (c) only III (d) I and III both

150. For a certain gas which deviates a little from ideal behaviour, a plot between $P/$ *vs* P was found to be non-linear. The intercept on y-axis will be :
(a) $\frac{RT}{M}$ (b) $\frac{M}{RT}$ (c) $\frac{MZ}{RT}$ (d) $\frac{R}{TM}$

151. At low pressure, the van der Waal's equation become :
(a) $PV_m \quad RT$ (b) $P(V_m \quad b) \quad RT$
(c) $P \quad \frac{a}{V_m^2} \quad V_m \quad RT$ (d) $P \quad \frac{RT}{V_m} \quad \frac{a}{V_m^2}$

152. At low pressure, if $RT \quad 2\sqrt{a\,P}$, then the volume occupied by a real gas is :
(a) $\frac{2RT}{P}$ (b) $\frac{2P}{RT}$ (c) $\frac{RT}{2P}$ (d) $\frac{RT}{P}$

153. For a gas deviation from ideal behaviour is maximum at :
(a) 0°C and 1.0 atm (b) 100°C and 2.0 atm
(c) – 13°C and 1.0 atm (d) – 13°C and 2.0 atm

154. At low pressures, van der Waals' equation is written as $P \quad \frac{a}{V^2} \quad V \quad RT$. The compressibility factor is then equal to:
(a) $1 \quad \frac{a}{RTV}$ (b) $1 \quad \frac{RTV}{a}$
(c) $1 \quad \frac{a}{RTV}$ (d) $1 \quad \frac{RTV}{a}$

For More Material Join: @JEEAdvanced_2024

155. The compressibility factor for a real gas at high pressure is :

(a) 1 (b) 1 $\frac{Pb}{RT}$

(c) 1 $\frac{Pb}{RT}$ (d) 1 $\frac{RT}{Pb}$

156. The compressibility of a gas is greater than unity at 1 atm and 273 K. Therefore :

(a) V_m 22.4 L (b) V_m 22.4 L

(c) V_m 22.4 L (d) V_m 44.8 L

157. At 273 K temperature and 9 atm pressure, the compressibility for a gas is 0.9. The volume of 1 milli-mole of gas at this temperature and pressure is :

(a) 2.24 litre (b) 0.020 mL

(c) 2.24 mL (d) 22.4 mL

158. The compressibility factor for nitrogen at 330 K and 800 atm is 1.90 and at 570 K and 200 atm is 1.10. A certain mass of N_2 occupies a volume of 1 dm^3 at 330 K and 800 atm. Calculate volume occupied by same quantity of N_2 gas at 570 K and 200 atm :

(a) 1 L (b) 2 L

(c) 3 L (d) 4 L

159. Consider the equation Z $\frac{PV}{RT}$. Which of the following statements is correct?

(a) When Z 1, real gases are easier to compress than the ideal gas

(b) When Z 1, real gases get compressed easily

(c) When Z 1, real gases are difficult to compress

(d) When Z 1, real gases are difficult to compress

160. What is the compressibility factor (Z) for 0.02 mole of a van der Waals' gas at pressure of 0.1 atm. Assume the size of gas molecules is negligible.

Given : RT 20 L atm mol^{-1} and a 1000 atm L^2 mol^{-2}

(a) 2 (b) 1

(c) 0.02 (d) 0.5

161. Consider the following statements. If the van der Waal' s parameters of two gases are given as

	a/dm^6 bar mol^{-2}	b/dm^3 mol^{-1}
Gas A	6.5	0.055
Gas B	2	0.01

then :

1. Critical volume of A Critical volume of B
2. Critical pressure A Critical pressure of B
3. Critical temperature of A Critical temperature of B

Which of the above statements is/are incorrect?

(a) 1 alone (b) 1 and 2

(c) 1, 2 and 3 (d) 2 and 3

For More Material Join: @JEEAdvanced_2024

162. The van der Waals parameters for gases *W, X, Y* and *Z* are

Gas	a (atm L^2 mol^{-2})	b (L mol^{-1})
W	4.0	0.027
X	8.0	0.030
Y	6.0	0.032
Z	12.0	0.027

Which one of these gases has the highest critical temperature?
(a) *W* (b) *X* (c) *Y* (d) *Z*

163. Pressure remaining the constant, the volume of a given mass of an ideal gas increases for every degree centigrade rise in temperature by definite fraction of its volume at:
(a) 0°C
(b) its critical temperature
(c) absolute zero
(d) its Boyle temperature

164. The critical temperature of a substance is:
(a) the temperature above which the substance undergoes decomposition
(b) the temperature above which a substance can exist only in gaseous state
(c) boiling point of the substance
(d) all are wrong

165. The excluded volume of a gas will be larger, if $\frac{T_C}{P_C}$ is :
(a) small
(b) large
(c) equal to 1
(d) less than unity

166. Select incorrect statement :
(a) we can condense vapour simply by applying pressure
(b) to liquefy a gas one must lower the temperature below T_C and also apply pressure
(c) at T_C, there is no distinction between liquid and vapour state hence density of the liquid is nearly equal to density of the vapour
(d) However great the pressure applied, a gas cannot be liquified below it's critical temp.

167. The correct order of temperature of a real gas is :
(I) Boyle's temperature
(II) Critical temperature
(III) Inversion temperature
(a) III > I > II
(b) I > II > III
(c) II > I > III
(d) I > III > II

168. The temperature at which the second virial coefficient of real gas is zero is called :
(a) Critical temperature
(b) Triple point
(c) Boiling point
(d) Boyle's temperature

169. The van der Waals' equation for one mole may be expressed as

$$V_m^3 \quad b \quad \frac{RT}{p} \quad V_m^2 \quad \frac{aV_m}{p} \quad \frac{ab}{p} \quad 0$$

For More Material Join: @JEEAdvanced_2024

where V_m is the molar volume of the gas. Which of the following is incorrect?
(a) For a temperature less than T_C, V has three real roots
(b) For a temperature less than T_C, V has three imaginary roots
(c) For a temperature equal to T_C all three roots of V are real and identical
(d) On increasing the temp. (T T_C), the three roots become closer to one another

170. For a gas obeying the van der Waals' equation, at the critical point

(a) both $\left(\frac{P}{V}\right)_T$ and $\left(\frac{^2P}{V^2}\right)_T$ are zero
(b) only $\left(\frac{^2P}{V^2}\right)_T$ is not zero
(c) $\left(\frac{P}{V}\right)_T$ is zero but $\left(\frac{^2P}{V^2}\right)_T$ is non-zero
(d) $\left(\frac{P}{V}\right)_T$ is non-zero but $\left(\frac{^2P}{V^2}\right)_T$ is zero

171. However great the pressure, a gas cannot be liquified above its :
(a) Boyle temperature
(b) Inversion temperature
(c) Critical temperature
(d) Room temperature

172. The temperature at which real gases obey the ideal gas laws over a wide range of low pressure is called:
(a) Critical temperature
(b) Inversion temperature
(c) Boyle temperature
(d) Reduced temperature

173. Inversion temperature T_i $\frac{2a}{Rb}$ is defined as the temperature above which if a gas is expanded adiabatically it gets warmed up but if temperature of gas is lower than T_i then it will cool down. What will happen to a gas if it is adiabatically expanded at 600 K if its Boyle's temperature is 290 K?
(a) Heating
(b) Cooling
(c) Constant
(d) None

174. The van der Waals' equation of law of corresponding states for 1 mole of gas is :


(a) P_r $\frac{3}{V_r^2}$ $(3V_r$ 1) 8 T_r
(b) P_r $\frac{3}{V_r^2}$ $(3V_r$ 1) 8 T_r
(c) P_r $\frac{3}{V_r^2}$ $(3V_r$ 1) 8 T_r
(d) P_r $\frac{3}{V_r^2}$ $(3V_r$ 1) 8

175. Calculate the volume occupied by 16 gram O_2 at 300 K and 8.31 MPa if $\frac{P_cV_c}{RT_c}$ 3/8 and $\frac{P_rV_r}{T_r}$ 2.21 (Given : R 8.314 J/K-mol)
(a) 125.31 mL
(b) 124.31 mL
(c) 248.62 mL
(d) none of these

For More Material Join: @JEEAdvanced_2024


Level 2

1. Consider the composite system, which is held at 300 K, shown in the following figure. Assuming ideal gas behavior, calculate the total pressure if the barriers separating the compartments are removed. Assume that the volume of the barriers is negligible. (Given : R 0.082 atm L/mol.K)



(a) 1 atm (b) 2 atm (c) 2.3 atm (d) 3.2 atm

2. 11 moles of N_2 and 12 moles of H_2 mixture reacted in 20 litre vessel at 800 K. After equilibrium was reached, 6 mole of H_2 was present. 3.58 litre of liquid water is injected in equilibrium mixture and resultant gaseous mixture suddenly cooled to 300K. What is the final pressure of gaseous mixture? Neglect vapour pressure of liquid solution. Assume (i) all NH_3 dissolved in water (ii) no change in volume of liquid (iii) At 300 K no reaction takes place between N_2 and H_2



(a) 18.47 atm (b) 60 atm (c) 22.5 atm (d) 45 atm

3. Two vessels connected by a valve of negligible volume. One container (I) has 2.8 g of N_2 at temperature T_1(K). The other container (II) is completely evacuated. The container (I) is heated to T_2 (K) while container (II) is maintained at $T_2/3$ (K). volume of vessel (I) is half that of vessel (II). If the valve is opened then what is the mass ratio of N_2 in both vessel (W_I/W_{II})?

(a) 1 : 2 (b) 1 : 3 (c) 1 : 6 (d) 3 : 1

4. A mixture of $NH_3(g)$ and $N_2H_4(g)$ is placed in a sealed container at 300 K. The total pressure is 0.5 atm. The container is heated to 1200 K at which time both substances decompose completely according to the equations

$$2NH_3(g) \qquad N_2(g) + 3H_2(g) \text{ and } N_2H_4(g) \qquad N_2(g) + 2H_2(g)$$

After decomposition is complete, the total pressure at 1200 K is found to be 4.5 atm. Find the mole% of N_2H_4 in the original mixture :

(a) 20% (b) 25% (c) 50% (d) 75%

For More Material Join: @JEEAdvanced_2024

5. Correct expression for density of an ideal gas mixture of two gases 1 and 2, where m_1 and m_2 are masses and n_1 and n_2 are moles and M_1 and M_2 are molar masses.

(a) $d \quad \frac{(m_1 \quad m_2)}{(M_1 \quad M_2)}$ (b) $d \quad \frac{(m_1 \quad m_2)}{(n_1 \quad n_2)} \frac{P}{RT}$

(c) $d \quad \frac{(n_1 \quad n_2)}{(m_1 \quad m_2)} \quad \frac{P}{RT}$ (d) None of these

6. Two closed vessel A and B of equal volume containing air at pressure P_1 and temperature T_1 are connected to each other through a narrow open tube. If the temperature of one is now maintained at T_1 and other at T_2 (where $T_1 \quad T_2$) then that what will be the final pressure?

(a) $\frac{T_1}{2P_1T_2}$ (b) $\frac{2P_1T_2}{T_1 \quad T_2}$ (c) $\frac{2P_1T_1}{T_1 \quad T_2}$ (d) $\frac{2P_1}{T_1 \quad T_2}$

7. A balloon containing 1 mole air at 1 atm initially is filled further with air till pressure increases to 4 atm. The initial diameter of the balloon is 1 m and the pressure at each stage is proportional to diameter of the balloon. How many moles of air added to change the pressure from 1 atm to 4 atm.

(a) 80 (b) 257 (c) 255 (d) 256

8. If Pd *vs*. P(where P denotes pressure in atm and d denotes density in gm/L) is plotted for He gas (assume ideal) at a particular temperature. If $\frac{d}{dP}(Pd)_{P \quad 8.21 \text{ atm}} \quad 5$, then the temperature will be

(a) 160 K (b) 320 K (c) 80 K (d) none of these


9. What is the density of wet air with 75% relative humidity at 1 atm and 300 K? Given : vapour pressure of H_2O is 30 torr and average molar mass of air is 29 g mol ^{1}.

(a) 1.614 g/L (b) 0.96 g/L (c) 1.06 g/L (d) 1.164 g/L

10. 7 moles of a tetra-atomic non-linear gas 'A' at 10 atm and T K are mixed with 6 moles of another gas B at $\frac{T}{3}$ K and 5 atm in a closed, rigid vessel without energy transfer with surroundings. If final temperature of mixture was $\frac{5T}{6}$ K, then gas B is? (Assuming all modes of energy are active)

(a) monoatomic (b) diatomic (c) triatomic (d) tetra atomic

11. Three closed rigid vessels, A, B and C, which initially contain three different gases at different temperatures are connected by tube of negligible volume, without any energy transfer with surroundings. The vessel A contain 2 mole Ne gas, at 300 K, vessel 'B' contain 2 mole SO_2 gas at 400 K and vessel 'C' contain 3 mole CO_2 gas at temperature 500 K. What is the final pressure (in atm) attained by gases when all valves of connecting three vessels are opened and additional 15.6 kcal heat supplied to vessels through valve. The volume of A, B and C vessel is 2, 2 and 3 litre respectively



For More Material Join: @JEEAdvanced_2024

Given : R 2 calorie/mol-K; C_v (Ne) $3/2\,R$, C_v (CO) $5/2\ R$ and C_v (SO_2) $3\,R$

(a) 73.89 atm (b) 67.31 atm (c) 80 atm (d) none of these

12. Gas molecules each of mass 10^{26} kg are taken in a container of volume 1 dm^3. The root mean square speed of gas molecules is 1 km sec^{-1}. What is the temperature of gas molecules.

(Given : N_A 6 10^{23}; R 8 J/mol.K)

(a) 298 K (b) 25 K (c) 250 K (d) 2500 K

13. A balloon of diameter 21 meter weighs 100 kg. Calculate its pay-load, if it is filled with He at 1.0 atm and 27°C. Density of air is 1.2 kg m^{-3}. (Given : R 0.082 L atm K ^{1}mol ^{1})

(a) 4952.42 kg (b) 4932.42 kg (c) 493.242 kg (d) none of these

14. A given volume of ozonised oxygen (containing 60% oxygen by volume) required 220 sec to effuse while an equal volume of oxygen took 200 sec only under identical conditions. If density of O_2 is 1.6 g/L then find density of O_3.


(a) 1.936 g/L (b) 2.16 g/L (c) 3.28 g/L (d) 2.24 g/L

15. If 250 mL of N_2 over water at 30°C and a total pressure of 740 torr is mixed with 300 mL of Ne over water at 25°C and a total pressure of 780 torr, what will be the total pressure if the mixture is in a 500 mL vessel over water at 35°C.

(Given : Vapour pressure (Aqueous tension) of H_2O at 25°C, 30°C and 35°C are 23.8, 31.8 and 42.2 torr respectively. Assume volume of $H_2O(l)$ is negligible in final vessel)

(a) 760 torr (b) 828.4 torr (c) 807.6 torr (d) 870.6 torr

16. A bulb of constant volume is attached to a manometer tube open at other end as shown in figure. The manometer is filled with a liquid of density (1/3) rd that of mercury. Initially h was 228 cm.



Through a small hole in the bulb gas leaked assuming pressure decreases as $\frac{dP}{dt}$ kP.

If value of h is 114 cm after 14 minutes, what is the value of k (in hour ^{1})?

[Use : ln(4/3) 0.28 and density of Hg 13.6 g/mL.].

(a) 0.6 (b) 1.2 (c) 2.4 (d) none of these

17. A mixture of nitrogen and water vapours is admitted to a flask at 760 torr which contains a sufficient solid drying agent. After long time the pressure attained a steady value of 722 torr.

If the experiment is done at 27°C and drying agent increases in mass by 0.9 gm, what is the volume of the flask? Neglect any possible vapour pressure of drying agent and volume occupied by drying agent.


(a) 443.34 L (b) 246.3 L (c) 12.315 L (d) 24.63 L

For More Material Join: @JEEAdvanced_2024

18. At room temperature following reaction goes to completion

$$2AB(g) \quad B_2(g) \qquad 2AB_2(s)$$

AB_2 is solid with negligible vapour pressure below 0°C. At 300 K, the AB in the smaller flask exerts a pressure of 3 atm and in the larger flask B_2 exerts a pressure of 1 atm at 400 K when they are separated out by a close valve. The gases are mixed by opening the stop cock and after the end of the reaction the flask are cooled to 250 K




The final pressure is :

(a) 0.156 atm (b) 0.3125 atm

(c) 0.625 atm (d) 3.2 atm

19. A vessel of uniform cross-section of length 500 cm as shown in figure is divided in two parts by a weightless and frictionless piston one part contains 5 moles of He(g) and other part 2 moles of $H_2(g)$ and 4 moles of $O_2(g)$ added at the same temperature and pressure in which reaction takes place finally vessel cooled to 300 K and 1 atm. What is the length of He compartment? (Assume volume of piston and vol. of $H_2O(l)$ formed are negligible)



(a) 187.5 cm (b) 300 cm (c) 312.5 cm (d) none of these

20. For a real gas (mol. mass = 60) if density at critical point is 0.80 g/cm^3 and its T_c $\frac{4 \quad 10^5}{821}$ K, then van der Waals' constant a (in atm L^2 mol ^{2}) is

(a) 0.3375 (b) 3.375 (c) 1.68 (d) 0.025

21. The van der Waals' constant 'b' of a gas is 4 10^{-4} L/mol. How near can the centres of the two molecules approach each other? [Use : N_A 6 10^{23}]

(a) 10^{-7} m (b) 10^{-10} m (c) 5 10^{-11} m (d) 5 10^{-9} m

22. The density of vapour of a substance (X) at 1 atm pressure and 500 K is 0.8 kg/m^3. The vapour effuses through a small hole at a rate of 4/5 times slower than oxygen under the same condition. What is the compressibility factor (Z) of the vapour?

(a) 0.974 (b) 1.35 (c) 1.52 (d) 1.22

For More Material Join: @JEEAdvanced_2024

23. van der Waal's gas equation can be reduced to virial equation and virial equation (in terms of volume) is $Z \quad A \quad \frac{B}{V_m} \quad \frac{C}{V_m^2} \quad \ldots\ldots$

where A first virial coefficient, B second virial coefficient, C third virial coefficient. The third virial coefficient of Hg(g) is 625 $(cm^2/mol)^2$. What volume is available for movement of 10 moles He(g) atoms present in 50 L vessel?


(a) 49.75 L (b) 49.25 L (c) 25 L (d) 50 L

24. If the slope of 'Z' (compressibility factor) vs. 'P' curve is constant slope $\frac{}{492.6}$ atm ^{1} at a particular temperature (300 K) and very high pressure, then calculate diameter of the molecules.

(Given : $N_A \quad 6.0 \quad 10^{23}, R \quad 0.0821$ atm. L mol ^{1} K ^{1})

(a) 7.5 Å (b) 5 Å (c) 2.5 Å (d) 1.25 Å

25. A graph is plotted between P (atm) *vs* t C for 10 mol of an ideal gas as follows:




Then slope of curve and volume of container (L) respectively, are:

(a) 0.1, 8.21 (b) 8.21, 0.1

(c) 27.3, 8.21 (d) 8.21, 27.3

26. For two samples of ideal gases A and B curves are plotted n *vs* V (volume of container) at 16.42 atm pressure. Then temperature of A and B respectively are:



(a) $\frac{200}{\sqrt{3}}$ K, $200\sqrt{3}$ K (b) $\frac{200}{\sqrt{3}}$ C, $(200\sqrt{3})$ C

(c) $200\sqrt{3}$ K, $\frac{200}{\sqrt{3}}$ K (d) 200 K, $\frac{\sqrt{3}}{200}$ K

27. At a constant pressure, what should be the percentage increase in the temperature in kelvin for a 10% increase in volume:

(a) 10% (b) 20% (c) 5% (d) 50%

28. 6 litre H_2O is placed in a closed room of volume 827 L at the temperature of 300 K. If vapour pressure of liquid water is 22.8 mm of Hg at 300 K and its density is 1 g/cm^3:

[Given : $R \quad 0.0821$ atm. L mol ^{1} K ^{1}, Assuming volume of liquid water to be constant]

Column-I		Column-II	
(P)	Mass of H_2O in gaseous form (in g)	**(1)**	6
(Q)	Moles of H_2O in gaseous state	**(2)**	18
(R)	Approximate mass of water left in liquid state (in kg)	**(3)**	3
(S)	Total number of moles of all atoms in vapour form	**(4)**	1

For More Material Join: @JEEAdvanced_2024

Codes :

	P	Q	R	S
(a)	1	2	4	3
(b)	4	3	2	1
(c)	2	4	1	3
(d)	1	2	3	4

29. Match the items of columns I and II.

Column-I		**Column-II**	
(P)	Z for ideal gas behaviour	**(1)**	3/8
(Q)	Z for real gas at low pressure	**(2)**	$1 \quad \frac{Pb}{RT}$
(R)	Z for real gas at high pressure	**(3)**	1
(S)	Z for critical state	**(4)**	$1 \quad \frac{a}{RTV}$

Codes :

	P	Q	R	S
(a)	1	2	4	3
(b)	3	4	2	1
(c)	2	1	4	3
(d)	1	2	3	4

30.

List-I (Conditions for real gas)		**List-II**	
(P)	If force of attraction among gas particles are negligible	**(1)**	$PV_m \quad RT$
(Q)	At 1 atm and 273 K	**(2)**	$PV_m \quad RT \quad \frac{a}{V_m}$
(R)	If the volume of gas particles is negligible	**(3)**	$P \quad \frac{a}{V_m^2} \quad (V_m \quad b) \quad RT$
(S)	At low pressure and high temperature	**(4)**	$PV_m \quad RT \quad Pb$

Codes :

	P	Q	R	S
(a)	4	1	3	2
(b)	4	3	2	1
(c)	2	1	4	3
(d)	1	2	3	4

For More Material Join: @JEEAdvanced_2024

Level 3

Passage 1

van der Waals' equation for calculating the pressure of a non ideal gas is

$$P \quad \frac{an^2}{V^2} \quad (V \quad nb) \quad n\,RT$$

van der Waals' suggested that the pressure exerted by an ideal gas, P_{ideal}, is related to the experimentally measured pressure, P_{real} by the equation

$$P_{ideal} \quad P_{real} \qquad \frac{an^2}{V^2}$$

observed pressure — correction term

Constant **'a'** is measure of intermolecular interaction between gaseous molecules that gives rise to nonideal behavior. It depends upon how frequently any two molecules approach each other closely. Another correction concerns the volume occupied by the gas molecules. In the ideal gas equation, V represents the volume of the container. However, each molecule does occupy a finite, although small, intrinsic volume, so the effective volume of the gas becomes $(V \quad nb)$, where n is the number of moles of the gas and b is a constant. The term nb represents the volume occupied by gas particles present in n moles of the gas.

Having taken into account the corrections for pressure and volume, we can rewrite the ideal gas equation as follows :

$$P \quad \frac{an^2}{V^2} \quad (V \quad nb) \quad nRT$$

corrected pressure — corrected volume

1. At relatively high pressures, the van der Waals' equation of state reduces to

(a) $PV \quad RT \quad a/V_m$ (b) $PV \quad aRT/V_m^2$ (c) $P \quad RT \quad a/V_m^2$ (d) $PV_m \quad RT \quad Pb$

2. For non-zero value of force of attraction between gas molecules at large volume, gas equation will be :

(a) $PV \quad nRT \quad \frac{n^2 a}{V}$ (b) $PV \quad nRT \quad nbP$ (c) $P \quad \frac{nRT}{V \quad b}$ (d) $PV \quad nRT$

3. The van der Waals' constant 'a' for CO_2 gas is greater than that of H_2 gas. Its mean that the

(a) strength of van der Waals' force of CO_2 gas is less than that of H_2 gas

(b) strength of van der Waals' force of CO_2 gas is equal to that of H_2 gas

(c) CO_2 gas can be more easily liquified

(d) H_2 gas can be more easily liquified

For More Material Join: @JEEAdvanced_2024

4. Using van der Waals' equation, find the constant 'a' (in atm L^2 mol^{-2}) when two moles of a gas confined in 4 L flask exerts a pressure of 11.0 atmospheres at a temperature of 300 K. The value of b is 0.05 L mol^{-1}. (R 0.082 atm.L/K mol)


(a) 2.62 (b) 2.64 (c) 6.24 (d) 6.46



Passage 2

Collision cross-section is an area of an imaginary sphere of radius around the molecule within which the centre of another molecule cannot penetrate.

The volume swept by a single molecule in unit time is



V (^{2}) $\bar{u}$ where $\bar{u}$ is the average speed


If N^* is the number of molecules per unit volume, then the number of molecules within the volume V is

$$N \quad VN^* \quad (\quad ^2\bar{u})N^*$$

Hence, the number of collision made by a single molecule in unit time will be

$$Z \quad N \quad (\quad ^2\bar{u})N^*$$

In order to account for the movements of all molecules, we must consider the average velocity along the line of centres of two colliding molecules instead of the average velocity of a single molecule. If it is assumed that, on an average, molecules collide while approaching each other perpendicularly, then the average velocity along their centres is $\sqrt{2}\,\bar{u}$ as shown below.



Number of collision made by a single molecule with other molecules per unit time is given by

$$Z_1 \qquad ^2(\bar{u}_{rel})N^* \quad \sqrt{2} \quad ^2\,\bar{u}N^*$$

The total number of bimolecular collisions Z_{11} per unit volume per unit time is given by

$$Z_{11} \quad \frac{1}{2}(Z_1N^*) \text{ or } Z_{11} \quad \frac{1}{2}(\sqrt{2} \quad ^2\bar{u}N^*)\,N^* \quad \frac{1}{\sqrt{2}} \quad ^2\bar{u}N^{*2}$$

If the collision involve two unlike molecules then the number of collisions Z_{12} per unit volume per unit time is given as

$$Z_{12} \qquad ^2_{12} \sqrt{\frac{8kT}{}} \quad N_1N_2$$

For More Material Join: @JEEAdvanced_2024

where N_1 and N_2 are the number of molecules per unit volume of the two types of molecules, $_{12}$ is the average diameter of the two molecules and is the reduced mass. The mean free path is the average distance travelled by a molecule between two successive collisions. We can express it as follows :

$$\frac{\text{Average distance travelled per unit time}}{\text{No. of collisions made by a single molecule per unit time}} \quad \frac{\bar{u}}{Z_1}$$

or

$$\frac{\bar{u}}{\sqrt{2} \quad ^2\bar{u}N^*} \qquad \frac{1}{\sqrt{2} \quad ^2N^*}$$

1. For a given gas the mean free path at a particular pressure is :
(a) independent to temperature (b) decreases with rise in temperature
(c) increases with rise in temperature (d) directly proportional to T^2

2-3 Three ideal gas samples in separate equal volume containers are taken and following data is given :

	Pressure	Temperature	Mean free paths	Mol. wt.
Gas *A*	1 atm	1600 K	0.16 nm	20
Gas *B*	2 atm	200 K	0.16 nm	40
Gas *C*	4 atm	400 K	0.04 nm	80


2. Calculate ratio of collision frequencies (Z_{11}) $(A : B : C)$ of following for the three gases.
(a) 1 : 2 : 4 (b) 4 : 2 : 1 (c) 1 : 4 : 16 (d) 16 : 4 :1

3. Calculate number of collision by one molecule per sec (Z_1).
(a) 4 : 1 : 4 (b) 1 : 4 : 4 (c) 4 : 3 : 2 (d) 1 : 2 : 4



Passage 3

A manometer contains a liquid of density 5.44 g/cm^3 is attached to a flask containing gas 'A' as follows



For More Material Join: @JEEAdvanced_2024

1. If the same liquid is used in barometer to measure the atmospheric pressure, then what will be the length of the liquid column, which exerts pressure equal to 1 atm ? (density of Hg = 13.6 g/cm^3)

(a) 190 cm (b) 76 cm (c) 30.4 cm (d) 266 cm

2. The initial pressure of gas A in the flask is:

(a) 1.5 atm (b) 1 atm (c) 1.3 atm (d) 1.2 atm


3. If gas A undergoes 30% trimerisation [$3A(g) \rightleftharpoons A_3(g)$] then the difference in height of the liquid level in two columns is:


(a) 38 cm (b) 7.6 cm (c) 3.04 cm (d) 15.1 cm


One or More Answers is/are Correct




1. Which of the following curves represent(s) Boyle's law?

(a) 

(b) 

(c) 

(d) 

2. If a gas expands at constant temperature:

(a) the pressure decreases

(b) the kinetic energy of the molecules remains the same

(c) the K.E. of the molecules decrease

(d) the number of molecules of the gas increase

3. Which of the following statements are correct?


(a) It is not possible to compress a gas at a temperature below T_C

(b) At a temperature below T_C, the molecules are close enough for the attractive forces to act, and condensation occurs

(c) No condensation takes place above T_C

(d) Due to higher kinetic energy of the gas molecules above T_C, it is considered as super critical fluid

4. What conclusion would you draw from the following graphs for an ideal gas?



(a) As the temperature is reduced, the volume as well as the pressure increase

(b) As the temperature is reduced, the volume becomes zero and the pressure reaches infinity

For More Material Join: @JEEAdvanced_2024

(c) As the temperature is reduced, the pressure decrease
(d) A point is reached where, theoretically, the volume become zero

5. At Boyle temperature:
(a) the effects of the repulsive and attractive intermolecular forces just offset each other
(b) the repulsive intermolecular forces are stronger than the attractive intermolecular forces
(c) the repulsive intermolecular forces are weaker than the attractive intermolecular forces
(d) $b \quad \frac{a}{RT} \quad 0$

6. Indicate the correct statement for equal volumes of $N_2(g)$ and $CO_2(g)$ at 25°C and 1 atm.
(a) The average translational K.E. per molecule is the same for N_2 and CO_2
(b) The rms speed is same for both $N_2$and CO_2
(c) The density of N_2 is less than that of CO_2
(d) The total translational K.E. of both N_2 and CO_2 is the same

7. Which of the following is correct for critical temperature?
(a) It is the highest temperature at which liquid and vapour can coexist
(b) Beyond this temperature, there is no distinction between the two phases and a gas cannot be liquefied by compression.
(c) At this temperature, the gas and the liquid phases have different critical densities
(d) All are correct


8. Consider the following statement regarding Maxwell's distribution of speeds. The correct statement(s) is/are :
(a) as temperature increases, the peak (maxima) of a curve is shifted towards right side
(b) as temperature increases, the most probable speed of molecules increases but fraction of molecules of maximum speed decreases
(c) the area under the curve at all the temperatures is the same because it represents the number of gaseous molecules
(d) the fractions of molecules having different speeds are different at a given temperature

9. If a gas expands at a constant pressure by providing heat :
(a) the temperature increases
(b) the kinetic energy of the gaseous molecules remains same
(c) the kinetic energy of gaseous molecules decreases
(d) the number of molecules of the gas decreases

10. Select the incorrect statement(s) :
(a) The product of pressure and volume of fixed amount of a gas is independent of temperature
(b) The value of universal gas constant depends upon temperature, volume and number of gaseous molecules
(c) The gas constant also known as Boltzmann's constant
(d) The average kinetic energy of molecules depends only on temperature


11. Following represents the Maxwell distribution curve for an ideal gas at two temperature T_1 andT_2. Which of the following option(s) is/are true?

For More Material Join: @JEEAdvanced_2024



(a) Total area under the two curves is independent to moles of gas
(b) u_{mp} decreases as temperature decreases
(c) T_1 T_2 and hence higher the temperature, sharper the curve
(d) The fraction of molecules having speed u_{mp} decreases as temperature increases

12. Two container each containing liquid water are connected as shown in diagram.



Given that vapour pressure of $H_2O(l)$ at 300 K and 350 K are 22 torr and 40 torr, select correct statement(s) :
(a) The final pressure in each container if valve is opened while keeping the containers at the given temperature is 22 torr
(b) The final pressure in each container if valve is opened while keeping the containers at the given temperature is 40 torr
(c) Mass of $H_2O(l)$ is decreased in container *X*
(d) Mass of $H_2O(l)$ is decreased in container *Y*

13. Select the correct statement(s) :
(a) At Boyle's temperature a real gas behaves like an ideal gas at low pressure
(b) Above critical conditions, a real gas is called super critical fluid
(c) For hydrogen gas '*b*' domainates over '*a*' at all temperatures
(d) At high pressure van der Waals' constant '*b*' domainates over '*a*'


14. Select the correct statement :
(a) The value of compressibility factor '*Z*' for H_2 gas is greater than one at room temperature and pressure
(b) The real gas behaves as an ideal gas at Boyle's temperature.

For More Material Join: @JEEAdvanced_2024

(c) For a real gas following van der Waals' equation of state, the expression of critical temperature is $\frac{8a}{27R.b}$

(d) At low pressure, the compressibility factor 'Z' 1 $\frac{P.b}{RT}$ for a van der Waals' gas.

15. A open ended mercury manometer is used to measure the pressure exerted by a trapped gas as shown in the figure. Initially manometer shows no difference in mercury level in both columns as shown in diagram.



After sparking 'A' dissociates according to following reaction

$$A(g) \qquad B(g) \quad 3C(g)$$

If pressure of Gas "A" decreases to 0.9 atm, then (Assume temperature to be constant and is 300 K)

(a) total pressure increases to 1.3 atm

(b) total pressure increases by 0.3 atm

(c) total pressure increases by 22.3 cm of Hg

(d) difference in mercury level is 228 mm.

16. Select incorrect statements for real gases:

(a) In low pressure region repulsive forces dominates

(b) Volume of gas particles is not negligible in low pressure region

(c) Gases behaves as an ideal gas at low pressure and low temperature

(d) In high pressure region attractive forces dominates

17. Select correct statements:

(a) A real gas can be liquified at critical temperature

(b) Critical pressure is the maximum pressure at which a substance is present in its liquid state at T_C

(c) Ideal gas can be liquified below T_C

(d) Critical volume is the molar volume of substance in gaseous state at T_C and P_C

18. Which is/are correct for real gases?

(a) $\lim_{P \to 0}(PV_m)$ constant at constant high temperature

(b) $\lim_{V_m \to 0}(PV_m)$ constant at constant low temperature

For More Material Join: @JEEAdvanced_2024

(c) $\lim_{P \to 0} \frac{PV_m}{RT}$ 1 at high temperature

(d) $\lim_{V \to 0} \frac{PV_m}{RT}$ R

19. Select incorrect statement(s)
(a) At very low pressure real gases show minimum deviation from ideal behaviour.
(b) The compressibility factor for an ideal gas is zero.
(c) At Boyle temperature real gas behave as ideal gas in high pressure region.
(d) Real gas show maximum deviation at high pressure and low temperature.

20. If an ideal gas is heated at constant pressure:
(a) the volume increases
(b) the mass of gas remains same
(c) the kinetic energy of the molecules increases
(d) attraction forces between gas particles increases

Match the Column



Column-I and **Column-II** contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

1.

Column I	Column II
(A) Boyle's temperature	(P) a/Rb
(B) Inversion temperature	(Q) $8a/27Rb$
(C) Critical temperature	(R) $2a/Rb$
(D) Critical pressure	(S) $a/27b^2$

2.

Column I	Column II
(A) Root mean square speed	(P) $\frac{3}{2}RT$
(B) Most probable speed	(Q) $\sqrt{\frac{8P}{d}}$
(C) Average speed	(R) $\sqrt{\frac{3P}{d}}$
(D) K.E. of gas mol ^{1}	(S) $\sqrt{\frac{2RT}{M}}$

For More Material Join: @JEEAdvanced_2024





3.

Column I	Column II
(A) If force of attraction among the gas molecules be negligible	(P) $P \quad \frac{a}{V^2} \quad (V \quad b) \quad RT$
(B) If the volume of the gas molecules be negligible	(Q) $PV \quad RT \quad \frac{a}{V}$
(C) At STP	(R) $PV \quad RT \quad Pb$
(D) At low pressure and at high temperature	(S) $PV \quad RT$

4. Match gases under specified conditions listed in Column I with their properties/laws in Column II.


Column I	Column II
(A) Hydrogen gas ($P \quad 200$ atm, $T \quad 273$K)	(P) Compressibility factor 1
(B) Hydrogen gas ($P \sim 0, T \quad 273$ K)	(Q) Attractive forces are dominant
(C) CO_2 ($P = 1$ atm, $T = 273$ K)	(R) $PV \quad nRT$
(D) Real gas with very large molar volume	(S) $P(V \quad nb) \quad nRT$

5.

Column I	Column II
(A) $\frac{1}{V^2}$ $vs.$ P for ideal gas at constant T and n.	(P) 
(B) V $vs.$ $\frac{1}{T}$ for ideal gas at constant P and n	(Q) 
(C) PT $vs.$ T^2 for ideal gas at constant V and n.	(R) 
(D) V $vs.$ $\frac{1}{P^2}$ for ideal gas at constant T and n.	(S) 


For More Material Join: @JEEAdvanced_2024

6. Match the correct column from list-1 to list-2 on the basis of following Andrew's isotherm of real gas.



Column-I	Column-II
(A) Substance exist in both liquid and gas state	(P) At AB part
(B) Only liquid state exist	(Q) At BD part
(C) Substance exist in gas state only	(R) At DE part
(D) Real gas is called super critical fluid	(S) At point C
	(T) At GF curve

7.



Column-I	Column-II
(A) Speed at point 'A' is	(P) $\sqrt{\frac{8RT}{M}}$
(B) Speed at point 'B' is	(Q) $\sqrt{\frac{3RT}{M}}$
(C) Speed at point 'C' is	(R) U_{rms}
(D) The speed possessed by maximum fraction of the gas particles	(S) $\sqrt{\frac{2RT}{M}}$

For More Material Join: @JEEAdvanced_2024

Assertion-Reason Type Questions



Each question contains STATEMENT-1 (Assertion) and STATEMENT-2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below

(A) If both the statement are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1

(B) If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1

(C) If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

(D) If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

1. STATEMENT-1 : The heat absorbed during the isothermal expansion of an ideal gas against vacuum is zero.
STATEMENT-2 : The volume occupied by the molecules of an ideal gas is zero.

2. STATEMENT-1 : A lighter gas diffuses more rapidly than a heavier gas.
STATEMENT-2 : At a given temperature, the rate of diffusion of a gas is inversely proportional to the density.

3. STATEMENT-1 : The value of the van der Waals' constant 'a' is larger for ammonia than for nitrogen.
STATEMENT-2 : Hydrogen bonding is present in ammonia.

4. STATEMENT-1 : Helium shows only positive deviation from ideal behaviour at room temperature.
STATEMENT-2 : Helium is an inert gas.

5. STATEMENT-1 : CH_4, CO_2 has value of Z (compressibility factor) less than one at 0°C.
STATEMENT-2 : Z 1 is due to the attractive forces dominate among the molecules.

6. STATEMENT-1 : The Joule-Thomson coefficient for an ideal gas is zero.
STATEMENT-2 : There are no intermolecular attractive forces in an ideal gas.

7. STATEMENT-1 : The average translational kinetic energy per molecule of the gas per degree of freedom is 1/2 KT.
STATEMENT-2 : For every molecule there are three rotational degree of freedom.

8. STATEMENT-1 : On increasing the temperature, the height of the peak of the Maxwell distribution curve decreases.
STATEMENT-2 : The fraction of molecules is very less at the higher speeds.

9. STATEMENT-1 : The gases He and H_2 are very different in their behaviour at any temperature and pressure but their compressibility factors are nearly the same at the critical point.
STATEMENT-2 : They have nearly the same critical constant.

For More Material Join: @JEEAdvanced_2024

10. STATEMENT-1 : Most probable speed is the speed possessed by maximum fraction of molecules at the same temperature.

STATEMENT-2 : On collision, more and more molecules acquire higher speed at the same temperature.

11. STATEMENT-1 : Plot of P *vs* $1/V$ (volume) is a straight line for an ideal gas.

STATEMENT-2 : Pressure is directly proportional to volume for an ideal gas.

12. STATEMENT-1 : 1 mol of H_2 and O_2 each occupy 22.7 L of volume at 0°C and 1 bar pressure, considering ideal behaviour.

STATEMENT-2 : Molar volume for all ideal gases at the same temperature and pressure are equal.

13. STATEMENT-1 : Reacting gases react to form a new gas having pressure equal to the sum of their partial pressure.

STATEMENT-2 : Pressure exerted by a mixture of non-reacting gases present in a container is equal to the sum of their partial pressures.

14. STATEMENT-1 : $1/4^{th}$ of the initial mole of the air is expelled, if air present in an open vessel is heated from 27°C to 127°C.

STATEMENT-2 : Rate of diffusion of a gas is inversely proportional to the square root of its molecular mass.

15. STATEMENT-1 : Compressibility factor for hydrogen varies with pressure with positive slope at all pressures at 0°C.

STATEMENT-2 : Even at low pressures, repulsive forces dominate in hydrogen gas at 0°C.

16. STATEMENT-1 : Wet air is heavier than dry air.

STATEMENT-2 : The density of dry air is less than the density of water at 1 atm and 273K.

Subjective Problems


1. Collapsible balloon is inflated to a volume of 10 L at a pressure of 1 atm. When the balloon is immersed to the bottom of a lake, its volume reduce to 1.25 L. Assuming atmospheric pressure to be equivalent to 10 m column of water and no change in temperature, if the depth of the lake is $x \quad 10^3$ cm, what is the value of x ?

2. A gaseous mixture containing equal moles of H_2, O_2 and He is subjected to series of effusion steps. The composition (by moles) of effused mixture after 4 effusion steps is $x:1:y$ respectively. Then find the value of $\frac{x}{y}$.

3. One mole of a gas changed from its initial state (15 L, 2 atm) to final state (4L, 10 atm) reversibly. If this change can be represented by a straight line in $P \quad V$ curve maximum temperature (approximate), the gas attained is $x \quad 10^2$K. Then find the value of x.

4. Two moles of an ideal gas undergoes the following process. Given that $\left(\frac{P}{T}\right)_V$ is $x \quad 10^{\ y}$, then calculate the value of $(x \quad y)$


For More Material Join: @JEEAdvanced_2024



5. 1 mole of a diatomic gas present in 10 L vessel at certain temperature exert a pressure of 0.96 atm. Under similar conditions an ideal gas exerted 1.0 atm pressure. If volume of gas molecule is negligible, then find the value of van der Waals' constant ''a'' (in atm L^2/mol^2).

6. The graph of compressibility factor (Z) *vs.* P for one mole of a real gas is shown in following diagram. The graph is plotted at constant temperature 273 K. If the slope of graph at very high pressure $\frac{dZ}{dP}$ is $\frac{1}{2.8}$ atm ^{1}, then calculate volume of one mole of real gas molecules (in L /mol)

Given : N_A 6 10^{23} and R $\frac{22.4}{273}$ L atm K ^{1}mol ^{1}




7. Under the identical conditions of temperature, the density of a gas X is two times to that of gas Y while molecular mass of gas Y is three times that of X. Calculate the ratio of pressure of X and Y.

8. The time taken for a certain volume of a certain gas to diffuse through a small hole was 2 min. Under similar conditions an equal volume of oxygen took 5.65 min to pass. What is the molecular mass of gas (in amu) ?

9. Excess $F_2(g)$ reacts at 150 C and 1.0 atm pressure with $Br_2(g)$ to give a compound BrF_n. If 423 mL of $Br_2(g)$ at the same temperature and pressure produced 4.2 g of BrF_n, what is n ? [Atomic mass Br 80, F 19]


10. Initially bulb "*A*" contained oxygen gas at 27 C and 950 mm of Hg and bulb *"B"* contained neon gas at 27 C and 900 mm of Hg. These bulbs are connected by a narrow tube of negligible volume equipped with a stopcock and gases were allowed to mix-up freely. The pressure obtained in the combined system was found to be 910 mm of Hg.

For More Material Join: @JEEAdvanced_2024



If volume of bulb *B* was measured to be 10 L, then find the volume of oxygen gas present initially in bulb *"A"*.

11. Air is trapped in a horizontal glass tube by 36 cm mercury column as shown below :



If the tube is held vertical keeping the open end up, length of air column shrink to 19 cm. What is the length (in cm) by which the mercury column shifts down?

12. A flask containing air at 107°C and 722 mm of Hg is cooled to 100 K and 760 mm of Hg. If density in the initial condition 1 g/cm^3, then what is the final density (g/cm^3)?

13. If an ideal gas at 100 K is heated to 109 K in a rigid container, the pressure increases by X%. What is the value of *X*?

14. The van der Waal's constants for a gas are a 3.6 atm L^2 mol^{-2}, b 0.6 L mol^{-1}. If R 0.08 L atm K^{-1} mol^{-1} and the Boyle's temperature (K) is T_B of this gas, then what is the value of $\frac{T_B}{15}$?

15. A flask has 10 gas particles, out of which four particles are moving at 7 ms^{-1} and the remaining are moving at the same speed of '*X*' ms^{-1}. If the r.m.s. of the gas is 5 ms^{-1}, what is the value of *X* ?

For More Material Join: @JEEAdvanced_2024

Answers

Level 1

1.	(c)	**2.**	(c)	**3.**	(a)	**4.**	(d)	**5.**	(a)	**6.**	(b)	**7.**	(a)	**8.**	(d)	**9.**	(c)	**10.**	(c)
11.	(a)	**12.**	(a)	**13.**	(c)	**14.**	(c)	**15.**	(b)	**16.**	(c)	**17.**	(a)	**18.**	(a)	**19.**	(c)	**20.**	(c)
21.	(b)	**22.**	(b)	**23.**	(a)	**24.**	(d)	**25.**	(d)	**26.**	(c)	**27.**	(b)	**28.**	(a)	**29.**	(c)	**30.**	(b)
31.	(c)	**32.**	(b)	**33.**	(b)	**34.**	(b)	**35.**	(a)	**36.**	(d)	**37.**	(d)	**38.**	(c)	**39.**	(c)	**40.**	(a)
41.	(a)	**42.**	(b)	**43.**	(c)	**44.**	(a)	**45.**	(d)	**46.**	(a)	**47.**	(c)	**48.**	(d)	**49.**	(b)	**50.**	(a)
51.	(b)	**52.**	(d)	**53.**	(c)	**54.**	(b)	**55.**	(b)	**56.**	(d)	**57.**	(a)	**58.**	(a)	**59.**	(c)	**60.**	(a)
61.	(d)	**62.**	(d)	**63.**	(b)	**64.**	(a)	**65.**	(b)	**66.**	(c)	**67.**	(b)	**68.**	(b)	**69.**	(a)	**70.**	(a)
71.	(c)	**72.**	(a)	**73.**	(a)	**74.**	(b)	**75.**	(d)	**76.**	(c)	**77.**	(b)	**78.**	(b)	**79.**	(b)	**80.**	(a)
81.	(b)	**82.**	(d)	**83.**	(a)	**84.**	(a)	**85.**	(c)	**86.**	(a)	**87.**	(b)	**88.**	(b)	**89.**	(b)	**90.**	(b)
91.	(b)	**92.**	(a)	**93.**	(c)	**94.**	(b)	**95.**	(a)	**96.**	(b)	**97.**	(d)	**98.**	(d)	**99.**	(b)	**100.**	(d)
101.	(c)	**102.**	(c)	**103.**	(a)	**104.**	(c)	**105.**	(a)	**106.**	(b)	**107.**	(c)	**108.**	(d)	**109.**	(b)	**110.**	(c)
111.	(c)	**112.**	(b)	**113.**	(b)	**114.**	(d)	**115.**	(d)	**116.**	(b)	**117.**	(d)	**118.**	(b)	**119.**	(a)	**120.**	(b)
121.	(a)	**122.**	(b)	**123.**	(b)	**124.**	(b)	**125.**	(b)	**126.**	(c)	**127.**	(d)	**128.**	(a)	**129.**	(a)	**130.**	(c)
131.	(b)	**132.**	(d)	**133.**	(a)	**134.**	(b)	**135.**	(c)	**136.**	(c)	**137.**	(b)	**138.**	(a)	**139.**	(c)	**140.**	(b)
141.	(d)	**142.**	(b)	**143.**	(a)	**144.**	(c)	**145.**	(c)	**146.**	(c)	**147.**	(c)	**148.**	(d)	**149.**	(d)	**150.**	(a)
151.	(c)	**152.**	(c)	**153.**	(d)	**154.**	(a)	**155.**	(b)	**156.**	(a)	**157.**	(c)	**158.**	(d)	**159.**	(c)	**160.**	(d)
161.	(c)	**162.**	(d)	**163.**	(a)	**164.**	(b)	**165.**	(b)	**166.**	(d)	**167.**	(a)	**168.**	(d)	**169.**	(b)	**170.**	(a)
171.	(c)	**172.**	(c)	**173.**	(a)	**174.**	(a)	**175.**	(b)										

Level 2

1.	(b)	**2.**	(c)	**3.**	(c)	**4.**	(b)	**5.**	(b)	**6.**	(b)	**7.**	(c)	**8.**	(a)	**9.**	(d)	**10.**	(b)
11.	(a)	**12.**	(c)	**13.**	(b)	**14.**	(d)	**15.**	(d)	**16.**	(b)	**17.**	(d)	**18.**	(c)	**19.**	(c)	**20.**	(b)
21.	(b)	**22.**	(c)	**23.**	(a)	**24.**	(b)	**25.**	(a)	**26.**	(a)	**27.**	(a)	**28.**	(c)	**29.**	(b)	**30.**	(b)

For More Material Join: @JEEAdvanced_2024



Level 3

Passage–1	**1.** (d)	**2.** (a)	**3.** (c)	**4.** (d)
Passage–2	**1.** (c)	**2.** (c)	**3.** (a)	
Passage–3	**1.** (a)	**2.** (d)	**3.** (b)	

One or More Answers is/are correct

1. (a,b,c,d) **2.** (a,b) **3.** (b,c,d) **4.** (c,d) **5.** (a) **6.** (a,c,d) **7.** (a,b) **8.** (a,b,c,d)
9. (a) **10.** (a,b,c) **11.** (a,b,d) **12.** (a,d) **13.** (a,b,d) **14.** (a,b,c) **15.** (a,b,d) **16.** (a,b,c,d)
17. (a,d) **18.** (a,c) **19.** (b,c) **20.** (a,b,c)

Match the Column

1. A P; B R; C Q; D S
2. A R; B S; C Q; D P
3. A R; B Q; C P; D S
4. A P, S; B R; C P, Q; D R
5. A R; B S; C P; D Q
6. A Q, S; B R; C P, T; D T
7. A S; B P; C Q, R; D S

Assertion-Reason Type Questions

1. (B) **2.** (C) **3.** (A) **4.** (B) **5.** (A) **6.** (A) **7.** (C) **8.** (B) **9.** (C) **10.** (C)
11. (C) **12.** (A) **13.** (D) **14.** (B) **15.** (A) **16.** (D)

Subjective Problems

1. 7	**2.** 4	**3.** 7	**4.** 3	**5.** 4	**6.** 2	**7.** 6	**8.** 4	**9.** 5	**10.** 4
11. 9	**12.** 4	**13.** 9	**14.** 5	**15.** 3					

For More Material Join: @JEEAdvanced_2024

Hints and Solutions

Level 1

2. (c) Boyle's Law

Temperature and mass constant

$PV = K$

$\log P = \log K - \log V$

3. (a) $\frac{V_1}{T_1} = \frac{V_2}{T_2}$; pressure and mass constant

$\frac{0.2}{273} = \frac{V_2}{546}$

$V_2 = 0.4\,\text{L}$

4. (d) Charle's law

Pressure and mass constant

$\frac{V}{T}$ constant $= K$

$V = KT$

$\log V = \log K + \log T$

5. (a) $\frac{P_1V_1}{T_1} = \frac{P_2V_2}{T_2}$

$\frac{P \quad V}{348} = \frac{(2P) \quad (0.85\,V)}{T_2}$

$T_2 = 592\,\text{K} = 319\,^\circ\text{C}$

6. (b) $V = KT$

(i) $\frac{V}{T} = K$ (ii) $VT = KT^2$

(iii) $\log V = \log K + \log T$

8. (d) V constant, mass constant

$\frac{P}{T}$ constant

$P = KT$

$\log P = \log K + \log T$

10. (c) Boyle's law $PV = K$

$$\frac{P}{V} = \frac{K}{V^2} \quad \ldots(i)$$

$$PV^2 = KV \quad \ldots(ii)$$

$$\log P = \log K + \log V \quad \ldots(iii)$$

12. (a) P n, initial mole $n_1 = \frac{PV}{RT} = 0.1$ mole

$\frac{P_1}{P_2} = \frac{n_1}{n_2}$

$\frac{1}{0.75} = \frac{0.1}{n_2}$ $n_2 = 0.075$

Escape mole $= 0.025$

13. (c) For ideal gas , $PV = nRT$

15. (b) $\frac{P_1V_1}{T_1} = \frac{P_2V_2}{T_2}$

$\frac{P \quad 0.1}{T} = \frac{1.5\,P \quad V_2}{\frac{4T}{3}}$

$V_2 = 88.9\,\text{mL}$

16. (c) $\frac{V_1}{T_1} = \frac{V_2}{T_2}$

17. (a) $P = KT$

$P = K(t\,^\circ\text{C} + 273)$

18. (a) $PM = dRT$

$M = dT$

$\frac{M_1}{M_2} = \frac{d_1T_1}{d_2T_2} = \frac{1}{2} \cdot \frac{2}{1} = \frac{1}{1}$

19. (c) $PM = dRT$

$\frac{P_A}{P_B} = \frac{d_A}{d_B} = \frac{M_B}{M_A}$


$\frac{4}{1}$

20. (c) $V = T$

Because temperature by 10 K

expelled air $10\,\text{cm}^3$

21. (b)



$PV = nRT$

n, V constant

$P = T$

$T_2 = T_1$

$P_2 = P_1$

22. (b) $PV = \frac{W}{M}RT$

For More Material Join: @JEEAdvanced_2024

24. (d) $PV = \frac{W}{M}RT$

$V = \frac{1}{M}$

25. (d) $P_1V_1 = P_2V_2$

26. (c) $d = \frac{PM}{RT}$

27. (b) The moles of the gas in the bubble remains constant, so that $n_1 = n_2$. To calculate the final volume, V_2,

$V_2 = V_1 \frac{P_1}{P_2} \frac{T_2}{T_1}$

$2.0\text{ mL} \frac{6.0\text{ atm}}{1.0\text{ atm}} \frac{298\text{ K}}{281\text{ K}}$

12.72 mL

29. (c) $n_{C_2H_2} = \frac{PV}{RT} = \frac{1 \quad 2.64}{0.0821 \quad 273} = 0.118\text{ mole}$

$n_{O_2} = \frac{5}{2}n_{C_2H_2} = \frac{5}{2} \quad 0.118 = 0.294$

$V = \frac{nRT}{P} = 6.6\text{ L}$

30. (b) $d = \frac{PM}{dRT}$


31. (c) $P_1V_1 = P_2V_2$

$V_1 = V$, $V_2 = V/10$

32. (b) $\frac{P_1}{T_1} = \frac{P_2}{T_2}$

33. (b)

(a)




$P = 1/V$

$V_1 \quad V_2 \quad V_3 \quad V_4$

$PV = nRT$

(b)



$P \quad T$

$P \quad T$

$P_1 \quad P_2 \quad P_3 \quad P_4$

34. (b) $PV = nRT$

$n = \frac{1 \quad 10}{76 \quad 0.0821 \quad 1273} = 1.26 \quad 10^{-3}$

No. of Na atoms $= 1.26 \quad 10^{-3} \quad 6.02 \quad 10^{23}$

$= 7.6 \quad 10^{19}$

35. (a) Moles of C_2H_6 and C_2H_4 are a and b respectively.

$a \quad b = \frac{28}{22.4} = 1.25$

$\underset{a}{C_2H_6}(g) \quad \underset{3.5a}{3.5\,O_2}(g) \quad 2\,CO_2(g) \quad 3H_2O(l)$

$\underset{b}{C_2H_4}(g) \quad \underset{3b}{3O_2}(g) \quad 2\,CO_2(g) \quad 2H_2O(l)$

Moles of $O_2 \quad 3.5a \quad 3b = 4$

$a \quad b = 1.25$

$a = 0.5$

$b = 0.75$

36. (d) $n_{H_2O} = \frac{PV}{RT} = 1.54$

$\frac{n_{MgSO_4 \cdot H_2O}}{n_{H_2O}} = \frac{1}{x} = \frac{54.2}{(120 \quad 18x) \quad 1.54}$

$x = 7$

37. (d) Volume of alveoli $= \frac{4}{3} \quad R^3$

$\frac{4}{3} \quad 3.14 \quad (5 \quad 10^{-3})^3\text{mL}$

$5.23 \quad 10^{-7}\text{mL} \quad 5.23 \quad 10^{-10}\text{ L}$

$n_2 = \frac{14}{100} \quad n_{air} = 0.14 \quad \frac{PV}{RT}$

$0.14 \quad 1 \quad \frac{5.23 \quad 10^{-10}}{0.0821 \quad 310}$

$2.87 \quad 10^{-12}$

Molecules $= 2.87 \quad 6.02 \quad 10^{23} \quad 10^{-12}$

$= 1.7 \quad 10^{12}$

38. (c) $PM = dRT$

$M = 46$

39. (c) Because the number of moles is constant.

$\frac{P_iV_i}{T_i} = \frac{P_fV_f}{T_f}$; $P_f = \frac{P_iV_iT_f}{V_fT_i}$

$P_f = \frac{P_iV_iT_f}{V_fT_i}$

$3.21 \quad 10^5\text{ Pa} \quad \frac{V_i}{1.03\ V_i} \quad \frac{(273 \quad 28.0)}{(273 \quad 5.00)}$

$3.50 \quad 10^5\text{ Pa}$

For More Material Join: @JEEAdvanced_2024

40. (a) n_{N_2} 53.57 mole

$\frac{P_1}{P_2}$ $\frac{n_1}{n_2}$

n_2 $\frac{1.8}{200}$ 53.57 0.48213

n_{N_2} escape 53.087

volume $\frac{nRT}{p}$ 1264 L

41. (a) $\frac{P_1V_1}{T_1}$ $\frac{P_2V_2}{T_2}$

V_2 $\frac{745 \quad 1.41 \quad 10^4 \quad 225}{294 \quad 63.1}$

$1.274 \quad 10^5$ L

42. (b) P in mm of Hg $\frac{0.61 \quad 10^3}{1.013 \quad 10^5}$ 760

4.6 mm of Hg

43. (c) $h_{Gallium}$ $\frac{74 \quad 13.6}{6.095}$ 165.1

44. (a) Volume of He at 0.1 atm

P_1V_1 P_2V_2

50 100 V_2 1

V_2 5000 L

Volume available for balloon

5000 50 4950 L at 1 atm

Volume of balloon 100 m^3

100000 L 49500 L at 0.1 atm

So, 2 cylinder required.

45. (d) P_1V_1 P_2V_2 , P_2 $\frac{76 \quad 30.40 \quad 10}{13.6}$

V_2 $\frac{76 \quad 14}{98.41}$ 10.8 L

46. (a) $\frac{P_1}{T_1}$ $\frac{P_2}{T_2}$

T_2 3 $\frac{76}{80}$ 273

47. (c) n_1 n_2

$\frac{1}{32}$ $\frac{2.375}{M_2}$

M_2 76

48. (d) P_i $\frac{6.81}{4}$ $\frac{0.0821 \quad 250}{1.1 \quad 104}$

$3.012 \quad 10^{-3}$

P_f 4 10^{-3}

P 1 10^{-3}

$n_{He\text{-added}}$ $\frac{10^{-3} \quad 1.16 \quad 10^4}{0.0821 \quad 250}$

wt n_{He} 4 2.24 gm.

49. (b) P $\frac{nRT}{V}$ 2.46 atm

50. (a) V $\frac{nRT}{P}$ $\frac{10 \quad 0.0821 \quad T}{0.821}$

V T

$\log V$ $\log T$

slope 1

45

51. (b) Intercept on y-axis

$\log_{10}\frac{nR}{V}$ $\log_{10}\frac{10 \quad 0.0821}{8.21}$ 1.0

$\frac{P}{T}$ v/s curve $\frac{P}{T}$ $\frac{nR}{V}$

Intercept $\frac{nR}{V}$

$\frac{10 \quad 0.0821}{8.21}$ 0.1, slope 0

52. (d) $\frac{3}{2}KT$ mgh

h $\frac{3}{2}$ $\frac{1.38 \quad 10^{-23} \quad 300}{9.81 \quad 4 \quad 1.66 \quad 10^{-27}}$

$9.53 \quad 10^4$ m

53. (c) d_A $2d_B$; $3M_A$ M_B; PM dRT

$\frac{P_A}{P_B}$ $\frac{M_A}{M_B}$ $\frac{d_A}{d_B}$ $\frac{RT}{RT}$

$\frac{P_A}{P_B}$ $\frac{1}{2}$ 2

$\frac{P_A}{P_B}$ $\frac{4}{1}$

54. (b) Case I—Suppose inner balloon burst first

$\frac{600}{300}$ $\frac{800}{T_2}$ T_2 400 K

Case II—Suppose outer balloon burst first

$\frac{1500}{300}$ $\frac{1800}{T_2}$; T_2 360 K

55. (b) V_1 V, T_1 300 K, T_2 500 K, V_2 ?

At constant pressure V_1T_2 V_2T_1

V_2 $\frac{V_1T_2}{T_1}$ $\frac{V \quad 500}{300}$ $\frac{5V}{3}$

Volume of air escaped

final volume – initial volume

For More Material Join: @JEEAdvanced_2024

$\frac{5V}{3}$ V $\frac{2V}{3}$

% of air escaped $\frac{2V/3}{5V/3}$ 100 40%

56. (d) R $\frac{PV}{nT}$

57. (a)

	$2NH_3(g)$	$N_2(g)$	$3H_2(g)$
Before sparking	76	0	0
After sparking at eqm	76 $2x$	x	$3x$


Increase in pressure $2x$ 6; x 3 cm Hg

Partial pressure of H_2 3 3 9 cm Hg

58. (a) P_{Atm} $\frac{h}{13.6}$ P_{Gas} aq Tension

P_{Gas} P_{Atm} aq tension $\frac{h}{13.6}$

59. (c)



P_1V_1 P_2V_2

P $4V$ P 1 $\frac{h}{10}$ V

4 1 $\frac{h}{10}$

h 30 m

60. (a) $h_g d_g$ $h_{Hg} d_{Hg}$; 5 2.72 13.6 h_{Hg}

h_{Hg} 1 m

P_{gas} 0.76 1 1.76 m of Hg

= 1760 mm of Hg

$\frac{1760}{760}$ 10 n R 300


n 0.94 mole

61. (d) n_T n_1 n_2

$\frac{P \quad 1.25}{R \quad 287.5}$ $\frac{611 \quad 10^3 \quad 0.5}{293.5\,R}$ $\frac{433 \quad 10^3 \quad 0.75}{204.5\,R}$

P $4.84 \quad 10^5$ Pa

62. (d)



$\frac{0.2P \quad (100 \quad V)}{RT}$ $\frac{100P}{RT}$

0.2 (100 V) 100

V 400 cm^3

63. (b) Moles of C_2H_2 and C_3H_8 are a and b

C_2H_2 2.5 O_2 $2CO_2$ H_2O

a $2a$

C_3H_8 $5O_2$ $3CO_2$ $4H_2O$

b $3b$

$2a$ $3b$ 230

a b 80

b 70

a 10

mole fraction $\frac{7}{8}$ 0.875

64. (a) p n

$2H_2(g)$ $O_2(g)$ $2H_2O(g)$

a b

a $2b$ b b

a b 1

a $2b$ 0.4

$3b$ 0.6

b 0.2

a 0.8

65. (b) P_1 $\frac{5 \quad RT}{V}$; p.pr. of He $\frac{2}{5}$ $\frac{5RT}{V}$

$\frac{2 \quad 0.0821 \quad 400}{8.21}$ 8 atm

66. (c) P_{O_2} $\frac{3}{10}$ P_T;

After removing 2 mole of O_2,

P_{O_2} $\frac{1}{8}$ P_T

Decrease in partial pressure of O_2

$\frac{\frac{3P_T}{10} \quad \frac{P_T}{8}}{\frac{3P_T}{10}}$ 100

58.33

67. (b) P_{N_2} 760 30 730 torr.

n_{N_2} $\frac{730 \quad 0.821}{760 \quad 0.0821 \quad 300}$ 0.032

68. (b) % Relative humidity

$\frac{\text{Partial pressure of } H_2O(g)}{\text{Vapour pressur of } H_2O}$ 100

at constant temperature

For More Material Join: @JEEAdvanced_2024

69. (a) Volume of O_2 1 L
and p_{O_2} 760 355 405 mm Hg
Since, the temperature is constant, using Boyle's law we get
1 L 405 mm 0.3 L p_{O_2}
p_{O_2} 1350 mm Hg

70. (a) Dalton's law is applicable for a non-reacting gaseous mixture.
SO_2 Cl_2 SO_2Cl_2

71. (c) n_{CH_4} $\frac{x}{16}$ n_{O_2} $\frac{x}{32}$
mole fraction O_2 1/3

72. (a) n_T n_1 n_2 n_1 $\frac{P}{R}\frac{1}{T}$ 1
$\frac{P}{RT}\frac{2\,L}{}$ 2 P RT
P RT P

73. (a) n $\frac{PV}{RT}$ $\frac{3170}{8.314}\frac{10^{-3}}{300}$ 1.27 10^{-3} mole.

74. (b) Dalton's Law is applicable on a non-reacting gas mixture.

75. (d) mole N_2 $\frac{56}{28}$ 2, mole O_2 $\frac{96}{32}$ 3
P_{O_2} $\frac{2}{5}$ 10 4 atm
P_{N_2} $\frac{3}{5}$ 10 6 atm

76. (c) P n
$\frac{P_{H_2}}{P_{CO_2}}$ $\frac{n_{H_2}}{n_{CO_2}}$ $\frac{22}{1}$
P_{H_2} 22 atm

77. (b) P_{gas} $P_{dry\ gas}$ $P_{moisture}$ at T K
or P_{dry} 830 30 800
Now at T_2 0.99 T_1;
at constant volume $\frac{P_1}{T_1}$ $\frac{P_2}{T_2}$
P_{dry} $\frac{800\quad 0.99\,T}{T}$ 792 mm
P_{gas} P_{dry} $P_{moisture}$
792 25 817 mm

78. (b) We have P_{O_2} X_{O_2} P_{total}
0.60 $\frac{n_{O_2}}{n_{O_2}\quad n_{SO_2}}$ P_{total}
Let no. of moles of O_2 are x;
0.60 $\frac{x/32}{x/32\quad 3x/64}$ P_{total}
0.60 0.4 P_{total}
P_{total} $\frac{0.60}{0.4}$ 1.5 atm
$\because$ P_{O_2} P_{SO_2} 1.5 atm
P_{SO_2} 1.5 P_{O_2} 1.5 0.60 0.9 atm
Concentration of SO_2 $\frac{0.9}{0.0821\quad 300}$
0.036

79. (b) Given P 10 atm,
total numbers of moles; n_A n_B n_C 10
P_A 3 atm, P_B 1 atm
$\because$ P_A x_A $P_{(total)}$ $\frac{n_A}{n_A\quad n_B\quad n_C}$ 10
$\frac{n_A}{10}$ 10, n_A 3
Similarly, P_B x_B $P_{(total)}$
So, n_B 1
n_C 10 $(n_A$ $n_B)$ 10 4 6
Mass of C 6 2 12 g

80. (a) PV $\frac{10}{M}RT$ …(1)
Let x g of the gas remain in the flask when final pressure $\frac{P}{2}, \frac{2T}{3}$
$\frac{P}{2}$ V $\frac{x}{M}$ R $\frac{2T}{3}$ …(2)
(1)/(2) 2 $\frac{10}{x}$ $\frac{3}{2}$
x $\frac{3\quad 10}{4}$ 7.5 g

81. (b) $\frac{P_x}{P_y}$ $\frac{n_x}{n_y}$
$\frac{2}{1}$ $\frac{1/m_x}{2/m_y}$
m_y $4m_x$

82. (d) P $\frac{nRT}{V}$
P $\frac{0.05\quad 0.0821\quad 303}{44\quad 4.6}$ 6.14 10^{-3} atm


For More Material Join: @JEEAdvanced_2024

83. (a) $P_T \quad P_{He} \quad P_{Ne} \quad V.P\,H_2O$

$745 \quad 368 \quad P_{Ne} \quad 28.3$

$P_{Ne} \quad 348.7$ mm of Hg

84. (a)



$$n_{He} \quad \frac{1.2 \quad 0.63}{0.082 \quad 289}, \quad n_{Ne} \quad \frac{3.4 \quad 2.8}{0.0821 \quad 289}$$

$$0.032 \qquad 0.4$$

$$P_f \quad \frac{0.431 \quad 0.0821 \quad 289}{4.6} \quad 2.223$$

$$P_{He} \quad \frac{0.032}{0.432} \quad 2.223 \quad 0.164 \text{ atm}$$

85. (c) From the total pressure and the vapour pressure of water we can calculate the partial pressure of O_2.

$$P_{O_2} \quad P_T \quad P_{H_2O}$$

$$760 \quad 22.4$$

$$737.6 \text{ mm Hg}$$

From the ideal gas equation we write

$$m \quad \frac{PVM}{RT}$$

$$\frac{(0.974 \text{ atm})\,(0.128 \text{ L})\,(32.0 \text{ g/mol})}{(0.0821 \text{ L atm/K mol})\,(273 + 24)\text{ K}}$$

$$0.163 \text{ g}$$

86. (a) $PV \quad nRT \qquad R \quad K \quad N_A$

$PV \quad n \quad N_A KT$

$n \quad N_A \quad \frac{PV}{KT}$ no. of particles

87. (b) $KE \quad \frac{1}{2}Mv^2$

88. (b) K.E is temperature dependent

same moles would have same volume at same P and T

$PV \quad nRT$

$d \quad \frac{PM}{RT}$

89. (b) According to Boyle's law at constant temperature and mass

$$PV \quad K$$

$$P \quad \frac{1}{V}$$

90. (b) $KE_T \quad \frac{3}{2}nRT$ at constant P and V

$PV \quad K$ at constant T and mass

$V \quad KT$ at constant P and mass

91. (b) $KE_{avg} \quad \frac{3}{2}KT$

92. (a) $PV \quad nRT$

$PV \quad T$ at given mass.

93. (c) According to kinetic theory of gases collision between molecules are perfectly elastic.

94. (b) $KE_T \quad \frac{3}{2}nRT \quad \frac{3}{2}\frac{W}{M}RT$

$KE \quad \frac{T}{M}$

$$\frac{KE_1}{KE_2} \quad \frac{T_1}{M_1}\frac{M_2}{T_2} \quad \frac{300}{2} \quad \frac{44}{600} \quad \frac{11}{1}$$

95. (a) $PV \quad \frac{1}{3}mu^2$

$P \quad 1 \quad \frac{2}{3} \quad \frac{1}{2}mu^2$

$P \quad \frac{2}{3}E.$

96. (b) Moles ratio $\frac{n_{He}}{n_{N_2}} \quad \frac{w/4}{w/28} \quad \frac{7}{1}$

Ratio of translational K.E. $\frac{n_{He}\ T_{He}}{n_{N_2}\ T_{N_2}}$

$\frac{7}{1} \quad \frac{300}{700} \quad 3:1$

97. (d) Due to phase transfer distance between particle changes, because temperature remains constant, so KE remains constant.

98. (d) $K \quad \frac{3}{2}n_T RT \qquad n_T \quad \frac{3000 \quad 2}{3 \quad 250 \quad 8.314}$

$3 \quad 10^3 \quad \frac{3}{2}n_T \quad 8.314 \quad 250$

$\frac{x}{20} \quad \frac{30 \quad x}{40} \quad \frac{2}{8.314}$

$\%Ne \quad 28.3$

99. (b) $KE \quad \frac{3}{2}RT$

$U_{rms} \quad \sqrt{\frac{3RT}{M}}$

100. (d) $U_{rms} \quad \sqrt{\frac{U_1^2 \quad U_2^2 \quad U_3^2 \quad U_4^2}{4}} \quad \sqrt{\frac{54}{4}} \quad \frac{\sqrt{54}}{2}$

101. (c) $U_{ms} \quad \sqrt{\frac{4 \quad (6)^2 \quad 5(2)^2 \quad 10(3)^2}{19}}$

3.6 cm/sec.

For More Material Join: @JEEAdvanced_2024

102. (c) $U_{rms} = \sqrt{\frac{3RT}{Mw}}$

$\frac{U_{H_2}}{U_{O_2}} = \sqrt{\frac{50}{2} \cdot \frac{32}{800}} = 1$

104. (c) $U_{rms} = \sqrt{\frac{3RT}{M}}$ $\frac{(U_{rms})_{H_2}}{(U_{rms})_{N_2}} = \sqrt{\frac{T_{H_2}}{M_{H_2}} \cdot \frac{M_{N_2}}{T_{N_2}}}$;

$(U_{rms})_{H_2} = \sqrt{5}\,(U_{rms})_{N_2}$

$\frac{(U_{rms})_{H_2}}{(U_{rms})_{N_2}} = \sqrt{5} = \sqrt{\frac{T_{H_2}}{T_{N_2}} \cdot \frac{28}{2}}$

$\frac{\sqrt{5}}{1} = \sqrt{\frac{T_{H_2}}{T_{N_2}} \cdot 14}$

$5 = \frac{T_{H_2}}{T_{N_2}} \cdot 14$

$T_{N_2} \cdot 5 = T_{H_2} \cdot 14$

$T_{N_2} = T_{H_2}$

105. (a) $U_{rms} = \sqrt{\frac{2RT}{M}}$ $U_{avg} = \sqrt{\frac{8}{} \cdot \frac{RT}{M}}$ $U_{mp} = \sqrt{\frac{2RT}{M}}$

$U_{rms} : U_{avg} : U_{mp}$

1.224 : 1.128 : 1

$U_{rms} > U_{avg} > U_{mp}$

106. (b) $200 = \sqrt{\frac{2RT}{2 \times 10^{-3}}}$ $RT = 40$

Average K.E. $= \frac{3}{2}nRT = \frac{3}{2} \cdot \frac{8}{2} \cdot 40$

$= 240$ J

107. (c) Second member of C_nH_{2n} series

C_3H_6 42

$\sqrt{\frac{8RT_1}{M_1}} = \sqrt{\frac{8RT_2}{M_2}}$ $\frac{900}{71} = \frac{T_2}{42}$

$T_2 = 532.4$ K

108. (d) $U_{rms} = \sqrt{\frac{3RT}{M}}$

109. (b) $P = \frac{1}{3}\frac{nmc^2}{V}$

$\frac{1}{3} \cdot \frac{6 \times 10^{22} \times 10^{-24} \times (100)^2}{10 \times 10^{-3}}$

$= 2 \times 10^4$ Pa

110. (c) Second member is C_3H_4;

$\sqrt{\frac{2RT_1}{M_1}} = \sqrt{\frac{2RT_2}{M_2}}$

$T_1 = T_2 \cdot \frac{M_1}{M_2} = 800 \cdot \frac{40}{64}$ K

$= 500$ K or 227°C

111. (c) $300 = \sqrt{\frac{3RT}{4 \times 10^{-3}}}$; $RT = 120$

Total K.E. of He gas $= \frac{3}{2}nRT$

$= \frac{3}{2} \cdot \frac{8}{4} \cdot 120$ J

$= 360$ J

112. (b) $U_{mp} = \sqrt{\frac{2RT}{M}}$

$U_{avg} = \sqrt{\frac{8}{} \cdot \frac{RT_2}{M}}$

$U_{rms} = \sqrt{\frac{RT_1}{M}}$

$T_3 > T_2 > T_1$

113. (b) $U_1 = \sqrt{\frac{3P_1}{d_1}}$

$U_{rms} = \sqrt{\frac{3}{d}}\,(\sqrt{P_2} - \sqrt{P_1})$

$= \sqrt{\frac{3}{0.75}}\,(300 - 200)$

$= \sqrt{4} \cdot 100 = 200$

114. (d) Rate of diffusion $\propto \frac{1}{\sqrt{\text{density}}}$

at constant P and T

$d = \frac{PM}{RT}$, $d \propto M$

115. (d) $: v : u = \sqrt{2} : \sqrt{\frac{8}{}} : \sqrt{3}$

116. (b) $\sqrt{\frac{8RT}{M}} = \sqrt{\frac{28}{88}} = \sqrt{\frac{7}{22}}$

$\frac{8 \times 8.314 \times T}{16 \times 10^{-3}} = \frac{7}{22} \times 10^6$

$T = \frac{1000 \times 2}{8.314} = 240.55$ K

T °C $= 240.55 - 273 = -32.45$ °C

117. (d) $\sqrt{\frac{2RT}{M_{SO_2}}} = \sqrt{\frac{3R \times T_1}{M_{O_2}}}$ $\frac{2 \times T}{64} = \frac{3 \times 300}{32}$

$T = 900$ K

For More Material Join: @JEEAdvanced_2024

118. (b) $U_{rms} \quad \sqrt{\frac{3RT}{M}}$

119. (a) Rate of diffusion $\quad \frac{1}{\sqrt{M}} \quad P$

Rate of diffusion $\quad \frac{P}{\sqrt{M}}$

120. (b) According to Grahm's law

$r \quad \frac{1}{\sqrt{d}}, \qquad \frac{r_1}{r_2} \quad \sqrt{\frac{d_2}{d_1}}$ at constant P and T

121. (a) $\frac{r_1}{r_2} \quad \sqrt{\frac{M_2}{M_1}}$ at constant P and T

122. (b) $\frac{r_1}{r_{H_2}} \quad \sqrt{\frac{2}{M_1}} \quad \frac{1}{6}$

$M \quad 72$

123. (b) $\frac{r_1}{r_2} \quad \sqrt{\frac{M_2}{M_1}}$

$\frac{V_1}{t_1} \quad \frac{t_2}{V_2} \quad \sqrt{\frac{M_2}{M_1}}$

124. (b) $\frac{r_{H_2}}{r_{H.C}} \quad \sqrt{\frac{M}{2}} \quad 3\sqrt{3}$

$\frac{M}{2} \quad 9 \quad 3$

$M \quad 54$

$n \quad 4$

125. (b) $\frac{r_{O_2}}{r_{CH_4}} \quad \frac{n_{O_2}}{n_{CH_4}} \quad \sqrt{\frac{M_{CH_4}}{M_{O_2}}}$

$\frac{3}{2} \quad \frac{16}{32} \quad \sqrt{\frac{16}{32}} \quad \frac{3}{4\sqrt{2}}$

126. (c) $\frac{\frac{n_{SO_2}}{n_{CH_4}}}{\frac{8}{1}} \quad \frac{16}{64}^{3/2} \quad \frac{1}{1};$

$\frac{r_{SO_2}}{r_{CH_4}} \quad \frac{1}{1}\sqrt{\frac{16}{64}} \quad \frac{1}{2}$

127. (d) $\frac{n_{He}}{n_{CH_4}} \quad \frac{1}{2}\sqrt{\frac{16}{4}} \quad \frac{1}{1}$


$\frac{n_{He}}{n_{SO_2}} \quad \frac{1}{3}\sqrt{\frac{64}{4}} \quad \frac{4}{3}$

$n_{He} : n_{CH_4} : n_{SO_2} \quad 4 : 4 : 3$

128. (a) $\frac{r_1}{r_2} \quad \frac{V_1/t_1}{V_2/t_2} \quad \frac{V_1}{V_2}\frac{t_2}{t_1} \quad \sqrt{\frac{M_2}{M_1}};$

$\frac{80}{V_2}\frac{3}{2} \quad \sqrt{\frac{64}{32}} \quad \sqrt{2}; \quad V_2 \quad \frac{120}{\sqrt{2}}$

129. (a) $\frac{r_{NH_3}}{r_{HCl}} \quad \sqrt{\frac{36.5}{17}} \quad \frac{x}{87.5 \quad x}$



$x \quad 51.7$ cm

130. (c) $\frac{r_{CH_3OCH_3}}{r_{CH_4}} \quad \sqrt{\frac{M_{CH_4}}{M_{CH_3OCH_3}}} \quad \frac{P_{CH_3OCH_3}}{P_{CH_4}}$

$\sqrt{\frac{16}{46}} \quad \frac{0.8}{0.2} \quad 2.36 : 1$

131. (b) $\frac{n_1}{n_2}\frac{t_2}{t_1} \quad \sqrt{\frac{M_2}{M_1}}; \quad \frac{w_1}{M_1} \quad \frac{M_2}{w_2} \quad \sqrt{\frac{M_2}{M_1}}$

$\frac{w_1}{w_2} \quad \sqrt{\frac{M_1}{M_2}}; \quad \frac{w_1}{4} \quad \sqrt{\frac{4}{64}}$

$w_1 \quad 1$ gm

132. (d) $\frac{r_{mix}}{r_x} \quad \frac{4/5}{4/10} \quad 2 \quad \sqrt{\frac{M_x}{M_{Mix}}}$

$M_{mix} \quad 9$

$M_{mix} \quad M_{H_2}X_{H_2} \quad M_{CH_4}X_{CH_4}$

$2X_{H_2} \quad 16(1 \quad X_{H_2}) \quad 9$

$X_{H_2} \quad 0.5$

133. (a) Mass of the filled balloon $\quad 50 \quad 685$

735 kg

Pay load $\quad$ Mass of displaced air $\quad$ Mass of balloon

$5108 \quad 735 \quad 4373$ kg

134. (b) Energy associated with each degree of freedom is $\frac{1}{2}k_B T$

135. (c) $C_v \quad \frac{3}{2}R \quad R \quad 4R \quad \frac{3}{4} \quad \frac{11}{2}R;$

$C_p \quad \frac{11}{2}R \quad R \quad \frac{13}{2}R$

$\frac{C_p}{C_v} \quad \frac{13}{11} \quad 1.18$

For More Material Join: @JEEAdvanced_2024

136. (c) $C_v = \frac{\frac{3}{2}R + \frac{5}{2}R}{2} = 2R$

$C_p = \frac{\frac{5}{2}R + \frac{7}{2}R}{2} = 3R$

$\frac{C_p}{C_v} = 1.5$

137. (b) $C_v = \frac{3}{2}RT$ for monoatomic gas;

$C_p = \frac{5}{2}RT$

$C_v = \frac{5}{2}2RT$ for diatomic gas

$C_p = \frac{7}{2}RT$

Thus, for mixture of 1 mole each,

$$C_v = \frac{\frac{3}{2}RT + \frac{5}{2}RT}{2}$$

and $$C_p = \frac{\frac{5}{2}RT + \frac{7}{2}RT}{2}$$

Thus, $\frac{C_p}{C_v} = \frac{3RT}{2RT} = 1.5$

138. (a) Internal energy of a substance is a function of temperature and volume (except ideal gas). For ideal gas it is only function of temperature.

139. (c) For ideal gas $PV = nRT$

volume of 1 mole gas at 1 atm pressure and 273 K is 22.4 L and at 300 K and 1 atm pressure volume = 24.63 L.

140. (b) If attraction are disappear than pressure of real gas

141. (d) $P_{Ideal} - P_{Real} = \frac{n^2a}{V^2}$

142. (b) Real gas behaves like an ideal gas at low pressure and high temperature at this condition $Z = 1$.

143. (a) $P_{Ideal} - P_r = \frac{n^2a}{V^2}$

144. (c) $P_{Real} - P_i = \frac{n^2a}{V^2}$

$a \propto P_r$

$P_2 > P_3 > P_1$

145. (c) $P(V - b) = RT$; $P = \frac{RT}{(V - b)}$;

$$P = \frac{R}{(V - b)}T > 0$$

146. (c) $b = 4$ volume occupied by molecules of 1 mol of a gas,

or $b = 4N_A \frac{4}{3}r^3$

or $r = \left(\frac{3b}{16N_A}\right)^{1/3}$

$$\left(\frac{3 \times 24}{16 \times 6.023 \times 10^{23} \times 3.14}\right)^{1/3} \text{cm}$$

$= 1.355 \times 10^{-8}$ cm

Now, molecular diameter, $d = 2r$

$= 2.67 \times 10^{-8}$ cm

147. (c) Liquifaction I.M.A.F. $\propto a$

148. (d) $V_i = V - V_{ex}$

$V_{ex} = n.b$

$b \propto$ size

149. (d) A real gas can behave like an ideal gas at low pressure and high temperature.

150. (a) $P(V - nb) = nRT$

$PV - Pnb = nRT \quad d = \frac{\text{Mass}}{\text{Volume}}$

$PV = nRT + Pnb \quad n = \frac{\text{Mass}}{M}$

$$P\frac{\text{Mass}}{d} = nRT + Pnb$$

$$\frac{P}{d} = \frac{n}{\text{Mass}}RT + \frac{Pnb}{\text{Mass}}$$

$$\frac{P}{d} = \frac{RT}{M} + \frac{Pb}{M}$$

151. (c) $P + \frac{a}{V_m^2}(V_m - b) = RT$

152. (c) At low pressure,

$$P + \frac{a}{V^2}(V) = RT$$

i.e., $PV^2 - RTV + a = 0$

$$V = \frac{RT - \sqrt{R^2T^2 - 4Pa}}{2P} \approx \frac{RT}{2P}$$

($\because 4aP < R^2T^2$)

153. (d) Real gas shows ideal behaviour at low pressure and high temperature.

For More Material Join: @JEEAdvanced_2024

154. (a) $\frac{PV}{RT}$ Z Z 1 $\frac{a}{VRT}$

155. (b) van der Waals' equation for one mole of a real gas is

$$P \quad \frac{a}{V^2} \quad (V \quad b) \quad RT$$

or $PV \quad RT \quad Pb \quad \frac{ab}{V^2} \quad \frac{a}{V}$

At high pressures, the van der Waals' gas equation reduces to $PV \quad RT \quad Pb$

or $\frac{PV}{RT} \quad 1 \quad \frac{Pb}{RT}$

Compressibility factor $Z \quad \frac{PV}{RT} \quad 1 \quad \frac{Pb}{RT}$

156. (a) $\frac{PV}{RT} \quad 1 \quad RT$ constant

$PV \quad 1 \quad$ at 1 atm

$V_m \quad 22.4$ L

157. (c) $Z \quad \frac{PV}{nRT} \quad \frac{V}{n} \quad \frac{0.9 \quad 0.0821 \quad 273}{9}$

2.24 litre/mol

Volume of 1 milli-mole of gas 2.24 mL

158. (d) $Z \quad \frac{PV}{nRT}$; $1.90 \quad \frac{1 \quad 800}{n \quad R \quad 330}$;

$$n \quad \frac{1 \quad 800}{1.90 \quad R \quad 330}$$

$$Z \quad 1.10 \quad \frac{V \quad 200}{n \quad R \quad 570};$$

$$1.10 \quad \frac{V \quad 200 \quad 1.90 \quad R \quad 330}{800 \quad R \quad 570}$$

$V \quad 4$ L

159. (c) $Z \quad 1$ real gas shows positive deviation from ideal behaviour when repulsive forces dominate.

160. (d) $0.1 \quad \frac{1000 \quad (0.02)^2}{V^2} \quad V \quad 20 \quad 0.02$

$$0.1\,V^2 \quad 0.4\,V \quad 0.4 \quad 0$$

$$V^2 \quad 4V \quad 4 \quad 0$$

$V \quad 2$ L

$Z \quad \frac{PV}{nRT} \quad \frac{0.1 \quad 2}{20 \quad 0.02} \quad 0.5$

161. (c) $T_C \quad \frac{8a}{27\,Rb} \quad P_C \quad \frac{a}{27\,b^2} \quad V_C \quad 3b$

163. (a) $V \quad K(t \quad C \quad 273)$

1 C $V_1 \quad K(273 \quad 1) \quad K \quad 273 \quad K$

$V \quad (K \quad 273) \quad t \quad 0$ C

2 C $V_2 \quad K(273 \quad 2) \quad K \quad 273 \quad 2\,K$

164. (b) It is a temperature above which a gas can not be liquified.

165. (b) $\frac{T_C}{P_C} \quad \frac{8a \quad 27b^2}{27\,Rb \quad a} \quad \frac{8b}{R}$

$V_{\text{Excluded}} \quad nb$

166. (d) $T_{\text{Liquifaction}} \quad \frac{1}{\text{I.M.A.F}} \quad \frac{1}{a} \quad \frac{1}{T_C}$

Critical temperature is a temperature above which a gas can't be liquify.

167. (a) $T_C \quad \frac{8a}{27\,Rb}$, $T_b \quad \frac{a}{Rb}$, $T_i \quad \frac{2a}{Rb}$

$T_i \quad T_b \quad T_c$

168. (d) At low pressure $Z \quad 1 \quad b \quad \frac{a}{RT} \quad \frac{1}{V_m}$ at

Boyles temperature $b \quad \frac{a}{RT} \quad 0$

169. (b) Below critical temperature a gas can be liquified.

So V_m has three real roots.

170. (a) At critical temperature slope of pressure volume curve is maximum and it is zero.

171. (c) Above critical temperature a gas can't be liquified.

172. (c) At Boyle's temperature and low pressure $Z \quad 1$.

At low pressure $Z \quad 1 \quad b \quad \frac{a}{RT} \quad \frac{1}{V_m}$

At $T_B \quad a/Rb$, $Z \quad 1$

173. (a) Because $T \quad T_i$ and if gas expanded adiabatically then T

174. (a) The van der Waals' equation for 1 mole of gas is

$$P \quad \frac{a}{V^2} \quad (V \quad b) \quad RT$$

if we put, $P_r \quad \frac{P}{P_C}, V_r \quad \frac{V_m}{V_C}$ and $T_r \quad \frac{T}{T_C}$;

$$\because \quad P_C \quad \frac{a}{27b^2}, V_C \quad 3b \text{ and } T_C \quad \frac{8a}{27Rb}$$

$$P_r \quad \frac{a^2}{27b^2} \quad \frac{a}{V_r^2\,(3b)^2} \quad (V_r(3b) \quad b) \quad RT_r \quad \frac{8a}{27Rb}$$

This equation is called van der Waals' equation of law of corresponding states.

For More Material Join: @JEEAdvanced_2024

175. (b) $\frac{P_r V_r}{T_r}$ 2.21

$\frac{P}{P_C}\ \frac{V}{V_C}\ \frac{T_C}{T}$ 2.21

$\frac{PV}{T}\ \frac{T_C}{P_C V_C}$ 2.21

0.1243 L 124.31 mL

Level 2

1. (b) The number of moles of He, Ne and Xe is given by

n_{He} $\frac{PV}{RT}\ \frac{3}{RT}$; n_{Ne} $\frac{PV}{RT}\ \frac{10}{RT}$;

n_{Xe} $\frac{PV}{RT}\ \frac{1}{RT}$

n n_{He} n_{Ne} n_{Xe} $\frac{(3\ 10\ 1)}{RT}$

The total pressure is given by

P $\frac{(n_{He}\ n_{Ne}\ n_{Xe})\,RT}{V}$

P_{total} $\frac{(3\ 10\ 1)}{RT}\ \frac{RT}{7}$ 2 atm

2. (c) $N_2 + 3H_2 \rightleftharpoons 2NH_3$

Initial moles 11 12 0

at equilibrium 9 6 4

moles of N_2 and H_2 present at equilibrium

15

after addition of water


$NH_3(g)$ $H_2O(l)$ $NH_4OH(l)$

Volume of vessel available for gaseous mixture of N_2 and H_2 20 3.58

16.42 L

Pressure exerted by gaseous mixture at 300 K $\frac{15\ 0.821\ 300}{16.42}$ 22.5 atm

3. (c) I II



Let x mole of N_2 present into vessel II and P is final pressure of N_2

$P(2V)$ $xR(T_2/3)$ and $P(V)$ $(0.1\ x)RT_2$

2 $\frac{x}{3(0.1\ x)}$

x 0.6/7 mole;

$\frac{0.6}{7}$ 28 2.4 g N_2

II has 2.4 g N_2 and I has 0.4 g of N_2;

$\frac{W_I}{W_{II}}\ \frac{0.4}{2.4}$ 1 : 6

4. (b) Let initial mixture contains n_1 and n_2 moles of NH_3 and N_2H_4 respectively

Total moles of gases originally present

n_1 n_2

Total moles of gases after decomposition of gases $2n_1$ $3n_2$

0.5 V $(n_1\ n_2)\,R$ 300

4.5 V $(2n_1\ 3n_2)\,R$ 1200

$\frac{2n_1\ 3n_2}{n_1\ n_2}\ \frac{9}{4}$

$\frac{n_1}{n_2}\ \frac{1}{3}$

$\frac{n_2}{n_1\ n_2}$ 100 25%

6. (b) Let T_1 T_2; final pressure will be same, Let x mole transfer from A to B vessel.

$P_A V$ $(n\ x)\,RT$

and $P_A V$ $(n\ x)\,RT_2$

x $\frac{n(T_1\ T_2)}{T_1\ T_2}$;

finally P_1 $2V$ $2nRT_1$; V $\frac{nRT_1}{P_1}$

P_A $\frac{nRT_1}{P_1}$ n $\frac{n(T_1\ T_2)}{(T_1\ T_2)}$ RT_1

P_A $\frac{2P_1T_2}{T_1\ T_2}$

7. (c) P d; P kd and k $\frac{1\text{ atm}}{1\text{ metre}}$

PV nRT; kd $\frac{1}{6}$ d^3 nRT;

$\frac{d_1^4}{d_2^4}$ $\frac{n_1}{n_2}$; $\frac{1}{4^4}$ n_1/n_2; n_2 256

no. of moles added 256 1 = 255

8. (a) PM dRT

Pd P^2 $\frac{M}{RT}$

$\frac{d(Pd)}{dP}$ $\frac{2PM}{RT}$ 5

For More Material Join: @JEEAdvanced_2024

$$\frac{0.2 \quad 8.21 \quad 4}{0.0821 \quad T} \quad 5$$

$$T \quad 160 \text{ K}$$

9. (d) % Relative humidity

$$\frac{\text{Partial pressure of } H_2O}{\text{Vapour pressure of } H_2O} \quad 100$$

$$75 \quad \frac{P_{H_2O}}{30} \quad 100$$

P_{H_2O} 22.5 torr;

% of H_2O vapour in air

$$\frac{(22.5)}{760} \quad 100 \quad 2.96$$

molar mass of wet air

$$\frac{29 \quad 97.04 \quad 2.96 \quad 18}{100}$$

$$\frac{2814.16 + 53.28}{100} \quad 28.67$$

density of wet air $\frac{PM}{RT}$ $\frac{1 \quad 28.67}{0.0821 \quad 300}$

1.164 g/L

10. (b) C_v of gas A $\frac{3}{2}R$ $\frac{3}{2}R$ $6R$ $9R$

Let C_v of gas B X

Heat lost by gas A $7 \quad 9R \quad T$ $\frac{5T}{6}$

Heat gained by gas B $6X$ $\frac{5T}{6}$ $\frac{T}{3}$

$$63R \quad \frac{T}{6} \quad 6X \quad \frac{T}{2} \quad X \quad \frac{7}{2}R;$$

gas is diatomic

11. (a) Total heat received by gas = heat provided

$$2 \quad \frac{3}{2}R(T_f \quad 300) \quad 2 \quad 3\,R(T_f \quad 400)$$

$$3 \quad \frac{5}{2}R\,(T_f \quad 500) \quad 15600$$

T_f 900 K

$$P_f \quad \frac{900 \quad 0.0821 \quad 7}{7}$$

P_f 73.89 atm

12. (c) $1000 \quad \sqrt{\frac{3RT}{M}}; 10^3 \quad \frac{3 \quad 8.314 \quad T}{10^{-26} \quad 6 \quad 10^{23}}^{1/2}$

$$T \quad \frac{10^6 \quad 10^{-26} \quad 6 \quad 10^{23}}{3 \quad 8} \quad 250 \text{ K}$$

13. (b) Mass of balloon 100 kg 10 10^4 g

Volume of balloon $\frac{4}{3} \quad r^3$

$$\frac{4}{3} \quad \frac{22}{7} \quad \frac{21}{2}^3$$

4851 m^3 4851 10^3 L

Mass of gas (He) in balloon $\frac{PVM}{RT}$

$$\frac{1 \quad 4851 \quad 10^3 \quad 4}{0.082 \quad 300}$$

$78.878 \quad 10^4$

Total mass of gas and balloon

$78.878 \quad 10^4 \quad 10 \quad 10^4$

$88.878 \quad 10^4$ g or 888.78 kg

Mass of air displaced 1.2 4851

5821.2 kg

Pay load = mass of air displaced

mass of (balloon + gas)

Pay load 5821.2 888.78 = 4932.42 kg

14. (d) Let V mL of gas effused

$$\frac{V/220}{V/200} \quad \sqrt{\frac{d_{O_2}}{d_{mix}}} \quad d_{mix}1.6 \quad (1.1)^2 \quad 1.936 \text{ g/L}$$

Let density of ozone is d; In 100 volume ozonised oxygen, 60% O_2 and 40% by volume O_3 is present

Mass of mixture = mass of ozone

+ mass of oxygen

$$100 \quad 1.936 = 40 \quad d \quad 60 \quad 1.6$$

density of O_3 is 2.44 g/L

15. (d) n_{N_2} $\frac{\frac{708.2}{760} \quad 0.25}{0.0821 \quad 303}$ $9.36 \quad 10^{-3}$

$$n_{O_2} \quad \frac{\frac{756.2}{760} \quad 0.3}{(0.0821) \quad 298}$$

0.0122

n_{total} moles 0.02156

pressure in final vessel P

$$\frac{(n_{total})\,RT}{V} \quad \frac{0.02156 \quad 00821 \quad 308}{0.5}$$

P 1.09 atm or 828.4 torr

P_{total} $P_{(O_2 + N_2)}$ V.pr. of H_2O

828.4 42.2 = 870.6 torr

16. (b) $P \quad P_0 e^{-kt}$; P_0 $\frac{228}{3}$ 76 152 cm Hg

For More Material Join: @JEEAdvanced_2024

At t 14 min; P $\frac{114}{3}$ 76 114 cm Hg

$152e^{-kt}$ 114

k $\frac{1}{t}\ln\frac{152}{114}$ $\frac{1}{14}$ 0.28

0.02 60 hr $^{-1}$ 1.2 hr $^{-1}$

17. (d) P_{H_2O} 760 722 $\frac{38}{760}$ atm; n_{H_2O} $\frac{0.9}{18}$

V $\frac{0.9}{18}$ $\frac{0.821 \quad 300}{38}$ 760; V 24.63 L

18. (c) n_{AB} $\frac{3 \quad 8.21}{0.0821 \quad 300}$ 1;

n_{B_2} $\frac{1 \quad 8.21}{0.0821 \quad 400}$ 0.25

	$2AB(g)$	$B_2(g)$	$2AB_2(s)$
Initial moles	1	0.25	
After reaction	0.5	0	

P_{AB} $(8.21 \quad 8.21) = 0.5 \quad 0.0821 \quad 250$

P_{AB} 0.625 atm

19. (c) Moles of O_2 remaining after reaction

4 1 3

V n is due to same T and P in each compartment after reaction.

Length of He compartment

$\frac{5}{5 \quad 3}$ 500 312.5 cm

20. (b) V_c $\frac{60}{0.80}$ 75 cm ^{3} mol $^{-1}$;

b $\frac{V_c}{3}$ 25 cm ^{3} mol $^{-1}$ 0.025 L mol $^{-1}$

T_c $\frac{8a}{27Rb}$

$\frac{4 \quad 10^5}{821}$ $\frac{8 \quad a}{27 \quad 0.0821 \quad 0.025}$

a 3.375

21. (b) b 4 $\frac{4}{3}$ r^3 N_A;

4 10^{-4} 1000

4 $\frac{4}{3}$ r^3 $6 \quad 10^{23}$

r $5 \quad 10^{-9}$ cm

Distance of closest approach $2r$

10^{-8} cm or 10^{-10} m

22. (c) $\frac{r_x}{r_{O_2}}$ $\sqrt{\frac{M_{O_2}}{M_x}}$ $\frac{4}{5}^2$ $\frac{32}{M_x}$ M_x 50

d_x 0.80 kg / m ^{3},

V_m $\frac{1000}{800}$ 50 62.5 L

Z $\frac{PV_m}{RT}$ $\frac{1 \quad 62.5}{0.0821 \quad 500}$ 1.52

23. (a) b 25 cm ^{3}/mol

b 0.025 L/mol

V_{Real} V_i nb 50 10 0.025 49.75 L

24. (b) Z 1 $\frac{Pb}{RT}$ at high pressure

slope $\frac{b}{RT}$ $\frac{}{492.6}$

b $\frac{}{492.6}$ 0.0821 300 and

b $\frac{4}{3}$ r^3 $4N_A$

$\frac{}{492.6}$ 0.0821 300 10^{-3}

r 2.5 Å or d 5 Å

26. (a) n $\frac{P}{RT}$ V

Slope of line $\frac{P}{RT}$ tan

T $\frac{P}{R\tan}$ $\frac{16.42}{0.0821\tan}$

For B, T $200\sqrt{3}$ K

For A, T $\frac{200}{\sqrt{3}}$ K

Level 3

Passage-1

1. (d) p $\frac{a}{V_m^2}$ $(V_m \quad b)$ RT

P V_m P $\frac{a}{V_m^2}$ p

$P(V_m \quad b)$ RT

PV_m RT Pb

2. (a) V P

V nb V

P $\frac{n^2a}{V^2}$ V nRT

PV $\frac{n^2a}{V}$ nRT

PV nRT $\frac{n^2a}{V}$

For More Material Join: @JEEAdvanced_2024

3. (c) van der Waal's constant 'a' I.M.A.F.
So CO_2 gas can be more easily liquified.

4. (d) $P \quad \frac{n^2a}{V^2} \quad (V \quad nb) \quad nRT$

Passage-2

2 & 3. Let $_A$, $_B$ and $_C$ be collision diameters of A, B and C

$$\frac{kT}{\sqrt{2} \quad {}^2P}, \text{ i.e., } \quad \sqrt{\frac{T}{P}}$$

$$\sqrt{\frac{1600}{0.16 \quad 1}}^{A} : \sqrt{\frac{200}{0.16 \quad 2}}^{B} : \sqrt{\frac{400}{0.04 \quad 4}}^{C}$$

$4 : 1 : 2$

2. (c) $Z_{11} \quad \frac{{}^2P^2}{\sqrt{m}T^{3/2}}$

$$A:B:C \quad \frac{16 \quad 1}{\sqrt{20} \quad (1600)^{3/2}} : \frac{1 \quad 4}{\sqrt{40} \quad (200)^{3/2}} : \frac{4 \quad 16}{\sqrt{80} \quad (400)^{3/2}}$$

$= 1 : 4 : 16$

3. (a) $Z_1 \quad \frac{({}^2)P}{\sqrt{MT}}$

$$A : B : C \quad \frac{0.6 \quad 1}{\sqrt{20 \quad 1600}} : \frac{1 \quad 2}{\sqrt{40 \quad 200}} : \frac{4 \quad 4}{\sqrt{80 \quad 400}}$$

$4 : 1 : 4$

Passage-3

1. (a) $h_L d_L \quad h_{Hg} d_{Hg}$

$$h_L \quad \frac{76 \quad 13.6}{5.44} \quad 190 \text{ cm}$$

2. (d) $P_{Gas} \quad P_{Atm} \quad P_L \quad 1 \quad \frac{38}{190} \quad 1.2$

3. (b)

$$3A(g) \rightleftharpoons A_3g$$

$t \quad 0$	1.2 atm	$A_3(g)$	
$t \quad t_{eq.}$	$1.2 \quad 0.36$	$\frac{1}{3}(0.36)$	0.12 atm

$P_T \quad 1.2 \quad 0.36 \quad 0.12 \quad 0.96$ atm

Pressure difference in column

$1 \quad 0.96 \quad 0.04$ atm

The difference in height of the liquid level in two columns $0.04 \quad 190 \quad 7.6$ cm

One or More Answers is/are Correct

1. (a, b, c, d)

For Boyle's law

$T \quad$ constant

$PV \quad K \quad$ constant

2. (a, b)

$PV \quad nRT$

$T \quad$ constant

$PV \quad$ constant

$V \quad P$

KE is a function of temperature.

3. (b, c, d)

T_c is a temperature above which a gas can not be liquified at this point, slope of $P \quad V$ curve is maximum and attraction force is dominating at this point. Both liquid vapour are exist in equilibrium.

4. (c, d)

at constant volume	at constant P
$P \quad T$	$V \quad T$
$P \quad KT$	$V \quad KT$
$P \quad K(t \quad C \quad 273.15)$	$V \quad K(t \quad C \quad 273.15)$

5. (a)

T_B is a temperature at which gas behave like an ideal gas mean attractive and repulsing forces just offset each other.

6. (a, c, d)


$KE \quad \frac{3}{2} RT \,(1 \text{ mole}) \quad U_{rms} \quad \sqrt{\frac{3RT}{M_w}}$

$d \quad \frac{PM_w}{RT}$

7. (a, b)

Refer. Q.3

8. (a, b, c, d)



9. (a) Refer Q.2

10. (a, b, c)

$PV \quad nRT$

R is constant and independent of P and T

$R \quad K \quad N_A$

K Boltzmann's constant

For More Material Join: @JEEAdvanced_2024

11. (a,b,d)

(*A*) ∵ area under the curve gives fraction of molecules and total area is constant.

(*B*) u_{mp} decreases with decrease in temperature.

(*C*) T_2 is higher temperature

(*D*) As seen from graph; *A, B, D*

12. (a, d)

Vapour pressure of liquid is only and only temperature dependent.

13. (a, b, d)

For H_2 gas IMAF 0

Z 1 Z 1 $\frac{Pb}{RT}$

at low pressure Z 1 Z 1 $\frac{a}{RTV_m}$

Real gas behave like an ideal gas at low pressure and high temperature.

14. (a, b, c)

T_C $\frac{8a}{27Rb}$ Refer. Q.14

15. (a,b,d)

P_T (1 3) 1 3 0.1 1.3 atm.

P 0.2 atm or 76 0.3 cm of Hg

or 760 0.3 mm of Hg

16. (a, b, c, d)

Refer. Q. No. 13

17. (a, d)

Critical pressure is the minimum pressure at which substance convert into liquid state at critical point.

18. (a,c)

Low P and high temperature gas behaves as an ideal gas.

PV constant and $\frac{PV_m}{RT}$ 1 .

19. (b, c)

For ideal gas Z 1

at low pressure and Boyle's temperature real gas behave like an ideal gas.

20. (a, b, c)

V T at constant P

KE $\frac{3}{2}RT$ (1 mole)

Match the Column

5. y $\frac{1}{V^2}$ or $\sqrt{y}$ $\frac{1}{V}$

P x and P $\frac{\text{constant}}{V}$

(A) x $(k)\sqrt{y}$ y k^1x^2

(B) V kT; y V & $\frac{1}{T}$ x y $\frac{k}{x}$

(C) P kT; PT kT^2 or y kx

(D) v $\frac{c}{p}$ y $c\sqrt{x}$; y^2 cx

Subjective Problems

1. Refer Q.No. 59 Level-1

2. For H_2 and He.

f^{4th} $\frac{x/y}{1}$ $\sqrt{\frac{M_{He}}{M_{H_2}}}^4$

$\frac{x}{y}$ $(\sqrt{2})^4$ 4


3. Equation of line

P 2 $\frac{10\ \ 2}{4\ \ 15}(V\ \ 15)$

P 2 $\frac{8}{11}(V\ \ 15)$

P 2 $\frac{8V}{11}$ $\frac{15\ \ 8}{11}$

P $\frac{142}{11}$ $\frac{8V}{11}$



$f(T)$ $\frac{1}{nR}$ $\frac{142V}{11}$ $\frac{8V^2}{11}$

$\frac{d(F(T))}{dV}$ $\frac{1}{nR}$ $\frac{142}{11}$ $\frac{8V^2}{11}$ 0

V $\frac{142}{11\ \ 16}$ 8.875


P $\frac{142}{11}$ $\frac{8}{11}$ $\frac{8.875}{16}$ $\frac{71}{11}$

T_{Max} $\frac{PV}{nR}$ $\frac{71}{11}$ $\frac{142}{16\ \ 0.0821}$ 700

7 10^2 K

For More Material Join: @JEEAdvanced_2024

4. $T \quad 300 \quad \frac{4 \quad 3}{100}(P \quad 3)$



$T \quad 300 \quad \frac{P}{100} \quad \frac{3}{100}$

Multiply by 100

$T \quad 100 \quad 3 \quad 10^4 \quad P \quad 3$

$P \quad 3 \quad 3 \quad 10^4 \quad 100\,T$

$\frac{dP}{dT} \quad 0 \quad 100 \quad 1 \quad 10^2$

$x \quad y \quad 3$

5. $P \quad \frac{n^2a}{V^2} \quad P_i$

$\frac{n^2a}{V^2} \quad 0.04$

$a \quad 4$

6. $Z \quad 1 \quad \frac{Pb}{RT}$ high pressure

$\frac{dZ}{dP} \quad \frac{b}{RT} \quad \frac{1}{2.8}$

$b \quad \frac{RT}{2.8} \quad \frac{22.4}{2.8} \quad 4 \quad N_A \quad \frac{4}{3} \quad R^3$

$N_A \quad \frac{4}{3} \quad R^3$ Volume of 1 mole gas

$\frac{5.6}{2.8} \quad 2$

7. $P \quad \frac{dRT}{M_w}; \quad \frac{p_x}{p_y} \quad \frac{d_x}{d_y} \quad \frac{M_y}{M_x} \quad 6$

8. $\frac{r_x}{r_{O_2}} \quad \sqrt{\frac{M_{O_2}}{M_x}}$

$\frac{V}{2} \quad \frac{5.65}{V} \quad \sqrt{\frac{32}{M}}$

$8 \quad \frac{32}{M}$

$M \quad 4$

9. $n_{Br_2} \quad \frac{1 \quad 0.423}{0.0821 \quad 423} \quad \frac{1}{82.1}$ mole

POAC on 'Br'

$\frac{1}{82.1} \quad 2 \quad \frac{4.2}{80 \quad 19n} \quad 1$

$n \quad 5$

10. $n_T \quad n_1 \quad n_2$

$\frac{910 \quad (10 \quad V)}{RT} \quad \frac{950 \quad V}{RT} \quad \frac{900 \quad 10}{RT}$

$V \quad 4\,L$

11. $P_f \quad 1 \quad \frac{36}{76} \quad \frac{112}{76}$ atm. Final height $\quad 19$ cm

$P_i \quad 1$ atm, initial length $\quad h_i$ cm

Boyle's law $\quad P_iV_i \quad P_fV_f$

$1 \quad h_iA \quad \frac{112}{76} \quad 19A$

$h_i \quad 28$ cm

The length by which the Hg column shifts down $\quad h_i \quad h_f$

$28 \quad 19 \quad 9$ cm

12. $P_i \quad 722$ mm $\qquad P_f \quad 760$ mm

$T_i \quad 107 \quad 273 \quad 380$ K $\qquad T_f \quad 100$ K

$d_i \quad 1_g/\text{cm}^3 \qquad d_f \quad ?$

$\frac{d_iT_iR}{P_i} \quad \frac{d_fT_fR}{P_f}$

$d_f \quad \frac{P_f}{P_i} \quad \frac{T_i}{T_f} \quad d_i \quad \frac{760}{722} \quad \frac{380}{100} \quad 1\,g/\text{cm}^3$

$4\,g/\text{cm}^3$

13. V, n constant.

$\frac{P_i}{T_i} \quad \frac{P_f}{T_f} \qquad P_f \quad \frac{T_f}{T_i}P_i \quad \frac{109}{100} \quad P_i$

$P_{\text{increases}} \quad P \quad P_f \quad P_i \quad \frac{9}{100}P_i$

% Pressure increases

$\frac{P_f}{P_i} \quad 100 \quad \frac{9P_i}{100P_i} \quad 100\%$

$X\% \quad 9\%$

$X \quad 9$

14. $T_B \quad \frac{a}{Rb} \quad \frac{3.6}{0.08 \quad 0.6} \quad 75$ K , $\frac{75}{15} \quad 5$ K

15. $3\,\text{ms}^{-1}$

$u_{\text{rms}} \quad \sqrt{\frac{u_1^2N_1 \quad u_2^2N_2}{N_1 \quad N_2}} \qquad u_{\text{rms}}^2 \quad \frac{u_1^2N_1 \quad u_2^2N_2}{N_1 \quad N_2}$


$25 \quad \frac{4 \quad 7^2 \quad u_2^2}{10} \qquad u_2 \quad \sqrt{\frac{54}{6}} \quad 3\,\text{ms}^{-1}$

For More Material Join: @JEEAdvanced_2024

4

THERMODYNAMICS

- The subject of **Thermodynamics** deals basically with the interaction of one body with another in terms of quantities of heat & work.
- A **System** is defined as that macroscopic part of the universe which is at the moment under investigation.
- **Surroundings** are the rest of the universe outside the system.
- **Boundary :** The actual or imaginary surface that separates the system from the surroundings.



For More Material Join: @JEEAdvanced_2024

Thermodynamic Properties

Intensive Properties
do not depend on the amount of matter present in system

Examples : Pressure, Temperature, Density, Melting point, Boiling point etc.

Extensive Properties
depend on the amount of matter present in the system.

Examples : Mass, Volume, Heat Capacity, Enthalpy, Entropy etc.

Thermodynamic Process : The change of a system from one state to another state.

Thermodynamic Processes

Isothermal	Isobaric	Isochoric	Adiabatic	Polytropic
temperature remains constant throughout the process *i.e.*, : $dT=0$	pressure remains constant throughout the process *i.e.*, : $dP=0$	volume remains constant throughout the process *i.e.*, : $dV=0$	system does not exchange heat with surrounding *i.e.*, : $dq=0$	PV^n = constant

- **Cyclic Process :** System undergoes a number of different processes and finally returns to its initial state.
- **REVERSIBLE PROCESS (QUASI-STATIC) & IRREVERSIBLE PROCESS :**
  A process which is carried out so slowly that the system and the surroundings are nearly in equilibrium is known as a **reversible Process.** If this condition does not hold good, the process is said to be, **irreversible.**
  In a reversible process the driving force is infinitesimally larger than the opposing force. If the driving force is made infinitely smaller than opposing force, the system can be brought back without producing any permanent change.
- **All natural processes are irreversible process.**
  Spontaneous Process : Proceeds on its own *i.e.*, without any external help.
- **INTERNAL ENERGY (*U*) :** The sum of all microscopic forms of energy.
  $U \quad U_{\text{Translational}} \quad U_{\text{Rotational}} \quad U_{\text{Vibrational}} \quad U_{\text{Bonding}} \quad U_{\text{Electronic}} \quad \ldots\ldots$
- U is a state function & an extensive property.
- $U \quad U_{\text{Final}} \quad U_{\text{Initial}}$; $U \quad q_v$
- $dU \quad \left(\frac{U}{T}\right)_v dT \quad \left(\frac{U}{V}\right)_T dV$, $dU \quad nC_{Vm}(T_2 \quad T_1)$ at constant volume

For More Material Join: @JEEAdvanced_2024

Enthalpy

Chemical reactions are generally carried out at constant pressure (atmospheric pressure) so it has been found useful to define a new state function called Enthalpy (H) as :

- H U PV ; H U (PV) ; at constant pressure H U P V ; combining with first law. H q_p
- The difference between H & U becomes significant only when gases are involved (insignificant in solids and liquids)


$$H \quad U \quad (PV) \quad U \quad nR \quad T$$
$$H \quad U \quad (\ n_g)\,RT$$

- For a given system consider; H $f(T,P)$; dH $\left(\frac{H}{T}\right)_P dT$ $\left(\frac{H}{P}\right)_T dP$
- For isobaric process : dP 0 ; dH $\left(\frac{H}{T}\right)_P dT$; dH $C_P.dT$; H $C_P.dT$
- For an ideal gas H $f(T)$ only

$\left(\frac{H}{P}\right)_T$ 0 or dH $C_P.dT$; H $C_P.dT$; for n moles of ideal gas H $nC_P\,dT$

Kirchoff's Equation

- For chemical reactions d $_rH$ $_rC_p.dT$
- $_rH_{T_2}$ $_rH_{T_1}$ $_rC_p(T_2$ $T_1)$
- Similarly $_rU_{T_2}$ $_rU_{T_1}$ $_rC_v(T_2$ $T_1)$



- **ZEROTH LAW OF THERMODYNAMICS**

Two systems in thermal equilibrium with a third system, are also in thermal equilibrium with each other.

- **FIRST LAW OF THERMODYNAMICS**


It is law of conservation of energy. Mathematically for **closed system** at rest in absence of external fields this law is written as :

$$\boldsymbol{U \quad q \quad w}$$

Where U is change in internal energy of the system, q is the transfer of heat from surroundings to the system and w is the work involved (either done on the system or by the system).

For More Material Join: @JEEAdvanced_2024

❖ **Sign convention according to IUPAC :**



❖ **HEAT :** The energy that flow into or out of a system because of a difference in temperature between the system and its surrounding.

❖ **TYPES OF WORK :** Two types of work normally come across in chemistry.

1. Mechanical work (*e.g.*, PV work) **2.** Electrical work (Non PV work)

❖ *Mechanical work* is involved when a system changes its volume in the presence of an externally applied pressure (*i.e.*, pressure volume work). It is especially important in system containing gases

$$dw \quad P_{ext} . dV \; ; \quad w \quad \int_{V_1}^{V_2} P_{ext.} \, dV$$

❖ **CALCULATION OF WORK DONE IN VARIOUS PROCESS**

❖ **Isochoric process:**

Since $dV \quad 0$ so $w \quad 0$

from 1st law $U \quad q_v \quad w \quad q_v$

❖ **Isobaric process:**

$$w \quad P_{ext}(V_2 \quad V_1)$$

& $H \quad q_p$

❖ **Isothermal process:**

Since $dT \quad 0$ so $dU \quad 0$ for an ideal gas

from I[st] law $q \quad w \quad 0$ or $q \quad w$

If process is reversible

$$w \quad nRT \ln \frac{V_2}{V_1} \quad nRT \ln \frac{P_1}{P_2} \quad \text{or} \quad w \quad 2.303 nRT \log \frac{V_2}{V_1} \quad 2.303 nRT \log \frac{P_1}{P_2}$$

If process is irreversible

$$w \quad P_{ext} \quad \frac{nRT}{P_2} \quad \frac{nRT}{P_1}$$

In case of free expansion $p_{ext} \quad 0$ so $w \quad 0$

❖ **Adiabatic process :** $q \quad 0$

$$U \quad w \quad nC_v(T_2 \quad T_1) \quad \frac{P_2V_2 \quad P_1V_1}{1}, \text{ where } \quad \frac{C_P}{C_V}$$


If process is reversible $T_1V_1^{\quad 1} \quad T_2V_2^{\quad 1}$

For More Material Join: @JEEAdvanced_2024

If process is irreversible $\frac{T_2 \quad T_1}{1} \quad P_{ext} \cdot \frac{T_2}{P_2} \quad \frac{T_1}{P_1}$

❖ **Cyclic process:**

Work done Area enclosed in PV-diagram
For clockwise it is ve
For anti-clockwise it is ve



❖ Electrical work ($w \quad V.I.t$) is a type of non PV-work.

Heat Capacity : Between any two temperatures, the quantity of heat required to raise the temperature of the system from the lower to the higher temperature divided by the temperature difference.

❖ $C \quad dq/dT$.

❖ $q_v \quad U$

❖ $C_v \quad \frac{q}{T_2 \quad T_1}_v \quad \frac{U}{T_2 \quad T_1}_v$ or $C_v \quad (U/T)_v$

❖ $C_P \quad \frac{H}{T}_P$

❖ $C_{p,m} \quad C_{v,m} \quad R$ (for ideal gas)

❖ $C_{p,m}$ and $C_{v,m}$ depends on temperature even for an ideal gas. ($C \quad a \quad bT \quad cT^2$)

❖ $_{mix} \quad \frac{n_1 C_{p_1} \quad n_2 C_{p_2} \quad \ldots\ldots}{n_1 C_{v_1} \quad n_2 C_{v_2} \quad \ldots\ldots}$

Reversible Adiabatic process involving ideal gas

❖ $T_1 V_1^{\ 1} \quad T_2 V_2^{\ 1}$

❖ $TV^{\ 1}$ constant

❖ PV constant

❖ $T \ P^{1}$ constant

❖ **Molar Specific heat (C_m) of ideal gas in any Reversible Polytropic Process :**

$C_m \quad C_{v,m} \quad \frac{R}{1 \quad x} \quad \frac{R}{1} \quad \frac{R}{1 \quad x}$

Second Law of Thermodynamics

Statements :

(i) No cyclic engine is possible which take heat from one single source and in a cycle completely convert it into work without producing any change in surroundings.

(ii) In an irreversible process, entropy of universe increases but it remains constant :

in a reversible process


$S_{sys} \quad S_{surr} \quad 0$ for reversible process

$S_{sys} \quad S_{surr} \quad 0$ for irreversible process

$S_{sys} \quad S_{surr} \quad 0$ (In general)

For More Material Join: @JEEAdvanced_2024

(iii) Efficiency of Carnot engine working reversibly is maximum.



Carnot cycle

AB Isothermal Reversible Expansion $w_{AB} \quad nRT_2 \ln \frac{V_2}{V_1}$

BC Adiabatic Reversible Expansion $w_{BC} \quad C_V(T_1 \quad T_2)$

CD Isothermal Reversible Expansion $w_{CD} \quad nRT_1 \ln \frac{V_4}{V_3} \quad T_2 \quad T_1$

DA Adiabatic Reversible Expansion $w_{DA} \quad C_V(T_2 \quad T_1)$

for Carnot efficiency ( ) calculation $\frac{w_{\text{Total}}}{q_2} \quad \frac{T_2 \quad T_1}{T_2} \quad \frac{q_2 \quad q_1}{q_2}$

Calculation of Entropy change (S or S_{sys})

Ideal gas in any Reversible/Irreversible process from (P_1, V_1, T_1) to (P_2, V_2, T_2)

❖ $S \quad nC_v \ln \frac{T_2}{T_1} \quad nR \ln \frac{V_2}{V_1} \quad nC_p \ln \frac{T_2}{T_1} \quad nR \ln \frac{P_1}{P_2}$

❖ Reversible & irreversible isothermal expansion / contraction of an ideal gas

$$S \quad nR \ln \frac{V_2}{V_1}$$

❖ Isobaric heating or cooling : $S \quad nC_P \ln \frac{T_2}{T_1}$

❖ Isochoric heating or cooling : $S \quad nC_V \ln \frac{T_2}{T_1}$

❖ Adiabatic process :

❖ $S \quad nC_V \ln \frac{T_2}{T_1} \quad nR \ln \frac{V_2}{V_1}$ for irreversible process

$S \quad 0$ for reversible adiabatic compression and expansion.

❖ General reversible heating or cooling of any substance

$$S \int_{T_1}^{T_2} \frac{dq_{\text{rev}}}{T}$$

❖ Reversible Phase Transformation

(a) $S_{\text{Fusion}} \quad H_{\text{Fusion}}/T_m$, T_m is normal melting point (K)

(b) $S_{\text{vap}} \quad H_{\text{vap}}/T_B$, T_B is normal boiling point (K)

(c) $S_{\text{sub}} \quad H_{\text{sub}}/T_{\text{sub}}$, T_{sub} is sublimation temperature (K)

(d) $S_{\text{Trans}} \quad H_{\text{Trans}}/T_{\text{Trans}}$, T_{Trans} is the phase transition temperature (K)

❖ For chemical reaction

$$_rS \quad \nu_P \quad _fS \text{ (products)} \quad \nu_R \quad _fS \text{ (reactants)}$$

For More Material Join: @JEEAdvanced_2024

Third Law of Thermodynamics

"At absolute zero temperature, the entropy of a perfectly crystalline substance is taken as zero", which means that at absolute zero every crystalline solid is in a state of perfect order and its entropy should be zero.

By virtue of the third law, the absolute value of entropy (unlike absolute value of enthalpy) for any pure substance can be calculated at room temperature.

$$S_T \quad S_{0K} \quad \int_0^T \frac{q_{rev}}{T}, \text{ since } S_{0K} \quad 0 \; ; \; S_T \quad \int_0^T \frac{q_{rev}}{T}$$

Absolute entropies of various substances have been tabulated and these value are used to calculate entropy changes for the reactions by the formula;

$$S \quad \nu_P S^\circ_m \text{ (products)} \quad \nu_R S^\circ_m \text{ (reactants)}$$

Variation of S_r with temperature & pressure : $dS_r \quad \frac{{}_rC_{p,m}.dT}{T}$

$${}_rS_{T_2} \quad {}_rS_{T_1} \quad {}_rC_{p,m} \ln \frac{T_2}{T_1}$$

Gibbs Free Energy (G) and Spontaneity

- A new thermodynamic state function G, the Gibbs free energy is defined as :
  $G \quad H \quad TS$ or $G \quad H \quad T \; S$ (at constant temperature and pressure)
- $G \quad 0$ for spontaneous process
- $G \quad 0$ for equilibrium.

Criteria for Spontaneity

For a feasible process $S_T \quad S_{sys} \quad S_{surr.} \quad 0$

In absence of non PV work at constant T and P for a feasible process $G \quad 0$.

Standard Free Energy Change (G)

The free energy change for a process at a specified temperature in which the reactants in their standard state are converted to the products in their standard state. It is denoted by G .

- **Standard states :**
  (a) Standard temperature (any specified temperature)
  (b) For gases/solid/liquid: Standard pressure $P \quad 1$ bar
  (c) For ion/substance in solution: Standard concentration (C) $\quad 1M$

Like the standard enthalpy of formation of an element "the standard free energy of formation of an element in its standard state is zero". And so;

- ${}_rG \quad \nu_P \quad {}_fG$ (products) $\quad \nu_R \quad {}_fG$ (reactants)

A Thermodynamic Relationship

For any reversible reaction : $G \quad G \quad RT \ln Q$.

where Q is reaction quotient.

At equilibrium $G \quad 0, \quad Q \quad K$

$G \quad RT \ln K, \quad G \quad 2.303 \, RT \log K_{eq.}$


For More Material Join: @JEEAdvanced_2024

Effect of Temperature : Van't Hoff Equation

(a) $\frac{d(\ln K)}{dT}$ $\frac{H}{RT^2}$ (b) $\frac{d(\ln K)}{d\ \frac{1}{T}}$ $\frac{H}{R}$

G H $T\ S$ $\ln K$ $\frac{H}{RT}$ $\frac{S}{R}$

Integrated form $\ln \frac{K_2}{K_1}$ $\frac{H}{R}$ $\frac{1}{T_2}$ $\frac{1}{T_1}$



❖ **RELATIONSHIP BETWEEN G & $w_{\text{non-}PV}$**

dU dq dw_{PV} $dw_{\text{non-}PV}$

for reversible process at constant T & P

dU pdV TdS $dw_{\text{non-}PV}$

dH TdS $dw_{\text{non-}PV}$

$(dG_{\text{system}})_{T,P}$ $dw_{\text{non-}PV}$

$(dG_{\text{system}})_{T,P}$ $(dw_{\text{non-}PV})_{\text{by system}}$

Means free energy change for a process is equal to the maximum possible work that can be derived from the process *i.e.*,

❖ G $w_{\max}$ (for a reversible change at constant pressure and temperature)

❖ In case of a galvanic cell. G nFE_{cell}, where E_{cell} e.m.f. of the cell ; F Faraday constant and n number of electrons being transferred in the chemical process

❖ So G $nF\,E_{\text{cell}}$, where E°_{Cell} is the standard cell potential.

THERMOCHEMISTRY

❖ **Enthalpy of Reaction :**

The amount of heat evolved or absorbed when the number of moles of reactants react completely to give the products as given by the balanced chemical equation.

❖ **Enthalpy of formation :**

The change in heat enthalpy when one mole of a substance is formed directly from its constituent elements at standard states.

(i) For elements, ${}_fH$ 0 ; *e.g.*, ${}_fH$ (Zn, S) 0, ${}_fH$ (Br_2,*l*) 0 , ${}_rH$ (O_2, *g*) 0

(ii) If element exists in two or more allotropic forms then for most stable state ${}_fH$ 0

e.g., ${}_fH$ [C(*s*, graphite)] 0 , ${}_fH$ [S(*s*, rhombic)] 0

❖ **Enthalpy of combustion :**

The change in enthalpy when one mole of a substance is completely burnt in oxygen.

❖ **Enthalpy of Transition :**

The enthalpy change when one mole of one allotropic form changes to another.

For example : C(graphite) C(diamond) ; H_{trs} 1.90 kJ mol ^{1}.

❖ **Enthalpy of fusion :**

The change in enthalpy when one mole of a solid is completely converted into liquid state at its melting point.

For More Material Join: @JEEAdvanced_2024

❖ **Enthalpy of vaporisation :**

The change in enthalpy when one mole of a liquid is completely converted into its vapours at its boiling point.

❖ **Enthalpy of sublimation :**

The change in enthalpy when one mole of a solid is directly converted into vapours at sublimation temperature.

❖ **Enthalpy of solution :**

The change in enthalpy when one mole of a substance is dissolved in excess of water so that further dilution does not involve any heat change.

❖ **Enthalpy of Dilution :**

The change in enthalpy when a solution containing 1 mole of solute is diluted from one concentration to another.

❖ **Enthalpy of Hydration : (For anhydrous salt)**

The enthalpy change when 1 mole an hydrous salt combines with the requisite amount of water to form a new hydrated stable salt. For example, the hydration of anhydrous cupric sulphate is represented by

$$CuSO_4(s) \quad 5H_2O(l) \qquad CuSO_4 \quad 5H_2O(s)$$

❖ **Enthalpy of neutralisation :**

Enthalpy of neutralisation is the change in enthalpy when one gram equivalent of an acid is completely neutralised by one gram equivalent of a base in dilute solutions.

Strong acid Strong base Salt Water, H_{neut} 56 kJ mol ^{1}

❖ **Enthalpy of formation of ions :**

The enthalpy change when one mole of aqueous ions is obtained from element in its standard state as.

$$\frac{1}{2}Cl_2(g) \quad aq \qquad Cl \ (aq); \ H_r \quad H_r(Cl \ , aq)$$

By convention, the standard enthalpy of formation of H (aq) is taken to be zero.

❖ **Hess's law of constant heat summation :**

The total heat change in a particular reaction is the same whether the reaction takes place in one step or in a number of steps.

e.g., (i) $A \xrightarrow{H} B$

(ii) $A \xrightarrow{H_1} C \xrightarrow{H_2} B$

According to Hess's Law $H \quad H_1 \quad H_2$.

❖ **Enthalpy of atomisation :**

The change in enthalpy when one mole of substance converts into gaseous atoms.

❖ **Bond enthalpies :**

The average of enthalpies required to dissociate the bond present in different gaseous compounds into free atoms or radicals in the gaseous state.

The bond enthalpy given for any particular pair of atoms is the average value of the dissociation enthalpies of the bond for a number of molecules in which the pair of atoms appears.

For More Material Join: @JEEAdvanced_2024

❖ **Estimation of Enthalpy of a reaction from bond Enthalpies :**

For the gaseous reaction; $C_2H_4(g)$ $HCl(g)$ $C_2H_5Cl(g)$

H Energy required to break reactants into gaseous atoms — Energy released to from products from the gaseous atoms

❖ **Resonance Enthalpy :**

$H_{\text{Resonance}}$ $H_{f,\text{experimental}}$ $H_{f,\text{calculated}}$

$H_{c,\text{calculated}}$ $H_{c,\text{experimental}}$

For More Material Join: @JEEAdvanced_2024



Level 1

1. Out of molar entropy (I), specific volume (II), heat capacity (III), volume (IV), extensive properties are :
(a) I, II (b) I, II, IV (c) II, III (d) III, IV

2. Out of internal energy (I), boiling point (II), pH (III) and E.M.F. of the cell (IV) intensive properties are :
(a) I, II (b) II, III, IV (c) I, III, IV (d) All of these

3. Thermodynamic equilibrium involves
(a) chemical equilibrium (b) mechanical equilibrium
(c) thermal equilibrium (d) all the above simultaneously

4. Which has maximum internal energy at 290 K?
(a) Neon gas (b) Nitrogen gas (c) Ozone gas (d) Equal for all

5. A 10 g piece of iron (C 0.45 J/g°C) at 100°C is dropped into 25 g of water (C 4.2 J/g°C) at 27°C. Find temperature of the iron and water system at thermal equilibrium.
(a) 30°C (b) 33°C (c) 40°C (d) None of these

6. When freezing of a liquid takes place in a system it:
(a) may have q 0 or q 0 depending on the liquid
(b) is represented by q 0
(c) is represented by q 0
(d) has q 0

7. Mechanical work is specially important in systems that contain
(a) gas-liquid (b) liquid-liquid (c) solid-solid (d) amalgam

8. Determine which of the following reactions taking place at constant pressure represents system that do work on the surrounding environment
I. Ag (aq) Cl (aq) AgCl(s) II. $NH_4Cl(s)$ $NH_3(g)$ HCl(g)
III. $2NH_3(g)$ $N_2(g)$ $3H_2(g)$
(a) I (b) III (c) II and III (d) I and II

9. Determine which of the following reactions taking place at constant pressure represent surrounding that do work on the system environment
I. $4NH_3(g)$ $7O_2(g)$ $4NO_2(g)$ $6H_2O(g)$
II. $CO(g)$ $2H_2(g)$ $CH_3OH(l)$
III. C$(s,$ graphite$)$ $H_2O(g)$ $CO(g)$ $H_2(g)$
IV. $H_2O(s)$ $H_2O(l)$
(a) III, IV (b) II and III (c) II, IV (d) I and II, IV

10. A sample of liquid in a thermally insulated container (a calorimeter) is stirred for 2 hr. by a mechanical linkage to a motor in the surrounding. For this process :
(a) w 0; q 0; U 0 (b) w 0; q 0; U 0
(c) w 0; q 0; U 0 (d) w 0; q 0; U 0

11. A system undergoes a process in which U 300 J while absorbing 400 J of heat energy and undergoing an expansion against 0.5 bar. What is the change in the volume (in L)?
(a) 4 (b) 5 (c) 2 (d) 3

For More Material Join: @JEEAdvanced_2024

12. An ideal gas expands against a constant external pressure of 2.0 atmosphere from 20 litre to 40 litre and absorbs 10 kJ of heat from surrounding. What is the change in internal energy of the system? (Given : 1 atm-litre 101.3 J)

(a) 4052 J (b) 5948 J (c) 14052 J (d) 9940 J

13. One mole of an ideal gas at 25°C expands in volume from 1.0 L to 4.0 L at constant temperature. What work (in J) is done if the gas expands against vacuum ($P_{external}$ 0)?

(a) 4.0 10^2 (b) 3.0 10^2 (c) 1.0 10^2 (d) Zero

14. At 25°C, a 0.01 mole sample of a gas is compressed from 4.0 L to 1.0 L at constant temperature. What is the work done for this process if the external pressure is 4.0 bar?

(a) 1.6 10^3 J (b) 8.0 10^2 J (c) 4.0 10^2 J (d) 1.2 10^3 J

15. Calculate the work done (in J) when 4.5 g of H_2O_2 reacts against a pressure of 1.0 atm at 25°C

$$2H_2O_2(l) \quad O_2(g) \quad 2H_2O(l)$$

(a) 1.63 10^2 (b) 4.5 10^2 (c) 3.2 10^2 (d) 6.1 10^2

16. 2 mole of zinc is dissolved in HCl at 25°C. The work done in open vessel is :

(a) 2.477 kJ (b) 4.955 kJ (c) 0.0489 kJ (d) None


17. Temperature of 1 mole of a gas is increased by 2°C at constant pressure. Work done is :

(a) R (b) $2R$ (c) $R/2$ (d) $3R$

18. A sample of an ideal gas is expanded 1 m^3 to 3 m^3 in a reversible process for which P KV^2, with K 6 bar/m^6. Work done by the gas is :


(a) 5200 kJ (b) 15600 kJ (c) 52 kJ (d) 5267.6 kJ

19. A given mass of gas expands reversibly from the state A to the state B by three paths 1, 2 and 3 as shown in the figure. If w_1, w_2 and w_3 respectively be the work done by the gas along three paths then:



(a) w_1 w_2 w_3
(b) w_1 w_2 w_3
(c) w_1 w_2 w_3
(d) w_2 w_3 w_1


20. Heat absorbed by a system in going through a cyclic process shown in figure is :



(a) 10^7 J (b) 10^6 J
(c) 10^2 J (d) 10^4 J

For More Material Join: @JEEAdvanced_2024


21. In the cyclic process shown in *P–V* diagram, the magnitude of the work done is :



(a) $\frac{(P_2 - P_1)^2}{2}$ (b) $\frac{(V_2 - V_1)^2}{2}$

(c) $\frac{1}{4}(P_2 - P_1)(V_2 - V_1)$ (d) $(V_2 - V_1)^2$

22. An ideal gas is taken around the cycle *ABCA* as shown in *P–V* diagram. The net work done during the cycle is equal to :



(a) $12P_1V_1$ (b) $6P_1V_1$


(c) $5P_1V_1$ (d) P_1V_1

23. An ideal gas is at pressure *P* and temperature *T* in a box, which is kept in vacuum with in a large container. The wall of the box is punctured. What happens as the gas occupies entire container?

(a) It's temperature falls (b) Its temperature rises

(c) Its temperature remains the same (d) Unpredictable

24. In diagrams (1 to 4), variation of volume with changing pressure is shown. A gas is taken along the path *ABCD*. The change in internal energy of the gas will be :



(a) Positive in all the cases (1) to (4)

(b) Positive in cases (1), (2), (3) but zero in case (4)

(c) Negative in cases (1), (2), (3) but zero in case (4)

(d) Zero in all the cases

For More Material Join: @JEEAdvanced_2024


25. If the door of a refrigerator is kept open, the room in which the refrigerator is kept
(a) gets cooled
(b) gets heated
(c) neither gets cooled nor gets heated
(d) gets cooled or heated depending on the initial temperature of the room


26. The temperature of an ideal gas increases in an :
(a) adiabatic expansion (b) isothermal expansion
(c) adiabatic compression (d) isothermal compression


27. For two mole of an ideal gas :
(a) $C_v \quad C_p \quad R$ (b) $C_p \quad C_v \quad 2R$ (c) $C_p \quad C_v \quad R$ (d) $C_v \quad C_p \quad 2R$


28. Which of the following expressions is true for an ideal gas ?
(a) $\left(\frac{V}{T}\right)_P \quad 0$ (b) $\left(\frac{P}{T}\right)_V \quad 0$ (c) $\left(\frac{U}{V}\right)_T \quad 0$ (d) $\left(\frac{U}{T}\right)_V \quad 0$

29. Liquefied oxygen at 1 atmosphere is heated from 50 K to 300 K by supplying heat at a constant rate. The graph that correctly shows the relationship between temperature and time is :

(a) 


(b) 


(c) 


(d) 


30. If w_1, w_2, w_3 and w_4 for an ideal gas are magnitude of work done in isothermal, adiabatic, isobaric and isochoric reversible expansion processes, the correct order will be :
(a) $w_1 \quad w_2 \quad w_3 \quad w_4$ (b) $w_3 \quad w_2 \quad w_1 \quad w_4$
(c) $w_3 \quad w_2 \quad w_4 \quad w_1$ (d) $w_3 \quad w_1 \quad w_2 \quad w_4$

31. For a closed container containing 100 mole of an ideal gas fitted with movable, frictionless, weightless piston operating such that pressure of gas remains constant at 8.21 atm. Which graph represents correct variation of log V *vs.* log T where V is in litre and T in kelvin?

(a) 

(b) 

(c) 

(d) 


32. A gas expands against a variable pressure given by $P \quad \frac{20}{V}$ (where P in atm and V in L). During expansion from volume of 1 litre to 10 litre, the gas undergoes a change in internal energy of 400 J. How much heat is absorbed by the gas during expansion?
(a) 46 J (b) 4660 J (c) 5065.8 J (d) 4260 J

33. 2 mole of an ideal gas at 27°C expands isothermally and reversibly from a volume of 4 litre to 40 litre. The work done (in kJ) by the gas is :
(a) $w \quad 28.72$ kJ (b) $w \quad 11.488$ kJ
(c) $w \quad 5.736$ kJ (d) $w \quad 4.988$ kJ

For More Material Join: @JEEAdvanced_2024

34. 10 mole of ideal gas expand isothermally and reversibly from a pressure of 10 atm to 1atm at 300 K. What is the largest mass which can lifted through a height of 100 meter?
(a) 31842 kg (b) 58.55 kg (c) 342.58 kg (d) None of these

35. A heat engine carries one mole of an ideal mono-atomic gas around the cycle as shown in the figure. Select the correct option:



(a) q_{AB} 450 R and q_{CA} 450 R
(b) q_{AB} 450 R and q_{CA} 225 R
(c) q_{AB} 450 R and q_{CA} 375 R
(d) q_{AB} 375 R and q_{CA} 450 R

36. What is the final temperature of 0.10 mole monoatomic ideal gas that performs 75 cal of work adiabatically if the initial temperature is 227°C? (use R 2 cal/K-mol)
(a) 250 K (b) 300 K (c) 350 K (d) 750 K


37. The work done by the gas in reversible adiabatic expansion process is :
(a) $\frac{P_2V_2 \quad P_1V_1}{1}$ (b) $\frac{nR\,(T_1 \quad T_2)}{1}$ (c) $\frac{P_2V_2 \quad P_1V_1}{}$ (d) None of these

38. During an adiabatic process, the pressure of gas is found to be proportional to the cube of its absolute temperature. The ratio of $(C_{p,m}/C_{v,m})$ for gas is :
(a) $\frac{3}{2}$ (b) $\frac{5}{3}$ (c) $\frac{7}{2}$ (d) $\frac{4}{3}$

39. A gas expands adiabatically at constant pressure such that $T \quad V^{1/2}$
The value of $(C_{p,m}/C_{v,m})$ of the gas will be :
(a) 1.30 (b) 1.50 (c) 1.70 (d) 2

40. For a reversible adiabatic ideal gas expansion $\frac{dP}{P}$ is equal to :
(a) $\frac{dV}{V}$ (b) $\frac{dV}{V}$ (c) $\frac{}{1}\frac{dV}{V}$ (d) $\frac{dV}{V}$

41. P–V plot for two gases (assuming ideal)during adiabatic processes are given in the Fig. Plot A and plot B should correspond respectively to:



(a) He and H_2
(b) H_2 and He
(c) He and Ne
(d) H_2 and Cl_2

42. Calculate the final temperature of a monoatomic ideal gas that is compressed reversible and adiabatically from 16 L to 2 L at 300 K :
(a) 600 K (b) 1044.6 K (c) 1200 K (d) 2400 K

For More Material Join: @JEEAdvanced_2024

43. 5 mole of an ideal gas expand isothermally and irreversibly from a pressure of 10 atm to 1 atm against a constant external pressure of 1 atm. w_{irr} at 300 K is :
(a) 15.921 kJ (b) 11.224 kJ (c) 110.83 kJ (d) None of these

44. With what minimum pressure (in kPa), a given volume of an ideal gas ($C_{p,m}$ 7/2 R), originally at 400 K and 100 kPa pressure can be compressed irreversibly adiabatically in order to raise its temperature to 600 K :
(a) 362.5 kPa (b) 275 kPa (c) 437.5 kPa (d) 550 kPa

45. The work done in adiabatic compression of 2 mole of an ideal monoatomic gas against constant external pressure of 2 atm starting from initial pressure of 1 atm and initial temperature of 300 K (R 2 cal/mol-degree)
(a) 360 cal (b) 720 cal (c) 800 cal (d) 1000 cal


46. One mole of an ideal gas $C_{v,m}$ $\frac{5}{2}R$ at 300 K and 5 atm is expanded adiabatically to a final pressure of 2 atm against a constant pressure of 2 atm. Final temperature of the gas is :
(a) 270 K (b) 273 K (c) 248.5 K (d) 200 K


47. 10 litre of a non linear polyatomic ideal gas at 127°C and 2 atm pressure is suddenly released to 1 atm pressure and the gas expanded adiabatically against constant external pressure. The final temperature and volume of the gas respectively are.
(a) T 350 K ; V 17.5 L (b) T 300 K ; V 15 L
(c) T 250 K ; V 12.5 L (d) None of these

48. Calculate average molar heat capacity at constant volume of gaseous mixture contained 2 mole of each of two ideal gases A $C_{v,m}$ $\frac{3}{2}R$ and B $C_{v,m}$ $\frac{5}{2}R$:
(a) R (b) $2R$ (c) $3R$ (d) $8R$

49. 0.5 mole each of two ideal gases A $C_{v,m}$ $\frac{5}{2}R$ and B ($C_{v,m}$ $3R$) are taken in a container and expanded reversibly and adiabatically, during this process temperature of gaseous mixture decreased from 350 K to 250 K. Find H (in cal/mol) for the process :
(a) 100 R (b) 137.5 R
(c) 375 R (d) None of these

50. A cyclic process $ABCD$ is shown in P–V diagram for an ideal gas. Which of the following diagram represents the same process?





For More Material Join: @JEEAdvanced_2024

51. 36 mL of pure water takes 100 sec to evaporate from a vessel and heater connected to an electric source which delivers 806 watt. The $H_{vaporization}$ of H_2O is :
(a) 40.3 kJ/mol (b) 43.2 kJ/mol (c) 4.03 kJ/mol (d) None of these

52. For the reaction : $PCl_5(g)$ $PCl_3(g)$ $Cl_2(g)$:
(a) H U (b) H U (c) H U (d) None of the above

53. Consider the reaction at 300 K

$$H_2(g) \quad Cl_2(g) \quad 2HCl(g); \quad H \quad 185\,kJ$$

If 2 mole of H_2 completely react with 2 mole of Cl_2 to form HCl, what is U for this reaction?
(a) 0 (b) 185 kJ (c) 370 kJ (d) None of these

54. Which of the indicated relationship is correct for the following exothermic reaction carried out at constant pressure?

$$CO(g) \quad 3H_2(g) \quad CH_4(g) \quad H_2O(g)$$

(a) U H (b) U H (c) w 0 (d) q 0

55. One mole of an ideal gas undergoes a change of state (2.0 atm, 3.0 L) to (2.0 atm, 7.0 L) with a change in internal energy (U) 30 L-atm. The change in enthalpy (H) of the process in L-atm :
(a) 22 (b) 38 (c) 25 (d) None of these

56. What is the change in internal energy when a gas contracts from 377 mL to 177 mL under a constant pressure of 1520 torr, while at the same time being cooled by removing 124 J heat?
(a) 40.52 J (b) 83.48 J (c) 248 J (d) None of these

57. For the real gases reaction $2CO(g)$ $O_2(g)$ $2CO_2(g)$; H 560 kJ. In 10 litre rigid vessel at 500 K the initial pressure is 70 bar and after the reaction it becomes 40 bar.
The change in internal energy is :
(a) 557 kJ (b) 530 kJ (c) 563 kJ (d) None of these

58. One mole of a non-ideal gas undergoes a change of state from (1.0 atm, 3.0 L, 200 K) to (4.0 atm, 5.0 L, 250 K) with a change in internal energy (U) 40 L-atm. The change in enthalpy of the process in L-atm :
(a) 43 (b) 57 (c) 42 (d) None of these

59. Consider the reaction at 300 K

$$C_6H_6(l) \quad \frac{15}{2}O_2(g) \quad 6\,CO_2(g) \quad 3H_2O(l); \quad H \quad 3271\,kJ$$

What is U for the combustion of 1.5 mole of benzene at 27°C?
(a) 3267.25 kJ (b) 4900.88 kJ (c) 4906.5 kJ (d) 3274.75 kJ

60. For the reaction; $FeCO_3(s)$ $FeO(s)$ $CO_2(g)$; H 82.8 kJ at 25°C, what is (E or U) at 25°C?
(a) 82.8 kJ (b) 80.32 kJ (c) 2394.77 kJ (d) 85.28 kJ

61. At 5 10^5 bar pressure density of diamond and graphite are 3 g/cc and 2 g/cc respectively, at certain temperature 'T'. Find the value of U H for the conversion of 1 mole of graphite to 1 mole of diamond at temperature 'T' :
(a) 100 kJ/mol (b) 50 kJ/mol (c) 100 kJ/mol (d) None of these

62. Predict which of the following reaction(s) has a positive entropy change?
I. $Ag\ (aq)$ $Cl\ (aq)$ $AgCl(s)$
II. $NH_4Cl(s)$ $NH_3(g)$ $HCl(g)$

For More Material Join: @JEEAdvanced_2024

III. $2NH_3(g)$ $N_2(g)$ $3H_2(g)$

(a) I and II (b) III (c) II and III (d) II

63. Predict which of the following reaction(s) has a negative entropy change?

I. $CH_4(g)$ $2O_2(g)$ $CO_2(g)$ $2H_2O(l)$

II. $NH_3(g)$ $HCl(g)$ $NH_4Cl(s)$

III. $2KClO_4(s)$ $2KClO_3(s)$ $O_2(g)$

(a) III (b) II (c) I and II (d) I

64. Which of the following reactions is associated with the most negative change in entropy?

(a) $2SO_2(g)$ $O_2(g)$ $2SO_3(g)$

(b) $C_2H_4(g)$ $H_2(g)$ $C_2H_6(g)$

(c) $C(s, \text{graphite})$ $O_2(g)$ $CO_2(g)$

(d) $3C_2H_2(g)$ $C_6H_6(l)$

65. When two mole of an ideal gas $C_{p,m}$ $\frac{5}{2}R$ heated from 300 K to 600 K at constant pressure, the change in entropy of gas (S) is :

(a) $\frac{3}{2}R\ln 2$ (b) $\frac{3}{2}R\ln 2$ (c) $5R\ln 2$ (d) $\frac{5}{2}R\ln 2$

66. Which of the following expression for entropy change of an irreversible process ?

(a) dS $\frac{dq}{T}$ (b) dS $\frac{dq}{T}$ (c) dS $\frac{dq}{T}$ (d) dS $\frac{dU}{T}$

67. Which of the following expressions is known as Clausius inequality ?

(a) $\circ\frac{dq}{T}$ 0 (b) $\circ\frac{ds}{T}$ 0 (c) $\circ\frac{T}{dq}$ 0 (d) $\circ\frac{dq}{T}$ 0

68. In problem 65, calculate S_{gas} if process is carried out at constant volume :

(a) $5R\ln 2$ (b) $\frac{3}{2}R\ln 2$ (c) $3R\ln 2$ (d) $3R\ln 2$

69. If one mole of an ideal gas $C_{p,m}$ $\frac{5}{2}R$ is expanded isothermally at 300 K until it's volume is tripled, then change in entropy of gas is :

(a) zero (b) infinity (c) $\frac{5}{2}R\ln 3$ (d) $R\ln 3$

70. In previous problem if expansion is carried out freely (P_{ext} 0), then S is :

(a) zero (b) infinity (c) $R\ln 3$ (d) None

71. When one mole of an ideal gas is compressed to half of its initial volume and simultaneously heated to twice its initial temperature, the change in entropy of gas (S) is :

(a) $C_{p,m}\ln 2$ (b) $C_{v,m}\ln 2$ (c) $R\ln 2$ (d) $(C_{v,m}$ $R)\ln 2$

72. What is the change in entropy when 2.5 mole of water is heated from 27°C to 87°C? Assume that the heat capacity is constant. ($C_{p,m}$ (H_2O) 4.2 J/g-K and $\ln(1.2) = 0.18$)

(a) 16.6 J/K (b) 9 J/K (c) 34.02 J/K (d) 1.89 J/K

73. Calculate standard entropy change in the reaction

$Fe_2O_3(s)$ $3H_2(g)$ $2Fe(s)$ $3H_2O(l)$

Given : S_m (Fe_2O_3, S) 87.4 , S_m (Fe, S) 27.3,

S_m (H_2, g) 130.7, S_m (H_2O, l) 69.9 JK ^{1} mol ^{1}

For More Material Join: @JEEAdvanced_2024

(a) 212.5 JK ^{1} mol ^{1} (b) 215.2 JK ^{1} mol ^{1}
(c) 120.9 JK ^{1} mol ^{1} (d) None of these

74. Calculate the entropy change (J/mol K) of the given reaction. The molar entropies [J/K-mol] are given in brackets after each substance.

$2PbS(s)$ [91.2] $3O_2(g)$ [205.1] $2PbO(s)$ [66.5] $2SO_2(g)$ [248.2]

(a) 113.5 (b) 168.3 (c) +72.5 (d) 149.2

75. Given $_rS$ 266 and the listed [S_m values] calculate S for $Fe_3O_4(s)$:

$4Fe_3O_4(s)$ [ ] $O_2(g)$ [205] $6Fe_2O_3(s)$ [87]

(a) +111.1 (b) +122.4 (c) 145.75 (d) 248.25

76. The entropy change for a phase transformation is :

(a) $\frac{U}{dT}$ (b) $\frac{T}{H}$ (c) $\frac{H}{T}$ (d) $\frac{H \quad G}{T}$

77. What is the melting point of benzene if H_{fusion} 9.95 kJ/mol and S_{fusion} 35.7 J/K-mol?

(a) 278.7°C (b) 278.7 K (c) 300 K (d) 298 K

78. S for freezing of 10 g of $H_2O(l)$ (enthalpy of fusion is 80 cal/g) at 0°C and 1 atm is :

(a) 12.25 J/K (b) 0.244 J/K (c) 2.93 J/K (d) 12.25 J/K

79. Chloroform has $H_{\text{vaporization}}$ 29.2 kJ/mol and boils at 61.2°C. What is the value of $S_{\text{vaporization}}$ for chloroform?

(a) 87.3 J/mol-K (b) 477.1 J/mol-K (c) 87.3 J/mol-K (d) 477.1 J/mol-K

80. The entropy of vaporization of benzene is 85 JK ^{1} mol ^{1}. When 117g benzene vaporizes at it's normal boiling point, the entropy change of surrounding is :

(a) 85 JK ^{1} (b) 85 1.5 JK ^{1} (c) 85 1.5 JK ^{1} (d) None of these

81. Identify the correct statement regarding entropy

(a) At absolute zero temperature, the entropy of perfectly crystalline substances is ve
(b) At absolute zero temperature entropy of perfectly crystalline substance is taken to be zero
(c) At 0 C the entropy of a perfectly crystalline substance is taken to be zero
(d) At absolute zero temperature, the entropy of all crystalline substances is taken to be zero

82. Calculate S for following process :

$$\underset{\text{at 100 K}}{X(s)} \qquad \underset{\text{at 200 K}}{X(l)}$$

Given : Melting point of $X_{(s)}$ 100 K ; H_{Fusion} 20 kJ/mol ; $C_{p,m}(X,l)$ 10 J/mol.K

(a) 26.93 J/K (b) 206.93 J/K (c) 203 J/K (d) 206.93 kJ/K

83. For a perfectly crystalline solid $C_{p,m}$ aT^3, where a is constant. If $C_{p,m}$ is 0.42 J/K mol at 10 K, molar entropy at 20 K is :

(a) 0.42 J/K mol (b) 0.14 J/K mol (c) 1.12 J/K mol (d) zero

84. Consider the following spontaneous reaction $3X_2(g)$ $2X_3(g)$. What are the sign of H, S and G for the reaction?

(a) +ve, +ve, +ve (b) ve, ve, ve (c) ve, ve, ve (d) ve, ve, ve

85. For the reaction $2H(g)$ $H_2(g)$, the sign of H and S respectively are :

(a) , (b) +, + (c) , (d) ,

For More Material Join: @JEEAdvanced_2024

86. Consider the following reaction.

$$C_6H_6(l) \quad \frac{15}{2}O_2(g) \qquad 6CO_2(g) \quad 3H_2O(g)$$

signs of H, S and G for the above reaction will be

(a) , , (b) , , (c) , , (d) , ,

87. Consider the following reaction at temperature T :

$$CH_2 \quad CH_2(g) \quad Cl_2(g) \qquad ClCH_2CH_2Cl(g)$$

$_rH$ 217.5 kJ/mol, $_rS$ 233.9 J/K-mol

Reaction is supported by :

(a) entropy (b) enthalpy (c) both (a) & (b) (d) neither

88. For a process to be spontaneous at constant T and P :

(a) $(G)_{system}$ must be negative (b) $(G)_{system}$ must be positive

(c) $(S)_{system}$ must be positive (d) $(S)_{system}$ must be negative

89. For a reaction to occur spontaneously :

(a) S must be negative (b) $(H \quad T \quad S)$ must be positive

(c) $H \quad T \quad S$ must be negative (d) H must be negative

90. Which of the following conditions regarding a chemical process ensures its spontaneity at all temperature?

(a) $H \quad 0, \; S \quad 0$ (b) $H \quad 0, \; S \quad 0$ (c) $H \quad 0, \; S \quad 0$ (d) $H \quad 0, \; S \quad 0$

91. The free energy change $G \quad 0$, when

(a) the system is at equilibrium

(b) catalyst is added

(c) reactants are initially mixed thoroughly

(d) the reactants are completely consumed

92. Which of the following conditions will always lead to a non-spontaneous change?

(a) H and S both +ve (b) H is ve and S is +ve

(c) H and S both ve (d) H is +ve and S is ve

93. Suppose that a reaction has $H \quad 40$ kJ and $S \quad 50$ J/K. At what temperature range will it change from spontaneous to non-spontaneous?

(a) 0.8 K to 1 K (b) 799 K to 800 K (c) 800 K to 801 K (d) 799 K to 801 K

94. For isothermal expansion in case of an ideal gas :

(a) $G \quad S$ (b) $G \quad H$ (c) $G \quad T. \; S$ (d) None of these

95. What is the normal boiling point of mercury?

Given : H_f (Hg, l) 0; S (Hg, l) 77.4 J/K-mol

H_f (Hg, g) 60.8 kJ/mol; S (Hg, g) 174.4 J/K-mol

(a) 624.8 K (b) 626.8 K (c) 636.8 K (d) None of these

96. 18 gm of ice is converted into water at 0°C and 1 atm. The entropies of $H_2O(s)$ and $H_2O(l)$ are 38.2 and 60 J/mol K respectively. The enthalpy change for this conversion is :

(a) 5951.4 J/mol (b) 595.14 J/mol (c) 5951.4 J/mol (d) None of these

For More Material Join: @JEEAdvanced_2024

97. Using the listed [G_f values] calculate G for the reaction :

$3H_2S(g)$ [33.6] $2HNO_3(l)$ [80.6] $2NO(g)$ [86.6] $4H_2O(l)$ [237.1] $3S(s)$ [0.0]

(a) 513.2 (b) 1037.0 (c) +433.4 (d) +225.0

98. From the following H and S values, predict which of reactions I, II, and III would be spontaneous at 25°C.

	H (kJ)	S (J/K)
I.	+10.5	+30
II.	+1.8	113
III.	126	+84

(a) III (b) I (c) II and III (d) I and II

99. Calculate H_f for UBr_4 from the G of reaction and the $S°$ values at 298 K.

$U(s)$ $2Br_2(l)$ $UBr_4(s)$; G 788.6 kJ; S (J/K-mol) 50.3, 152.3, 242.6

(a) 822.1 kJ/mol (b) 841.2 kJ/mol (c) 775.6 kJ/mol (d) 804.3 kJ/mol

100. The entropies of $H_2(g)$ and $H(g)$ are 130.6 and 114.6 J mol ^{1} K ^{1} respectively at 298 K. Using the data given below calculate the bond energy of H_2 (in kJ/mol) :

$H_2(g)$ $2H(g)$; G 406.6 kJ

(a) 377.2 (b) 436.0 (c) 425.5 (d) 430.5

101. Consider the G_f and H_f (kJ/mol) for the following oxides. Which oxide can be most easily decomposed to form the metal and oxygen gas?

(a) ZnO (G 318.4, H 348.3) (b) Cu_2O (G 146.0, H 168.8)

(c) HgO (G 58.5, H 90.8) (d) PbO (G 187.9, H 217.3)

102. Which of the following option is correct ?

(a) $\frac{\ln K_p}{T}$ $\frac{H}{RT^2}$ (b) $\frac{\ln k}{T}$ $\frac{E_a}{RT^2}$

(c) $\frac{\ln K_p}{T}$ $\frac{U}{RT^2}$ (d) All of these

103. Calculate G (kJ/mol) at 127°C for a reaction with $K_{\text{equilibrium}}$ 10^5 :

(a) 38.294 (b) 16.628 (c) 9.16 (d) None of these

104. When reaction is carried out at standard states then at the equilibrium :

(a) H 0 (b) S 0

(c) equilibrium constant K 0 (d) equilibrium constant K 1

105. At 25°C, G for the process $H_2O(l) \rightleftharpoons H_2O(g)$ is 8.6 kJ. The vapour pressure of water at this temperature, is nearly :

(a) 24 torr (b) 285 torr (c) 32.17 torr (d) 100 torr

106. For the auto-ionization of water at 25°C, $H_2O(l) \rightleftharpoons H\ (aq)$ $OH\ (aq)$ equilibrium constant is 10^{14}.

What is G for the process?

(a) ~8 10^4 J (b) ~3.5 10^4 J (c) ~10^4 J (d) None of these

For More Material Join: @JEEAdvanced_2024

107. The molar entropies of HI(g), H(g) and I(g) at 298 K are 206.5, 114.6, and 180.7 J mol ^{1} K ^{1} respectively. Using the G given below, calculate the bond energy of HI.

HI(g) H(g) I(g); G 271.8 kJ

(a) 282.4 kJ mol ^{1} (b) 298.3 kJ mol ^{1} (c) 290.1 kJ mol ^{1} (d) 315.4 kJ mol ^{1}

108. Hess's law states that :

(a) the standard enthalpy of an overall reaction is the sum of the enthalpy changes in individual reactions.

(b) enthalpy of formation of a compound is same as the enthalpy of decomposition of the compound into constituent elements, but with opposite sign.

(c) at constant temperature the pressure of a gas is inversely proportional to its volume

(d) the mass of a gas dissolved per litre of a solvent is proportional to the pressure of the gas in equilibrium with the solution

109. An imaginary reaction X Y takes place in three steps

X A, H q_1; B A, H q_2; B Y, H q_3

If Hess' law is applicable, then the heat of the reaction (X Y) is :

(a) q_1 q_2 q_3 (b) q_2 q_3 q_1 (c) q_1 q_2 q_3 (d) q_3 q_2 q_1

110. The enthalpy change for a reaction does not depend upon :

(a) the physical states of reactants and products

(b) use of different reactants for the same product

(c) the number of intermediate reaction steps

(d) the differences in initial or final temperatures of involved substances

111. The standard enthalpy of formation of gaseous H_2O at 298 K is 241.82 kJ/mol. Calculate H at 373 K given the following values of the molar heat capacities at constant pressure :

$H_2O(g)$ 33.58 JK ^{1} mol ^{1}; $H_2(g)$ 29.84 JK ^{1} mol ^{1}; $O_2(g)$ 29.37 JK ^{1} mol ^{1}

Assume that the heat capacities are independent of temperature :

(a) 242.6 kJ/mol (b) 485.2 kJ/mol (c) 121.3 kJ/mol (d) 286.4 kJ/mol

112. Which of the following value of H_f represent that the product is least stable ?

(a) 94.0 kcal mol ^{1} (b) 231.6 kcal mol ^{1} (c) 21.4 kcal mol ^{1} (d) 64.8 kcal mol ^{1}

113. For which of the following substances the enthalpy of formation in the standard state is zero ?

(a) Sucrose (b) Ethanol (c) Aluminium (d) Calcium chloride

114. Calculate the standard enthalpy of reaction for the following reaction using the listed enthalpies of reaction :

$3\,Co(s)$ $2O_2(g)$ $Co_3O_4(s)$

$2\,Co(s)$ $O_2(g)$ $2\,CoO(s)$; H_1 475.8 kJ

$6CoO(s)$ $O_2(g)$ $2\,Co_3O_4(s)$; H_2 355.0 kJ

(a) 891.2 kJ (b) 120.8 kJ (c) +891.2 kJ (d) 830.8 kJ

115. From given following equations and H values, determine the enthalpy of reaction at 298 K for the reaction :

$C_2H_4(g)$ $6F_2(g)$ $2\,CF_4(g)$ $4HF(g)$

$H_2(g)$ $F_2(g)$ $2HF(g)$; H_1 537 kJ

For More Material Join: @JEEAdvanced_2024

$C(s)\quad 2F_2(g)\qquad CF_4(g);\quad H_2\quad 680\ kJ$

$2C(s)\quad 2H_2(g)\qquad C_2H_4(g);\quad H_3\quad 52\ kJ$

(a) 1165 (b) 2486 (c) +1165 (d) +2486

116. Given :

$C(diamond)\quad O_2\qquad CO_2;\quad H\quad 395\ kJ$

$C(graphite)\quad O_2\qquad CO_2;\quad H\quad 393\ kJ$

The enthalpy of formation of diamond from graphite is

(a) 2.0 kJ (b) 1.5 kJ (c) 788 kJ (d) 788 kJ

117. Which of the following equations represents a reaction that provides the enthalpy of formation of CH_3Cl?

(a) $C(s)\quad HCl(g)\quad H_2(g)\qquad CH_3Cl(g)$

(b) $C(s)\quad 3H(g)\quad Cl(g)\qquad CH_3Cl(g)$

(c) $C(s)\quad 3/2H_2(g)\quad 1/2Cl_2(g)\qquad CH_3Cl(g)$

(d) $CH_4(g)\quad Cl_2(g)\qquad CH_3Cl(g)\quad HCl(g)$

118. Use the given standard enthalpies of formation (in kJ/mol) to determine the enthalpy of reaction of the following reaction :

$NH_3(g)\quad 3F_2(g)\qquad NF_3(g)\quad 3HF(g)$

$H_f\ (NH_3, g)\quad 46.2;\qquad H_f\ (NF_3, g)\quad 113.0;\qquad H_f\ (HF, g)\quad 269.0$

(a) 335.8 kJ/mol (b) 873.8 kJ/mol (c) 697.2 kJ/mol (d) 890.4 kJ/mol

119. The standard enthalpy of formation of octane (C_8H_{18}) is 250 kJ/mol. Calculate the enthalpy of combustion of C_8H_{18}. The enthalpy of formation of $CO_2(g)$ and $H_2O(l)$ are 394 kJ/mol and 286 kJ/mol respectively :

(a) – 5200 kJ/mol (b) – 5726 kJ/mol (c) – 5476 kJ/mol (d) – 5310 kJ/mol

120. Determine the enthalpy of formation of $B_2H_6(g)$ in kJ/mol of the following reaction :

$B_2H_6(g)\quad 3O_2(g)\qquad B_2O_3(s)\quad 3H_2O(g);$

Given : $_rH\quad 1941$ kJ/mol; $H_f\ (B_2O_3, s)\quad 1273$ kJ/mol;

$H_f\ (H_2O, g)\quad 241.8$ kJ/mol

(a) 75.6 (b) +75.6 (c) 57.4 (d) 28.4

121. Consider the following reactions :

$C(s)\quad O_2(g)\qquad CO_2(g)\quad x\ kJ$

$CO(g)\quad \frac{1}{2}O_2(g)\qquad CO_2(g)\quad y\ kJ$

The heat of formation of $CO(g)$ is :

(a) $(x\quad y)$ kJ/mol (b) $(x\quad y)$ kJ/mol (c) $(y\quad x)$ kJ/mol (d) None of these

122. If $_fH\ (C_2H_4)$ and $_fH\ (C_2H_6)$ are x_1 and x_2 kcal mol ^{1}, then heat of hydrogenation of C_2H_4 is :

(a) $x_1\quad x_2$ (b) $x_1\quad x_2$ (c) $x_2\quad x_1$ (d) $x_1\quad 2x_2$

123. What amount of energy (kJ) is released in the combustion of 5.8 g of $C_4H_{10}(g)$?

$2C_4H_{10}(g)\quad 13\,O_2(g)\qquad 8CO_2(g)\quad 10H_2O(l);\quad H\quad 5756\ kJ$

(a) 575.6 (b) 287.8 (c) 182 (d) 57.56

For More Material Join: @JEEAdvanced_2024

124. The enthalpy of the reaction forming PbO according to the following equation is 438 kJ. What heat energy (kJ) is released in formation of 22.3 g PbO(*s*)?
(Atomic masses : Pb 207, O 16.0)

$$2Pb(s) \quad O_2(g) \qquad 2PbO(s)$$

(a) 21.9 (b) 28.7 (c) 14.6 (d) 34.2

125. The fat, $C_{57}H_{104}O_6(s)$, is metabolized via the following reaction. Given the enthalpies of formation, calculate the $_rH$ (kJ) when 1.0 g of this fat reacts.

$$C_{57}H_{104}O_6(s) \quad 80\,O_2(g) \qquad 57CO_2(g) \quad 52H_2O(l)$$

$_fH$ ($C_{57}H_{104}O_6$, *s*) 70870 kJ/mol; $_fH$ (H_2O, *l*) 285.8 kJ/mol;

$_fH$ (CO_2, *g*) 393.5 kJ/mol

(a) H 37.98 (b) H 40.4 (c) H 33.4 (d) H 30.2

126. The $_fH$ of $NH_3(g)$ is 46 kJ mol^{-1}. The H (in kJ mol^{-1}) of the reaction, $2NH_3(g) \qquad N_2(g) \quad 3H_2(g)$ is :

(a) 46 (b) 46 (c) 92 (d) 92

127. Consider the following reaction:

$$H_2O(l) \qquad H_2O(g); \quad H_1 \quad 44 \text{ kJ}$$

$$2CH_3OH(l) \quad 3O_2(g) \qquad 4H_2O(l) \quad 2CO_2(g); \quad H_2 \quad 1453 \text{ kJ}$$

What is the value of H for second reaction if water vapour instead of liquid water is formed as product?

(a) 1409 kJ (b) 1629 kJ (c) 1277 kJ (d) None of these

128. The standard enthalpy change for the following reaction is 436.4 kJ :

$$H_2(g) \qquad H(g) \quad H(g)$$

What is the $_fH$ of atomic hydrogen (H)?

(a) 872.8 kJ/mol (b) 218.2 kJ/mol (c) 218.2 kJ/mol (d) 436.9 kJ/mol

129. Determine enthalpy of formation for $H_2O_2(l)$, using listed enthalpies of reaction :

$$N_2H_4(l) \quad 2H_2O_2(l) \qquad N_2(g) \quad 4H_2O(l); \quad _rH_1 \quad 818 \text{ kJ/mol}$$

$$N_2H_4(l) \quad O_2(g) \qquad N_2(g) \quad 2H_2O(l); \quad _rH_2 \quad 622 \text{ kJ/mol}$$

$$H_2(g) \quad 1/2\,O_2(g) \qquad H_2O(l); \quad _rH_3 \quad 285 \text{ kJ/mol}$$

(a) 383 kJ/mol (b) 187 kJ/mol (c) 498 kJ/mol (d) None of these

130. Heat of combustion of ethanol at constant pressure and at temperature T K is found to be q J mol^{-1}. Hence, heat of combustion (in J mol^{-1}) of ethanol at the same temperature at constant volume will be :

(a) $RT \quad q$ (b) $(q \quad RT)$ (c) $q \quad RT$ (d) $q \quad RT$

131. Stearic acid [$CH_3(CH_2)_{16}CO_2H$] is a fatty acid, the part of fat that stores most of the energy. 1.0 g of stearic acid was burned in a bomb calorimeter. The bomb had a heat capacity of 652 J/°C. If the temperature of 500 g water (c 4.18 J/g °C) increase from 25.0 to 39.3°C, how much heat was released when the stearic acid was burned?
[Given C_p (H_2O) 4.18 J/g°C]

(a) 39.21 kJ (b) 29.91 kJ (c) 108 kJ (d) 9.32 kJ

For More Material Join: @JEEAdvanced_2024

132. Gasoline has an enthalpy of combustion 24000 kJ/ gallon. When gasoline burns in an automobile engine, approximately 30% of the energy released is used to produce mechanical work. The remainder is lost as heat transfer to the engine's cooling system. As a start on estimating how much heat transfer is required, calculate what mass of water could be heated from 25°C to 75°C by the combustion of 1.0 gallon of gasoline in an automobile?

(Given : C (H_2O) 4.18 J/g°C)

(a) 34.45 kg (b) 80.383 kg (c) 22 kg (d) 224 kg

133. A 0.05 L sample of 0.2 *M* aqueous hydrochloric acid is added to 0.05 L of 0.2 *M* aqueous ammonia in a calorimeter. Heat capacity of entire calorimeter system is 480 J/K. The temperature increase is 1.09 K. Calculate $_rH$ in kJ/mol for the following reaction :

$$HCl(aq.) \quad NH_3(aq.) \quad NH_4Cl(aq.)$$

(a) 52.32 (b) 61.1 (c) 55.8 (d) 58.2

134. A coffee cup calorimeter initially contains 125 g of water, at a temperature of 24.2°C. 8 g of ammonium nitrate (NH_4NO_3), also at 24.2°C, is added to the water, and the final temperature is 18.2°C. What is the heat of solution of ammonium nitrate in kJ/mol? The specific heat capacity of the solution is 4.2 J/°C g.

(a) 33.51 kJ/mol (b) 39.5 kJ/mol (c) 32.2 kJ/mol (d) 37.3 kJ/mol

135. Read following statement(s) carefully and select the right option :

(I) The enthalpy of solution of $CaCl_2$ $6H_2O$ in a large volume of water is endothermic to the extent of 3.5 kcal/ mol. If H 23.2 kcal for the reaction,

$$CaCl_2(s) + 6H_2O(l) \quad CaCl_2.6H_2O(s)$$

then heat of solution of $CaCl_2$ (anhydrous) in a large volume of water is – 19.7 kcal/mol

(II) For the reaction $2Cl(g)$ $Cl_2(g)$; the sign of H and S are negative.

(a) statement I and II both are wrong (b) both are correct

(c) only I is correct (d) only II is correct

136. If the enthalpy of formation and enthalpy of solution of $HCl(g)$ are 92.3 kJ/mol and 75.14 kJ/mol respectively then find enthalpy of formation of Cl (aq) :

(a) 17.16 kJ/mol (b) 167.44 kJ/mol (c) 17.16 kJ/mol (d) None of these

137. At 25°C, when 1 mole of $MgSO_4$ was dissolved in water, the heat evolved was found to be 91.2 kJ. One mole of $MgSO_4$ $7H_2O$ on dissolution gives a solution of the same composition accompanied by an absorption of 13.8 kJ. The enthalpy of hydration, *i.e.,* H_h for the reaction

$$MgSO_4(s) \quad 7H_2O(l) \quad MgSO_4 \quad 7H_2O(s) \text{ is :}$$

(a) 105 kJ/mol (b) 77.4 kJ/mol (c) 105 kJ/mol (d) None of these

138. The enthalpies of neutralization of a weak base *A*OH and a strong base *B*OH by HCl are 12250 cal/mol and 13000 cal/mol respectively. When one mole of HCl is added to a solution containing 1 mole of *A*OH and 1 mole of *B*OH, the enthalpy change was 12500 cal/ mol. In what ratio is the acid distribution between *A*OH and *B*OH?

(a) 2 : 1 (b) 2 : 3 (c) 1 : 2 (d) None of these

139. The magnitude of enthalpy change for neutralization of the reaction ;

$MgO(s)$ $2HCl(aq)$ $MgCl_2(aq)$ $H_2O(l)$ (Given H (aq) OH (aq) $H_2O(l)$; 57 kJ/mol) will be :

For More Material Join: @JEEAdvanced_2024

(a) less that 57 kJ mol ^{1} (b) 57 kJ mol ^{1}

(c) greater than 57 kJ mol ^{1} (d) 57 kJ mol ^{1}

140. Enthalpy of neutralization of HCl by NaOH is 55.84 kJ/mol and by NH_4OH is 51.34 kJ/mol. The enthalpy of ionization of NH_4OH is :

(a) 107.18 kJ/mol (b) 4.5 kJ/mol (c) 4.5 kJ/mol (d) None of these

141. Which of the following reaction is endothermic ?

(a) $CaCO_3$ CaO CO_2 (b) Fe S FeS

(c) NaOH HCl NaCl H_2O (d) CH_4 $2O_2$ CO_2 $2H_2O$

142. Which of the following is not correct ?

(a) Dissolution of a salt in excess of water may be endothermic process

(b) Neutralisation is always exothermic

(c) The absolute value of enthalpy (H) can be determined by calorimeter

(d) The heat of reaction at constant volume is denoted by U

143. Substance $A_2B(g)$ can undergoes decomposition to form two set of products :

$A_2B(g) \rightarrow A_2(g)$ $B(g)$; H 40 kJ/mol

$A_2B(g) \rightarrow A(g)$ $AB(g)$; H 50 kJ/mol

If the molar ratio of $A_2(g)$ to $A(g)$ is 5 : 3 in a set of product gases, then the energy involved in the decomposition of 1 mole of $A_2B(g)$ is :

(a) 48.75 kJ/mol (b) 43.73 kJ/mol (c) 46.25 kJ/mol (d) None of these

144. Boron can undergo the following reactions with the given enthalpy changes :

$$2B(s) \quad \frac{3}{2}O_2(g) \qquad B_2O_3(s); \quad H \quad 1260 \text{ kJ}$$

$$2B(s) \quad 3H_2(g) \qquad B_2H_6(g); \quad H \quad 30 \text{ kJ}$$

Assume no other reactions are occurring.

If in a container (operating at constant pressure) which is isolated from the surrounding, mixture of H_2 (gas) and O_2 (gas) are passed over excess of $B(s)$, then calculate the molar ratio $(O_2 : H_2)$ so that temperature of the container do not change :

(a) 15 : 3 (b) 42 : 1 (c) 1 : 42 (d) 1 : 84

145. The enthalpy change for the following reaction is 368 kJ. Calculate the average O—F bond energy.

$$OF_2(g) \qquad O(g) \quad 2F(g)$$

(a) 184 kJ/mol (b) 368 kJ/mol (c) 536 kJ/mol (d) 736 kJ/mol

146. The enthalpy change for the reaction, $C_2H_6(g)$ $2C(g)$ $6H(g)$ is X kJ. The bond energy of C—H bond is :

(a) $\frac{X}{2}$ (b) $\frac{X}{3}$ (c) $\frac{X}{6}$ (d) data insufficient

For More Material Join: @JEEAdvanced_2024

147. The table given below lists the bond dissociation energy (E_{diss}) for single covalent bonds formed between C and atoms A, B, D, E.

Bond	E_{diss} (kcal mol^{-1})
C — A	240
C — B	382
C — D	276
C — E	486

Which of the atoms has smallest size ?

(a) D (b) E (c) A (d) D

148. Calculate P—Cl bond enthalpy

Given : $_fH$ (PCl_3, g) 306 kJ/mol; $H_{atomization}$ (P, s) 314 kJ/mol;
$_fH$ (Cl, g) 121 kJ/mol

(a) 123.66 kJ/mol (b) 371 kJ/mol (c) 19 kJ/mol (d) None of these

149. Calculate the enthalpy for the following reaction using the given bond energies (kJ/mol) :
(C—H 414; H—O 463; H—Cl 431, C—Cl 326; C—O 335)

CH_3 $OH(g)$ $HCl(g)$ CH_3 $Cl(g)$ $H_2O(g)$

(a) 23 kJ/mol (b) 42 kJ/mol (c) 59 kJ/mol (d) 511 kJ/mol

150. Based on the values of B.E. given, $_fH$ of $N_2H_4(g)$ is :

Given : N—N = 159 kJ mol^{-1}; H—H = 436 kJ mol^{-1}
N N = 941 kJ mol^{-1}; N—H = 398 kJ mol^{-1}

(a) 711 kJ mol^{-1} (b) 62 kJ mol^{-1} (c) 98 kJ mol^{-1} (d) 711 kJ mol^{-1}

For More Material Join: @JEEAdvanced_2024

Level 2

1. A heating coil is immersed in a 100 g sample of $H_2O(l)$ at 1 bar and 100°C in a closed vessel. In this heating process, 60% of the liquid is converted in to gaseous form at constant pressure of 1 bar. Densities of liquid and gaseous water under these conditions are 1000 kg/m^3 and 0.60 kg/m^3 respectively. Magnitude of the work done for the process is :
(a) 4997 J (b) 4970 J (c) 9994 J (d) None of these

2. A rigid and insulated tank of 3 m^3 volume is divided into two compartments. One compartment of volume of 2 m^3 contains an ideal gas at 0.8314 MPa and 400 K while the second compartment of volume 1 m^3 contains the same gas at 8.314 MPa and 500 K. If the partition between the two compartments is ruptured, the final temperature of the gas is :
(a) 420 K (b) 450 K (c) 480 K (d) None of these

3. What is the value of change in internal energy at 1 atm in the process?

$$H_2O(l, 323\ K) \qquad H_2O(g, 423\ K)$$

Given : $C_{v,m}(H_2O, l)$ 75.0 JK ^{1} mol ^{1}; $C_{p,m}(H_2O, g)$ 33.314 JK ^{1} mol ^{1}

H_{vap} at 373 K 40.7 kJ/mol


(a) 42.91 kJ/mol (b) 43086 kJ/mol (c) 42.6 kJ/mol (d) 49.6 kJ/mol

4. For an ideal gas $\frac{C_{p,m}}{C_{v,m}}$. The molecular mass of the gas is M, its specific heat capacity at constant volume is :
(a) $\frac{R}{M(\quad 1)}$ (b) $\frac{M}{R(\quad 1)}$ (c) $\frac{RM}{1}$ (d) $\frac{R}{M(\quad 1)}$

5. 1 mole of an ideal gas A ($C_{v,m}$ $3R$) and 2 mole of an ideal gas B are $C_{v,m}$ $\frac{3}{2}R$ taken in a container and expanded reversible and adiabatically from 1 litre to 4 litre starting from initial temperature of 320 K. E or U for the process is :
(a) 240 R (b) 240 R (c) 480 R (d) 960 R


6. Calculate the work done by the system in an irreversible (single step) adiabatic expansion of 2 mole of a polyatomic gas (4/3) from 300 K and pressure 10 atm to 1 atm :
(a) 227 R (b) 205 R (c) 405 R (d) None of these

7. One mole of an ideal gas is carried through the reversible cyclic process as shown in figure. The max. temperature attained by the gas during the cycle :
(a) $\frac{7}{6R}$ (b) $\frac{12}{49R}$
(c) $\frac{49}{12R}$ (d) None of these



For More Material Join: @JEEAdvanced_2024

8. Two moles of a triatomic linear gas (neglect vibration degree of freedom) are taken through a reversible process ideal starting from A as shown in figure.



The volume ratio $\frac{V_B}{V_A}$ 4. If the temperature at A is 73 C, then:

(i) Work done by the gas in AB process is :

(a) 6.16 kJ (b) 308.3 kJ

(c) 9.97 kJ (d) 0 J

(ii) Total enthalpy change in both steps is :

(a) 3000 R (b) 4200 R (c) 2100 R (d) 0

9. A gas $C_{v,m}$ $\frac{5}{2}R$ behaving ideally was allowed to expand reversibly and adiabatically from 1 litre to 32 litre. Its initial temperature was 327°C. The molar enthalpy change (in J/mol) for the process is :

(a) 1125 R (b) 675 (c) 1575 R (d) None of these

10. Two mole of an ideal gas is heated at constant pressure of one atmosphere from 27°C to 127°C. If $C_{v,m}$ 20 10 ^{2} T JK ^{1} mol ^{1}, then q and U for the process are respectively :

(a) 6362.8 J, 4700 J (b) 3037.2 J, 4700 J (c) 7062.8 J, 5400 J (d) 3181.4 J, 2350 J

11. 10 mole of an ideal gas is heated at constant pressure of one atmosphere from 27°C to 127°C. If $C_{v,m}$ 21.686 10 ^{3} T (JK ^{1} mol ^{1}), then H for the process is :

(a) 3000 J (b) 3350 J (c) 3700 J (d) 30350 J

12. For polytropic process PV^{n} constant, molar heat capacity (C_m) of an ideal gas is given by :

(a) $C_{v,m}$ $\frac{R}{(n \quad 1)}$ (b) $C_{v,m}$ $\frac{R}{(1 \quad n)}$ (c) $C_{v,m}$ R (d) $C_{p,m}$ $\frac{R}{(n \quad 1)}$

13. 2 mole of an ideal monoatomic gas undergoes a reversible process for which PV^{2} C. The gas is expanded from initial volume of 1 L to final volume of 3 L starting from initial temperature of 300 K. Find H for the process :

(a) 600 R (b) 1000 R (c) 3000 R (d) None of these

14. Calculate S for 3 mole of a diatomic ideal gas which is heated and compressed from 298 K and 1 bar to 596 K and 4 bar: [Given: $C_{v,m}$ (gas) $\frac{5}{2}R$; ln (2) 0.70; R 2 cal K ^{1} mol ^{1}]

(a) 14.7 cal K ^{1} (b) +14.7 cal K ^{1} (c) 4.9 cal K ^{1} (d) 6.3 cal K^{-1}

15. One mole of an ideal monoatomic gas at 27°C is subjected to a reversible isoentropic compression until final temperature reaches to 327°C. If the initial pressure was 1.0 atm then find the value of (ln P_2): (Given : ln 2 0.7)

(a) 1.75 atm (b) 0.176 atm (c) 1.0395 atm (d) 2.0 atm

16. Two mole of an ideal gas is expanded irreversibly and isothermally at 37°C until its volume is doubled and 3.41 kJ heat is absorbed from surrounding. S_{total} (system + surrounding) is :

(a) 0.52 J/K (b) 0.52 J/K (c) 22.52 J/K (d) 0

17. For a perfectly crystalline solid $C_{p,m}$ aT^{3} bT, where a and b are constant. If $C_{p,m}$ is 0.40 J/K mol at 10 K and 0.92 J/K mol at 20 K, then molar entropy at 20 K is :

(a) 0.92 J/K mol (b) 8.66 J/K mol (c) 0.813 J/K mol (d) None of these

For More Material Join: @JEEAdvanced_2024

18. Which of the following statement(s) is/are correct?

Statement (i) : The entropy in isolated system with P–V work only, is always maximized at equilibrium.

Statement (ii) : It is possible for the entropy of a closed system to decrease substantially in an irreversible process.

Statement (iii) : Entropy can be created but not destroyed.

Statement (iv) : S_{system} is zero for reversible process in an isolated system.

(a) Statement i, ii, iii (b) Statement ii, iv

(c) Statement i, ii, iv (d) All of these

19. Combustion of sucrose is used by aerobic organisms for providing energy for the life sustaining processes. If all the capturing of energy from the reaction is done through electrical process (non P–V work) then calculate maximum available energy which can be captured by combustion of 34.2 gm of sucrose

Given : $H_{combustion}$ (sucrose) 6000 kJ mol ^{1}

$S_{combustion}$ 180 J/K-mol and body temperature is 300 K

(a) 600 kJ (b) 594.6 kJ (c) 5.4 kJ (d) 605.4 kJ

20. For the hypothetical reaction

$$A_2(g) \quad B_2(g) \rightleftharpoons 2AB(g)$$

$_rG$ and $_rS$ are 20 kJ/mol and 20 JK ^{1} mol ^{1} respectively at 200 K.

If $_rC_p$ is 20 JK ^{1} mol ^{1} then $_rH$ at 400 K is :

(a) 20 kJ/mol (b) 7.98 kJ/mol (c) 28 kJ/mol (d) None of these

21. Calculate $_fG$ for (NH_4Cl, s) at 310 K.

Given : $_fH$ (NH_4Cl, s) 314.5 kJ/mol; $_rC_p$ 0

$S_{N_2(g)}$ 192 JK ^{1} mol ^{1}; $S_{H_2(g)}$ 130.5 JK ^{1} mol ^{1};

$S_{Cl_2(g)}$ 233 JK ^{1} mol ^{1}; $S_{NH_4Cl(s)}$ 99.5 JK ^{1} mol ^{1}

All given data at 300 K.

(a) 198.56 kJ/mol (b) 426.7 kJ/mol (c) 202.3 kJ/mol (d) None of these

22. Using listed informations, calculate $_rG$ (in kJ/mol) at 27°C

$$Co_3O_4(s) \quad 4CO(g) \quad 3Co(s) \quad 4CO_2(g)$$

Given : At 300 K H_f (kJ/mol) 891, 110.5, 0.0, 393.5

S (J/K-mol) 102.5, 197.7, 30.0, 213.7

(a) 214.8 (b) 195.0 (c) 200.3 (d) 256.45

23. Fixed mass of an ideal gas contained in a 24.63 L sealed rigid vessel at 1 atm is heated from 73 C to 27 C. Calculate change in Gibb's energy if entropy of gas is a function of temperature as S 2 10 ^{2} T (J/K): (Use 1 atm L 0.1 kJ)

(a) 1231.5 J (b) 1281.5 J (c) 781.5 J (d) 0

24. The molar heat capacities at constant pressure (assumed constant with respect to temperature) of A, B and C are in ratio of 3 : 1.5 : 2.0. The enthalpy change for the exothermic reaction A $2B$ $3C$ at 300 K and 310 K is H_{300} and H_{310} respectively then :

For More Material Join: @JEEAdvanced_2024

(a) H_{300} H_{310}
(b) H_{300} H_{310}
(c) H_{300} H_{310}
(d) if T_2 T_1 then H_{310} H_{300} and if T_2 T_1 then H_{310} H_{300}

25. Determine U at 300 K for the following reaction using the listed enthalpies of reaction :

$$4CO(g) \quad 8H_2(g) \quad 3CH_4(g) \quad CO_2(g) \quad 2H_2O(l)$$

C (graphite) $1/2\ O_2(g)$ $CO(g)$; H_1 110.5 kJ

$CO(g)$ $1/2\ O_2(g)$ $CO_2(g)$; H_2 282.9 kJ

$H_2(g)$ $1/2\ O_2(g)$ $H_2O(l)$; H_3 285.8 kJ

C (graphite) $2H_2(g)$ $CH_4(g)$; H_4 74.8 kJ

(a) 653.5 kJ (b) 686.2 kJ (c) 747.4 kJ (d) None of these

26. Calculate $_fH$ (in kJ/mol) for Cr_2O_3 from the $_rG$ and the S values provided at 27°C

$$4Cr(s) \quad 3O_2(g) \quad 2Cr_2O_3(s); \quad {}_rG \quad 2093.4 \text{ kJ/mol}$$

S (J/K mol) : S (Cr, s) 24; S (O_2, g) 205; S (Cr_2O_3, s) 81

(a) 2258.1 kJ/mol (b) 1129.05 kJ/mol (c) 964.35 kJ/mol (d) None of these

27. Calculate the heat produced (in kJ) when 224 gm of CaO is completely converted to $CaCO_3$ by reaction with CO_2 at 27°C in a container of fixed volume.

Given : H_f ($CaCO_3$, s) 1207 kJ/mol; H_f (CaO, s) 635 kJ/mol

H_f (CO_2, g) 394 kJ/mol; [Use R 8.3 JK^{-1} mol^{-1}]

(a) 702.04 kJ (b) 721.96 kJ (c) 712 kJ (d) 721 kJ

28. When 1.0 g of oxalic acid ($H_2C_2O_4$) is burned in a bomb calorimeter whose heat capacity is 8.75 kJ/K, the temperature increases by 0.312 K. The enthalpy of combustion of oxalic acid at 27°C is :

(a) 245.7 kJ/mol (b) 244.452 kJ/mol (c) 241.5 kJ/mol (d) None of these

29. Enthalpy of neutralization of H_3PO_3 acid is 106.68 kJ/mol using NaOH. Enthalpy of neutralization of HCl by NaOH is 55.84 kJ/mol. Calculate $H_{\text{ionization}}$ of H_3PO_3 into its ions :

(a) 50.84 kJ/mol (b) 5 kJ/mol (c) 2.5 kJ/mol (d) None of these

30. The enthalpy of neutralization of a weak monoprotic acid (H*A*) in 1 *M* solution with a strong base is 55.95 kJ/mol. If the unionized acid required 1.4 kJ/mol heat for its complete ionization and enthalpy of neutralization of the strong monobasic acid with a strong monoacidic base is 57.3 kJ/mol, what is the % ionization of the weak acid in molar solution ?

(a) 1% (b) 3.57% (c) 35.7% (d) 10%

31. Determine C—C and C—H bond enthalpy (in kJ/mol)

Given : $_fH$ (C_2H_6, g) 85 kJ/mol, $_fH$ (C_3H_8, g) 104 kJ/mol

$_{sub}H$ (C, s) 718 kJ/mol, B.E. (H—H) 436 kJ/mol

(a) 414, 345 (b) 345, 414 (c) 287, 404.5 (d) None of these

32. Consider the following data : $_fH$ (N_2H_4, l) 50 kJ/mol, $_fH$ (NH_3, g) 46 kJ/mol

B.E. (N—H) 393 kJ/mol and B.E. (H—H) 436 kJ/mol,

$_{vap}H$ (N_2H_4, l) 18 kJ/mol

The N—N bond energy in N_2H_4 is :

(a) 226 kJ/mol (b) 154 kJ/mol (c) 190 kJ/mol (d) None of these

For More Material Join: @JEEAdvanced_2024

33. What is the bond enthalpy of Xe—F bond?

$XeF_4(g)$ Xe (g) F (g) $F_2(g)$ F(g) ; $_rH$ 292 kcal/mol

Given : Ionization energy of Xe 279 kcal/mol

B.E. (F—F) 38 kcal/mol, Electron affinity of F 85 kcal/mol

(a) 24 kcal/mol (b) 34 kcal/mol (c) 8.5 kcal/mol (d) None of these

34. If enthalpy of hydrogenation of $C_6H_6(l)$ into $C_6H_{12}(l)$ is 205 kJ/mol and resonance energy of $C_6H_6(l)$ is 152 kJ/ mol then enthalpy of hydrogenation of  is

Assume H_{vap} of $C_6H_6(l), C_6H_8(l), C_6H_{12}(l)$ all are equal :

(a) 535.5 kJ/mol (b) 238 kJ/mol (c) –357 kJ/mol (d) 119 kJ/mol

35. The enthalpy of combustion of propane (C_3H_8) gas in terms of given data is :

Bond energy (kJ/mol)


C H	O=O	C=O	O—H	C—C
x_1	x_2	x_3	x_4	x_5

Resonance energy of CO_2 is z kJ/mol and $H_{vaporization}$ $[H_2O(l)]$ is y (kJ/mol)

(a) $8x_1$ $2x_5$ $5x_2$ $6x_3$ $8x_4$ $4y$ $3z$ (b) $6x_1$ x_5 $5x_2$ $3x_3$ $4x_4$ $4y$ $3z$

(c) $8x_1$ $2x_5$ $5x_2$ $6x_3$ $8x_4$ y z (d) $8x_1$ x_5 $5x_2$ $6x_3$ $8x_4$ $4y$ $3z$


36. The efficiency of an ideal gas with adiabatic exponent ' ' for the shown cyclic process would be



(a) $\frac{(\quad 1)(2\ln 2 \quad 1)}{1 \quad (\quad 1)2\ln 2}$ (b) $\frac{(\quad 1)(1 \quad 2\ln 2)}{(\quad 1)2\ln 2 \quad 1}$

(c) $\frac{(2\ln 2 \quad 1)(\quad 1)}{(\quad 1)2\ln 2 \quad 1}$ (d) $\frac{(2\ln 2 \quad 1)}{/(\quad 1)}$

37. Initially one mole of ideal gas $C_v \quad \frac{5}{2}R$ at 1.0 atm and 300 K is put through the following cycle :



Step-I : Heating to twice its initial pressure at constant volume.

Step-II : Adiabatic expansion to its initial temperature.

For More Material Join: @JEEAdvanced_2024

Step-III : Isothermal compression back to 1.00 atm.

What is the volume at state X ?

(a) 40.4 L (b) 65.0 L (c) 139 L (d) 4.35 L

38. What is the enthalpy of neutralization of HF against a strong base ?

Given : H (aq) OH (aq) $H_2O(l)$; $_rH$ 56 kJ/mol

$_sH$ (HF, aq) 329 kJ/mol ; $_sH$ (H_2O, l) 285 kJ/mol

$_sH$ (F , aq) 320 kJ/mol

(a) 17 kJ/mol (b) 38 kJ/mol (c) 47 kJ/mol (d) 43 kJ/mol

39. The molar heat capacities of Iodine vapour and solid are 7.8 and 14 cal/mol respectively. If enthalpy of sublimation of iodine is 6096 cal/mol at 200°C, then what is U (internal energy change) at 250°C in cal/mol

(a) 5360 (b) 4740 (c) 6406 (d) None of these

40. For the reaction taking place at certain temperature $NH_2COONH_4(s) \rightleftharpoons 2NH_3(g)$ $CO_2(g)$, if equilibrium pressure is $3X$ bar then $_rG$ would be

(a) $RT \ln 9$ $3RT \ln X$ (b) $RT \ln 4$ $3RT \ln X$

(c) $3RT \ln X$ (d) None of these

For More Material Join: @JEEAdvanced_2024

Level 3

Passage 1

The first law of thermodynamics for a closed system is $dU = dq + dw$, where $dw = dw_{pv} + dw_{non\text{-}pv}$. The most common type of $w_{non\text{-}pv}$ is electrical work. As per IUPAC convention work done on the system is positive.

1. A system generates 50 J of electrical energy, has 150 J of pressure-volume work done on it by the surroundings while releasing 300 J of heat energy. What is the change in the internal energy of the system?
(a) 500 (b) 100 (c) 300 (d) 200

2. A system generates 50 J of electrical energy and delivers 150 J of pressure-volume work against the surroundings while releasing 300 J of heat energy. What is the change in the internal energy of the system?
(a) 100 (b) 400 (c) 300 (d) 500

Passage 2

If the boundary of system moves by an infinitesimal amount, the work involved is given by $dw = P_{ext}\, dV$

for irreversible process $w = P_{ext}\, V$ (where $V = V_f - V_i$)

for reversible process $P_{ext} = P_{int} - dP \sim P_{int}$

so for reversible isothermal process $w = nRT \ln \dfrac{V_f}{V_i}$

2 mole of an ideal gas undergoes isothermal compression along three different paths :

(i) reversible compression from $P_i = 2$ bar and $V_i = 8$ L to $P_f = 20$ bar

(ii) a single stage compression against a constant external pressure of 20 bar, and

(iii) a two stage compression consisting initially of compression against a constant external pressure of 10 bar until $P_{gas} = P_{ext}$, followed by compression against a constant pressure of 20 bar until $P_{gas} = P_{ext}$.

1. Work done (in bar-L) on the gas in reversible isothermal compression is :
(a) 9.212 (b) 36.848 (c) 18.424 (d) None of these

2. Work done on the gas in single stage compression is :
(a) 36 (b) 72 (c) 144 (d) None of these

3. Total work done on the gas in two stage compression is :
(a) 40 (b) 80 (c) 160 (d) None of these

4. Order of magnitude of work is :
(a) $w_1 = w_2 = w_3$ (b) $w_3 = w_2 = w_1$ (c) $w_2 = w_3 = w_1$ (d) $w_1 = w_2 = w_3$

For More Material Join: @JEEAdvanced_2024



Passage 3

Standard Gibb's energy of reaction ($_rG$) at a certain temperature can be computed as $_rG$ $_rH$ T. $_rS$ and the change in the value of $_rH$ and $_rS$ for a reaction with temperature can be computed as follows :

$$_rH_{T_2} \quad _rH_{T_1} \quad _rC_p\,(T_2 \quad T_1)$$

$$_rS_{T_2} \quad _rS_{T_1} \quad _rC_p \ln \frac{T_2}{T_1}$$

$$_rG \quad _rH \quad T. \ _rS$$

and by $_rG$ $RT \ln K_{eq}$.

Consider the following reaction : $CO(g)$ $2H_2(g) \rightleftharpoons CH_3OH(g)$

Given : $_fH$ (CH_3OH, g) 201 kJ/mol; $_fH$ (CO, g) 114 kJ/mol

S (CH_3OH, g) 240 J/K-mol ; S (H_2, g) 29 JK^{-1} mol^{-1}

S (CO, g) 198 J/mol-K ; $C_{p,m}$ (H_2) 28.8 J/mol-K

$C_{p,m}$ (CO) 29.4 J/mol-K; $C_{p,m}$ (CH_3OH) 44 J/mol-K

and $\ln \frac{320}{300}$ 0.06, all data at 300 K

1. $_rS$ at 300 K for the reaction is :
(a) 152.6 J/K-mol (b) 181.6 J/K-mol (c) 16 J/K-mol (d) None of these

2. $_rH$ at 300 K for the reaction is :
(a) 87 kJ/mol (b) 87 kJ/mol (c) 315 kJ/mol (d) 288 kJ/mol

3. $_rS$ at 320 K is :
(a) 155.18 J/mol-K (b) 150.02 J/mol-K (c) 172 J/mol-K (d) None of these

4. $_rH$ at 320 K is :
(a) 288.86 kJ/mol (b) 289.1 kJ/mol (c) 87.86 kJ/mol (d) None of these

5. $_rG$ at 320 K is :
(a) 48295.2 kJ/mol (b) 240.85 kJ/mol (c) 240.85 kJ/mol (d) 81.91 kJ/mol



Passage 4

Enthalpy of neutralization is defined as the enthalpy change when 1 mole of acid/base is completely neutralized by base/acid in dilute solution.

For strong acid and strong base neutralization net chemical change is

$$H \ (aq) \quad OH \ (aq) \quad H_2O(l); \quad _rH \quad 55.84 \text{ kJ/mol}$$

$H_{\text{ionization}}$ of aqueous solution of strong acid and strong base is zero.

When a dilute solution of a weak acid or base is neutralized, the enthalpy of neutralization is some what less because of the absorption of heat in the ionization of the weak acid or base,

for weak acid/base

$$H_{\text{neutralization}} \quad H_{\text{ionization}} \quad _rH \quad (H \quad OH \quad H_2O)$$

For More Material Join: @JEEAdvanced_2024

1. If enthalpy of neutralization of CH_3COOH by HCl is 49.86 kJ/mol then enthalpy of ionization of CH_3COOH is :
(a) 5.98 kJ/mol (b) 5.98 kJ/mol (c) 105.7 kJ/mol (d) None of these

2. What is H for complete neutralization of strong diacidic base $A(OH)_2$ by HNO_3?
(a) 55.84 kJ/mol (b) 111.68 kJ/mol (c) 55.84 kJ/mol (d) None of these

3. Under the same conditions how many mL of 0.1 M NaOH and 0.05 M H_2A (strong diprotic acid) solution should be mixed for a total volume of 100 mL produce the highest rise in temperature :
(a) 25 : 75 (b) 50 : 50 (c) 75 : 25 (d) 66.66 : 33.33



Passage 5

Gibbs Helmholtz equation relates the enthalpy, entropy and free energy change of the process at constant pressure and temperature as

$G \quad H \quad T \; S$ (at constant P, T)

In General the magnitude of H does not change much with the change in temperature but the term $T \; S$ changes appreciably. Hence in some process spontaneity is very much dependent on temperature and such processes are generally known as entropy driven process.

1. For the reaction at 298 K; $A_2B_4 \quad 2AB_2$
$H \quad 2$ kJ and $S \quad 20$ J/K at constant P and T, the reaction will be
(a) spontaneous and entropy driven (b) spontaneous and enthalpy driven
(c) non-spontaneous (d) at equilibrium

2. When $CaCO_3$ is heated to a high temperature it decomposes into CaO and CO_2, however it is quite stable at room temperature. It can be explained by the fact that
(a) $_rH$ dominates the term $T \; S$ at high temperature
(b) the term $T \; S$ dominates the $_rH$ at high temperature
(c) at high temperature both $_rS$ and $_rH$ becomes negative
(d) thermodynamics can not say anything about spontaneity

3. The Dissolution of $CaCl_2 \; 6H_2O$ in a large volume of water is endothermic to the extent of 3.5 kcal mol ^{1} and H for the reaction is 23.2 kcal mol ^{1}.

$CaCl_2(s) \quad 6H_2O(l) \qquad CaCl_2 \; 6H_2O(s)$

Select the correct statement :
(a) $H_{Solution}$ for anhydrous $CaCl_2$ is –19.7 kcal/mol and the process is enthalpy driven
(b) $H_{Solution}$ for anhydrous $CaCl_2$ is –19.7 kcal/mol and the process is entropy driven
(c) Dissolution of $CaCl_2 \; 6H_2O$ in water is enthalpy driven process
(d) The $_rS$ the reaction $CaCl_2(s) \quad 6H_2O(l) \qquad CaCl_2 \; 6H_2O(s)$ is negative

For More Material Join: @JEEAdvanced_2024

One or More Answers is/are Correct

1. Identify the intensive quantities from the following :
(a) Enthalpy (b) Temperature (c) Pressure (d) Mass

2. Identify the extensive quantities from the following :
(a) Gibb's energy (b) Entropy (c) Refractive index (d) Specific heat

3. Identify the state functions from the following :
(a) Heat (b) Work (c) Enthalpy (d) Enthalpy change

4. Which of the following statements is/are correct as per IUPAC sign convention?
(a) The work done by the system on the surrounding is negative
(b) The work done by the surrounding on the system is positive
(c) The heat absorbed by the system from the surrounding is positive
(d) The heat absorbed by the surrounding from the system is positive


5. In an isothermal irreversible expansion of an ideal gas as per IUPAC sign convention :
(a) U 0 (b) H 0 (c) w $nRT \ln \frac{P_1}{P_2}$ (d) w q

6. In reversible isothermal expansion of an ideal gas :
(a) w 0 (b) U_1 U_2 (c) H_1 H_2 (d) q $nRT \ln \frac{V_2}{V_1}$

7. An adiabatic process is that process in which :
(a) energy is transferred as heat
(b) no energy is transfer red as heat
(c) U w
(d) the temp. of gas increases in a reversible adiabatic compression

8. In adiabatic process, the work involved during expansion or compression of an ideal gas is given by :
(a) nC_v T (b) $\frac{nR}{1}(T_2 \quad T_1)$
(c) nRP_{ext} $\frac{T_2P_1 \quad T_1P_2}{P_1P_2}$ (d) $2.303\ RT \log \frac{V_2}{V_1}$

9. One mole of an ideal gas is subjected to a two step reversible process (A—B and B—C). The pressure at A and C is same. Mark the correct statement(s) :
(a) Work involved in the path AB is zero
(b) In the path AB work will be done on the gas by the surrounding
(c) Volume of gas at C 3 volume of gas at A
(d) Volume of gas at B is 16.42 litres



10. Assume ideal gas behaviour for all the gases considered and neglect vibrational degrees of freedom. Separate equimolar samples of Ne, O_2, CO_2 and SO_2 were subjected to a two process as mentioned. Initially all are at same state of temperature and pressure.

For More Material Join: @JEEAdvanced_2024

Step I All undergo reversible adiabatic expansion to attain same final volume, which is double the original volume thereby causing the decreases in their temperature.

Step II After step I all are given appropriate amount of heat isochorically to restore the original temperature.

Mark the correct option(s) :

(a) Due to step I only, the decrease in temperature will be maximum for Ne

(b) During step II, heat given will be minimum for SO_2

(c) There will be no change in internal energy for any of the gas after both the steps of process are competed

(d) The P–V graph of O_2 and CO_2 will be same

11. Which of the following is/are correct?

(a) H U (PV) when P and V both changes

(b) H U P V when pressure is constant

(c) H U V P when volume is constant

(d) H U P V V P when P and V both changes

12. H U for the reaction(s) :

(a) $N_2(g)$ $3H_2(g)$ $2NH_3(g)$

(b) $Ag_2O(s)$ $2Ag(s)$ $\frac{1}{2}O_2(g)$

(c) $CO(g)$ $\frac{1}{2}O_2(g)$ $CO_2(g)$

(d) $C(s)$ $O_2(g)$ $CO_2(g)$

13. Which of the following conditions may lead to a non-spontaneous change?

(a) H and S both + ve

(b) H ve; S ve

(c) H ve; S ve

(d) H ve; S ve

14. For a process to be spontaneous :

(a) $(G_{system})_{T,p}$ 0

(b) S_{system} $S_{surrounding}$ 0

(c) S_{system} $S_{surrounding}$ 0

(d) $(G_{system})_{T,p}$ 0

15. The normal boiling point of a liquid 'X' is 400 K. Which of the following statement is true about the process $X(l)$ $X(g)$?

(a) at 400 K and 1 atm pressure G 0

(b) at 400 K and 2 atm pressure G ve

(c) at 400 K and 0.1 atm pressure G ve

(d) at 410 K and 1 atm pressure G ve

16. When ice melts at 1°C :

(a) an increase in entropy

(b) a decrease in enthalpy

(c) a decrease in free energy

(d) process is spontaneous

17. The value of $H_{transition}$ of C (graphite) C (diamond) is 1.9 kJ/mol at 25°C. Entropy of graphite is higher than entropy of diamond. This implies that :

(a) C (diamond) is more thermodynamically stable than C (graphite) at 25°C

(b) C (graphite) is more thermodynamically stable than C (diamond) at 25°C

(c) diamond will provide more heat on complete combustion at 25°C

(d) $G_{transition}$ of C (diamond) C (graphite) is ve

18. Which of the following statement(s) is/are false?

(a) All adiabatic processes are isoentropic (or isentropic) processes

(b) When $(G_{system})_{T,p}$ 0; the reaction must be exothermic

For More Material Join: @JEEAdvanced_2024

(c) $dG \quad VdP \quad SdT$ is applicable for closed system, both PV and non-PV work

(d) The heat of vaporisation of water at 100°C is 40.6 kJ/mol. When 9 gm of water vapour condenses to liquid at 100°C and 1 atm, then $S_{system} \quad 54.42$ J/K

19. Which of the following statement(s) is/are true?

(a) $U \quad 0$ for combustion of $C_2H_6(g)$ in a sealed rigid adiabatic container

(b) ${}_fH$ (S, monoclinic) $\quad 0$

(c) If dissociation energy of $CH_4(g)$ is 1656 kJ/mol and $C_2H_6(g)$ is 2812 kJ/mol, then value of C—C bond energy will be 328 kJ/mol

(d) If $H_f(H_2O, g) \quad 242$ kJ/mol; $H_{vap}(H_2O, l) \quad 44$ kJ/mol then, ${}_fH$ (OH , $aq.$) will be $\quad 142$ kJ/mol

20. From the following data, mark the option(s) where H is correctly written for the given reaction.

Given : H (aq) OH $(aq) \quad H_2O(l)$; $H \quad 57.3$ kJ

$H_{solution}$ of $HA(g) \quad 70.7$ kJ/mol

$H_{solution}$ of $BOH(g) \quad 20$ kJ/mol

$H_{ionization}$ of $HA \quad 15$ kJ/mol and BOH is a strong base.

Reaction	${}_rH$ (kJ/mol)
(a) $HA(aq) \quad BOH(aq) \quad BA(aq) \quad H_2O$	42.3
(b) $HA(g) \quad BOH(g) \quad BA(aq) \quad H_2O$	93
(c) $HA(g) \quad$ H $(aq) \quad A \ (aq)$	55.7
(d) $B \ (aq) \quad$ OH $(aq) \quad BOH(aq)$	20

21. Select correct statement(s)

(a) An adiabatic system can exchange energy with its surroundings.

(b) A thermodynamic property which is intensive is additive.

(c) Work done may be zero in a cyclic process.

(d) For a simple compressible substance, the relation $dq \quad P.dv \quad 0$ is true for any cycle involving mechanical work only.

22. For an isolated system, the entropy :

(a) either increases or remains constant (b) either decreases or remains constant

(c) can never decrease (d) can never increase

23. The normal boiling point of a liquid X is 400 K. H_{vap} at normal boiling point is 40 kJ/mol. Select correct statement(s) :

(a) $S_{Vaporisation} \quad 100$ J/mol.K at 400 K and 2 atm

(b) $S_{Vaporisation} \quad 10$ J/mol.K at 400 K and 1 atm

(c) $G_{Vaporisation} \quad 0$ at 410 K and 1 atm

(d) $U \quad 43.32$ kJ/mol.K at 400 K and 1 atm

24. Select incorrect statement(s)

(a) A closed system with all adiabatic boundaries must be an isolated system

(b) Total heat exchange in a cyclic process may be zero

(c) Entropy of a closed system is maximum at equilibrium

(d) Molar Gibb's Energy is an extensive property

For More Material Join: @JEEAdvanced_2024

25. Select correct statement(s) for the reaction $H_2O(g)$ $CO(g)$ $H_2(g)$ $CO_2(g)$

Substance	$CO(g)$	$CO_2(g)$	$H_2O(g)$	$H_2(g)$
${}_fH_{400}$ (kcal mol^{-1})	25	95	55	0
S_{400} (cal mol^{-1}K^{-1})	45	50	40	30

(a) Reaction is enthalpy driven
(b) Reaction is entropy driven
(c) Reaction is spontaneous at 400 K
(d) Reaction is non-spontaneous at 400 K

Match the Column



Column-I and Column-II contains four entries each. Entries of Column-I are to be matched with some entries of column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

1.

Column-I	Column-II
(A) Reversible cooling of an ideal gas at constant volume	(P) w 0; q 0; U 0
(B) Reversible isothermal expansion of an ideal gas	(Q) w 0; q 0; U 0
(C) Adiabatic expansion of non-ideal gas into vacuum	(R) w 0; q 0; U 0
(D) Reversible melting of sulphur at normal melting point	(S) w 0; q 0; U 0

2.

Column-I	Column-II
(A) Adiabatic process	(P) q 0
(B) Isothermal process	(Q) H 0
(C) Isoenthalpic process	(R) T 0
(D) Isoentropic process	(S) S 0

3.

Column-I	Column-II
(A) Reversible isothermal expansion of an ideal gas	(P) w 2.303 $nRT \log \frac{V_2}{V_1}$
(B) Reversible adiabatic compression of an ideal gas	(Q) PV constant
(C) Irreversible adiabatic expansion of an ideal gas	(R) w $\frac{nR}{(\ 1)}(T_2\ T_1)$
(D) Irreversible isothermal compression of an ideal gas	(S) H 0

For More Material Join: @JEEAdvanced_2024

4.

Column-I	Column-II
(A) $PCl_5(g) \rightleftharpoons PCl_3(g)$ $Cl_2(g)$ in a closed system	(P) H U
(B) $2HI(g) \rightleftharpoons H_2(g)$ $I_2(g)$ in a closed system	(Q) H U 0
(C) $N_2(g)$ $3H_2(g) \rightleftharpoons 2NH_3(g)$ in a closed system	(R) H U
(D) $CO(g)$ $\frac{1}{2}O_2(g) \rightleftharpoons CO_2(g)$ in an isolated system	(S) U 0

5.

Column-I (Process)	Column-II (Entropy Change)
(A) Reversible isothermal compression of an ideal gas	(P) S_{system} 0
(B) Isothermal free expansion (P_{ext} 0) of an ideal gas	(Q) S_{system} 0
(C) Reversible adiabatic expansion of an ideal gas	(R) S_{system} 0
(D) Reversible ideal gas expansion	(S) Information insufficient

6.

Column-I (Process)	Column-II (Entropy change)
(A) Reversible isothermal ideal gas expansion	(P) $S_{surrounding}$ 0
(B) Reversible adiabatic ideal gas compression	(Q) $S_{surrounding}$ 0
(C) Adiabatic free expansion (P_{ext} 0) of an ideal gas	(R) $S_{surrounding}$ 0
(D) Irreversible isothermal ideal gas compression	(S) S_{system} 0

7.

Column-I	Column-II
(A) $(G_{system})_{T,P}$ 0	(P) Process is in equilibrium
(B) S_{system} $S_{surrounding} > 0$	(Q) Process is nonspontaneous
(C) S_{system} $S_{surrounding} < 0$	(R) Process is spontaneous
(D) $(G_{system})_{T,P}$ 0	(S) System is unable to do useful work

For More Material Join: @JEEAdvanced_2024

8.

Column-I	Column-II
(A) $(G_{system})_{T,P}$	(P) $nR \ln \frac{V_2}{V_1}$
(B) Work done in reversible isothermal ideal gas expansion	(Q) $nRT \ln \frac{P_2}{P_1}$
(C) G for reversible isothermal expansion of an ideal gas	(R) nFE
(D) S_{gas} for isothermal expansion of an ideal gas	(S) $nR \ln \frac{P_1}{P_2}$

9.

Column-I	Column-II
(A) Heating of an ideal gas at constant pressure	(P) H $nC_{p,m}$ T 0
(B) Compression of liquid at constant temperature	(Q) U 0
(C) Reversible process for an ideal gas at constant temperature	(R) G V P
(D) Adiabatic free expansion of an ideal gas	(S) G $nRT \ln \frac{P_2}{P_1}$

10.

Column-I	Column-II
(Sign of H and S respectively)	(Nature of reaction)
(A) &	(P) Spontaneous only at low temperature
(B) &	(Q) Spontaneous only at high temperature
(C) + & +	(R) Spontaneous at all temperature
(D) &	(S) Non-spontaneous at all temperature

11.

Column-I	Column-II
(A) Reversible adiabatic compression	(P) S_{system} 0
(B) Reversible vaporisation of liquid	(Q) S_{system} 0
(C) $2N(g)$ $N_2(g)$	(R) $S_{surrounding}$ 0
(D) $MgCO_3(s)$ $MgO(s)$ $CO_2(g)$	(S) $S_{surrounding}$ 0

For More Material Join: @JEEAdvanced_2024

12.

Column-I	Column-II
(A) $H^+(aq)$	(P) $\Delta_f H^\circ = 0$
(B) $H(g)$	(Q) $\Delta_f H^\circ < 0$
(C) $H_2(g)$	(R) $\Delta_f G^\circ = 0$
(D) $C(s, \text{diamond})$	(S) $\Delta_f S^\circ = 0$

13.

Column-I (Partial derivative)	Column-II (Thermodynamic variable)
(A) $\left(\frac{\partial U}{\partial T}\right)_V$	(P) C_P
(B) $\left(\frac{\partial H}{\partial T}\right)_P$	(Q) C_V
(C) $\left(\frac{\partial G}{\partial T}\right)_P$	(R) S
(D) $\left(\frac{\partial G}{\partial P}\right)_T$	(S) V

14.

Column-I	Column-II
(A) $C(s, \text{graphite}) + O_2(g) \rightarrow CO_2(g)$	(P) $\Delta H_{\text{combustion}}$
(B) $C(s, \text{graphite}) \rightarrow C(g)$	(Q) $\Delta H_{\text{formation}}$
(C) $CO(g) + \frac{1}{2}O_2(g) \rightarrow CO_2(g)$	(R) $\Delta H_{\text{atomization}}$
(D) $CH_4(g) \rightarrow C(g) + 4H(g)$	(S) $\Delta H_{\text{sublimation}}$

15.

Column-I	Column-II
(A) $O_2(g)$	(P) $\Delta_f H^\circ$ ve; $\Delta_f S^\circ$ ve
(B) $O_3(g)$	(Q) $\Delta_f H^\circ$ ve; $\Delta_f S^\circ$ ve
(C) $Br_2(g)$	(R) $\Delta_f H^\circ$ ve; $\Delta_f S^\circ$ ve
(D) $H_2O(l)$	(S) $\Delta_f H^\circ = 0$; $\Delta_f S^\circ = 0$

For More Material Join: @JEEAdvanced_2024

Assertion-Reason Type Questions



Each question contains STATEMENT-1 (Assertion) and STATEMENT-2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below :

(A) If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1

(B) If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1

(C) If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

(D) If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

1. **STATEMENT-1 :** Heat and work are "definite quantities".
   STATEMENT-2 : Heat and work are not properties of a system. Their values depend on the path of the process and vary accordingly.
2. **STATEMENT-1 :** There is no change in internal energy for an ideal gas at constant temperature.
   STATEMENT-2 : Internal energy of an ideal gas is a function of temperature only.
3. **STATEMENT-1 :** The heat absorbed during the isothermal expansion of an ideal gas against vacuum is zero.
   STATEMENT-2 : The volume occupied by the molecules of an ideal gas is zero.
4. **STATEMENT-1 :** H and U are the same for the reaction, $N_2(g)$ $O_2(g) \rightleftharpoons 2NO(g)$,
   STATEMENT-2 : All reactants and products are gases where all gases are ideal.
5. **STATEMENT-1 :** The magnitude of the work involved in an isothermal expansion is greater than that involved in an adiabatic expansion.
   STATEMENT-2 : P V curve (P on y-axis and V on x-axis) decrease more rapidly for reversible adiabatic expansion compared to reversible isothermal expansion starting from same initial state.
6. **STATEMENT-1 :** Entropy change in reversible adiabatic expansion of an ideal gas is zero.
   STATEMENT-2 : The increase in entropy due to volume increase just componsate the decrease in entropy due to fall in temperature.
7. **STATEMENT-1 :** There cannot be chemical equilibrium in an open system.
   STATEMENT-2 : There is no fixed mass in an open system.
8. **STATEMENT-1 :** The standard free energy changes of all spontaneously occuring reactions are negative.
   STATEMENT-2 : The standard free energies of the elements in their standard states at 1 bar and 298 K are taken as zero.
9. **STATEMENT-1 :** Enthalpy and entropy of any elementary substance in the standard states are taken as zero.
   STATEMENT-2 : At absolute zero, particles of the perfectly crystalline substance become completely motionless.
10. **STATEMENT-1 :** A reaction which is spontaneous and accompanied by decrease of randomness must be exothermic.
    STATEMENT-2 : All exothermic reactions are accompanied by decrease of randomness.

For More Material Join: @JEEAdvanced_2024

11. STATEMENT-1 : Many endothermic reactions that are not spontaneous at room temperatures become spontaneous at high temperature.
STATEMENT-2 : H of the endothermic reaction may increases with increase in temperature.

12. STATEMENT-1 : Decrease of free energy during the process under constant temperature and pressure provides a measure of its spontaneity.
STATEMENT-2 : A spontaneous change must have +ve sign of S_{system}.

13. STATEMENT-1 : $|\ _fH|$ of (H_2O, l) $|\ _fH|$ of (H_2O, g)
STATEMENT-2 : $H_{condensation}$ is negative.

14. STATEMENT-1 : All combustion reactions are exothermic.
STATEMENT-2 : Enthalpies of products are greater than enthalpies of reactants ($v_p\ _fH(P)$ $v_R\ _fH(R)$)

15. STATEMENT-1 : Enthalpy of neutralization of CH_3COOH by NaOH is less than that of HCl by NaOH.
STATEMENT-2 : Enthalpy of neutralization of CH_3COOH is less because of the absorption of heat in the ionization process.

16. STATEMENT-1 : Internal energy of a real gas may change during expansion at constant temperature.
STATEMENT-2 : Internal energy of a real gas is a function of temperature and volume.

17. STATEMENT-1 : Work is a state function which is expressed in joule.
STATEMENT-2 : Work appears only at the boundary of the system.

18. STATEMENT-1 : The expansion of a gas into an evacuated space takes place non-spontaneously.
STATEMENT-2 : A process in which all steps cannot be retraced by themselves is called a spontaneous process.

Subjective Problems



1. A perfect gas undergoes a reversible adiabatic expansion from (300 K, 200 atm) to (90 K, 10 atm). Find the atomicity of gas.

2. 5 mole of an ideal gas at temp. T are compressed isothermally from 12 atm. to 24 atm. Calculate the value of $10r$

Where, r $\dfrac{\text{Work done along reversible process}}{\text{Work done along single step irreversible process}}$ (Given : $\ln 2$ 0.7)

3. A diatomic ideal gas is expanded according to PV^3 constant, under very high temperature (Assume vibration mode active). Calculate the molar heat capacity of gas (in cal / mol K) in this process.

4. A heat engine is operating between 500 K to 300 K and it absorbs 10 kcal of heat from 500 K reservoir reversibly per cycle. Calculate the work done (in kcal) per cycle.

5. In a carnot cycle involving ideal non-linear triatomic gas, if during adiabatic expansion volume increases from 2L to 16 L and heat absorbed during isothermal expansion is 8 kcal, then calculate magnitude of work done by carnot engine (in kcal).

For More Material Join: @JEEAdvanced_2024

6. Molar heat capacities at constant pressure for A, B and C are 3, 1.5 and 2 J/K mol. The enthalpy of reaction and entropy of reaction, $A \quad 2B \quad 3C$ are 20 kJ/mol and 20 J/K mol at 300 K. Calculate G (in kJ / mol) for the reaction ,

$$\frac{1}{2}A \quad B \quad \frac{3}{2}C$$

7. Standard molar enthalpy of combustion of glucose is 2880 kJ. If only 25% of energy is available for muscular work and 1.0 km walk consumes 90 kJ of energy, what maximum distance (in km) a person can walk after eating 90 g of glucose.

8. Given $C_2H_2(g) \quad H_2(g) \quad C_2H_4(g)$: $H \quad 175\,kJ\,mol^{-1}$

$H_{f(C_2H_4,g)} \quad 50\,kJ\,mol^{-1}$; $H_{f(H_2O,l)} \quad 280\,kJ\,mol^{-1}$; $H_{f(CO_2g)} \quad 390\,kJ\,mol^{-1}$

If H is enthalpy of combustion (in $kJ\,mol^{-1}$) of $C_2H_2(g)$, then calculate the value of $\left|\frac{H}{257}\right|$.

9. The integral enthalpies of solution of anhydrous $CuSO_4(s)$ and hydrated $CuSO_4 \cdot 5H_2O(s)$ are 70 kJ per mol and 10 kJ per mol respectively. Determine the magnitude of enthalpy of hydration of 0.1 mole anhydrous $CuSO_4(s)$ as

$CuSO_4(s) \quad 5H_2O(l) \quad CuSO_4 \cdot 5H_2O(s)$

10. If enthalpy of neutralisation of HCl by NaOH is $57\,kJ\,mol^{-1}$ and with NH_4OH is $50\,kJ\,mol^{-1}$. Calculate enthalpy of ionisation of $NH_4OH(aq)$.


11. Lattice energy of NaCl(s) is $790\,kJ\,mol^{-1}$ and enthalpy of hydration is $785\,kJ\,mol^{-1}$. Calculate enthalpy of solution of NaCl(s).

12. x g sample of NH_4NO_3 is decomposed in a Bomb calorimeter. The temperature of calorimeter increase by 4 C. The heat capacity of the system is 1.25 kJ/ C. Calculate the value of x.
Given molar heat of decomposition of NH_4NO_3 at constant volume is $400\,kJ\,mol^{-1}$.

13. A heat engine operating between 227°C and 77°C absorbs 10 kcal of heat from the 227°C reservoir reversibly per cycle. Calculate total work done (in kcal) in two cycles.

14. Calculate work done in chemical reaction (in kcal)
$A(s) \quad 3B(g) \quad C(l)$ at 227°C at 1 atm in closed vessel.

15. One mole ideal monoatomic gas is heated according to path AB and AC.
If temperature of state B and state C are equal.
Calculate $\frac{q_{AC}}{q_{AB}}$ 10.



For More Material Join: @JEEAdvanced_2024

Answers

Level 1

1.	(d)	**2.**	(b)	**3.**	(d)	**4.**	(c)	**5.**	(a)	**6.**	(c)	**7.**	(a)	**8.**	(c)	**9.**	(d)	**10.**	(d)
11.	(c)	**12.**	(b)	**13.**	(d)	**14.**	(d)	**15.**	(a)	**16.**	(b)	**17.**	(b)	**18.**	(a)	**19.**	(b)	**20.**	(c)
21.	(c)	**22.**	(c)	**23.**	(c)	**24.**	(d)	**25.**	(b)	**26.**	(c)	**27.**	(b)	**28.**	(c)	**29.**	(c)	**30.**	(d)
31.	(a)	**32.**	(c)	**33.**	(b)	**34.**	(b)	**35.**	(c)	**36.**	(a)	**37.**	(a)	**38.**	(a)	**39.**	(b)	**40.**	(b)
41.	(b)	**42.**	(c)	**43.**	(b)	**44.**	(b)	**45.**	(b)	**46.**	(c)	**47.**	(a)	**48.**	(b)	**49.**	(c)	**50.**	(c)
51.	(a)	**52.**	(b)	**53.**	(d)	**54.**	(b)	**55.**	(b)	**56.**	(b)	**57.**	(b)	**58.**	(b)	**59.**	(b)	**60.**	(b)
61.	(a)	**62.**	(c)	**63.**	(c)	**64.**	(d)	**65.**	(c)	**66.**	(a)	**67.**	(a)	**68.**	(c)	**69.**	(d)	**70.**	(c)
71.	(d)	**72.**	(c)	**73.**	(b)	**74.**	(b)	**75.**	(c)	**76.**	(c)	**77.**	(b)	**78.**	(d)	**79.**	(a)	**80.**	(b)
81.	(b)	**82.**	(b)	**83.**	(c)	**84.**	(d)	**85.**	(d)	**86.**	(b)	**87.**	(b)	**88.**	(a)	**89.**	(b)	**90.**	(b)
91.	(a)	**92.**	(d)	**93.**	(d)	**94.**	(c)	**95.**	(b)	**96.**	(a)	**97.**	(a)	**98.**	(a)	**99.**	(a)	**100.**	(b)
101.	(c)	**102.**	(d)	**103.**	(a)	**104.**	(d)	**105.**	(a)	**106.**	(a)	**107.**	(b)	**108.**	(a)	**109.**	(b)	**110.**	(c)
111.	(a)	**112.**	(d)	**113.**	(c)	**114.**	(a)	**115.**	(b)	**116.**	(a)	**117.**	(c)	**118.**	(b)	**119.**	(c)	**120.**	(c)
121.	(c)	**122.**	(c)	**123.**	(b)	**124.**	(a)	**125.**	(a)	**126.**	(c)	**127.**	(c)	**128.**	(b)	**129.**	(b)	**130.**	(a)
131.	(a)	**132.**	(b)	**133.**	(a)	**134.**	(a)	**135.**	(b)	**136.**	(b)	**137.**	(a)	**138.**	(a)	**139.**	(c)	**140.**	(b)
141.	(a)	**142.**	(c)	**143.**	(b)	**144.**	(d)	**145.**	(a)	**146.**	(d)	**147.**	(b)	**148.**	(a)	**149.**	(a)	**150.**	(b)

For More Material Join: @JEEAdvanced_2024



Level 2

1.	(c)	**2.**	(c)	**3.**	(c)	**4.**	(a)	**5.**	(d)	**6.**	(c)	**7.**	(c)	**8.**	**i** (c), **ii** (b)	**9.**	(c)	**10.**	(a)
11.	(d)	**12.**	(b)	**13.**	(b)	**14.**	(d)	**15.**	(a)	**16.**	(b)	**17.**	(c)	**18.**	(d)	**19.**	(d)	**20.**	(a)
21.	(a)	**22.**	(d)	**23.**	(c)	**24.**	(c)	**25.**	(d)	**26.**	(b)	**27.**	(a)	**28.**	(d)	**29.**	(b)	**30.**	(b)
31.	(b)	**32.**	(c)	**33.**	(b)	**34.**	(d)	**35.**	(a)	**36.**	(a)	**37.**	(c)	**38.**	(c)	**39.**	(b)	**40.**	(d)



Level 3

Passage–1	**1.** (d)	**2.** (d)							
Passage–2	**1.** (b)	**2.** (c)	**3.** (b)	**4.** (c)					
Passage–3	**1.** (c)	**2.** (a)	**3.** (d)	**4.** (c)	**5.** (d)				
Passage–4	**1.** (a)	**2.** (b)	**3.** (b)						
Passage–5	**1.** (a)	**2.** (b)	**3.** (a)						

One or More Answers is/are Correct

1. (b,c)	**2.** (a,b)	**3.** (c)	**4.** (a,b,c)	**5.** (a,b,d)
6. (b,c,d)	**7.** (b,c,d)	**8.** (a,b,c)	**9.** (c,d)	**10.** (a,c,d)
11. (a,b,c)	**12.** (a,c)	**13.** (a,c,d)	**14.** (b,d)	**15.** (a,b,c)
16. (a,c,d)	**17.** (b,c,d)	**18.** (a,b,c,d)	**19.** (a,b,c)	**20.** (a,b,c)
21. (a,c,d)	**22.** (a,c)	**23.** (a,c)	**24.** (a,c,d)	**25.** (a,c)

Match the Column

1.	A	P;	B	S;	C	R;	D Q
2.	A	P;	B	R;	C	Q;	D S
3.	A	P, S;	B	Q,R;	C	R;	D S
4.	A	R;	B	Q;	C	P;	D S
5.	A	Q;	B	P;	C	R;	D S
6.	A	Q;	B	P, S;	C	P;	D R
7.	A	P, S;	B	R;	C	Q,S;	D Q,S
8.	A	R;	B	Q;	C	Q;	D P, S
9.	A	P;	B	R;	C	Q, S;	D Q,S
10.	A	P;	B	R;	C	Q;	D S

For More Material Join: @JEEAdvanced_2024

11. A S; B P, R; C Q; D P, R
12. A P, R; B Q; C P, R; D Q, S
13. A Q; B P; C R; D S
14. A P, Q; B Q,R, S; C P; D R
15. A S; B R; C P; D Q

Assertion-Reason Type Questions

1. (D) **2.** (A) **3.** (B) **4.** (B) **5.** (A) **6.** (A) **7.** (A) **8.** (B) **9.** (D) **10.** (C)
11. (B) **12.** (C) **13.** (A) **14.** (C) **15.** (A) **16.** (A) **17.** (D) **18.** (D)

Subjective Problems

1.	1	**2.**	7	**3.**	6	**4.**	4	**5.**	4	**6.**	7	**7.**	4	**8.**	5	**9.**	8	**10.**	7
11.	5	**12.**	1	**13.**	6	**14.**	3	**15.**	8										

For More Material Join: @JEEAdvanced_2024

Hints and Solutions

Level 1

1. (d) Heat capacity and volume are extensive properties.

2. (b) Boiling point, pH and emf are intensive property.

3. (d) If system is subjected to chemical equilibrium, mechanical equilibrium and thermal equilibrium then system is consider in thermodynamic equilibrium.

4. (c) Ozone have maximum internal energy

$E \quad \frac{3RT}{2} \quad \frac{3RT}{2} \quad (3N \quad 6)RT$. For non-linear molecule

$E \quad \frac{3RT}{2} \quad \frac{2RT}{2} \quad (3N \quad 5)RT$

For linear molecule.

5. (a) Heat lost by iron + Heat gained by water 0

10 0.45 $(T \quad 373) \quad 25 \quad 4.2 \quad (T \quad 300) \quad 0$

$T \quad 303$ K or 30°C

6. (c) In freezing heat released so $q \quad 0$

7. (a) $w \quad \int_{V_1}^{V_2} PdV$

8. (c) $W \quad n_g RT$ ve

II. $n_g \quad 2$, III. $n_g \quad 2$

9. (d) $w \quad P_{ext}.\ V \quad n_g RT$ and n_g is ve for I and III

also V ve for IV

10. (d) $q \quad 0$, $U \quad q \quad w$

$w \quad 0$ so $U \quad 0$

11. (c) $U \quad q \quad w$

300 400 0.5 $V \quad 100$

$V \quad 2$L

12. (b) $U \quad q \quad w$

10 1000 2 (20) 101.3 5948 J

13. (d) $w \quad PdV \quad 0$

14. (d) $w \quad 4 \int_1^4 dV \quad 4 \quad 3 \quad 100, w \quad 1.2 \quad 10^3$ J

15. (a) $w \quad n_g RT \quad \frac{4.5}{34 \quad 2} \quad 8.314 \quad 298,$

$w \quad 163.9$ J

16. (b) $w \quad P.\ V \quad n_{H_2}$

$RT \quad \frac{2 \quad 8.314 \quad 298}{1000} \quad 4.955$ kJ

17. (b) $w \quad nR(T_2 \quad T_1), w \quad 1R(2) \quad 2R$

18. (a) $dw \quad P.dV$

$w_{rev} \quad 6.V^2 dV$

$6 \ \frac{V_2^3}{3} \quad \frac{V_1^3}{3}$ bar.m^3;

$w \quad 5200$ kJ

so, work done by the gas is 5200 kJ.

19. (b) $w \quad$ Area of P-V curve

20. (c) For cyclic process : $dU \quad 0$

$w \quad q$

Net work done by system Area of the circle

21. (c) Process direction in V–P diagram is clockwise so process direction in P–V diagram is anti-clockwise.

Net work done by system Area

$\frac{(P_2 \quad P_1)}{2} \ \frac{(V_2 \quad V_1)}{2}$

22. (c) Work done in the cyclic process

Area bounded $(ABCA) \quad 5P_1V_1$

23. (c) $w \quad P \ V \quad 0$

$U \quad q \quad w \quad 0$

Temperature remains constant.


24. (d) Zero

Internal energy is state function.


27. (b) $C_P \quad C_V \quad nR \quad 2R$

28. (c) No intramolecular force act between gaseous molecule.

29. (c)



For More Material Join: @JEEAdvanced_2024

30. (d) 

w_3 w_1 w_2 w_4

31. (a) $V \quad \frac{nRT}{P} \quad \frac{100 \quad 0.0821 \quad T}{8.21} \quad T$

$\log V \quad \log T$

32. (c) $dw \quad P.dV$

$w \quad 20 \frac{dV}{V} \quad 20 \ln \frac{V_2}{V_1}$

$w \quad 46.06$ L-atm $\quad 4665.8$ J

$U \quad q \quad w \quad 400 \quad q \quad 4665.8$

$q \quad 5065.8$ J

33. (b) $w \quad nRT \ln \frac{V_2}{V_1}$

$w \quad 2 \quad 8.314 \quad 10^{\ 3} \quad 300 \ln \frac{40}{4},$

$w \quad 11.488$ J

34. (b) $w \quad nRT \ln \frac{P_1}{P_2}$

$10 \quad 8.314 \quad 300 \ln \frac{10}{1}$

57441.42 J

$w \quad mgh$

$m \quad 9.81 \quad 100 \quad 57441.42$

$m \quad 58.55$ kg

35. (c) At constant volume $U \quad q_v \quad nC_{vm} \quad T$
at constant pressure $H \quad q_p \quad nC_{p,m} \quad T$

36. (a) $U \quad w$

$75 \quad 0.1 \quad \frac{3}{2} \quad 2 [T_2 \quad 500]$

$T_2 \quad 250$ K

37. (a) $U \quad w \quad \frac{P_2V_1 \quad P_1V_1}{1}$

38. (a) $T^{\overline{1}} .P$ constant or $P \quad T^{\overline{1}}$

$\because \quad P \quad T^{3}$

$\overline{1} \quad 3; \quad \frac{3}{2}$

39. (b) $T \quad \frac{1}{\sqrt{V}}; \quad TV^{1/2}$ constant

For adiabatic process,

$TV^{\ 1}$ constant

$1 \quad \frac{1}{2}, \quad \frac{3}{2}$

40. (b) $PV \quad$ constant, $P \quad V^{\ 1} \ dV \quad V \quad dP \quad 0$

$\frac{dP}{P} \quad \frac{dV}{V}$

41. (b) $PV \quad$ constant for isothermal process

$PV \quad$ constant for adiabatic process so more is the value of , more is decrease in pressure.

42. (c) $T.V^{\ 1} \quad$ constant

$\frac{T_2}{T_1} \quad \frac{V_1}{V_2}^{\ 1}$

$T_2 \quad 300 \quad \frac{16}{2}^{\frac{5}{3} \ 1}$

1200 K

43. (b) $dw \quad P_{ext} \ dV$

$w_{irr} \quad P_{ext} \quad \frac{nRT}{P_2} \quad \frac{nRT}{P_1}$

$w_{irr} \quad 1 \quad (5 \quad 8.314 \quad 300) \quad 1 \quad \frac{1}{10}$

$w_{irr} \quad 11.224$ kJ

44. (b) For an adiabatic irreversible compression,

$U \quad w$

$nC_{v,\,m} (T_2 \quad T_1) \quad P_{ext} (V_2 \quad V_1)$

here $P_{ext} \quad P_2$

$nC_{v,\,m} (T_2 \quad T_1) \quad P_2 \quad \frac{nRT_2}{P_2} \quad \frac{nRT_1}{P_1}$

$\frac{5}{2} R (T_2 \quad T_1) \quad P_2 \quad R \quad \frac{T_2}{P_2} \quad \frac{T_1}{P_1}$

$P_2 \quad 275$ kPa

45. (b) $w \quad U \quad n \int_{T_1}^{T_2} C_{vm} \, dT$

$w \quad 2 \quad \frac{R}{1} (T_2 \quad T_1)$

$\frac{T_2 \quad T_1}{1} \quad P_{ext} \quad \frac{T_2}{P_2} \quad \frac{T_1}{P_1}$

$\frac{T_2 \quad 300}{\frac{5}{3} \quad 1} \quad 2 \quad \frac{T_2}{2} \quad \frac{300}{1}$

$1.5 T_2 \quad 450 \quad T_2 \quad 600 \quad T_2 \quad 420$ K

For More Material Join: @JEEAdvanced_2024

$w \quad 2 \quad \frac{2}{\frac{5}{3} \quad 1}(420 \quad 300) \quad 720$

46. (c) $\because \quad q \quad 0 \qquad U \quad w$

$$nC_{v,m}(T_2 \quad T_1) \quad P_{ext} \quad \frac{nRT_2}{P_2} \quad \frac{nRT_1}{P_1}$$

$$C_{v,m}[T_2 \quad T_1] \quad P_{ext} \quad R \quad \frac{T_1}{P_1} \quad \frac{T_2}{P_2}$$

$$\frac{5}{2}R[T_2 \quad 300] \quad 2 \quad R \quad \frac{300}{5} \quad \frac{T_2}{2}$$

$$T_2 \quad 248.5 \text{ K}$$

47. (a) $U \quad w$

$$n \quad \frac{R}{1}(T_2 \quad T_1) \quad P_{ext} \quad \frac{nRT_2}{P_2} \quad \frac{nRT_1}{P_1}$$

$$3(T_2 \quad T_1) \quad 1 \quad \frac{T_2}{1} \quad \frac{T_1}{2}$$

$$3T_2 \quad 3T_1 \quad T_2 \quad \frac{T_1}{2}$$

$$4T_2 \quad \frac{7T_1}{2}$$

$$T_2 \quad \frac{7T_1}{8} \quad \frac{7 \quad 400}{8} \quad 350 \text{ K}$$

$$\frac{P_1V_1}{T_1} \quad \frac{P_2V_2}{T_2} \quad \frac{2 \quad 10}{400} \quad \frac{1 \quad V_2}{350} \quad V_2 \quad 17.5 \text{ L}$$

48. (b) Average $C_{v,m} \quad \frac{n_1C_{v,m_1} \quad n_2C_{v,m_2}}{n_1 \quad n_2}$

$$\frac{2 \quad \frac{3}{2}R \quad 2 \quad \frac{5}{2}R}{2 \quad 2} \quad 2R$$

49. (c) $H \quad (n_1C_{p,m_1} \quad n_2C_{p,m_2}) \quad T$

$$0.5 \quad \frac{7}{2}R \quad 0.5 \quad 4R \quad (\ 100)$$

$$375R$$

50. (c) *AB* : Isobaric expansion
BC : Isothermal expansion
CD : Isochoric
DA : Isothermal compression

51. (a) 1 watt 1 J/sec
Total heat supplied for 36 mL H_2O

$$806 \quad 100$$

$$80600 \text{ J}$$

$$H_{vap} \quad \frac{80600}{36} \quad 18$$

$$40300 \text{ J or } 40.3 \text{ kJ/mol}$$

52. (b) $H \quad U \quad n_gRT, \quad n_g \quad 1, \quad H \quad U$

53. (d) $n_g \quad 0 \qquad H \quad U$
For 2 mole $U \quad 370$ kJ

54. (b) $H \quad U \quad n_gRT, \quad n_g \quad 2, \quad H \quad U$
$w \quad n_gRT, \qquad w \quad 0, \quad q \quad \text{ve}$

55. (b) At constant pressure, $H \quad U \quad P \quad V$
$30 \quad 2 \quad 4$
38 atm-L

56. (b) $H \quad U \quad P \quad V$

$$124 \quad U \quad \frac{1520}{760} \quad (177 \quad 377)$$

$$10^{3} \quad 101.3$$

$$U \quad 83.48 \text{ J}$$

57. (b) At constant volume $H \quad U \quad V \quad P$
$560 \quad U \quad 10 \quad (\ 30) \quad 0.1$
$U \quad 530$ kJ

58. (b) When both *P* and *V* are changing
$H \quad U \quad (PV)$
$U \quad (P_2V_2 \quad P_1V_1)$
$H \quad 40 \quad (20 \quad 3)$
57 L-atm

59. (b) For combustion of 1 mole of benzene
$n_g \quad 1.5$
$H \quad U \quad n_gRT$

$$3271 \quad U \quad \frac{1.5 \quad 8.314 \quad 300}{1000}$$

$$U \quad 3267.25 \text{ kJ}$$

For 1.5 mole of combustion of benzene
$U \quad 3267.25 \quad 1.5$
4900.88 kJ

60. (b) $H \quad E \quad n_gRT$

$$82.8 \quad U \quad \frac{1 \quad 8.314 \quad 298}{1000}$$

$$U \quad 80.32$$

61. (a) C (graphite) C (diamond)
$H \quad U \quad P \quad V$

$$V_m \text{ (diamond)} \quad \frac{12}{3} \text{ mL}$$

$$V_m \text{ (graphite)} \quad \frac{12}{2} \text{ mL}$$

$$H \quad U \quad (500 \quad 10^3 \quad 10^5 \text{ N/m}^2)$$

$$\frac{12}{3} \quad \frac{12}{2} \quad 10^{6}$$

$$-100 \text{ kJ/mol}$$

$$U \quad H \quad 100 \text{ kJ/mol}$$

For More Material Join: @JEEAdvanced_2024

62. (c) n_g is +ve

63. (b) n_g ve , S ve

64. (d) n_g ve , S ve

(a) n_g 1 (b) n_g 1

(c) n_g 0 (d) n_g 3

65. (c) S $nC_{p,m} \ln \frac{T_2}{T_1}$

2 $\frac{5}{2} R \ln \frac{600}{300}$

5 R ln 2

68. (c) S_{gas} $nC_{v,m} \ln \frac{T_2}{T_1}$

2 $\frac{5}{2}$ 1 R ln 2

3 R ln 2

69. (d) S $nR \ln \frac{V_2}{V_1}$ 1$R \ln \frac{3V}{V}$ R ln 3

70. (c) S will be same because entropy is a state function.

71. (d) S $nC_{v,m} \ln \frac{T_2}{T_1}$ $nR \ln \frac{V_2}{V_1}$

$C_{v,m}$ ln 2 R ln $\frac{1}{2}$

S $(C_{v,m}$ $R)$ ln 2

72. (c) S $nC_{p,m} \ln \frac{T_2}{T_1}$

2.5 18 4.2 ln $\frac{360}{300}$

34.02 J/K

73. (b) $_rS$ $\nu_p S$ (Products) $\nu_R S$ (Reactants)

(2 27.3 3 69.9) (87.4 3 130.7)

215.2 JK ^{1} mol ^{1}

74. (b) $_rS$ (2 248.2) (2 66.5) (2 91.2) 3(205.1)

168.3 J

75. (c) 266 6 87 205 4x

x 145.75

76. (c) S $\frac{H}{T}$

77. (b) at M.P. S_{fusion} $\frac{H_{fusion}}{T}$

T $\frac{H}{S}$ $\frac{9.95 \quad 1000}{35.7}$ 278.7 K

78. (d) $S_{freezing}$ $\frac{H_{fusion}}{T}$

S $\frac{80 \quad 10}{273}$ 4.18

12.25 J/K

79. (a) S $\frac{H}{T_{B.P.}}$

80. (b) $n_{C_6H_6}$ $\frac{117}{78}$ 1.5


S_{system} 1.5 85 J/K

$S_{surrounding}$ 1.5 85 J/K

81. (b) According to 3rd law of thermodynamics at absolute zero temperature entropy of perfectly crystalline substance is taken to be zero.

$X(s)$ ^{S} $X(l)$

82. (b)



S $\frac{20 \quad 1000}{100}$ 1 $\int_{100}^{200} C_{pm} \frac{dT}{T}$

S 200 10 ln 2 200 10 0.693,

S 206.93 JK ^{1}

83. (c) 0.42 $a(10)^3$ a 0.42 10 ^{3}

S_m $\int_0^{20} \frac{C_{p,m}}{T} dT$

$\int_0^{20} aT^2 dT$ $\frac{a}{3}[20^3$ 0]

1.12 J/K-mol

84. (d) For spontaneous reaction G ve

S ve(n_g ve)

H ve

G H T S

85. (d) H ve (H — H bond formed)

S ve(n_g 1)

86. (b) H ve (Combustion reaction)

S ve(n_g ve)

G H T S ve

87. (b) For $_rG$ ve, $_rH$ should be ve and S should be +ve

88. (a) For spontaneous reaction

$(G_{System})_{T,P}$ ve

89. (b) $(G)_{system}$ H T S

and H T S 0

90. (b) G H T S, H ve , S ve

91. (a) At equilibrium G 0

92. (d) G H T S ve

for spontaneous change H 0, S 0

For More Material Join: @JEEAdvanced_2024

93. (d) G_{799} 40 1000 799 (50),
G_{799} 50 J spontaneous
G_{801} 40 1000 801(50)
G_{801} 50 J non-spontaneous

94. (c) For ideal gas isothermal expansion H 0
G T S
where S $nR \ln \frac{V_2}{V_1}$

95. (b) $Hg(l) \rightleftharpoons Hg(g)$,
$_rS$ 174.4 77.4 97 J/K-mol
$\because$ G H T S 0
T $\frac{H}{S}$
$\frac{60.8 \quad 1000}{97}$ 626.8 K

96. (a) At equilibrium H T S
$H_2O(s)$ $H_2O(l)$
S $S_{H_2O(l)}$ $S_{H_2O(s)}$ = 21.8 J/mol-K
H 273 (21.8) = 5951.4 J/mol

97. (a) G 2 86.6 4 (237.1)
3 0 3(33.6)
2(80.6) 513.2

98. (a) G H T S
G_I 10.5 1000 298 30 1560
G_{II} 1.8 1000 298 113 35474
G_{IV} 126 1000 298 84 ve
III is spontaneous

99. (a) G H T S
S 242.6 50.3 (2 152.3), S 112.3
G 788.6 1000 H 298 112.3
H 788.6 1000 298 112.3
H 822.1 kJ mol ^{1}

100. (b) G H T S
406.6 1000 H 298(2 114.6 130.6)
H 406.6 1000 298(98.6)
H 436 kJ mol ^{1}

101. (c) G $RT \ln K$; K stability of reactant

102. (d) $\frac{}{T} \ln K_p$ $\frac{H}{RT^2}$
$\frac{}{T} \ln K$ $\frac{E_a}{RT^2}$

103. (a) G $RT \ln K$ 8.314 10 ^{3} 400 $\ln 10^5$
G 38.294 kJ

104. (d) At standard state on equilibrium G 0
G 0 $RT \ln K$ K 1

105. (a) At equilibrium G $RT \ln K_{eq}$
where K_{eq} $(P_{H_2O}/P$) and P 1 bar

106. (a) G $RT \ln K_w$
8.314 298 2.303 (14)
~ 80000 J/mol

107. (b) S 206.5 114.6 180.7 88.8
G H T S
H 271.8 298 88.8 10 ^{3}
H 298.3

108. (a) According to Hess's law the standard enthalpy of overall reaction is sum of the enthalpy change in individual reaction.

109. (b) X A q_1
q_2 A B
B Y q_3
q_2 X Y q_1 q_3, H q_2 q_1 q_3

110. (c) The change in enthalpy does not depend upon number of intermediate steps.

111. (a) $_rC_p$ 10.945 for
$H_2(g)$ $\frac{1}{2}O_2(g)$ $H_2O(g)$
According to Kirchhoff's law
$_rH_{T_2}$ $_rH_{T_1}$ $_rC_p[T_2 \quad T_1]$
$_rH_{T_2}$ (241.82) [10.94 J (373 298)/100]
$_rH_{T_2}$ 242.6 kJ/mol

112. (d) $_fH$ stability

113. (c) $(_fH)_{Al(s)}$ 0

114. (a) H $\frac{3}{2}$ H_1 $\frac{H_2}{2}$
713.7 177.5
891.2 kJ

115. (b) H 2 H_1 2 H_2 H_3

116. (a) C(graphite) C(diamond)
H 395 393 2

117. (c) $C(s)$ $\frac{3}{2}H_2(g)$ $\frac{1}{2}Cl_2(g)$ $CH_3Cl(g)$

118. (b) $_rH$ $(_fH)_P$ $(_fH)_R$
$_rH$ 113 3 (269) 46.2 873.8 kJ

119. (c) $C_8H_{18}(g)$ $\frac{25}{2}O_2(g)$ $8CO_2(g)$ $9H_2O(l)$

For More Material Join: @JEEAdvanced_2024

$_rH$ 8 (394) 9 (286) (250)
– 5476 kJ/mol

120. (c) 1941 1273 3(241.8) ($_fH$)B_2H_6
($_fH$)$_{B_2H_6}$ 1273 725.4 1941 57.4

121. (c) $C(s)$ $O_2(g)$ $CO_2(g)$;
$_rH_1$ x kJ/mol ...(1)
$CO(g)$ 1/2 $O_2(g)$ $CO_2(g)$
$_rH_2$ y kJ/mol ...(2)
Equation (1) (2)
$C(s)$ $\frac{1}{2}O_2(g)$ $CO(g)$
$_fH$ $(y \quad x)$ kJ/mol

122. (c) $C_2H_4(g)$ $H_2(g)$ $C_2H_6(g)$
$_rH$ x_2 x_1

123. (b) Q $\frac{5.8}{58}$ $\frac{5756}{2}$ 287.8 kJ

124. (a) Q $\frac{1}{2}$ 438 $\frac{22.3}{223}$ 21.9

125. (a) $_rH$ 57 (393.5) 52 (285.8)
(70870)
33578.9 kJ/mol
$\frac{33578.9}{884}$ 37.98 kJ/mol

126. (c) $_rH$ 46 2 92

127. (c) $2CH_3OH$ $3O_2(g)$ $4H_2O(l)$ $2CO_2(g)$
1453
4 44 $4H_2O(l)$ $4H_2O(g)$
$2CH_3OH$ $3O_2(g)$ $4H_2O(g)$ $2CO_2(g)$
1277
H 1277

128. (b) ($_fH$) $\frac{436.4}{2}$ 218.2 kJ/mol

129. (b) For $H_2(g)$ $O_2(g)$ $H_2O_2(l)$
$_fH$ (H_2O_2, l) $_rH_3$ $\frac{_rH_2}{2}$ $\frac{_rH_1}{2}$

130. (a) $C_2H_5OH(l)$ $3O_2(g)$ $2CO_2(g)$ $3H_2O(l)$
n_g 2 3 1
H U n_gRT
U H RT RT q

131. (a) $q_{reaction}$ q_{bomb} q_{water}
$q_{reaction}$ (C (bomb) $(m_{water}$ $c))$ T
(652 500 4.18) 14.3
39210 J or 39.21 kJ

132. (b) q m c T, m $q/(c$ $T)$
(24 10^6 0.7)/(4.18 50)
80383 g or 80.383 kg

133. (a) m mole of acid 0.05 0.2 0.01
$_rH$ $\frac{480 \quad 1.09}{0.01 \quad 1000}$ 52.32 kJ/mol

134. (a) $H_{solution}$ $\frac{(125 \quad 8) \quad 4.2 \quad 6}{8}$ 80
33516 J/mol or 33.51 kJ/mol

135. (b) (I) $CaCl_2(s)$ aq $CaCl_2(aq)$ q
$CaCl_2(aq)$ $6H_2O(l)$ $CaCl_2$ $6H_2O(s)$
23.2
$CaCl_2(s)$ $6H_2O$ $CaCl_2$ $6H_2O(s)$ q
23.2
3.5 q 23.2
q 19.7
(II) $2Cl(g)$ $Cl_2(g)$
H ve, S ve.

136. (b) $HCl(g)$ aq H $(aq.)$ Cl $(aq.)$;
$_rH$ 75.14
75.14 $_fH$ (H , aq) $_fH$ (Cl , g)
$_fH$ (HCl, g)
$\because$ $_fH$ (H , aq) 0
$_fH$ (Cl , aq) 75.14 92.3
167.44 kJ/mol

137. (a) Given that
$MgSO_4(s)$ nH_2O $MgSO_4(nH_2O)$;
$_rH_1$ 91.2 kJ/mol ...(i)
$MgSO_4$ $7H_2O(s)$ $(n$ $7)H_2O$
$MgSO_4(nH_2O)$
$_rH_2$ 13.8 kJ/mol ...(ii)
Equation (i) (ii)
or H_{hyd} $_rH_1$ $_rH_2$
91.2 kJ/mol 13.8 kJ/mol
105 kJ/mol

138. (a) $12250x$ $13000(1$ $x)$ 12500
$750x$ 500 x 2/3 and y 1/3
So, $\frac{x}{y}$ $\frac{2}{1}$

139. (c) |Enthalpy of change| 57
2H participate in reaction.

140. (b) $H_{neut.}$
$H_{ionization}$ $_rH$ (H OH H_2O)

For More Material Join: @JEEAdvanced_2024

51.34 x 55.84

So, $H_{\text{ionization}}$ or x 4.5 kJ/mol

141.(a) $CaCO_3(s)$ Heat $CaO(s)$ $CO_2(g)$

142. (c) By calorimetry we can determine change in enthalpy.

143. (b) $_rH$ $\frac{5}{8}$ 40 $\frac{3}{8}$ 50 = 43.75 kJ/mol

144. (d) No. of moles of O_2 required to supplied 30 kJ heat to second reaction

$$\frac{30}{1260} \quad \frac{3}{2} \quad \frac{1}{28}$$

So $n_{O_2} : n_{H_2}$ $\frac{1}{28} : 3$ or 1 : 84

145.(a) (O — F) $\frac{368}{2}$ 184

146.(d) C — C bond enthalpy is not given.

147.(b) B.E. size

148. (a) $P(s)$ $\frac{3}{2}Cl_2(g)$ $PCl_3(g)$

306 (314 3 121) [B.E. (P Cl) 3];

B.E. (P—Cl) 123.66 kJ/mol

149.(a) $CH_3OH(g)$ $HCl(g)$ $CH_3Cl(g)$ $H_2O(g)$

$_rH$ $(B.E.)_R$ $(B.E.)_P$

3 (C — H) (C — O) (O — H) (H — Cl)

3 (C — H) (C — Cl)) 2 (O — H)

335 463 431 326 2 463 23 kJ

150. (b) $N_2(g)$ $2H_2(g)$ $N_2H_4(g)$

$_fH$ (N_2H_4, g)

(941 2 436) (159 4 398)

1813 1751 62 kJ mol ^{1}

Level 2

1. (c) w $P_{\text{ext}}(V_f \quad V_i)$

$$10^5 \quad \frac{60 \quad 10^{-3}}{0.60} \quad \frac{40 \quad 10^{-3}}{1000} \quad \frac{100 \quad 10^{-3}}{1000}$$

$10^5 (100 \quad 10^{-3} \quad 0.04 \quad 10^{-3} \quad 0.1 \quad 10^{-3})$

$|w|$ 9994 J

2. (c) Mole of the gas in the first compartment

$$n_1 \quad \frac{P_1V_1}{RT_1} \quad \frac{0.8314 \quad 10^6 \quad 2}{8.314 \quad 400} \quad 500$$

Similarly, n_2 2000

The tank is rigid and insulated hence w 0 and q 0 therefore U 0

Let T_f and P_f denote the final temperature and pressure respectively

U $n_1C_{V,m}[T_f \quad T_1]$ $n_2C_{V,m}[T_f \quad T_2]$

0

500 $(T_f \quad 400)$ 2000 $(T_f \quad 500)$ 0

T_f 480 K

3. (c) H_2O $(l, 323\text{ K})$ $\xrightarrow{U_1}$ H_2O $(l, 373\text{ K})$

$\downarrow U_2$

H_2O (g,323 K) $\xrightarrow{U_3}$ H_2O (g,373 K)

$C_{V,m}(H_2O, g)$ 33.314 8.314

= 25 J/K mol

U_2 H_2 n_gRT 37.6

U_{total} U_1 U_2 U_3

$C_{v,m}(l)$ T $V_{\text{vap.}}$ $C_{v,m}(g)$ T

$$\frac{75 \quad 50}{1000} \quad 37.6 \quad \frac{25 \quad 50}{1000}$$

42.6 kJ/mol

4. (a) $\because$ $\frac{C_{p,m}}{C_{V,m}}$ and $C_{p,m} \quad C_{V,m} \quad R$

$C_{V,m}$ $\frac{R}{1}$

$C_{V,m}$ $\frac{C_V}{n}$ and C_V m c_V

$\frac{R}{1}$ $\frac{m \quad c_V}{m}$ M

c_V $\frac{R}{(\quad 1)M}$

5. (d) Av. $C_{V,m}$ $\frac{n_1C_{V,m_1} \quad n_2C_{V,m_2}}{n_1 \quad n_2}$ $2R$

for adiabatic process dU dW

$$\frac{dT}{T} \quad \frac{R}{C_{V,m}} \quad \frac{dV}{V}$$

$n_1C_{V,m_1}\,dT$ $n_2C_{V,m_2}\,dT$

$(n_1RT \quad n_2RT)$ $\frac{dV}{V}$

$\ln \frac{T_2}{T_1}$ $\frac{1}{2} \ln \frac{V_2}{V_1}$

T_2 320 $\left(\frac{1}{4}\right)^{1/2}$

= 160 K

U $(n_1C_{V,m_1} \quad n_2C_{V,m_2})$ T 960 R

For More Material Join: @JEEAdvanced_2024

6. (c) 2 $C_{V,m}(T_2 \quad T_1) \quad 1 \quad \frac{2RT_2}{1} \quad \frac{2RT_1}{10}$

$$\frac{4}{3} \quad \frac{C_{p,m}}{C_{V,m}}$$

and $C_{p,m} \quad C_{V,m} \quad R \qquad C_{V,m} \quad 3R;$

$$3R(T_2 \quad 300) \quad RT_2 \quad \frac{R \quad 300}{10}$$

$T_2 \quad 232.5$ K

$w \quad U = 2 \quad 3R(300 \quad 232.5)$

$w \quad 405R$

7. (c) Max. temp. attained by gas in between B to C

According to equation of straight line

$$\frac{P \quad 4}{1 \quad 4} \quad \frac{V \quad 1}{2 \quad 1}$$

$$P \quad 4 \quad 3V \quad 3$$

$$P \quad 7 \quad 3V$$

For 1 mole gas

$$\frac{RT}{V} \quad 7 \quad 3V; \; RT \quad 7V \quad 3V^2 \qquad \ldots(1)$$

$$R\frac{dT}{dV} \quad 7 \quad 6V \quad 0$$

$V \quad \frac{7}{6}$ put in Eq. (1)

$$RT \quad 7 \quad 3 \quad \frac{7}{6} \quad \frac{7}{6}$$

$$T \quad \frac{49}{12R}$$

8. (i)(c) $w \quad P \quad V \quad nR \quad T \quad 2 \quad 8.314 \quad 600$

9.97 kJ

(ii)(b) $H_{\text{total}} \quad H_{AB} \quad H_{BC} \quad nC_{p,m} \quad T \quad 0$

$$2 \quad \frac{7}{2} \quad R \quad (800 \quad 200)$$

$$4200R$$

9. (c) $\frac{T_2}{T_1} \quad \frac{V_1}{V_2}^{1}$

$$T_2 \quad T_1 \quad \frac{1}{32}^{\frac{7}{5} \quad 1} \quad 600 \quad \frac{1}{2^5}^{\frac{2}{5}}$$

$$600(0.5)^2 \quad 150 \text{ K}$$

$$H_m \quad \frac{7}{2}R \quad (150 \quad 600) \quad 1575R$$

10. (a) $w \quad nR \quad T \quad 2 \quad 8.314 \quad 100$

1662.8 J

$$U \quad n \quad C_{v,m}\,dT$$

$$2 \quad (20 \quad 10^{\;2}T)\,dT$$

$$2 \quad 20 \quad (T_2 \quad T_1) \quad 2 \quad 10^{\;2} \quad \frac{(T_2^2 \quad T_1^2)}{2}$$

4700 J

4700 q 1662.8

q 6362.8 J

11. (d) $C_{p,m} \quad (21.686 \quad 8.314 \quad 10^{\;3}T)$

$30 \quad 10^{\;3}T$

$$H \quad nC_{p,m} \quad T$$

$$10 \quad (30 \quad 10^{\;3}T)\,dT$$

$$H \quad 10 \quad 30(T_2 \quad T_1) \quad \frac{10^{\;3}}{2} \quad [T_2^2 \quad T_1^2] \quad 10$$

30350 J

12. (b) $dU \quad dq \quad dw$

$$nC_{v,m}\,dT \quad nC_m\,dT \quad P\,dV$$

$$C_m \quad C_{v,m} \quad \frac{P\,dV}{n\,dT} \qquad \ldots(1)$$

$PV^n \quad K$ and $PV \quad nRT$

$KV^{1 \quad n} \quad nRT$

$$K(1 \quad n)V^{\;n}\,dV \quad nRdT$$

$$\frac{dV}{dT} \quad \frac{nR}{K(1 \quad n)V^{\;n}} \qquad \ldots(2)$$

from Eqs. (1) and (2)

$$C_m \quad C_{v,m} \quad \frac{R}{(1 \quad n)}$$

13. (b) $P_2V_2^2 \quad P_1V_1^2$

$$\frac{P_2}{P_1} \quad \frac{V_1}{V_2}^{\;2}$$

Now, $\frac{T_2}{T_1} \quad \frac{P_2V_2}{P_1V_1} \quad \frac{V_1}{V_2} \quad \frac{1}{3}$

$$T_2 \quad \frac{300}{3} \quad 100 \text{ K}$$

$$H \quad nC_{p,m} \quad T$$

$$2 \quad \frac{5}{2}R \quad (\ 200 \text{ K}) = 1000 \text{ R}$$

14. (d) $S \quad nC_{p,m} \ln \frac{T_2}{T_1} \quad nR \ln \frac{P_1}{P_2}$

$$3 \quad \frac{7}{2}R \ln \frac{596}{298} \quad 3R \ln \frac{1}{4}$$

6.3 cal K ^{1}

15. (a) For isoentropic process $S_{\text{system}} \quad 0$

$$nC_{p,m} \ln \frac{T_2}{T_1} \quad nR \ln \frac{P_1}{P_2} \quad 0$$

$$\ln(P_2) \quad \frac{5}{2} \quad \ln \frac{600}{300}$$

1.75 atm

16. (b) $S_{\text{system}} \quad nR \ln \frac{V_2}{V_1}$

$$2 \quad R \quad \ln 2$$

For More Material Join: @JEEAdvanced_2024

11.52 J/K

$S_{surrounding}$ $\frac{3.41 \quad 1000}{310}$

11 J/K

S_{total} 11.52 11

0.52 J/K

17. (c) 0.40 aT_1^3 bT_1

0.40 a (1000) b 10

0.4 1000a 10b ...(1)

0.92 aT_2^3 bT_2

0.92 a 8000 20b ...(2)

from Eqs. (1) and (2)

a 2 10^{-5}, b 0.038

S_m $\frac{aT^3 \quad bT}{T}\,dT$

$\frac{a[T_2^3 \quad T_1^3]}{3}$ $b[T_2 \quad T_1]$

0.813 J/K-mol

19. (d) No. of moles of sucrose $\frac{34.2}{342}$ 0.1

$(\ G)_{T,P}$ useful work done by the system

G H T S

(6000 0.1) $\frac{180 \quad 0.1 \quad 300}{1000}$

605.4 kJ

20. (a) G_{200} H_{200} T S_{200}

H_{200} 20 4 16 kJ/mol

H_{T_2} H_{T_1} $C_P[T_2 \quad T_1]$

H_{400} H_{200} $\frac{20 \quad 200}{1000}$ kJ/mol

16 4 20 kJ/mol

21. (a) $_fS$ (NH_4Cl, s) at 300 K

$S_{NH_4Cl(s)}$ $\frac{1}{2}S_{N_2}$ $2S_{H_2}$ $\frac{1}{2}S_{Cl_2}$

374 JK $^{-1}$ mol $^{-1}$

$\because$ $_rC_P$ 0

$_fS_{310}$ $_fS_{300}$

374 JK $^{-1}$ mol $^{-1}$

$_fH_{310}$ $_fH_{300}$ 314.5

$_fG_{310}$ $_fH$ 310 S

314.5 $\frac{310(\ 374)}{1000}$

198.56 kJ/mol

22. (d) $_rH$ 3 $_fH$ (CO, s) 4 $_fH$ (CO_2, g)

$_fH$ (Co_3O_4, s) 4 $_fH$ (CO, g)

$_rH$ 241 kJ/mol

$_rS$ 3 30 4 213.7 102.5

4 197.7

51.5 J/K-mol

$_rG$ $_rH$ T $_rS$

241 $\frac{300 \quad 51.5}{1000}$

256.45 kJ/mol

23. (c) At constant volume, $\frac{P_1}{T_1}$ $\frac{P_2}{T_2}$

P_2 1 $\frac{300}{200}$ $\frac{3}{2}$

and V_1 24.63 L

for single phase

$\because$ dG Vdp $S\,dT$

G V P $(2 \quad 10^{-2}T)\,dT$

1231.5 200 $\frac{10^{-2} \quad 50{,}000}{2}$

781.5 J

24. (c) $\because$ $_rC_p$ 0, H_{300} H_{310}

25. (d) H 3 H_1 H_2 2 H_3 3 H_4

747.4 kJ

H U $n_g RT$; where n_g 8

747.4 U $\frac{8 \quad 8.314 \quad 300}{1000}$

U 727.44 kJ

26. (b) $_rG$ $_rH$ T $_rS$

$_rS$ 2 81 4 24 3 205 J/mol

$_rH$ 2258.1 kJ/mol

$_rH$ 2 $_fH$ (Cr_2O_3, s)

$_fH$ (Cr_2O_3, s) $\frac{2258.1}{2}$

1129.05 kJ/mol

27. (a) $CaO(s)$ $CO_2(g)$ $CaCO_3(s)$

$_fH$ $H_f(CaCO_3)$ $H_f(CaO)$

$H_f(CO_2)$

1207 (635) (394)

178 kJ/mol

U H $n_g RT$

U 178 $\frac{(\ 1) \quad 8.3 \quad 300}{1000}$

175.51 kJ

For More Material Join: @JEEAdvanced_2024

n_{CaO} $\frac{224}{56}$ 4

q_v n $_rU$ 4 (175.51)

702.04 kJ

28. (d) $H_2C_2O_4(l)$ $\frac{1}{2}O_2(g)$ $H_2O(l)$ $2CO_2(g)$;

n_g 3/2

U_c $-\frac{0.312 \quad 8.75}{1}$ 90

245.7 kJ/mol

H U n_gRT

245.7 $\frac{3}{2}$ $\frac{8.314 \quad 300}{1000}$

241.95 kJ/mol

29. (b) H_3PO_3 $2H^+$ HPO_3^{2-} ; $_rH$?

2H 2OH $2H_2O$;

$_rH$ 55.84 2 111.68

106.68 $_{ion}H$ 55.84 2

$_{ion}H$ 5 kJ/mol

30. (b) HA H A^- ; $_rH$ 1.4 kJ/mol

$H_{neutralization}$ $H_{ionization}$ $_rH$

(H OH H_2O)

55.95 $H_{ionization}$ 57.3

$H_{ionization}$ for 1 M HA 1.35 kJ/mol

% heat utilized by 1 M acid for ionization

$\frac{1.35}{1.4}$ 100 96.43%

so, acid is 100 96.43 3.57% ionized

31. (b) 2 C(s) $3H_2(g)$ $C_2H_6(g)$

$_fH$ 2 $_{sub}$ H (C, s)

3 B.E. (H H)

B.E. (C C)

+ 6 B.E. (C H)

85 2 718 3 436 (x 6y)

x 6y 2829 ...(1)

Similarly, for $C_3H_8(g)$

2x 8y 4002 ...(2)

from Eqs. (1) and (2)

x 345 kJ/mol; y 414 kJ/mol

32. (c) $\frac{1}{2}N_2(g)$ $\frac{3}{2}H_2(g)$ $NH_3(g)$

Let B.E. of N N is x

46 $\frac{x}{2}$ $\frac{3}{2}$ 436 3 393 x 958

$N_2H_4(l)$ $N_2(g)$ $2H_2(g)$;

$_rH$ 50 kJ/mol

$_{vap}H$ (N_2H_4, l)

$_rH$ 4 B.E. (N H)

B.E. (N N)

B.E. (N N)

2 B.E. (H H)

50 (18 4 393 y) (958 2 436)

50 (1590 y) (1830)

B.E. (N N) or y 190 kJ/mol


33. (b) $_rH$ [Heat supplied] [Heat evolved]

292 [4x 279] [38 85]

x 34 kcal/mol

34. (d) Theoretical heat of hydrogenation of benzene (Actual heat of hydrogenation)

+ (Resonance energy)



205 152 357

Enthalpy of hydrogenation of 

$\frac{357}{3}$ = **119 kJ/mol**

35. (a) $C_3H_8(g)$ 5 $O_2(g)$ 3 $CO_2(g)$ $4H_2O(l)$

8 B.E. (C H)

$_cH$ 2 B.E. (C C)

5 B.E. (O O)


6 B.E. (C O)

8 B.E. (O H)

3 |R.E.| of CO_2

4 $_{vap}H$ (H_2O)

36. (a)



For More Material Join: @JEEAdvanced_2024

AB process

0 q_1 w_1

q_1 $nR(2T_0)\ln 2$

BC process U_2 q_2 w_2

$\frac{nR}{(\ \ 1)}(T_0 \ \ 2T_0)$ q_2 $\frac{nRT_0}{V_0}$ $(V_0 \ \ 2V_0)$

CA process


U_3 q_3 w_3

$\frac{nR}{(\ \ 1)}(2T_0 \ \ T_0)$ q_3 0

Efficiency $\frac{\text{Total work done}}{\text{Total heat absorbed}}$ $\frac{w_1 \ \ w_2}{q_1 \ \ q_3}$

$\frac{(\ \ 2RT_0\ln 2) \ \ (nRT_0)}{(2RT_0\ln 2) \ \ \frac{nRT_0}{1}}$

37. (c) V_1 24.63 L T_1 300 K ; T_2 600 K



For path (II)

$T_1V_1^{\ 1}$ $T_2V_2^{\ 1}$

$\frac{T_1}{T_2}$ $\frac{V_2}{V_1}^{\ 1}$

$\frac{600}{300}$ $\frac{V_2}{V_1}^{\frac{7}{5}\ 1}$

$(2)^{5/2}$ $\frac{V_2}{V_1}$

V_2 $4\sqrt{2}$ 24.63 139.3 L 139 L

38. (c)

$_rH$ $_rH$ (H_2O, l) $_rH$ (OH^-, aq) 229 56 285 $_rH$ (OH^-, aq);

$_rH$ (OH^-, aq) 229

$HF(aq)$ $OH^-(aq)$ $F^-(aq)$ $H_2O(l)$

ΔH_{neut} (320 285) (329 229)

47 kJ/mol

39. (b) $I_{2(s)}$ $I_2(g)$

H_2 H_1 $C_P(T_2 \ \ T_1)$

C_P $C_P(I_{2,g})$ $C_P(I_{2,s})$ 6.2 cal/mole

H_2 6096 6.2(50) 5786 cal/mole

H U 1 2 523

5786 U 1 2 523

U 5786 1046 4740 cal

40. (d) G $RT\ln K_p$; K_p $(2x)^2X$ $4X^3$

G $RT\ln(4X^3)$

G $RT\ln 4$ $3RT\ln X$

Level 3

Passage-1

1. (d) U q w

U 300 (50 150) 200

2. (d) U 300 (50 150) 500

Passage-2

1. (b) PV nRT 2 8 2 0.080 T

T 100 K

w_{rev} 2.303 n R $T\log\frac{P_1}{P_2}$

2.303 2 0.08 100 $\log\frac{1}{10}$

= 36.848 bar-L

2. (c) w_{irr} $P_{ext}(V_2 \ \ V_1)$

20 $\frac{nRT}{P_2}$ $\frac{nRT}{P_1}$

= 144 bar-L

3. (b) $w_{irr\ (total)}$ w_1 w_2

10 $\frac{nRT}{10}$ $\frac{nRT}{2}$

20 $\frac{nRT}{20}$ $\frac{nRT}{10}$

5 nRT 80 bar-L

4. (c) In isothermal compression is

$w_{One\text{-}stage}$ $w_{Two\text{-}stage}$ $w_{Reversible}$

Passage-3

1. (c) $_rS$ 240 198 29 2 16

2. (a) $_rH$ 201 114 87

3. (d) $_rS$ 16 $\int_{300}^{320} {}_rC_p\frac{dT}{T}$

$_rS$ 16 (44 29.4 2 28.8)$\ln\frac{320}{300}$

$_rS$ 16 (43)$\ln\frac{320}{300}$ 18.78

4. (c) $_rH$ 87 $\int_{300}^{320} {}_rC_P\,dT$

For More Material Join: @JEEAdvanced_2024

87 C 4.31 70 10^{3}

87.86 kJ/mol

5. (d) $_rS$ S_{CH_3OH} S_{CO} 2 S_{H_2} – J/K-mol

$_rH$ $_fH$ (CH_3OH) $_fH$ (CO)

2 $_fH$ (H_2)

87 kJ/mol

$_rS_{320}$ $_rS_{300}$ $_rC_p$ [T_2 T_1]

where $_rC_p$ 44 29.4 2 28.8

43 J/K-mol

$$_rS_{320} \quad 16 \quad (\ 43)\ln\frac{320}{320}$$

18.58

$_rH_{320}$ $_rH_{300}$ $_rC_p$ [T_2 T_1]

$$87 \quad \frac{(\ 43)\ 20}{1000}$$

87.86 kJ/mol

$_rG_{320}$ $_rH_{320}$ T $_rS_{320}$

$$87.86 \quad \frac{320\ (\ 18.58)}{1000}$$

81.91 kJ/mol

Passage-4

1. (a) 49.86 ($_{Ion}H)_{CH_3COOH}$ 55.84

($_{Ion}H)_{CH_3COOH}$ 5.98 kJ mol ^{1}

2. (b) $_rH$ 2 (55.84) 111.68 kJ

3. (b) 2NaOH H_2A Na_2A $2H_2O$

$$\frac{0.1\,V_1}{1000} \qquad \frac{0.05\,V_2}{1000}$$

V_1 V_2 100 ...(1)

$$2\ \frac{0.05\,V_2}{1000} \quad \frac{0.1\,V_1}{1000}$$

V_1 V_2

V_1 V_2 50 mL

Passage-5

1. (a) ($G)_{PT}$ 2000 (20 298)

3960 J/mol

2. (b) $CaCO_3$ CaO CO_2 H ve

Reaction becomes spontaneous at high temperature because T S dominates over H_{rxn}.

3. (a)

$CaCl_2(s)$ aq $6H_2O$ $CaCl_2(aq)$ x

$CaCl_2(aq)$ $CaCl_2$ $6H_2O(s)$ aq 3.5

$CaCl_2(s)$ $6H_2O$ $CaCl_2$ $6H_2O(s)$

x 3.5

x 3.5 23.2

x 19.7

H x 19.7

enthalpy driven

One or More Answers is/are Correct

1. (b,c)

Intensive properties : Temperature, pressure.

2. (a, b)

Extensive properties : Gibb's energy, Entropy.

3. (c) State function : Enthalpy

4. (a, b, c)

Work done by system is negative.

Work done by surrounding is positive.

Heat absorbed by system is positive.

Heat absorbed by surrounding is negative.

5. (a, b, d)

U n $C_{vm}dT$ H $C_{pm}dT$

U q w

6. (b,c,d) U 0 U_2 U_1 0

Similarly, H 0

$$q \quad w \quad nRT\ln\frac{V_2}{V_1}$$

7. (b, c, d)

No heat transfer

U q w 0 w w

U w

$$T_2 \quad T_1 \quad \frac{w}{nC_{vm}}$$

w ve

T_2 T_1

8. (a, b, c)

U q w 0 w w

w n $C_{vm}dT$

$$w \quad \frac{nR}{1}(T_2 \quad T_1)$$

$$w \quad nR\,P_{ext}\ \frac{T_2}{P_2}\ \frac{T_1}{P_1}$$

For More Material Join: @JEEAdvanced_2024

9. (c,d)

At V_A $\frac{1 \quad R \quad 100}{1}$ $100\ R$

V_B $\frac{1 \quad R \quad 600}{3}$ $\mathbf{200\ R}$


$\because V_B \quad V_A$ so expansion of gas takes place

$V_B \quad 200 \quad 0.0821 \quad 16.42$ L

10. (a,c,d)

First step is adiabatic ($q \quad 0$) so $U_1 \quad w_1$

Second step is isochoric ($w \quad 0$)



So, $U_2 \quad q_2$

$\because$ initial and final temp. are same

$U_{total} \quad U_1 \quad U_2 \quad 0$

or $w_1 \quad q_2 \quad 0$

Max. work done by the gas, SO_2 is (area) under the curve

so, SO_2 absorbed

$\because$ $SO_2 \quad CO_2 \quad O_2 \quad Ne$

so, max. decrease in temp. of Ne due to step 1.

11. (a,b,c)

$H \quad U \quad P \quad V \quad V \quad P \quad P \quad V$

is correct relation.

12. (a, c)

$H \quad U \quad n_g RT$

(a) $n_g \quad 2 \quad U \quad H$

(b) $n_g \quad \frac{1}{2} \quad U \quad H$

(c) $n_g \quad \frac{1}{2} \quad U \quad H$

(d) $n_g \quad 0 \quad U \quad H$

13. (a, c, d)

$G \quad H \quad T \quad S$

(a) $H \quad$ ve, $S \quad$ ve at low temperature

$G \quad$ ve

(b) $H \quad$ ve, $S \quad$ ve ; $G \quad$ ve

at any temperature.

(c) $H \quad$ ve , $S \quad$ ve, $G \quad$ ve at any temperature

(d) $H \quad$ ve , $S \quad$ ve, $G \quad$ ve at high temperature


14. (b, d)

For spontaneous process

$S_{System} \quad S_{Surrounding} \quad 0$

$(\ G_{System})_{T,P} \quad 0$

15. (a, b, c)



$x(l) \rightleftharpoons x(g) \quad T_b^0 \quad 400$ K

(a) At 400 K and 1 atm pressure

$G \quad 0$

(b) at 2 atm and 400 K, $G \quad$ ve

(c) at 400 K and 0.1 atm pressure, $G \quad$ ve

(d) at 410 K and 1 atm pressure, $G \quad$ ve

16. (a, c, d)

Entropy intermolecular force

$H \quad 0$

Spontaneous process $G \quad 0$

17. (b, c, d)

C(graphite) C(diamond)

$G \quad H \quad T \quad S$

$1.9 \quad 298(\ ve) \quad ve$

Graphite is more stable than diamond thermodynamically

$_rH \quad (\ _CH)_G \quad (\ _CH)_D \ ,(\ _CH)_G \quad x \quad (\ _CH)_D \quad y$

$1.9 \quad x \quad y \quad y \quad 1.9 \quad x$

Diamond provides more heat on complete combustion in comparison of graphite.

18. (a, b, c, d)

(a) Reversible adiabatic process is iso-entropic process

(b) When $(\ G_{System})_{T,P} \quad 0$ then process may be endothermic / exothermic

(c) $dG \quad VdP \quad SdT$, only PV work consider

(d) $S \quad \frac{9}{18} \quad \frac{40.6 \quad 1000}{373} \quad 54.42$

19. (a, b, c)

(a) $E \quad q \quad W \quad 0 \quad 0 \quad 0$

(b) $(\ _FH \ )(S,$ rhombic$) \quad 0$

For More Material Join: @JEEAdvanced_2024

(c) (C — C) (C — H) 2812

(C — C) 2812 6 $\frac{1656}{4}$ 328

(d) $H_2O(l)$ $H_2O(g)$

44 $(\ _fH)_{H_2O(l)}$ 242

$(\ _fH)_{H_2O(l)}$ 286

$H\ (aq)$ $OH\ (aq)$ $H_2O(l)$


57.3 286 4$(\ _fH)_{OH}\ (aq)$

$(\ _fH)_{OH}$ 228.7


20. (a, b, c)

(a) $_rH$ 15 0 57.3 42.3 kJ

(b)



(c)



$_rH$ 70.7 15 55.7

(d) $B\ (aq)$ $OH\ (aq)$ $BOH(aq)$

$_rH$ 0

21. (a,c,d)

Work is a form of energy exchange between system and surrounding in adiabatic process.

Intensive property is not additive.

In cyclic process work may be zero.

For cyclic process

dU 0 dq dw dq Pdv 0

22. (a,c,)

Entropy of isolated system increases or remains constant but it can never decreases.

23. (a,c)

S_{vap} $\frac{H}{B.P.}$ $\frac{40\ \ 1000}{400}$ 100

(a) P 2 atm B.P. 400 K

P B.P S_{vap} 100

(c) G H T S

40 1000 410 100 ve

G 0

(d) H U n_gRT

O U 1 8.31 $10^{\ 3}$ 400

U 36.676 kJ/mol

24. (a, c, d)

(a) In isolated system no change possible

(b) U q w 0 q w

w may be zero so q may be zero.

(c) Entropy of a closed system is not maximum at equilibrium.

(d) Intensive properties.

25. (a, c)

H	ve	Enthalpy driven
S	ve	Entropy driven
G	ve	Spontaneous process
G	ve	Non spontaneous process

Subjective Problems

1. For reversible adiabatic process

P_1^1 T_1 P_2^1 T_2

$\frac{T_1}{T_2}$ $\frac{P_2}{P_1}^{\frac{1}{}}$; $\frac{300}{90}$ $\frac{10}{200}^{\frac{1}{}}$;

$\frac{10}{3}$ $\frac{1}{20}^{\frac{1}{}}$; $\log\frac{10}{3}$ $\frac{1}{x}$ $\log\frac{1}{20}$

$\frac{1}{}$ $\frac{0.5229}{1.301}$ 0.4

$\frac{1}{}$ 0.6 1.66

Atomicity 1

2. r $\frac{nRT\ln\frac{12}{24}}{P_{ext}nRT\ \frac{1}{P_2}\ \frac{1}{P_1}}$

r $\frac{0.7}{24\ \frac{1}{24}\ \frac{1}{12}}$ $\frac{0.7\ \ 24}{24(1)}$

$10r$ 10 0.7 7

3. C_m C_{VM} $\frac{R}{1\ \ x}$ C_{VM} $\frac{R}{1\ \ 3}$

C_m $\frac{3R}{2}$ $\frac{2R}{2}$ R $\frac{R}{2}$ $3R$ 6

4. $\frac{w}{10}$ 1 $\frac{300}{500}$ w 4 kcal

5. $\frac{|w|}{8}$ ...(i)

1 $\frac{T_1}{T_2}$

For More Material Join: @JEEAdvanced_2024

T_2 2 ^{1} T_1 16 ^{1} $\frac{T_1}{T_2}$ $\frac{1}{2}$

$\frac{|w|}{8}$ 1 $\frac{1}{2}$ $\frac{1}{2}$ $|w|$ $\frac{1}{2}$ 8 4

6. $\frac{1}{2}A$ B $\frac{3}{2}C$

G H T S

(10 1000 300 10) 10^{3}

G 7 kJ/mol

7. Energy available $\frac{25}{100}$ 2880 $\frac{1}{2}$ 360

Minimum distance $\frac{1}{90}$ 360 4 km

8. $C_2H_2(g)$ $H_2(g)$ $C_2H_4(g)$

175 $({}_{f}H)_{C_2H_2}$ 50

$({}_{f}H)_{C_2H_2}$ 225

C_2H_2 $\frac{5}{2}O_2$ $2CO_2$ H_2O

H 2 (390) 280 225 1285

$\left|\frac{H}{257}\right|$ $\left|\frac{1285}{257}\right|$ 5

9. $5H_2O$ $CuSO_4(s)$ aq $CuSO_4(aq)$ 70

$CuSO_4(aq)$ $CuSO_4$ $5H_2O(s)$ aq 10

$CuSO_4(s)$ $5H_2O$ $CuSO_4$ $5H_2O$ 80


$|H|$ 80

For 0.1 mol : 80 0.1 mol : 80 0.1 8 kJ

10. 50 $({}_{Ion}H)_{NH_4OH}$ 57

$({}_{Ion}H)_{NH_4OH}$ 7 kJ

11.



$_{Sol.}H$ 790 785 5

12. Q_v 1.25 4 5

Q_v 5 $\frac{x}{80}$ 400

x 1

13. $\frac{T_2\ T_1}{T_2}$ $\frac{|w_{Total}|}{q_2}$ $\frac{500\ 350}{500}$ $\frac{|w_{Total}|}{10}$

Work done in one cycle 3

Work done in two cycles 3 2 6

14. W n_gRT $\frac{(3)\ 2\ 500}{1000}$ 3

15. Process AC polytropic process (P KV)

Molar Heat capacity c_m c_V $R/2$ $2R$

Process AB Isobaric

c_m c_p $5R/2$

$$\frac{q_{AC}}{q_{AB}} \frac{\int_{T_C}^{T_A} nC_m\,dT}{\int_{T_B}^{T_A} n\,C_{p,m}.dT} \frac{2R}{\frac{5}{2}R} 0.8 :$$

$\frac{q_{AC}}{q_{AB}}$ 10 0.8 10 8

For More Material Join: @JEEAdvanced_2024



5

CHEMICAL EQUILIBRIUM

Types of Reactions

	Reversible	Irreversible
(i)	A reaction in which not only the reactants react to form products but also the products react to form reactants under the same conditions is called reversible reaction.	A reaction which cannot take place in the reverse direction is called an irreversible reaction
(ii)	A few examples are given below $CaCO_3(s) \rightleftharpoons CaO(s)\ \ CO_2(g)$ $N_2O_4(g) \rightleftharpoons 2NO_2(g)$	A few examples are given below $AgNO_3(aq)\ \ NaCl(aq)\ \ \ AgCl(s)\ \ NaNO_3(aq)$ $BaCl_2(aq)\ \ Na_2SO_4(aq)\ \ \ BaSO_4(s)\ \ 2NaCl(aq)$

Characteristics of Equilibrium State

- A reaction (or a process) is said to be in equilibrium when the rate of forward reaction (process) becomes equal to the rate of backward reaction (process).
- At equilibrium, the concentration of each of the reactants and products become constant.
- Chemical equilibrium is dynamic in nature and not static *i.e.*, even after equilibrium is attained, the forward as well as the backward reaction take place but at equal speeds.
- A chemical equilibrium can be established only if none of the products is allowed to escape out.
- Chemical equilibrium can be attained from either direction.
- When reaction attains equilibrium at certain temperature and pressure, $_rG \quad 0$
- Catalyst increases rate of reaction in forward as well as backward direction upto same extent.

For More Material Join: @JEEAdvanced_2024

Law of Mass Action

It was put forward by Guldberg and Waage. It sates that the rate at which a substance reacts is directly proportional to its active mass raised with suitable powers and hence the rate at which substances react together is directly proportional to the product of their active masses. Active mass means molar concentration.

Law of Chemical Equilibrium

For the reaction $aA(aq)$ $bB(aq) \rightleftharpoons xX(aq)$ $yY(aq)$

$$K_c \quad \frac{[X]^x[Y]^y}{[A]^a[B]^b}$$

Where K_c is called equilibrium constant at constant temperature.

Types of Equilibrium Constants

K_c Equilibrium constant in terms of molar concentration $\frac{[X]^x[Y]^y}{[A]^a[B]^b}$

K_p Equilibrium constant in terms of pressure $\frac{p_X^x \cdot p_Y^y}{p_A^a \cdot p_B^b}$

K_x Equilibrium constant in terms of mole fraction $\frac{X_x)^x(X_y)^y}{(X_A)^a(X_B)^b}$

Unit of Equilibrium Constant

Unit of K_p (atm) ^{n_g} (where n_g change in gas mole of reaction)

Unit of K_c M ^{n_g}

Unit of K_x Unitless

Relationship between K_p and K_c: K_p $K_c(RT)$ ^{n_g}

where R universal gas constant, T temperature

Relationship between K_p and K_x: K_p $K_x(P)$ ^{n_g}

where P equilibrium pressure in container.

Types of Chemical Equilibria

- Homogeneous equilibrium. When in an equilibrium reaction, all the reactants and products are present in the same phase (*i.e.* gaseous or liquid), it is called homogeneous equilibrium.

$PCl_5(g) \rightleftharpoons PCl_3(g)$ $Cl_2(g)$; $CH_3COOH(l)$ $C_2H_5OH(l) \rightleftharpoons CH_3COOC_2H_3(l)$ $H_2O(l)$ etc.

- Heterogeneous equilibrium. When in an equilibrium reaction, the reactants and the products are present in two or more than two phases, it is called a heterogeneous equilibrium.

$PbCl_2(s) \rightleftharpoons Pb^2$ (aq) $2Cl$ (aq) ; $CaO(s)$ $CO_2(g) \rightleftharpoons CaCO_3(s)$ etc.

Expression of *K* for Equilibrium Reaction

- **Homogeneous equilibria:** If reactants and products are present in same phase.

Example: (i) $NH_3(aq)$ $H_2O(l) \rightleftharpoons NH_4(aq)$ OH (aq)

For More Material Join: @JEEAdvanced_2024

(ii) $3O_2(g) \rightleftharpoons 2O_3(g)$

(iii) $C_3H_8(g) \quad 5O_2(g) \rightleftharpoons 3CO_2(g) \quad 4H_2O(g)$

Equilibrium constant expression for them are

(i) $K_c \quad \dfrac{[NH_4(aq)][OH\ (aq)]}{[NH_3(aq)]}$; [] represents concentration in mol/ litre at equilibrium.

(ii) $K_c \quad \dfrac{[O_3]^2}{[O_2]^3}$; $K_p \quad \dfrac{P_{O_3}^2}{P_{O_2}^3}$ (iii) $K_C \quad \dfrac{[CO_2]^3[H_2O]^4}{[C_3H_8][O_2]^5}$; $K_p \quad \dfrac{P_{CO_2}^3 \ P_{H_2O}^4}{P_{C_3H_8} \ P_{O_2}^5}$

Note: Equilibrium constant for gaseous homogeneous equilibrium can be expressed in two ways K_p and K_C. This means value of equilibrium constant depends upon choice of standard state in which concentration of reactants and products are expressed.

❖ **Heterogeneous equilibria**

If reactants and products are present in two or more phases, the equilibria describing them is called heterogeneous equilibrium.

Example: (i) $AgI(s) \rightleftharpoons Ag\ (aq)\ \ I\ (aq)$

(ii) $CaO(s) \quad CO_2(g) \rightleftharpoons CaCO_3(s)$

(iii) $Br_2(l) \rightleftharpoons Br_2(g)$

Equilibrium expression for them can be written as

(i) $K_C \quad [Ag\ (aq)][I\ (aq)]$ (ii) $K_P \quad \dfrac{1}{P_{CO_2}}$; $K_C \quad \dfrac{1}{[CO_2(g)]}$

(iii) $K_P \quad P_{Br_2}(g)$; $K_C \quad [Br_2(g)]$

Predicting the Extent of a Reaction

High value of equilibrium constant indicates that product(s) concentration is high and its low value indicates that concentration of the product(s) in equilibrium mixture is low.

Large values of K_P or K_C (larger than about 10^3), favour the products strongly. For intermediate values of K (approximately in the range of $10^{\ 3}$ to 10^3), the concentrations of reactants and products are comparable. Small values of equilibrium constant (smaller than $10^{\ 3}$), favour the reactants strongly.

Predicting the Direction of the Reaction

For this purpose, we calculate the reaction quotient, Q. The reaction quotient is defined in the same way as the equilibrium constant (with molar concentrations to give Q_C, or with partial pressure to give Q_P) at any stage of reaction. For a general reaction

$aA \quad bB \rightleftharpoons cC \quad dD$

$Q_C \quad \dfrac{[C]^c[D]^d}{[A]^a[B]^d}$

Then, if $Q_C \quad K_C$, the reaction will proceed towards direction of reactants.

If $Q_C \quad K_c$, the reaction will move towards direction of the products.

If $Q_C \quad K_c$, the reaction mixture is at equilibrium

For More Material Join: @JEEAdvanced_2024

Important Point about Equilibrium Constant

- If the reaction is reversed, the value of equilibrium constant is inversed *e.g.,*

$CH_3COOH(aq) \quad C_2H_5OH(aq) \rightleftharpoons CH_3COOC_2H_5(aq) \quad H_2O(l)$, Equilibrium constant K

then for $CH_3COOC_2H_5(aq) \quad H_2O(l) \rightleftharpoons CH_3COOH(aq) \quad C_2H_5OH(aq)$, Equilibrium constant $1/K$

- If the reaction is divided by 2, equilibrium constant is the square root of the original *e.g.,*

for $N_2(g) \quad 3H_2(g) \rightleftharpoons 2NH_3(g)$, Equilibrium constant K

then for $\frac{1}{2}N_2(g) \quad \frac{3}{2}H_2(g) \rightleftharpoons NH_3(g)$, Equilibrium constant $\sqrt{K}$

If the reaction is multiplied by 2, equilibrium constant is the square of the original.

- If the reaction is written in two steps, equilibrium constant is equal to the product of the equilibrium constants of the step reactions *e. g.*, if for

$N_2(g) \quad 2O_2(g) \rightleftharpoons 2NO_2(g)$, Equilibrium constant K

and for the same reaction taking place in steps *i.e.*,

$N_2(g) \quad O_2(g) \rightleftharpoons 2NO(g)$, Equilibrium constant K_1

and $NO(g) \quad O_2(g) \rightleftharpoons 2NO_2(g)$, Equilibrium constant K_2

then $K \quad K_1 \quad K_2$

- Effect of temperature: According to van't Hoff equation , $\log \frac{K_2}{K_1} \quad \frac{H}{2.303R} \quad \frac{T_2 \quad T_1}{T_1 T_2}$

where K_1 and K_2 are the equilibrium constants at temperatures T_1 and T_2 respectively and H is the molar enthalpy change in the temperature range T_1 to T_2.

For exothermic reaction, as temperature increases K decreases.

For endothermic reaction, as temperature increases K increases.

Relationship Between Degree of Dissociation ( ) and Vapour Density

- For dissociation reaction: $A_n(g) \rightleftharpoons nA(g)$

Degree of dissociation ( ) $\quad \frac{\text{No. of moles dissociated}}{\text{Total no. of moles taken}}$

$$\frac{D \quad d}{d(n \quad 1)} \quad \frac{M_t \quad M_0}{M_0(n \quad 1)}$$

where M_t theoretical (calculated) molecular mass

M_0 observed (experimental) molecular mass

D theoretical vapour density, d observed vapour density.

n number of moles of product formed from 1 mole reactant

Density of gas mixture $\frac{PM}{RT}$

- For polymerization reaction

$nA(g) \rightleftharpoons A_n(g)$, where $n \quad 2$

$$\frac{D \quad d}{d \quad \frac{1}{n} \quad 1} \quad \frac{M_t \quad M_0}{M_0 \quad \frac{1}{n} \quad 1}$$

For More Material Join: @JEEAdvanced_2024

Le Chatelier's Principle

"If a system in equilibrium is subjected to a change of concentration, temperature or pressure, the equilibrium shifts in a direction so as to undo the effect of the change imposed."

❖ Effect of change in concentration on equilibrium

As we add or remove reactant (or product) the ratio of new concentrations become '*Q*' (reaction quotient)

$Q < K$: equilibrium will shift in forward direction.

$Q > K$: equilibrium will shift in backward direction.

❖ Effect of change in pressure

If a system in equilibrium consists of gases, then the concentrations of all the components can be altered by changing the pressure. When the pressure on the system is increased, the volume decreases proportionately. The equilibrium will shift in the direction in which there is decrease in number of moles *i.e.*, towards the direction in which there is decrease in volume.

❖ **Effect of change in pressure on melting point:** There are two types of solids:

(i) Solids whose volume decreases on melting, *e.g.*, ice, diamond, carborundum, magnesium nitride and quartz.

Solid (higher volume) $\rightleftharpoons$ Liquid (lower volume)

The process of melting is facilitated at high pressure, thus melting point is lowered.

(ii) Solid whose volume increase on melting, *e.g.*, Fe, Cu, Ag, Au, etc.

Solid (lower volume) $\rightleftharpoons$ Liquid (higher volume)

In this case the process of melting becomes difficult at high pressure, thus melting point becomes high.

❖ **Solubility of substances :** When solid substances are dissolved in water, either heat is evolved (exothermic) or heat is absorbed (endothermic).

(i) For endothermic solubility process, solubility increase with increase in temperature.

(ii) For exothermic solubility process, decrease with increase in temperature.

❖ **Solubility of gases in liquids :** When a gas dissolves in liquid, there is decrease in volume. Thus increase of pressure will favour the dissolution of gas in liquid.

❖ **Effect of temperature:** For endothermic reaction as temperature increases reaction shift in forward direction. For exothermic reaction as temperature increases reaction shift in backward direction.

❖ **Addition of inert gas**

(i) For reactions in which $n_p \quad n_r$ there is no effect of adding an inert gas at constant volume or at constant pressure on the equilibrium.

(ii) For reactions in which $n_p \quad n_r$, there is no effect of adding inert gas on the equilibrium at constant volume but at constant pressure, equilibrium shifts towards larger mole side.

Applying Le Chatelier's principle, the favourable conditions for the formation of NH_3 by Haber's process

i.e., $\quad N_2(g) \quad 3H_2(g) \rightleftharpoons 2NH_3(g), \quad H \quad 92.5$ kJ

are (i) High temperature (ii) Low pressure (iii) Removal of N_2 and H_2 (iv) Addition of inert gas at constant pressure.

For More Material Join: @JEEAdvanced_2024

❖ Physical Equilibrium

Only physical changes are involved.

Example : (i) Solid liquid equilibria

$$H_2O(s) \rightleftharpoons H_2O(l)$$

At equilibrium: Net rate of conversion of ice into water = Net rate of conversion of water into ice.

For $H_2O(l)$ normal freezing point is 273 K.

(ii) Liquid vapour equilibria :

$$H_2O(l) \rightleftharpoons H_2O(g)$$

Equilibrium is characterized by constant value of vapour pressure of $H_2O(l)$ at certain temperature.

For $H_2O(l)$ normal boiling point is 373 K.

For More Material Join: @JEEAdvanced_2024



Level 1

1. A reversible reaction is one which :
(a) proceeds in one direction
(b) proceeds in both directions
(c) proceeds spontaneously
(d) all the statements are wrong

2. The equilibrium constant K_c for the reaction

$$P_4(g) \rightleftharpoons 2P_2(g)$$

is 1.4 at 400°C. Suppose that 3 moles of $P_4(g)$ and 2 moles of $P_2(g)$ are mixed in 2 litre container at 400°C. What is the value of reaction quotient (Q_c)?
(a) $\frac{3}{2}$ (b) $\frac{2}{3}$ (c) 1 (d) None of these

3. In a chemical reaction equilibrium is established when :
(a) opposing reaction ceases
(b) concentrations of reactants and product are equal
(c) velocity of opposing reaction is the same as that of forward reaction
(d) reaction ceases to generate heat

4. The equilibrium constant for a reaction is K, and the reaction quotient is Q. For a particular reaction mixture, the ratio $\frac{K}{Q}$ is 0.33. This means that :
(a) the reaction mixture will equilibrate to form more reactant species
(b) the reaction mixture will equilibrate to form more product species
(c) the equilibrium ratio of reactant to product concentrations will be 3
(d) the equilibrium ratio of reactant to product concentrations will be 0.33

5. Consider the reaction $2SO_2(g)\ \ O_2(g) \rightleftharpoons 2SO_3(g)$ for which $K_c\ \ 278\ M^{\ 1}$. 0.001 mole of each of the reagents $SO_2(g), O_2(g)$ and $SO_3(g)$ are mixed in a 1.0 L flask. Determine the reaction quotient of the system and the spontaneous direction of the system :
(a) $Q_c\ \ 1000$; the equilibrium shifts to the right
(b) $Q_c\ \ 1000$; the equilibrium shifts to the left
(c) $Q_c\ \ 0.001$; the equilibrium shifts to the left
(d) $Q_c\ \ 0.001$; the equilibrium shifts to the right

6. In Q. No. 5, if the mixture of gases was allowed to come to equilibrium. The volume of the reaction vessel was then rapidly increased by a factor of two. As a result of the change the reaction quotient (Q_c) would :
(a) increase because of the pressure decrease
(b) decrease because of the pressure decrease
(c) remain the same because the equilibrium constant is independent of volume
(d) increase because the reaction is endothermic

7. For the reaction $A(g)\ \ 3B(g) \rightleftharpoons 2C(g)$ at 27°C, 2 moles of A, 4 moles of B and 6 moles of C are present in 2 litre vessel. If K_c for the reaction is 1.2, the reaction will proceed in :
(a) forward direction
(b) backward direction
(c) neither direction
(d) none of these

For More Material Join: @JEEAdvanced_2024

8. For a reversible gaseous reaction $N_2 + 3H_2 \rightleftharpoons 2NH_3$ at equilibrium, if some moles of H_2 are replaced by same number of moles of T_2 (T is tritium, isotope of H and assume isotopes do not have different chemical properties) without affecting other parameters, then :

(a) the sample of ammonia obtained after sometime will be radioactive.


(b) moles of N_2 after the change will be different as compared to moles of N_2 present before the change

(c) the value of K_p or K_c will change


(d) the average molecular mass of new equilibrium will be same as that of old equilibrium

9. For the synthesis of ammonia by the reaction $N_2 \quad 3H_2 \rightleftharpoons 2NH_3$ in the Haber's process, the attainment of equilibrium is correctly predicted by the curve


(a)




(b)




(c)



(d)



10. The figure shows the change in concentration of species A and B as a function of time.




The equilibrium constant K_c for the reaction $A(g) \rightleftharpoons 2B(g)$ is:

(a) $K_c \quad 1$ (b) $K \quad 1$

(c) $K \quad 1$ (d) data insufficient

11. Attainment of the equilibrium $A(g) \rightleftharpoons 2C(g) \quad B(g)$ gave the following graph. Find the correct option. (% dissociation = fraction dissociated 100)



For More Material Join: @JEEAdvanced_2024

(a) At t 5 sec equilibrium has been reached and K_c 40 (mol/litre)^2

(b) At t 5 sec equilibrium has been reached and % dissociation of A is 20%

(c) At t 5 sec equilibrium has been reached and % dissociation of A is 30%

(d) None of these

12. Using molar concentrations, what is the unit of K_c for the reaction?

$$CH_3OH(g) \rightleftharpoons CO(g) \quad 2H_2(g)$$

(a) M^2 (b) M^2 (c) M^1 (d) M

13. What is the unit of K_p for the reaction?

$$CS_2(g) \quad 4H_2(g) \rightleftharpoons CH_4(g) \quad 2H_2S(g)$$

(a) atm (b) atm^2 (c) atm^2 (d) atm^1

14. What is the equilibrium expression for the reaction $P_4(s) \quad 5O_2(g) \rightleftharpoons P_4O_{10}(s)$

(a) $K_c \quad [O_2]^5$ (b) $K_c \quad [P_4O_{10}] / 5[P_4][O_2]$

(c) $K_c \quad [P_4O_{10}] / [P_4][O_2]^5$ (d) $K_c \quad 1 / [O_2]^5$

15. At 527°C, the reaction given below has K_c 4

$$NH_3(g) \rightleftharpoons \frac{1}{2}N_2(g) \quad \frac{3}{2}H_2(g)$$

What is the K_p for the reaction?

$$N_2(g) \quad 3H_2(g) \rightleftharpoons 2NH_3(g)$$

(a) 16 $(800\,R)^2$ (b) $\left(\frac{800R}{4}\right)^2$ (c) $\left(\frac{1}{4 \quad 800R}\right)^2$ (d) None of these

16. The equilibrium constant for the reaction $N_2(g) \quad O_2(g) \rightleftharpoons 2NO(g)$ at temperature (T) is 4 10^{-4}. The value of K_c for the reaction $NO(g) \rightleftharpoons \frac{1}{2}N_2(g) \quad \frac{1}{2}O_2(g)$ at the same temperature is :

(a) 4 10^{-4} (b) 50 (c) 2.5 10^2 (d) 0.02

17. The equilibrium constant K_c for the following reaction at 842°C is 7.90 10^{-3}. What is K_p at same temperature?

$$\frac{1}{2}F_2(g) \rightleftharpoons F(g)$$

(a) 8.64 10^{-5} (b) 8.26 10^{-4} (c) 7.90 10^{-2} (d) 7.56 10^{-2}

18. The equilibrium constant K_p for the following reaction at 191°C is 1.24. What is K_c?

$$B(s) \quad \frac{3}{2}F_2(g) \rightleftharpoons BF_3(g)$$

(a) 6.7 (b) 0.61 (c) 8.30 (d) 7.6

19. For the equilibrium $SO_2Cl_2(g) \rightleftharpoons SO_2(g) \quad Cl_2(g)$, what is the temperature at which $\frac{K_p\,(\text{atm})}{K_c\,(\text{M})} \quad 3$?

(a) 0.027 K (b) 0.36 K (c) 36.54 K (d) 273 K

For More Material Join: @JEEAdvanced_2024

20. For the reversible reaction,

$N_2(g) \quad 3H_2(g) \rightleftharpoons 2NH_3(g)$

at 500°C, the value of K_p is 1.44 10 ^{5} when partial pressure is measured in atmospheres. The corresponding value of K_c with concentration in mole litre ^{1}, is:

(a) 1.44 10 ^{5}/(0.082 500) ^{2} (b) 1.44 10 ^{5}/(8.314 773) ^{2}

(c) 1.44 10 ^{5}/(0.082 773)^2 (d) 1.44 10 ^{5}/(0.082 773) ^{2}

21. For the reaction $CO(g) \quad Cl_2(g) \rightleftharpoons COCl_2(g)$ the value of $\frac{K_c}{K_p}$ is equal to :

(a) $\sqrt{RT}$ (b) RT (c) $\frac{1}{RT}$ (d) 1.0

22. The concentration of a pure solid or liquid phase is not included in the expression of equilibrium constant because :

(a) density of solid and liquid are independent of their quantities.

(b) solids and liquids react slowly.

(c) solids and liquids at equilibrium do not interact with gaseous phase.

(d) the molecules of solids and liquids cannot migrate to the gaseous phase.

23. A catalyst is a substance which :

(a) increases the equilibrium concentration of the product.

(b) changes the equilibrium constant of the reaction.

(c) shortens the time to reach equilibrium.

(d) supplies energy to the reaction.

24. What will be the effect on the equilibrium constant on increasing temperature, if the reaction neither absorbs heat nor releases heat?

(a) Equilibrium constant will remain constant.

(b) Equilibrium constant will decrease.

(c) Equilibrium constant will increase.

(d) Can not be predicted.

25. The equilibrium constant for the reaction $N_2(g) \quad O_2(g) \rightleftharpoons 2NO(g)$ is 4 10 ^{4} at 200 K. In presence of a catalyst, equilibrium is attained ten times faster. Therefore, the equilibrium constant in presence of the catalyst at 200 K is :

(a) 40 10 ^{4} (b) 4 10 ^{4}

(c) 4 10 ^{3} (d) difficult to compute without more data

26. For the reaction $H_2(g) \quad I_2(g) \rightleftharpoons 2HI(g)$, the equilibrium constant changes with :

(a) total pressure (b) catalyst

(c) concentration of H_2 and I_2 (d) temperature

27. Consider the reactions

(i) $2CO(g) \quad 2H_2O(g) \rightleftharpoons 2CO_2(g) \quad 2H_2(g)$; Eqm. Constant K_1

(ii) $CH_4(g) \quad H_2O(g) \rightleftharpoons CO(g) \quad 3H_2(g)$; Eqm. Constant K_2

For More Material Join: @JEEAdvanced_2024

(iii) $CH_4(g)$ $2H_2O(g) \rightleftharpoons CO_2(g)$ $4H_2(g)$; Eqm. Constant K_3

Which of the following relation is correct?

(a) K_3 $\frac{K_1}{K_2}$ (b) K_3 $\frac{K_1^2}{K_2^2}$ (c) K_3 K_1K_2 (d) K_3 $\sqrt{K_1}$ K_2

28. For the reaction $2NO_2(g)$ $\frac{1}{2}O_2(g) \rightleftharpoons N_2O_5(g)$, if the equilibrium constant is K_p, then the equilibrium constant for the reaction $2N_2O_5(g) \rightleftharpoons 4NO_2(g)$ $O_2(g)$ would be :

(a) K_p^2 (b) $\frac{2}{K_p}$ (c) $\frac{1}{K_p^2}$ (d) $\frac{1}{\sqrt{K_p}}$

29. The equilibrium constant (K_c) for the reaction

$$2HCl(g) \rightleftharpoons H_2(g) \quad Cl_2(g)$$

is 4 10^{34} at 25°C. What is the equilibrium constant for the reaction?

$$\frac{1}{2}H_2(g) \quad \frac{1}{2}Cl_2(g) \rightleftharpoons HCl(g)$$

(a) 2 10^{17} (b) 2.5 10^{33} (c) 5 10^6 (d) None of these

30. At a certain temperature, the following reactions have the equilibrium constants as shown below:

$$S(s) \quad O_2(g) \rightleftharpoons SO_2(g); \quad K_c \quad 5 \quad 10^{52}$$

$$2S(s) \quad 3O_2(g) \rightleftharpoons 2SO_3(g); \; K_c \quad 10^{29}$$

What is the equilibrium constant K_c for the reaction at the same temperature ?

$$2SO_2(g) \quad O_2(g) \rightleftharpoons 2SO_3(g)$$

(a) 2.5 10^{76} (b) 4 10^{23} (c) 4 10^{77} (d) None of these

31. Consider the following gaseous equilibria given below :

(I) N_2 $3H_2 \rightleftharpoons 2NH_3$; Eqm. Constant K_1

(II) $N_2 + O_2 \rightleftharpoons 2NO$; Eqm. Constant K_2

(III) $H_2 + \frac{1}{2}O_2 \rightleftharpoons H_2O$; Eqm. Constant K_3

The equilibrium constant for the reaction, $2NH_3 + \frac{5}{2}O_2 \rightleftharpoons 2NO + 3H_2O$ in terms of K_1, K_2 and K_3 will be :

(a) $K_1K_2K_3$ (b) $\frac{K_1K_2}{K_3}$ (c) $\frac{K_1K_3^2}{K_2}$ (d) $\frac{K_2K_3^3}{K_1}$

32. In the reaction $X(g)$ $Y(g) \rightleftharpoons 2\,Z(g)$, 2 mole of X, 1 mole of Y and 1 mole of Z are placed in a 10 litre vessel and allowed to reach equilibrium. If final concentration of Z is 0.2 M, then K_c for the given reaction is :

(a) 1.60 (b) $\frac{80}{3}$ (c) $\frac{16}{3}$ (d) None of these

For More Material Join: @JEEAdvanced_2024

33. An equilibrium mixture of the reaction $2H_2S(g) \rightleftharpoons 2H_2(g)\ \ S_2(g)$ had 0.5 mole H_2S, 0.10 mole H_2 and 0.4 mole S_2 in one litre vessel. The value of equilibrium constant (K) in mol litre ^{1} is :

(a) 0.004 (b) 0.008 (c) 0.016 (d) 0.160

34. Given $[CS_2]$ 0.120 M, $[H_2]$ 0.10 M, $[H_2S]$ 0.20 M and $[CH_4]$ 8.40 10^{5} M for the following reaction at 900°C, at eq. Calculate the equilibrium constant (K_c).

$$CS_2(g)\ \ 4H_2(g) \rightleftharpoons CH_4(g)\ \ 2H_2S(g)$$

(a) 0.0120 (b) 0.0980 (c) 0.280 (d) 0.120

35. The equilibrium constant for the following reaction is 10.5 at 500 K. A system at equilibrium has [CO] 0.250 M and $[H_2]$ 0.120 M. What is the $[CH_3OH]$?

$$CO(g)\ \ 2H_2(g) \rightleftharpoons CH_3OH(g)$$

(a) 0.03618 (b) 0.0435 (c) 0.546 (d) 0.0499

36. When sulphur (in the form of S_8) is heated at temperature T, at equilibrium, the pressure of S_8 falls by 30% from 1.0 atm, because $S_8(g)$ is partially converted into $S_2(g)$.

Find the value of K_p for this reaction.

(a) 2.96 (b) 6.14 (c) 204.8 (d) None of these

37. 9.2 grams of $N_2O_4(g)$ is taken in a closed one litre vessel and heated till the following equilibrium is reached $N_2O_4(g) \rightleftharpoons 2NO_2(g)$

At equilibrium, 50% $N_2O_4(g)$ is dissociated. What is the equilibrium constant (in mol litre ^{1}) (molecular mass of N_2O_4 92)

(a) 0.1 (b) 0.4 (c) 0.2 (d) 2

38. Two moles of NH_3 when put into a previously evacuated vessel (one litre), partially dissociated into N_2 and H_2. If at equilibrium one mole of NH_3 is present, the equilibrium constant is :

(a) 3/4 mol^{2}litre ^{2} (b) 27/64 mol^{2}litre ^{2} (c) 27/32 mol^{2}litre ^{2} (d) 27/16 mol^{2}litre ^{2}

39. In the presence of excess of anhydrous $SrCl_2$, the amount of water taken up is governed by K_p 10^{12} atm ^{4} for the following reaction at 273 K

$$SrCl_2\ \ 2H_2O(s)\ \ 4H_2O(g) \rightleftharpoons SrCl_2\ \ 6H_2O(s)$$

What is equilibrium vapour pressure (in torr) of water in a closed vessel that contains $SrCl_2\ \ 2H_2O(s)$?

(a) 0.001 torr (b) 10^{3} torr (c) 0.76 torr (d) 1.31 torr

40. $CuSO_4\ \ 5H_2O(s) \rightleftharpoons CuSO_4\ \ 3H_2O(s)\ \ 2H_2O(g)$; K_p 4 10^{4} atm^{2}. If the vapour pressure of water is 38 torr then percentage of relative humidity is : (Assume all data at constant temperature)

(a) 4 (b) 10 (c) 40 (d) None of these

41. $NH_4HS(s) \rightleftharpoons NH_3(g)\ \ H_2S(g)$

The equilibrium pressure at 25°C is 0.660 atm. What is K_p for the reaction?

(a) 0.109 (b) 0.218 (c) 1.89 (d) 2.18

For More Material Join: @JEEAdvanced_2024

42. For the reaction $2A(g) \rightleftharpoons B(g)\quad 3C(g)$, at a given temperature, $K_c\quad 16$. What must be the volume of the flask, if a mixture of 2 mole each of A, B and C exist in equilibrium?

(a) $\frac{1}{4}$ (b) $\frac{1}{2}$ (c) 1 (d) None of these

43. One mole of pure ethyl alcohol was treated with one mole of pure acetic acid at 25°C. One-third of the acid changes into ester at equilibrium. The equilibrium constant for the reaction will be :

(a) $\frac{1}{4}$ (b) 2 (c) 3 (d) 4

44. $I_2(aq)\quad I\ (aq) \rightleftharpoons I_3(aq)$. We started with 1 mole of I_2 and 0.5 mole of I in one litre flask. After equilibrium is reached, excess of $AgNO_3$ gave 0.25 mole of yellow precipitate. Equilibrium constant is :

(a) 1.33 (b) 2.66 (c) 2.0 (d) 3.0

45. At 87°C, the following equilibrium is established.

$$H_2(g)\quad S(s) \rightleftharpoons H_2S(g);\ K_c\quad 0.08$$

If 0.3 mole hydrogen and 2 mole sulphur are heated to 87°C in a 2 L vessel, what will be the concentration of H_2S at equilibrium?

(a) 0.011 M (b) 0.022 M (c) 0.044 M (d) 0.08 M

46. In the equilibrium $2SO_2(g)\quad O_2(g) \rightleftharpoons 2SO_3(g)$, the partial pressure of SO_2, O_2 and SO_3 are 0.662, 0.10 and 0.331 atm respectively. What should be the partial pressure of oxygen so that the equilibrium concentrations of SO_2 and SO_3 are equal?

(a) 0.4 atm (b) 1.0 atm (c) 0.8 atm (d) 0.25 atm

47. When heated, ammonium carbamate decomposes as follows :

$$NH_4COONH_2(s) \rightleftharpoons 2NH_3(g)\quad CO_2(g)$$

At a certain temperature, the equilibrium pressure of the system is 0.318 atm. K_p for the reaction is :

(a) 0.128 (b) 0.426 (c) $4.76\quad 10^{\ 3}$ (d) None of these

48. In a system $A(s) \rightleftharpoons 2B(g)\quad 3C(g)$, if the concentration of C at equilibrium is increased by a factor of 2, it will cause the equilibrium concentration of B to change to :

(a) two times the original value (b) one half of its original value

(c) $2\sqrt{2}$ times to the original value (d) $\frac{1}{2\sqrt{2}}$ times the original value

49. For $A\quad B \rightleftharpoons C\quad D$, if initially the concentration of A and B are both equal but at equilibrium concentration of D will be twice of that of A then what will be the equilibrium constant of reaction ?

(a) $\frac{4}{9}$ (b) $\frac{9}{4}$ (c) $\frac{1}{9}$ (d) 4

50. The equilibrium constant K_c for the reaction $SO_2(g)\quad NO_2(g) \rightleftharpoons SO_3(g)\quad NO(g)$ is 16. If 1 mole of each of all the four gases is taken in 1 dm^3 vessel, the equilibrium concentration of NO would be :

(a) 0.4 M (b) 0.6 M (c) 1.4 M (d) 1.6 M

For More Material Join: @JEEAdvanced_2024

51. On increasing the temperature, the rate of a reaction :
(a) always increases
(b) always decreases
(c) first increases and then decreases
(d) may increase or decrease depending upon the nature of the reaction

52. A catalyst increases the rate of a reaction by :
(a) increasing the activation energy of the reaction
(b) increasing the value of rate constants (k_f and k_b)
(c) increasing the enthalpy change of the reaction
(d) decreasing the enthalpy change of the reaction

53. At a certain temperature, only 50% HI is dissociated at equilibrium in the following reaction :

$$2HI(g) \rightleftharpoons H_2(g) \quad I_2(g)$$

The equilibrium constant for this reaction is :
(a) 0.25 (b) 1.0 (c) 3.0 (d) 0.5

54. The equilibrium constant K_p for the reaction

$$H_2(g) \quad CO_2(g) \rightleftharpoons H_2O(g) \quad CO(g)$$

is 4.0 at 1660°C. Initially 0.80 mole H_2 and 0.80 mole CO_2 are injected into a 5.0 litre flask. What is the equilibrium concentration of $CO_2(g)$?
(a) 0.533 M (b) 0.0534 M (c) 0.535 M (d) None of these

55. At 273 K and 1 atm, 10 litre of N_2O_4 decomposes to NO_2 according to equation

$$N_2O_4(g) \rightleftharpoons 2NO_2(g)$$

What is degree of dissociation ( ) when the original volume is 25% less than that of existing volume?
(a) 0.25 (b) 0.33 (c) 0.66 (d) 0.5

56. The equilibrium constant for the reaction $CO(g) \quad H_2O(g) \rightleftharpoons CO_2(g) \quad H_2(g)$ is 5. How many moles of CO_2 must be added to 1 litre container already containing 3 moles each of CO and H_2O to make 2M equilibrium concentration of CO?
(a) 15 (b) 19 (c) 5 (d) 20

57. A nitrogen-hydrogen mixture initially in the molar ratio of 1 : 3 reached equilibrium to form ammonia when 25% of the N_2 and H_2 had reacted. If the total pressure of the system was 21 atm, the partial pressure of ammonia at the equilibrium was :
(a) 4.5 atm (b) 3.0 atm (c) 2.0 atm (d) 1.5 atm

58. Ammonia under a pressure of 15 atm at 27°C is heated to 347°C in a closed vessel in the presence of a catalyst. Under the conditions, NH_3 is partially decomposed according to the equation,

$2NH_3 \rightleftharpoons N_2 \quad 3H_2$ the vessel is such that the volume remains effectively constant where as pressure increases to 50 atm. Calculate the percentage of NH_3 actually decomposed :
(a) 65% (b) 61.3% (c) 62.5% (d) 64%

For More Material Join: @JEEAdvanced_2024

59. 0.1 mole of $N_2O_4(g)$ was sealed in a tube under one atmospheric conditions at 25°C. Calculate the number of moles of $NO_2(g)$ present, if the equilibrium $N_2O_4(g) \rightleftharpoons 2NO_2(g)$ (K_p 0.14) is reached after some time :

(a) 1.8 10^2 (b) 2.8 10^2 (c) 0.034 (d) 2.8 10^{-2}

60. 5 moles of SO_2 and 5 moles of O_2 are allowed to react. At equilibrium, it was found that 60% of SO_2 is used up. If the pressure of the equilibrium mixture is one atmosphere, the partial pressure of O_2 is:

(a) 0.52 atm (b) 0.21 atm (c) 0.41 atm (d) 0.82 atm

61. $N_2(g)$ $3H_2(g) \rightleftharpoons 2NH_3(g)$

For the reaction initially the mole ratio was 1 : 3 of $N_2 : H_2$. At equilibrium 50% of each has reacted. If the equilibrium pressure is P, the partial pressure of NH_3 at equilibrium is :

(a) $\frac{P}{3}$ (b) $\frac{P}{4}$ (c) $\frac{P}{6}$ (d) $\frac{P}{8}$

62. 2.0 mole of PCl_5 were introduced in a vessel of 5.0 L capacity of a particular temperature. At equilibrium, PCl_5 was found to be 35% dissociated into PCl_3 and Cl_2. The value of K_c for the reaction

$$PCl_3(g) \quad Cl_2(g) \rightleftharpoons PCl_5(g)$$

(a) 1.89 (b) 0.377 (c) 1.33 (d) 13.3

63. At certain temperature compound $AB_2(g)$ dissociates according to the reaction

$$2AB_2(g) \rightleftharpoons 2AB(g) \quad B_2(g)$$

With degree of dissociation , which is small compared with unity, the expression of K_p, in terms of and initial pressure P is :

(a) $P\frac{^3}{2}$ (b) $\frac{P^{\ 2}}{3}$ (c) $P\frac{^3}{3}$ (d) $\frac{P^{\ 2}}{2}$

64. For the reaction

$H_2(g)$ $CO_2(g) \rightleftharpoons CO(g)$ $H_2O(g)$, if the initial concentration of $[H_2]$ $[CO_2]$ $1M$ and x moles/litre of hydrogen is consumed at equilibrium, the correct expression of K_p is :


(a) $\frac{x^2}{(1 \quad x)^2}$ (b) $\frac{(1 \quad x)^2}{(1 \quad x)^2}$ (c) $\frac{x^2}{(2 \quad x)^2}$ (d) $\frac{x^2}{1 \quad x^2}$


65. If D_T and D_O are the theoretical and observed vapour densities at a definite temperature and be the degree of dissociation of a substance, then, in the terms of D_O, D_T and n (number of moles of products formed from 1 mole reactant) is calculated by the formula :


(a) $\frac{D_O \quad D_T}{(1 \quad n)\, D_T}$ (b) $\frac{D_T \quad D_O}{(n \quad 1)\, D_T}$ (c) $\frac{D_T \quad D_O}{(n \quad 1)\, D_O}$ (d) $\frac{D \quad D_T}{(n \quad 1)\, D_T}$

66. For the dissociation of PCl_5 into PCl_3 and Cl_2 in gaseous phase reaction, if d is the observed vapour density and D the theoretical vapour density with ' ' as degree of dissociation, variation of D/d with ' ' is given by?

For More Material Join: @JEEAdvanced_2024

(a) 

(b) 

(c) 

(d) None of these

67. At 27°C and 1 atm pressure, N_2O_4 is 20% dissociation into NO_2. What is the density of equilibrium mixture of N_2O_4 and NO_2 at 27°C and 1 atm?
(a) 3.11 g/litre (b) 2.11 g/litre (c) 4.5 g/litre (d) None of these

68. $COCl_2$ gas dissociates according to the equation, $COCl_2(g) \rightleftharpoons CO(g)\ \ Cl_2(g)$. When heated to 700 K the density of the gas mixture at 1.16 atm and at equilibrium is 1.16 g/litre. The degree of dissociation of $COCl_2$ at 700 K is :
(a) 0.28 (b) 0.50 (c) 0.72 (d) 0.42

69. The degree of dissociation of I_2 molecule of 1000°C and under atmospheric pressure is 40% by volume. If the dissociation is reduced to 20% at the same temp., total equilibrium pressure on the gas is :
(a) 1.57 atm (b) 2.57 atm (c) 3.57 atm (d) 4.57 atm

70. Determine the value of equilibrium constant (K_c) for the reaction

$$A_2(g)\ \ B_2(g) \rightleftharpoons 2AB(g)$$

if 10 moles of A_2, 15 moles of B_2 and 5 moles of AB are placed in a 2 litre vessel and allowed to come to equilibrium. The final concentration of AB is 7.5 M :
(a) 4.5 (b) 1.5 (c) 0.6 (d) None of these

71. At 87°C, the following equilibrium is established

$$H_2(g)\ \ S(s) \rightleftharpoons H_2S(g);\ \ K_p\ \ 7\ \ 10^{\ 2}$$

If 0.50 mole of hydrogen and 1.0 mole of sulphur are heated to 87 C in 1.0 L vessel, what will be the partial pressure of H_2S at equilibrium?
(a) 0.966 atm (b) 1.38 atm (c) 0.0327 atm (d) 1 atm

72. Pure PCl_5 is introduced into an evacuated chamber and comes to equilibrium at 247°C and 2.0 atm. The equilibrium gaseous mixture contains 40% chlorine by volume.
Calculate K_p at 247°C for the reaction

$$PCl_5(g) \rightleftharpoons PCl_3(g)\ \ Cl_2(g)$$

(a) 0.625 atm (b) 4 atm (c) 1.6 atm (d) None of these

73. For the reaction

$$SnO_2(s)\ \ 2H_2(g) \rightleftharpoons 2H_2O(g)\ \ Sn(l)$$

calculate K_p at 900 K, where the equilibrium steam-hydrogen mixture was 45% H_2 by volume :
(a) 1.49 (b) 1.22 (c) 0.67 (d) None of these

For More Material Join: @JEEAdvanced_2024

74. For the reaction $XCO_3(s) \rightleftharpoons XO(s) \quad CO_2(g)$, K_p 1.642 atm at 727°C. If 4 moles of $XCO_3(s)$ was put into a 50 litre container and heated to 727°C.

What mole percent of the XCO_3 remains unreacted at equilibrium?

(a) 20 (b) 25 (c) 50 (d) None of these

75. $Fe_2O_3(s)$ may be converted to Fe by the reaction

$$Fe_2O_3(s) \quad 3H_2(g) \rightleftharpoons 2Fe(s) \quad 3H_2O(g),$$

for which K_c 8 at temp. 720°C.

What percentage of the H_2 remains unreacted after the reaction has come to equilibrium?

(a) ~22% (b) ~34% (c) ~66% (d) ~78%

76. $AB_3(g)$ is dissociates as $AB_3(g) \rightleftharpoons AB_2(g) \quad \frac{1}{2}B_2(g)$.

When the initial pressure of AB_3 is 800 torr and the total pressure developed at equilibrium is 900 torr, what fraction of $AB_3(g)$ is dissociated?

(a) 10% (b) 20% (c) 25% (d) 30%

77. At 1000 K, a sample of pure NO_2 gas decomposes as :

$$2NO_2(g) \rightleftharpoons 2NO(g) \quad O_2(g)$$

The equilibrium constant K_p is 156.25 atm. Analysis shows that the partial pressure of O_2 is 0.25 atm at equilibrium. The partial pressure of NO_2 at equilibrium is :

(a) 0.01 (b) 0.02 (c) 0.04 (d) None of these

78. Pure nitrosyl chloride (NOCl) gas was heated to 240°C in a 1.0 L container. At equilibrium the total pressure was 1.0 atm and the NOCl pressure was 0.64 atm. What would be the value of K_p?

(a) 1.02 atm (b) 16.875 10^{3} atm

(c) 16 10^{2} atm (d) None of these

79. At a certain temperature the equilibrium constant K_c is 0.25 for the reaction

$$A_2(g) \quad B_2(g) \rightleftharpoons C_2(g) \quad D_2(g)$$

If we take 1 mole of each of the four gases in a 10 litre container, what would be equilibrium concentration of $A_2(g)$?

(a) 0.331 M (b) 0.033 M (c) 0.133 M (d) 1.33 M

80. At 200°C PCl_5 dissociates as follows :

$$PCl_5(g) \rightleftharpoons PCl_3(g) \quad Cl_2(g)$$

It was found that the equilibrium vapours are 62 times as heavy as hydrogen. The % degree of dissociation of PCl_5 at 200°C is nearly :

(a) 10% (b) 42% (c) 50% (d) 68%

81. For the dissociation reaction $N_2O_4(g) \rightleftharpoons 2NO_2(g)$, the degree of dissociation ( ) in terms of K_p and total equilibrium pressure P is :

(a) $\sqrt{\frac{4P \quad K_P}{K_P}}$ (b) $\sqrt{\frac{K_P}{4P \quad K_P}}$


(c) $\sqrt{\frac{K_P}{4P}}$ (d) None of these


For More Material Join: @JEEAdvanced_2024


82. Consider the following equilibrium


$$N_2O_4(g) \rightleftharpoons 2NO_2(g)$$

Then the select the correct graph, which shows the variation in concentrations of NO_2 against concentrations of N_2O_4:

(a) 

(b) 

(c) 

(d) 

83. The vapour pressure of mercury is 0.002 mm Hg at 27°C. K_c for the process $Hg(l) \rightleftharpoons Hg(g)$ is:

(a) 0.002 (b) 8.12 10 ^{5} (c) 6.48 10 ^{5} (d) 1.068 10 ^{7}

84. Calculate the equilibrium constant (K_c) for the reaction given below, if at equilibrium mixture contains 5.0 mole of A_2, 3 mole of B_2 and 2 mole of AB_2 at 8.21 atm and 300 K

$$A_2(g) \quad 2B_2(g) \rightleftharpoons 2AB_2(g) \quad \text{Heat}$$

(a) 1.333 (b) 2.66 (c) 20 (d) None of these

85. For the reaction (1) and (2)

$$A(g) \rightleftharpoons B(g) \quad C(g) \qquad \ldots(1)$$

$$X(g) \rightleftharpoons 2Y(g) \qquad \ldots(2)$$

Given, $K_{P_1} : K_{P_2}$ 9 : 1

If the degree of dissociation of $A(g)$ and $X(g)$ be same then the total pressure at equilibrium (1) and (2) are in the ratio :

(a) 3 : 1 (b) 36 : 1 (c) 1 : 1 (d) 0.5 : 1

86. Given the following reaction at equilibrium, $N_2(g)$ $3H_2(g) \rightleftharpoons 2NH_3(g)$. At constant pressure if some inert gas added in to the system, then:

(a) More $NH_3(g)$ is produced (b) Less $NH_3(g)$ is produced

(c) No affect on the equilibrium (d) K_p of the reaction is decreased

87. In which of the following equilibrium, change in volume of the system does not alter the number of moles :

(a) $N_2(g)$ $O_2(g) \rightleftharpoons 2NO(g)$ (b) $PCl_5(g) \rightleftharpoons PCl_3(g)$ $Cl_2(g)$

(c) $N_2(g)$ $3H_2(g) \rightleftharpoons 2NH_3(g)$ (d) $SO_2Cl_2(g) \rightleftharpoons SO_2(g)$ $Cl_2(g)$

88. For the reaction

$$N_2(g) \quad 3H_2(g) \rightleftharpoons 2NH_3(g); \qquad H \qquad 93.6 \text{ kJ mol}^{1}$$

The number of moles of H_2 at equilibrium will increase if :

(a) volume is increased (b) volume is decreased

(c) argon gas is added at constant volume (d) NH_3 is removed

For More Material Join: @JEEAdvanced_2024

89. The volume of the reaction vessel containing an equilibrium mixture is increased in the following reaction

$$SO_2Cl_2(g) \rightleftharpoons SO_2(g) \quad Cl_2(g)$$

When equilibrium is re-established :

(a) the amount of $Cl_2(g)$ remains unchanged

(b) the amount of $Cl_2(g)$ increases

(c) the amount of $SO_2Cl_2(g)$ increases

(d) the amount of $SO_2(g)$ decreases

90. Some inert gas is added at constant volume to the following reaction at equilibrium

$$NH_4HS(s) \rightleftharpoons NH_3(g) \quad H_2S(g)$$

Predict the effect of adding the inert gas :

(a) The equilibrium shifts in the forward direction

(b) The equilibrium shifts in the backward direction

(c) The equilibrium remains unaffected

(d) The value of K_p is increased

91 Consider the reaction where K_p 0.497 at 500 K

$$PCl_5(g) \rightleftharpoons PCl_3(g) \quad Cl_2(g)$$

If the three gases are mixed in a rigid container so that the partial pressure of each gas is initially 1 atm, then which is correct observation ?


(a) More PCl_5 will be produced

(b) More PCl_3 will be produced

(c) Equilibrium will be established when 50% reaction is complete

(d) None of the above

92. The preparation of $SO_3(g)$ by reaction $SO_2(g) \quad \frac{1}{2}O_2(g) \rightleftharpoons SO_3(g)$ is an exothermic reaction. If the preparation follows the following temperature-pressure relationship for its % yield, then for temperatures T_1, T_2 and T_3. The correct option is :



(a) $T_3 \quad T_2 \quad T_1$


(b) $T_1 \quad T_2 \quad T_3$


(c) $T_1 \quad T_2 \quad T_3$


(d) Nothing could be predicted about temperature through given information


93. An equilibrium mixture at 700 K of $0.05\,M\ N_2(g), 0.3\,M\ H_2(g)$ and $0.2\,M\ NH_3(g)$ is present in a container. Now if this equilibrium is disturbed by adding $N_2(g)$ so that its concentration becomes $0.15\,M$ just after addition then which of the following graphs represents the above situation more appropriately:

For More Material Join: @JEEAdvanced_2024

(a)


(b)


(c)


(d)


94. In a vessel containing N_2, H_2 and NH_3 at equilibrium, some helium gas is introduced so that total pressure increase while temperature and volume remain constant. According to Le Chatelier's principle, the dissociation of NH_3 :

(a) increases (b) decreases

(c) remains unaltered (d) changes unpredictably

95. Le-Chatelier principle is not applicable to :

(a) $H_2(g)\quad I_2(g) \rightleftharpoons 2HI(g)$ (b) $Fe(s)\quad S(s) \rightleftharpoons FeS(s)$

(c) $N_2(g)\quad 3H_2(g) \rightleftharpoons 2NH_3(g)$ (d) $N_2(g)\quad O_2(g) \rightleftharpoons 2NO(g)$

96. Consider the following reactions. In which case the formation of product is favoured by decrease in pressure?

(1) $CO_2(g)\quad C(s) \rightleftharpoons 2\,CO(g);\qquad H\quad 172.5\,kJ$

(2) $N_2(g)\quad 3H_2(g) \rightleftharpoons 2NH_3(g);\qquad H\quad 91.8\,kJ$

(3) $N_2(g)\quad O_2(g) \rightleftharpoons 2NO(g);\qquad H\quad 181\,kJ$

(4) $2H_2O(g) \rightleftharpoons 2H_2(g)\quad O_2(g);\qquad H\quad 484.6\,kJ$

(a) 2, 3 (b) 3, 4 (c) 2, 4 (d) 1, 4

97. In which of the following reactions, the formation of product is favoured by decrease in temperature?

(1) $N_2(g)\quad O_2(g) \rightleftharpoons 2NO(g);\qquad H\quad 181$

(2) $2CO_2(g) \rightleftharpoons 2CO(g)\quad O_2(g);\qquad H\quad 566$

(3) $H_2(g)\quad I_2(g) \rightleftharpoons 2HI(g);\qquad H\quad 9.4$

(4) $H_2(g)\quad F_2(g) \rightleftharpoons 2HF(g);\qquad H\quad 541$

(a) 1, 2 (b) 2 only (c) 1, 2, 3 (d) 3, 4

For More Material Join: @JEEAdvanced_2024

98. For which of the following reactions is product formation favoured by low pressure and high temperature?

(a) $H_2(g)$ $I_2(g) \rightleftharpoons 2HI(g)$; H 9.4 kJ
(b) $CO_2(g)$ $C(s) \rightleftharpoons 2CO(g)$; H 172.5 kJ
(c) $CO(g)$ $2H_2(g) \rightleftharpoons CH_3OH(l)$; H 21.7 kJ
(d) $3O_2(g) \rightleftharpoons 2O_3(g)$; H 285 kJ

99. For which of the following reaction is product formation favoured by low pressure and low temperature?

(a) $CO_2(g)$ $C(s) \rightleftharpoons 2CO(g)$; H 172.5 kJ
(b) $CO(g)$ $2H_2(g) \rightleftharpoons CH_3OH$; H 21.7 kJ
(c) $2O_3(g) \rightleftharpoons 3O_2(g)$; H 285 kJ
(d) $H_2(g)$ $F_2(g) \rightleftharpoons 2HF(g)$; H 541 kJ

100. Consider the following reactions at equilibrium and determine which of the indicated changes will cause the reaction to proceed to the right.

(1) $CO(g)$ $3H_2(g) \rightleftharpoons CH_4(g)$ $H_2O(g)$ (add CH_4)
(2) $N_2(g)$ $3H_2(g) \rightleftharpoons 2NH_3(g)$ (remove NH_3)
(3) $H_2(g)$ $F_2(g) \rightleftharpoons 2HF(g)$ (add F_2)
(4) $BaO(s)$ $SO_3(g) \rightleftharpoons BaSO_4(s)$ (add BaO)

(a) 2, 3 (b) 1, 4 (c) 2, 4 (d) 2, 3, 4

101. If the pressure in a reaction vessel for the following reaction is increased by decreasing the volume, what will happen to the concentrations of CO and CO_2?

$$H_2O(g) \quad CO(g) \rightleftharpoons H_2(g) \quad CO_2(g) \quad \text{Heat}$$

(a) both the [CO] and [CO_2] will decrease
(b) neither the [CO] nor the [CO_2] will change
(c) the [CO] will decrease and the [CO_2] will increase
(d) both the [CO] and [CO_2] will increase

102. Consider the following reaction and determine which of the conditions will shift the equilibrium position to the right?

$$4NH_3(g) \quad 5O_2(g) \rightleftharpoons 4NO(g) \quad 6H_2O(g) \quad \text{Heat}$$

(a) increasing the temperature (b) increasing the pressure
(c) adding a catalyst (d) none of the above is correct

103. The conversion of ozone into oxygen is exothermic. Under what conditions is ozone the most stable?

$$2O_3(g) \rightleftharpoons 3O_2(g)$$

(a) At low pressure and low temperature (b) At high pressure and high temperature
(c) At high pressure and low temperature (d) At low pressure and high temperature

104. A system at equilibrium is described by the equation of fixed temperature *T*.

$$SO_2Cl_2(g) \rightleftharpoons SO_2(g) \quad Cl_2(g)$$

What effect will be the effect on equilibrium, if total pressure is increased by reducing volume?

For More Material Join: @JEEAdvanced_2024

(a) Concentration of $SO_2Cl_2(g)$ increases (b) Concentration of $SO_2(g)$ increases
(c) Concentration of $Cl_2(g)$ increases (d) Concentration of all gases increases

105. The reaction $2NO_2(g) \rightleftharpoons N_2O_4(g)$ is an exothermic equilibrium. This means that :
(a) equilibration of this gas mixture will be slower at high temperature
(b) a mole of N_2O_4 will occupy twice the volume of a mole of NO_2 at the same temperature
(c) the equilibrium will move to the right if an equilibrium mixture is cooled
(d) the position of equilibrium will move to the left with increasing gas pressure

106. Densities of diamond and graphite are 3.5 and 2.3 g/mL

C (diamond) $\rightleftharpoons$ C (graphite); $_rH$ 1.9 kJ/mol

favourable conditions for the formation of graphite are :
(a) high pressure and low temperature (b) low pressure and high temperature
(c) high pressure and high temperature (d) low pressure and low temperature


107. For an equilibrium $H_2O(s) \rightleftharpoons H_2O(l)$, which of the following statements is true?
(a) The pressure changes do not affect the equilibrium
(b) More of ice melts if pressure on the system is increased
(c) More of liquid freezes if pressure on the system is increased
(d) The pressure changes may increase or decrease the degree of advancement of the process


108. A pressure cooker reduces cooking time for food because:
(a) the higher pressure inside the cooker crushes the food material
(b) cooking involves chemical changes helped by a rise in temperature
(c) heat is more evenly distributed in the cooking space
(d) boiling point of water involved in cooking is increased


109. The vapour pressure of a liquid in a closed container depends on :
(1) temperature of liquid
(2) quantity of liquid
(3) surface area of the liquid
(a) 1 only (b) 2 only (c) 1 and 3 only (d) 1, 2 and 3


110. The pressure on a sample of water at its triple point is reduced while the temperature is held constant. Which phases changes are favoured?
(I) melting of ice
(II) sublimation of ice
(III) vaporization of liquid water
(a) I only (b) III only (c) II only (d) II and III

111. An exothermic reaction is represented by the graph :

(a) 


(b) 


(c) 


(d) 


For More Material Join: @JEEAdvanced_2024

112. An endothermic reaction is represented by the graph :


(a) 

(b) 

(c) 

(d) 

113. A schematic plot of ln K_{eq} *versus* inverse of temperature for a reaction is shown below



the reaction must be :

(a) Exothermic

(b) Endothermic

(c) One with negligible enthalpy change


(d) Highly spontaneous at ordinary temperature


114. What is the correct relationship between free energy change and equilibrium constant of a reaction?


(a) $G \quad RT \ln K$ (b) $G \quad RT \ln K$


(c) $G \quad RT \ln K$ (d) $G \quad RT \ln K$

115. For the chemical equilibrium $CaCO_3(s) \rightleftharpoons CaO(s) \quad CO_2(g)$, $_rH$ can be determined from which one of the following plots ?

(a) 

(b) 

(c) 

(d) 

116. K_p has the value of $10^{\ 6}$ atm^3 and $10^{\ 4}$ atm^3 at 298 K and 323 K respectively for the reaction

$$CuSO_4 \quad 3H_2O(s) \rightleftharpoons CuSO_4(s) \quad 3H_2O(g)$$

$_rH$ for the reaction is :

(a) 7.7 kJ/mol (b) 147.41 kJ/mol

(c) 147.41 kJ/mol (d) None of these

117. van't Hoff's equation shows the effect of temperature on equilibrium constants K_c and K_p. The K_p varies with temperature according to the relation :

(a) $\log \dfrac{K_{p_2}}{K_{p_1}} \quad \dfrac{H}{2.303\,R} \quad \dfrac{T_1 \quad T_2}{T_1 T_2}$ (b) $\log \dfrac{K_{p_2}}{K_{p_1}} \quad \dfrac{H}{2.303\,R} \quad \dfrac{T_2 \quad T_1}{T_1 T_2}$

For More Material Join: @JEEAdvanced_2024

(c) $\log \frac{K_{p_2}}{K_{p_1}} \quad \frac{E}{2.303\,R} \quad \frac{T_1 \quad T_2}{T_1 T_2}$ (d) None of these

118. For a reaction, the value of K_p increases with increase in temperature. The H for the reaction would be :

(a) positive (b) negative

(c) zero (d) cannot be predicted

119. The most stable oxides of nitrogen will be :

(a) $2NO_2(g) \rightleftharpoons N_2(g) \quad 2O_2(g); \qquad K \quad 6.7 \quad 10^{16}\ \text{mol L}^{1}$

(b) $2N_2O_5(g) \rightleftharpoons 2N_2(g) \quad 5O_2(g); \qquad K \quad 1.2 \quad 10^{24}\ \text{mol}^5\ \text{L}^{5}$

(c) $2NO(g) \rightleftharpoons N_2(g) \quad O_2(g); \qquad K \quad 2.2 \quad 10^{30}$

(d) $2N_2O(g) \rightleftharpoons 2N_2(g) \quad O_2(g); \qquad K \quad 3.5 \quad 10^{33}\ \text{mol L}^{1}$

120. When 1 mole of pure ethyl alcohol (C_2H_5OH) is mixed with 1 mole of acetic acid at 25°C, the equilibrium mixture contains 2/3 mole each of ester and water

$$C_2H_5OH(l) \quad CH_3COOH(l) \rightleftharpoons CH_3COOC_2H_5(l) \quad H_2O(l)$$

The G for the reaction at 298 K is :

(a) 3435 J (b) 4 J (c) 3435 J (d) Zero

121. The value of G for a reaction in aqueous phase having $K_c \quad 1$, would be :

(a) RT (b) 1 (c) 0 (d) RT

122. A plot of Gibbs energy of a reaction mixture against the extent of the reaction is :

(a) minimum at equilibrium (b) zero at equilibrium

(c) maximum at equilibrium (d) None of these

123. For the reaction at 300 K

$$A(g) \rightleftharpoons V(g) \quad S(g)$$

$$_rH \quad 30\ \text{kJ/mol}, \quad _rS \quad 0.1\,\text{kJ.K}^{1}\text{.mol}^{1}$$

What is the value of equilibrium constant?

(a) 0 (b) 1 (c) 10 (d) None of these

124. Solid $Ca(HCO_3)_2$ decomposes as

$$Ca(HCO_3)_2(s) \rightleftharpoons CaCO_3(s) \quad CO_2(g) \quad H_2O(g)$$

If the total pressure is 0.2 bar at 420 K, what is the standard free energy change for the given reaction ($_rG$)?

(a) 840 kJ/mol (b) 3.86 kJ/mol (c) 6.98 kJ/mol (d) 16.083 kJ/mol

125. The standard free energy change of a reaction is $G \quad 115$ kJ mol ^{1} at 298 K. Calculate the value of $\log_{10} K_p$ ($R \quad 8.314$ JK ^{1} mol ^{1})

(a) 20.16 (b) 2.303 (c) 2.016 (d) 13.83

For More Material Join: @JEEAdvanced_2024

Level 2

1. The following equilibrium constants were determined at 1120 K :

$$2CO(g) \rightleftharpoons C(s) \quad CO_2(g); \qquad K_{p_1} \quad 10^{14} \text{ atm}^{1}$$

$$CO(g) \quad Cl_2(g) \rightleftharpoons COCl_2(g); \qquad K_{p_2} \quad 6 \quad 10^{3} \text{ atm}^{1}$$

What is the equilibrium constant K_c for the following reaction at 1120 K :

$$C(s) \quad CO_2(g) \quad 2Cl_2(g) \rightleftharpoons 2COCl_2(g)$$

(a) $3.31 \quad 10^{11} \, M^{1}$ (b) $5.5 \quad 10^{10} \, M^{1}$ (c) $5.51 \quad 10^{6} \, M^{1}$ (d) None of these

2. One mole of $N_2(g)$ is mixed with 2 moles of $H_2(g)$ in a 4 litre vessel. If 50% of $N_2(g)$ is converted to $NH_3(g)$ by the following reaction :


$$N_2(g) \quad 3H_2(g) \rightleftharpoons 2NH_3(g)$$

What will be the value of K_c for the following equilibrium?

$$NH_3(g) \rightleftharpoons \frac{1}{2}N_2(g) \quad \frac{3}{2}H_2(g)$$

(a) 256 (b) 16 (c) $\frac{1}{16}$ (d) None of these

3.



The gas A_2 in the left flask allowed to react with gas B_2 present in right flask as $A_2(g) \quad B_2(g) \rightleftharpoons 2AB(g); K_c \quad 4$ at 27°C. What is the concentration of AB when equilibrium is established?

(a) 1.33 M (b) 2.66 M (c) 0.66 M (d) 0.33 M

4. Assume that the decomposition of HNO_3 can be represented by the following equation

$$4HNO_3(g) \rightleftharpoons 4NO_2(g) \quad 2H_2O(g) \quad O_2(g)$$

and the reaction approaches equilibrium at 400 K temperature and 30 atm pressure. At equilibrium partial pressure of HNO_3 is 2 atm.

Calculate K_c in $(\text{mol/L})^3$ at 400 K :

(Use : $R \quad 0.08$ atm-L/mol-K)

(a) 4 (b) 8 (c) 16 (d) 32

5. For the equilibrium

$$LiCl \quad 3NH_3(s) \rightleftharpoons LiCl \quad NH_3(s) \quad 2NH_3(g);$$

$K_p \quad 9 \text{ atm}^2$ at 37°C. A 5 litre vessel contains 0.1 mole of $LiCl \quad NH_3$. How many moles of NH_3 should be added to the flask at this temperature to derive the backward reaction for completion?

Use : $R \quad 0.082$ atm-L/mol K

(a) 0.2 (b) 0.59 (c) 0.69 (d) 0.79

For More Material Join: @JEEAdvanced_2024

6. Ammonium carbamate dissociates as

$$NH_2COONH_4(s) \rightleftharpoons 2NH_3(g) \quad CO_2(g)$$

In a closed vessel containing ammonium carbamate in equilibrium, ammonia is added such that partial pressure of NH_3 now equals to the original total pressure. Calculate the ratio of partial pressure of CO_2 now to the original partial pressure of CO_2 :

(a) 4 (b) 9 (c) $\frac{4}{9}$ (d) $\frac{2}{9}$

7. For the reaction $C_2H_6(g) \rightleftharpoons C_2H_4(g) \quad H_2(g)$

K_p is 5 10 ^{2} atm. Calculate the mole per cent of $C_2H_6(g)$ at equilibrium if pure C_2H_6 at 1 atm is passed over a suitable catalyst at 900 K :

(a) 20 (b) 33.33 (c) 66.66 (d) None of these

8. $2NOBr(g) \rightleftharpoons 2NO(g) \quad Br_2(g)$. If nitrosyl bromide (NOBr) is 40% dissociated at certain temp. and a total pressure of 0.30 atm. K_p for the reaction $2NO(g) \quad Br_2(g) \rightleftharpoons 2NOBr(g)$ is :

(a) 45 (b) 25 (c) 0.022 (d) 0.25

9. Consider the partial decomposition of A as

$$2A(g) \rightleftharpoons 2B(g) \quad C(g)$$

At equilibrium 700 mL gaseous mixture contains 100 mL of gas C at 10 atm and 300 K. What is the value of K_p for the reaction?

(a) $\frac{40}{7}$ (b) $\frac{1}{28}$ (c) $\frac{10}{28}$ (d) $\frac{28}{10}$

10. At a certain temperature and 2 atm pressure equilibrium constant (K_p) is 25 for the reaction

$$SO_2(g) \quad NO_2(g) \rightleftharpoons SO_3(g) \quad NO(g)$$

Initially if we take 2 moles of each of the four gases and 2 moles of inert gas, what would be the equilibrium partial pressure of NO_2?

(a) 1.33 atm (b) 0.1665 atm (c) 0.133 atm (d) None of these

11. 0.020 g of selenium vapour at equilibrium occupying a volume of 2.463 mL at 1 atm and 27°C. The selenium is in a state of equilibrium according to reaction

$$3Se_2(g) \rightleftharpoons Se_6(g)$$

What is the degree of association of selenium?

(At. mass of Se 79)

(a) 0.205 (b) 0.315 (c) 0.14 (d) None of these

12. Determine the degree of association (polymerization) for the reaction in aqueous solution, if observed (mean) molar mass of HCHO and $C_6H_{12}O_6$ is 150 :

$$6HCHO \rightleftharpoons C_6H_{12}O_6$$

(a) 0.50 (b) 0.833 (c) 0.90 (d) 0.96

13. A reaction system in equilibrium according to reaction $2SO_2(g) \quad O_2(g) \rightleftharpoons 2SO_3(g)$ in one litre vessel at a given temperature was found to be 0.12 mole each of SO_2 and SO_3 and 5 mole of O_2. In another vessel of one litre contains 32 g of SO_2 at the same temperature. What mass of O_2 must be added to this vessel in order that at equilibrium 20% of SO_2 is oxidized to SO_3?

(a) 0.4125 g (b) 11.6 g (c) 1.6 g (d) None of these

14. The equilibrium constant K_p for the following reaction is 4.5

$$N_2O_4(g) \rightleftharpoons 2NO_2(g)$$

For More Material Join: @JEEAdvanced_2024

What would be the average molar mass (in g/mol) of an equilibrium mixture of N_2O_4 and NO_2 formed by the dissociation of pure N_2O_4 at a total pressure of 2 atm?

(a) 69 (b) 57.5 (c) 80.5 (d) 85.5

15. A flask containing 0.5 atm pressure of $A_2(g)$, some solid AB added into flask which undergoes dissociation according to $2AB(s) \rightleftharpoons A_2(g) \quad B_2(g) \quad K_p \quad 0.06\ \text{atm}^2$

The total pressure (in atm) at equilibrium is :

(a) 0.70 (b) 0.6 (c) 0.10 (d) None of these

16. A vessel of 250 litre was filled with 0.01 mole of Sb_2S_3 and 0.01 mole of H_2 to attain the equilibrium at 440°C as

$$Sb_2S_3(s) \quad 3H_2(g) \rightleftharpoons 2Sb(s) \quad 3H_2S(g)$$

After equilibrium, the H_2S formed was analysed by dissolved it in water and treating with excess of Pb^{2} to give 1.19 g of PbS as precipitate. What is the value of K_c at 440°C?

(a) 1 (b) 2 (c) 4 (d) 8

17. For the reaction $2A(g) \quad B(g) \rightleftharpoons C(g) \quad D(g)$; $K_c \quad 10^{12}$. If the initial moles of A, B, C and D are 2, 1, 7 and 3 moles respectively in a one litre vessel, what is the equilibrium concentration of A?

(a) $4 \quad 10^{4}$ (b) $2 \quad 10^{4}$ (c) 10^{4} (d) $8 \quad 10^{4}$

18. The equilibrium constant for the following reaction in aqueous solution is 0.90.

$$H_3BO_3 + \text{glycerin} \rightleftharpoons (H_3BO_3 \quad \text{glycerin})$$

How many moles of glycerin should be added per litre of 0.10 M H_3BO_3 so that 80% of the H_3BO_3 is converted to the boric-acid-glycerin complex?

(a) 4.44 (b) 4.52 (c) 3.6 (d) 0.08

19. Rate of diffusion of ozonized oxygen is $0.4\sqrt{5}$ times that of pure oxygen. What is the per cent degree of association of oxygen assuming pure O_2 in the sample initially?

(a) 20 (b) 40 (c) 60 (d) None of these

20. One mole of SO_3 was placed in a two litre vessel at a certain temperature. The following equilibrium was established in the vessel

$$2SO_3(g) \rightleftharpoons 2SO_2(g) \quad O_2(g)$$

The equilibrium mixture reacts with 0.2 mole $KMnO_4$ in acidic medium. Hence, K_c is :

(a) 0.50 (b) 0.25

(c) 0.125 (d) None of these

21. At 800°C, the following equilibrium is established as

$$F_2(g) \rightleftharpoons 2F(g)$$

The composition of equilibrium may be determined by measuring the rate of effusion of the mixture through a pin hole. It is found that at 800°C and 1 atm mixture effuses 1.6 times as fast as SO_2 effuses under the similar conditions. (At. mass of F 19). What is the value of K_p (in atm)?

(a) 0.315 (b) 0.685

(c) 0.46 (d) 1.49

22. The equilibrium constant for the ionization of $RNH_2(g)$ in water as

$$RNH_2(g) \quad H_2O(l) \rightleftharpoons RNH_3^{+}(aq) \quad OH \quad (aq)$$

For More Material Join: @JEEAdvanced_2024

is 8 10 ^{6} at 25°C. Find the pH of a solution at equilibrium when pressure of $RNH_2(g)$ is 0.5 bar :

(a) 12.3 (b) 11.3 (c) 11.45 (d) None

23. Calculate $_rG$ for the reaction at 27°C

$$H_2(g) \quad 2Ag \ (aq) \rightleftharpoons 2Ag(s) \quad 2H \ (aq)$$

Given : P_{H_2} 0.5 bar; [Ag] 10 ^{5} M;

[H] 10 ^{3} M; $_rG$ [Ag (aq)] 77.1 kJ/mol

(a) 154.2 kJ/mol (b) 178.9 kJ/mol

(c) 129.5 kJ/mol (d) None of these

24. When N_2O_5 is heated at certain temperature, it dissociates as $N_2O_5(g) \rightleftharpoons N_2O_3(g) + O_2(g)$; K_c 2.5. At the same time N_2O_3 also decomposes as : $N_2O_3(g) \rightleftharpoons N_2O(g) + O_2(g)$. If initially 4.0 moles of N_2O_5 are taken in 1.0 litre flask and allowed to dissociate. Concentration of O_2 at equilibrium is 2.5 M. Equilibrium concentration of N_2O_5 is :

(a) 1.0 M (b) 1.5 M (c) 2.166 M (d) 1.846 M

25. Two solid compounds X and Y dissociates at a certain temperature as follows

$$X(s) \rightleftharpoons A(g) \quad 2B(g); \ K_{p_1} \quad 9 \quad 10^{-3} \text{ atm}^3$$

$$Y(s) \rightleftharpoons 2B(g) \quad C(g); \ K_{p_2} \quad 4.5 \quad 10^{-3} \text{ atm}^3$$

The total pressure of gases over a mixture of X and Y is :

(a) 4.5 atm (b) 0.45 atm (c) 0.6 atm (d) None of these

For More Material Join: @JEEAdvanced_2024

Level 3

Passage 1

For a gaseous reaction

$$aA(g) \quad bB(g) \rightleftharpoons cC(g) \quad dD(g)$$

equilibrium constants K_c, K_p and K_x are represented by the following relations

$$K_c \quad \frac{[C]^c\,[D]^d}{[A]^a\,[B]^b}; \quad K_p \quad \frac{p_C^c \;\; p_D^d}{p_A^a \;\; p_B^b} \text{ and } K_x \quad \frac{x_C^c \;\; x_D^d}{x_A^a \;\; x_B^b}$$

where $[A]$ represents molar concentration of A, p_A represents partial pressure of A and P represents total pressure, x_A represents mole fraction of A

1. On the basis of above work-up, select the write option

(a) $K_p \quad K_c\,(RT)^{\,n_g}; K_x \quad K_p\,(RT)^{\,n_g}$ (b) $K_c \quad K_p\,(RT)^{\,n_g}; K_p \quad K_x P^{\,n_g}$

(c) $K_c \quad K_x P^{\,n_g}; K_p \quad K_x P^{\,n_g}$ (d) $K_c \quad K_p\,(RT)^{\,n_g}; K_x \quad K_p\,(RT)^{\,n_g}$

2. For the reaction $SO_2Cl_2(g) \rightleftharpoons SO_2(g) \quad Cl_2(g)$, $K_p \quad K_x$ is obtained at :

(a) 0.5 atm (b) 0.8 atm (c) 1 atm (d) 2 atm

3. For the following equilibrium relation between K_c and K_x (in terms of mole fraction) is

$$PCl_3(g) \quad Cl_2(g) \rightleftharpoons PCl_5(g)$$

(a) $K_c \quad K_x(RT)^{\,1}$ (b) $K_c \quad K_x(RT)$ (c) $K_c \quad K_x \;\; \frac{RT}{P}$ (d) $K_c \quad K_x \;\; \frac{P}{RT}$

Passage 2

Variation of equilibrium constant K with temperature is given by van't Hoff equation

$$\ln K \quad \frac{{}_rS^\circ}{R} \quad \frac{{}_rH^\circ}{RT}$$

from this equation, ${}_rH^\circ$ can be evaluated if equilibrium constants K_1 and K_2 at two temperature T_1 and T_2 are known.

$$\log \frac{K_2}{K_1} \quad \frac{{}_rH^\circ}{2.303\,R} \;\; \frac{1}{T_1} \quad \frac{1}{T_2}$$

1. For an isomerization $X(g) \rightleftharpoons Y(g)$, the temperature dependency of equilibrium constant is given by :

$$\ln K \quad 2 \quad \frac{1000}{T}$$


The value of ${}_rS$ at 300 K is :

(a) $2R$ (b) $\frac{2}{R}$ (c) $1000\,R$ (d) None of these

For More Material Join: @JEEAdvanced_2024

2. Select the correct statement :
(a) Value of K_{eq} always increases with increasing temperature
(b) For exothermic reaction value of K_{eq} increases with decreasing in temperature
(c) For endothermic reaction value of K_{eq} increases with decreasing in temperature
(d) For exothermic reaction slope is ($\log K$ *vs*. $1/T$) negative

3. Variation of $\log_{10} K$ with $\frac{1}{T}$ is shown by the following graph in which straight line is at 45° hence H is :



(a) 4.606 kJ/mol (b) 19.147 kJ/mol (c) 8.314 kJ/mol (d) 10 kJ/mol

4. The equilibrium constant K_p for the following reaction is 1 at 27°C and 4 at 47°C.

$$A(g) \rightleftharpoons B(g) \quad C(g)$$

For the reaction calculate enthalpy change for the

$B(g) \quad C(g) \rightleftharpoons A(g)$ (Given : R 2 cal/mol-K)

(a) – 13.31 kcal/mol (b) 13.31 kcal/mol (c) – 19.2 kcal/mol (d) 55.63 kcal/mol



Passage 3

N_2O_3 is an unstable oxide of nitrogen and it decomposes into $NO(g)$ and $NO_2(g)$ where $NO_2(g)$ is further dimerise into N_2O_4 as

(i) $N_2O_3(g) \rightleftharpoons NO_2(g) \quad NO(g)$; K_{p_1} 2.5 bar

(ii) $2NO_2(g) \rightleftharpoons N_2O_4(g)$; K_{p_2}

A flask is initially filled with pure $N_2O_3(g)$ having pressure 2 bar and equilibria was established.

At equilibrium partial pressure of $NO(g)$ was found to be 1.5 bar.

1. The equilibrium partial pressure of $N_2O_3(g)$ is:
(a) 0.5 bar (b) 1.0 bar (c) 1.5 bar (d) 0.1 bar

2. The equilibrium partial pressure of $NO_2(g)$ is:
(a) 0.066 bar (b) 0.133 bar (c) 0.423 bar (d) 0.83 bar

3. The value of K_{p_2} is:
(a) 0.16 bar [1] (b) 0.32 bar [1] (c) 0.48 bar [1] (d) 0.64 bar [1]

For More Material Join: @JEEAdvanced_2024



Passage 4

If a system at equilibrium is subjected to a change of any one of the factors such as concentration, pressure or temperature, the system adjusts itself in such a way so as to minimise the effect of that change.

Effect of change in concentration on equilibrium:

As we add or remove reactant (or product) the ratio of equilibrium concentration become 'Q' (reaction quotient) and depending upon.

Q K : equilibrium will shift in forward direction

Q K : equilibrium will shift in backward direction

Effect of change in pressure :

If a system in equilibrium consists of gases, then the concentrations of all the components can be altered by changing the pressure. When the pressure on the system is increased, then equilibrium will shift in the direction in which there is decrease in number of moles *i.e.,* towards the direction in which there is decrease in volume.

Effect of change in pressure on melting point : There are two types of solids :

(a) Solids whose volume decreases on melting, *e.g.,* ice, diamond, carborundum, magnesium nitride and quartz.

Solid (higher volume) $\rightleftharpoons$ Liquid (lower volume)

The process of melting is facilitated at high pressure, thus melting point is lowered.

(b) Solid whose volume increase on melting, *e.g.,* Fe, Cu, Ag, Au, etc.

Solid (lower volume) $\rightleftharpoons$ Liquid (higher volume)

In this case the process of melting becomes difficult at high pressure, thus melting point becomes high.

(c) **Solubility of substances :** When solid substance are dissolved in water, either heat is evolved.

For endothermic solubility process solubility increase with increase in temperature.

For exothermic solubility decrease with increase in temperature.

(d) **Solubility of gases in liquids :** When a gas dissolves in liquid, there is decrease in volume. Thus increase of pressure will favour the dissolution of gas in liquid.

Effect of temperature : For endothermic reaction as temperature increases reaction shift in forward direction. For exothermic reaction as temperature increases reaction shift in backward direction.

1. A 'X' (g) solute when dissolved in water heat is evolved. Then solubility of 'X' will increase :

(a) high temperature, low pressure
(b) low temperature, high pressure
(c) high temperature, high pressure
(d) low temperature, low pressure

2. $Fe(l) \rightleftharpoons Fe(s)$

Above equilibrium is favoured at :

(a) high pressure, low temperature
(b) high pressure, high temperature
(c) low pressure, high temperature
(d) low pressure, low temperature

For More Material Join: @JEEAdvanced_2024

3. For the reaction

$N_2(g) \quad O_2(g) \rightleftharpoons 2NO(g)$

If pressure is increased by reducing the volume of the container then :

(a) total pressure at equilibrium will remain same

(b) concentration of all the component at equilibrium will change

(c) concentration of all the component at equilibrium will remain same

(d) equilibrium will shift in the backward direction

One or More Answers is/are Correct



1. A catalyst :

(a) increases the average kinetic energy of reacting molecules

(b) decreases the activation energy

(c) can alters the reaction mechanism

(d) can change pre-exponential factor

2. Which of the following is correct about the chemical equilibrium?

(a) $(\quad G)_{T,P} \quad 0$

(b) Equilibrium constant is independent of initial concentration of reactants

(c) Catalyst has no effect on equilibrium state

(d) Reaction stops at equilibrium

3. For the reaction

$$AB_2(g) \rightleftharpoons AB(g) \quad B(g)$$

If is negligiable w.r.t. 1 then degree of dissociation ( ) of AB_2 is proportional to :

(a) $\frac{1}{P}$ (b) $\frac{1}{V}$ (c) $\frac{1}{\sqrt{P}}$ (d) $\sqrt{V}$

4. Consider the reactions given below. In which cases will the reaction proceed toward right by increasing the pressure?

(a) $4HCl(g) \quad O_2(g) \quad 2Cl_2(g) \quad 2H_2O(g)$

(b) $Cl_2(g) \quad H_2O(g) \quad 2HCl(g) \quad \frac{1}{2}O_2(g)$

(c) $CO_2(g) \quad 4H_2(g) \quad CH_4(g) \quad 2H_2O(g)$

(d) $N_2(g) \quad O_2(g) \quad 2NO(g)$

5. Ammonia is a weak base that reacts with water according to the equation

$$NH_3(aq) \quad H_2O(l) \rightleftharpoons NH_4(aq) \quad OH \quad (aq)$$

Select the correct option(s) that can increase the moles of ammonium ion in water:

(a) Addition of HCl (b) Addition of NaOH

(c) Addition of NH_4Cl (d) Addition of H_2O

6. Consider the reaction $2CO(g) \quad O_2(g) \rightleftharpoons 2CO_2(g) \quad$ Heat

Under what conditions shift is undeterminable?

(a) Addition of O_2 and decrease in volume

(b) Addition of CO and removal of CO_2 at constant volume

For More Material Join: @JEEAdvanced_2024

(c) Increase in temperature and decrease in volume
(d) Addition of CO and increase in temperature at constant volume

7. What will be the effect of addition of catalyst at constant temperature?
(a) The equilibrium constant will remain constant
(b) H of the reaction will remain constant
(c) k_f and k_b will increase upto same extent
(d) equilibrium composition will change

8. For the reaction $PCl_5(g) \rightleftharpoons PCl_3(g)$ $Cl_2(g)$, the forward reaction at constant temperature is favoured by:
(a) introducing an inert gas at constant volume
(b) introducing chlorine gas at constant volume
(c) introducing an inert gas at constant pressure
(d) increasing the volume of the container

9. For the reaction : $Cl_2(g)$ $3F_2(g) \rightleftharpoons 2ClF_3(g)$; H 329 kJ, dissociation of $ClF_3(g)$ will be favoured by :
(a) increasing the temperature
(b) increasing the volume of the container
(c) adding of F_2 gas
(d) adding of inert gas at constant pressure

10. Increase in the pressure for the following equilibrium results in the:

$$H_2O(l) \rightleftharpoons H_2O(g)$$

(a) formation of more $H_2O(l)$ (b) formation of more $H_2O(g)$
(c) increase in b.p. of $H_2O(l)$ (d) decrease in b.p. of $H_2O(l)$

11. Heating a II group metal carbonate leads to decomposition as :

$$BaCO_3(s) \rightleftharpoons BaO(s) \quad CO_2(g)$$

Equilibrium will shift left
(a) by addition of BaO(s) (b) by addition of $CO_2(g)$
(c) by decreasing the temperature (d) by decreasing the volume of the vessel

12. $N_2(g)$ and $H_2(g)$ are allowed to react in a closed vessel at given temp. and pressure for the formation of $NH_3(g)$ [$N_2(g)$ $3H_2(g) \rightleftharpoons 2NH_3(g)$ 22.4 kcal]. If $He(g)$ is added at equilibrium at constant pressure then which is/are correct ?
(a) Concentration of $N_2(g)$, $H_2(g)$ and $NH_3(g)$ decrease.
(b) Moles of $NH_3(g)$ decreases.
(c) The extent of cooling depends on amount of $He(g)$ added.
(d) Concentration of N_2 and H_2 increases and concentration of NH_3 decreases.

For More Material Join: @JEEAdvanced_2024

Match the Column



Column-I and **Column-II** contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

1.

Column-I	Column-II
(A) $CaCO_3(s) \rightleftharpoons CaO(s) \quad CO_2(g)$	(P) $K_p \quad K_c$ above room temperature
(B) $CO(g) \quad Cl_2(g) \rightleftharpoons COCl_2(g)$	(Q) $K_p \quad K_c$ above room temperature
(C) $H_2(g) \quad I_2(g) \rightleftharpoons 2HI(g)$	(R) $K_p \quad K_c$ above room temperature
(D) $HCl(g) \rightleftharpoons H \ (aq) \quad Cl \ (aq)$	(S) K_p and K_c not defined

2.

Column-I	Column-II
(A) $3O_2(g) \rightleftharpoons 2O_3(g)$	(P) no unit
(B) $SO_2(g) \quad \frac{1}{2}O_2(g) \rightleftharpoons SO_3(g)$	(Q) atm $^{1/2}$
(C) $2HF(g) \rightleftharpoons H_2(g) \quad F_2(g)$	(R) atm ^{1}
(D) $CO(g) \quad 3H_2(g) \rightleftharpoons CH_4(g) \quad H_2O(g)$	(S) atm ^{2}

3.

Column-I (Reaction)	Column-II (If is negligiable w.r.t. 1)
(A) $2X(g) \rightleftharpoons Y(g) \quad Z(g)$	(P) $2\sqrt{K_c}$
(B) $X(g) \rightleftharpoons Y(g) \quad Z(g)$	(Q) $3\sqrt{K_c}$
(C) $3X(g) \rightleftharpoons Y(g) \quad Z(g)$	(R) $(2K_c)^{1/3}$
(D) $2X(g) \rightleftharpoons Y(g) \quad 2Z(g)$	(S) $\sqrt{K_c}$

4.

Column-I	Column-II
(A) $N_2(g) \quad 3H_2(g) \rightleftharpoons 2NH_3(g); \ H$ ve	(P) K increases with increase in temperature
(B) $N_2(g) \quad O_2(g) \rightleftharpoons 2NO(g); \ H$ ve	(Q) K decreases with increase in temperature
(C) $A(g) \quad B(g) \rightleftharpoons 2C(g) \quad D(g); \ H$ ve	(R) Pressure has no effect
(D) $PCl_5(g) \rightleftharpoons PCl_3(g) \quad Cl_2(g); \ H$ ve	(S) Moles of product increase due to addition of inert gas at constant pressure

For More Material Join: @JEEAdvanced_2024



5.

Column-I	Column-II
(A) $\frac{K_{10\ T\ C}}{K_{T\ C}}$ 2	(P) Endothermic
(B) $\frac{K_{10\ T\ C}}{K_{T\ C}}$ $\frac{1}{2}$	(Q) Not affected by pressure
(C) $A(g)$ $B(g) \rightleftharpoons C(g)$	(R) Exothermic
(D) $X(s)$ $Y(g) \rightleftharpoons Z(g)$	(S) Affected by volume


6.


Column-I	Column-II
(A) Pressure increased in $2NO(g) \rightleftharpoons N_2(g)$ $O_2(g)$	(P) Equilibrium shifted in forward direction
(B) Pressure increased in $CH_4(g)$ $H_2O(g) \rightleftharpoons CO(g)$ $3H_2(g)$	(Q) Equilibrium shifted in backward direction
(C) Temp. increased and pressure increased $3O_2(g) \rightleftharpoons 2O_3(g)$; H 285 kJ	(R) Equilibrium remains unaffected
(D) Pressure decreased and moles of N_2 increased $N_2(g)$ $2O_2(g) \rightleftharpoons 2NO_2(g)$; H 66.4 kJ	(S) Theoretically we cannot predict

7.

Column-I (Reactions)	Column-II (Equilibrium States)
(A) (A)  $(g) \rightleftharpoons$  (g)	(P) 
(B) $I_2(s) \rightleftharpoons I_2(aq)$	(Q) 

For More Material Join: @JEEAdvanced_2024

(C) $N_2O_4(g) \rightleftharpoons 2NO_2(g)$ (R) 

(D) $AgI(s) \rightleftharpoons Ag^+(aq) + I^-(aq)$ (S) 

Assertion-Reason Type Questions

Each question contains STATEMENT-1 (Assertion) and STATEMENT-2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below :

(A) If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1

(B) If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1

(C) If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

(D) If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

1. STATEMENT-1 : The endothermic reactions are favoured at lower temperature and the exothermic reactions are favoured at higher temperature.

STATEMENT-2 : When a system in equilibrium is disturbed by changing the temperature, it will tend to adjust itself so as to overcome the effect of change.

2. STATEMENT-1 : The melting point of ice decreases with increase of pressure.

STATEMENT-2 : Ice contracts on melting.

3. STATEMENT-1 : The equilibrium of $A(g) \rightleftharpoons B(g)$ $C(g)$ is not affected by changing the volume.

STATEMENT-2 : K_c for the reaction does not depend on volume of the container.

4. STATEMENT-1 : For a chemical reaction at initial stage rate of forward reaction (r_f) is greater than rate of reversed reaction (r_b)

STATEMENT-2 : When r_f r_b, chemical reaction is at equilibrium.

5. STATEMENT-1 : For the reaction $A(g) \rightleftharpoons B(g)$ $C(g)$, K_p 1 atm. If we start with equal moles of all gases at 9 atm of initial pressure, then at equilibrium partial pressure of A increases.

STATEMENT-2 : Reaction quotient Q_p K_p hence equilibrium shifts in backward direction.

For More Material Join: @JEEAdvanced_2024

6. STATEMENT-1 : The gas phase reaction $PCl_3(g)$ $Cl_2(g) \rightleftharpoons PCl_5(g)$ shifts to the right on increasing pressure.

STATEMENT-2 : When pressure increases, equilibrium shifts towards more number of moles.

7. STATEMENT-1 : For a reaction at equilibrium, the Gibb's free energy of reaction is minimum at constant temp. and pressure.

STATEMENT-2 : The Gibb's free energy of both reactants and products increases and become equal at equilibrium.

8. STATEMENT-1 : The physical equilibrium is not static but dynamic in nature.

STATEMENT-2 : The physical equilibrium is a state in which two opposing process are proceeding at the same rate.

9. STATEMENT-1 : Equilibrium constant for the reverse reaction is the inverse of the equilibrium constant for the reaction in the forward direction.

STATEMENT-2 : Eqilibrium constant depends upon the way in which the reaction is written.

10. STATEMENT-1 : If Q_p K_p reaction moves in direction of reactants.

STATEMENT-2 : Reaction quotient is defined in the same way as equilibrium constant at any stage of the reaction.

11. STATEMENT-1 : For the reaction $H_2(g)$ $I_2(g) \rightleftharpoons 2HI(g)$ if the volume of vessel is reduced to half of its original volume, equilibrium concentration of all gases will be doubled.

STATEMENT-2 : According to Le-Chatelier's principle, reaction shifts in a direction that tends to minimized the effect of the stress.

12. STATEMENT-1 : The equilibrium constant of the exothermic reaction at high temperature decreases.

STATEMENT-2 : Since $\ln \frac{K_2}{K_1}$ $\frac{H}{R}$ $\frac{1}{T_1}$ $\frac{1}{T_2}$ and for exothermic reaction,

H ve and thereby; $\frac{K_2}{K_1}$ 1

13. STATEMENT-1 : For the reaction at certain temperature

$$A(g) \quad B(g) \rightleftharpoons C(g)$$

there will be no effect by addition of inert gas at constant volume.

STATEMENT-2 : Molar concentration of all gases remains constant.

14. STATEMENT-1 : For the physical equilibrium $H_2O(s) \rightleftharpoons H_2O(l)$ on increasing temperature and increasing pressure more water will form.

STATEMENT-2 : Since forward reaction is endothermic in nature and volume of water is greater than that of the volume of ice.

15. STATEMENT-1 : The catalyst does not alter the equilibrium constant.

STATEMENT-2 : Because for the catalysed reaction and uncatalysed reaction H remains same and equilibrium constant depends on H.

For More Material Join: @JEEAdvanced_2024

Subjective Problems

1. If 50% of CO_2 converts to CO at the following equilibrium :

$$\frac{1}{2}C(s) \quad \frac{1}{2}CO_2(g) \rightleftharpoons CO(g)$$

and the equilibrium pressure is 12 atm. Calculate K_p.

2. Calculate partial pressure of B at equilibrium in the following equilibrium

$$A(s) \rightleftharpoons B(g) \quad 2C(g) \; ; \; K_p \quad 32\,\text{atm}^3.$$

3. In a gaseous reaction $A \quad 2B \rightleftharpoons 2C \quad D$ the initial concentration of B was 1.5 times that of A. At equilibrium the concentrations of A and D were equal. Calculate the equilibrium constant K_C.

4. For the reaction $A(g) \rightleftharpoons B(g); \quad K_C \quad 10$

$$B(g) \rightleftharpoons C(g); \quad K_C \quad 2$$

$$C(g) \rightleftharpoons D(g) \; ; \; K_C \quad 0.01$$

Calculate K_C for the reaction $D(g) \rightleftharpoons A(g)$.

5. 5 litre vessel contains 2 moles of each of gases A and B at equilibrium. If 1 mole each of A and B are removed. Calculate K_C for the reaction $A(g) \rightleftharpoons B(g)$.

6. Calculate K_p for the reaction $A(g) \rightleftharpoons B(s) \quad 2C(g) \; ; \; K_C \quad 0.2$ at 305 K.

7. A mixture of 3 moles of SO_2, 4 moles of NO_2, 1 mole of SO_3 and 4 moles of NO is placed in a 2.0 L vessel. $SO_2(g) \quad NO_2(g) \rightleftharpoons SO_3(g) \quad NO(g)$.

At equilibrium, the vessel is found to contain 1 mole of SO_2. Calculate the value of K_C.

8. The density of an equilibrium mixture of N_2O_4 and NO_2 at 1 atm and 373.5 K is 2.0 g/L. Calculate K_C for the reaction $N_2O_4(g) \rightleftharpoons 2NO_2(g)$

9. If chemical equilibrium is attained at standard states then what is the value of G ?


10. Calculate the equilibrium concentration ratio of C to A if equimolar ratio of A and B were allowed to come to equilibrium at 300 K.

$A(g) \quad B(g) \rightleftharpoons C(g) \quad D(g) \; ; \quad G \quad 830$ cal.

11. A definite amount of solid NH_4HS is placed in a flask already containing ammonia gas at a certain temperature and 0.1 atm pressure. NH_4HS decomposes to give NH_3 and H_2S and at equilibrium total pressure in flask is 1.1 atm. If the equilibrium constant K_p for the reaction $NH_4HS(s) \rightleftharpoons NH_3(g) \quad H_2S(g)$ is represented as $z \quad 10^{-1}$ then find the value of z.

For More Material Join: @JEEAdvanced_2024

12. The gaseous reaction : $A(g) \quad nB(g) \rightleftharpoons mC(g)$ is represented by following curves



What is the value of $n \quad m$?

For More Material Join: @JEEAdvanced_2024

Answers

Level 1

1. (b)	2. (b)	3. (c)	4. (a)	5. (b)	6. (a)	7. (a)	8. (a)	9. (a)	10. (a)
11. (b)	12. (b)	13. (b)	14. (d)	15. (c)	16. (b)	17. (d)	18. (d)	19. (c)	20. (d)
21. (b)	22. (a)	23. (c)	24. (a)	25. (b)	26. (d)	27. (d)	28. (c)	29. (d)	30. (c)
31. (d)	32. (c)	33. (c)	34. (c)	35. (a)	36. (a)	37. (c)	38. (d)	39. (c)	40. (c)
41. (a)	42. (b)	43. (a)	44. (a)	45. (a)	46. (a)	47. (c)	48. (d)	49. (d)	50. (d)
51. (a)	52. (b)	53. (a)	54. (b)	55. (b)	56. (b)	57. (b)	58. (b)	59. (c)	60. (c)
61. (a)	62. (d)	63. (a)	64. (a)	65. (c)	66. (a)	67. (a)	68. (c)	69. (d)	70. (a)
71. (a)	72. (c)	73. (a)	74. (d)	75. (b)	76. (c)	77. (b)	78. (b)	79. (c)	80. (d)
81. (b)	82. (b)	83. (d)	84. (b)	85. (b)	86. (b)	87. (a)	88. (a)	89. (b)	90. (c)
91. (a)	92. (b)	93. (a)	94. (c)	95. (b)	96. (d)	97. (d)	98. (b)	99. (c)	100. (a)
101. (d)	102. (d)	103. (b)	104. (d)	105. (c)	106. (d)	107. (b)	108. (d)	109. (a)	110. (d)
111. (c)	112. (b)	113. (a)	114. (b)	115. (a)	116. (c)	117. (b)	118. (a)	119. (a)	120. (c)
121. (c)	122. (a)	123. (b)	124. (d)	125. (a)					

Level 2

1. (a)	2. (c)	3. (c)	4. (d)	5. (d)	6. (c)	7. (c)	8. (a)	9. (c)	10. (c)
11. (b)	12. (d)	13. (b)	14. (b)	15. (a)	16. (a)	17. (a)	18. (b)	19. (c)	20. (c)
21. (d)	22. (b)	23. (c)	24. (d)	25. (b)					

For More Material Join: @JEEAdvanced_2024

Level 3

Passage–1:	**1.** (b)	**2.** (d)	**3.** (c)				
Passage–2:	**1.** (a)	**2.** (b)	**3.** (b)	**4.** (a)			
Passage–3:	**1.** (a)	**2.** (d)	**3.** (c)				
Passage–4:	**1.** (b)	**2.** (a)	**3.** (b)				

One or More Answers is/are correct

1. (b,c,d) **2.** (a,b,c) **3.** (c,d) **4.** (a,c) **5.** (a,c,d) **6.** (c,d) **7.** (a,b,c) **8.** (c,d)
9. (a,b,d) **10.** (a,c) **11.** (b,c,d) **12.** (a,b,c)

Match the Column

1. A P; B R; C Q; D S
2. A R; B Q; C P; D S
3. A P; B S; C Q; D R
4. A Q; B P, R; C P, S; D P, S
5. A P; B R; C S; D Q
6. A R; B Q; C P; D S
7. A Q; B S; C P; D S

Assertion-Reason Type Questions

1. (D) **2.** (A) **3.** (D) **4.** (B) **5.** (A) **6.** (C) **7.** (C) **8.** (A) **9.** (A) **10.** (D)
11. (B) **12.** (A) **13.** (A) **14.** (C) **15.** (A)

Subjective Problems

1. 4	**2.** 2	**3.** 4	**4.** 5	**5.** 1	**6.** 5	**7.** 9	**8.** 2	**9.** 0	**10.** 2
11. 3	**12.** 5								

For More Material Join: @JEEAdvanced_2024

Hints and Solutions

Level 1

2. (b) $Q_C = \frac{[P_2(g)]^4}{[P_4(g)]} = \frac{(1)^2}{(3/2)} = \frac{2}{3}$

4. (a) $\because \frac{K}{Q} = 0.33$

$K < Q$

i.e., reaction moves in backward direction to achieve equilibrium.

5-6. (b,a)

$Q_C = \frac{[SO_3(g)]^2}{[SO_2(g)]^2[O_2(g)]^1} = 1000 > K_c$

i.e., reaction moves backward

also if volume increases pressure decreases

Q_c

7. (a) $Q_C = \frac{\left(\frac{6}{2}\right)^2}{\frac{2}{2}\left(\frac{4}{2}\right)^3} = \frac{9}{8}$

$Q_c > K_c$ so reaction will proceed in forward direction.

8. (a) Since D_2 has similar chemical properties as H_2 so upon mixing, NH_2D and NHD_2 are formed.

9. (a) $[N_2]$ and $[H_2]$ start from non-zero value, decrease & then become constant.

$[NH_3]$ start from zero, increase & then become constant.

10. (a) From graph, $[A]_{eq} = 0.1\,M, [B]_{eq} = 0.4\,M$

$K_c = \frac{[B]^2}{[A]^1} = 1.6$

11. (b) $A(g) \rightleftharpoons 2C(g) + B(g)$

$t = 0 \quad 10M$

$t_{eq} \quad (10 - 10\ ) \quad 2(10\ ) \quad (\ )$

From graph

$[A]_{eq} = 10(1 - \ ) = 8$

0.2

$K_c = \frac{[C]^2[B]^1}{[A]^1}$

$4\,(\text{mol/L})^2$

12. (b) $K_c = \frac{[CO]^1[H_2]^2}{[CH_3OH]^1}$

So units $= M^2$

13. (b) $K_p = (\text{atm})^{n_g}$

$(\text{atm})^{2}$

14. (d) Activity of solid component is unit.

15. (c) If, $K_c = \frac{[N_2]^{1/2}[H_2]^{3/2}}{[NH_3]^1} = 4$

then $\frac{[NH_3]^2}{[N_2][H_2]^3} = (4)^{-2} = \frac{1}{16}$

also $K_p = K_c (RT)^{n_g}$

16. (b) For, $N_2 + O_2 \rightleftharpoons 2NO$, $n_g = 0$

$K_c = K_p = K_x = 4 \times 10^{-4}$

So for $NO \rightleftharpoons \frac{1}{2}N_2 + \frac{1}{2}O_2$

$K_c = (4 \times 10^{-4})^{-1/2}$

17-21. (d, d,c, d, b)

$K_p = K_c (RT)^{n_g}$

24. (a) $\ln \frac{K_2}{K_1} = \frac{H}{R}\left(\frac{1}{T_1} - \frac{1}{T_2}\right)$

25-26. (b,d)

Equilibrium constant is temperature dependant only.

27. (d) $CO(g) + H_2O(g) \rightleftharpoons CO_2(g) + H_2(g)$; $\sqrt{K_1}$

$CH_4(g) + H_2O(g) \rightleftharpoons CO(g) + 3H_2(g)$; K_2

Addition

$CH_4(g) + 2H_2O(g) \rightleftharpoons CO_2(g) + 4H_2(g)$; $(\sqrt{K_1} \times K_2)$

$K_3 = \sqrt{K_1} \times K_2$

28-29. (c,d) Similar to Q no. 16

30-31. (c,d) Similar to Q. no. 27

32. (c) $X(g) + Y(g) \rightleftharpoons 2Z(g)$

$t = 0 \quad 0.2M \quad 0.1M \quad 0.1M$

$t_{eq} \quad (0.2 - x) \quad (0.1 - x) \quad (0.1 + 2x)$

$\because [Z] = (0.1 + 2x) = 0.2$

$x = 0.05$

Now, $K_c = \frac{[Z]^2}{[X]^1[Y]^1}$

For More Material Join: @JEEAdvanced_2024

33. (c) $K_c = \dfrac{[H_2]^2[S_2]}{[H_2S]}$

34. (c) $K_c = \dfrac{[CH_4][H_2S]^2}{[CS_2][H_2]^4}$

35. (a) $K_c = \dfrac{[CH_3OH]}{[CO][H_2]^2}$

36. (a) $K_p = \dfrac{P^4_{S_2(g)}}{P_{S_8(g)}} = \dfrac{(4 \quad 0.3)^4}{0.70} = 2.96$

37. (c) $N_2O_4(g) \rightleftharpoons 2NO_2(g)$

$t \quad 0 \qquad 0.1M$

$t_{eq.} \qquad 0.1(1 \quad) \qquad 2(0.1 \quad)$

$\because \quad 0.5$

$K_c = \dfrac{[NO_2]^2}{[N_2O_4]}$

38. (d) $2NH_3(g) \rightleftharpoons N_2(g) \quad 3H_2(g)$

$t \quad 0 \qquad 2$ moles

$t_{eq.} \qquad 2 \quad 2x \qquad x \qquad 3x$

Given : $(2 \quad 2x) \quad 1$

$x \quad 0.5$

Now $K_c = \dfrac{[N_2][H_2]^3}{[NH_3]^2}$

39. (c) $K_p = \dfrac{1}{P^4_{H_2O(g)}}$

$P_{H_2O(g)} = \left(\dfrac{1}{K_p}\right)^{1/4}$

$(10^{\quad 12})^{1/4} \quad 10^{\quad 3}$ atm

$K_p \quad 10^{\quad 3} \quad 760 \quad 0.76$ torr

40. (c) $P_{H_2O} \quad 2 \quad 10^{\quad 2}$ atm

R.H. $= \dfrac{2 \quad 10^{\quad 2} \quad 760}{38} \quad 100$

40%

41. (a) $NH_4HS(s) \rightleftharpoons NH_3(g) \quad H_2S(g)$

$t \quad 0 \qquad a$ moles

$t_{eq.} \qquad (a \quad x) \qquad x \qquad x$

$\because \quad P_{Gas} \quad X_{Gas} \quad P_{Total}$

$P_{NH_3} = \dfrac{1}{2} \quad 0.66 \quad 0.33$ atm

$P_{H_2S} = \dfrac{1}{2} \quad 0.66 \quad 0.33$ atm

Now $K_p \quad (P_{NH_3})^1(P_{H_2S})^1$

42. (b) $K_c = \dfrac{n_B n_C^3}{n_A^2} = \dfrac{1}{V^2}$

$16 = \dfrac{2}{2^2} \dfrac{2^3}{V^2}$

$V = \dfrac{1}{2}$

43. (a) $C_2H_5OH(l) \quad CH_3COOH(l) \rightleftharpoons$

$t \quad 0 \qquad 1$ mole $\qquad 1$ mole

$t_{eq.} \qquad (1 \quad x) \qquad (1 \quad x)$

$CH_3COOC_2H_5(l) \quad H_2O(l)$

$\quad - \qquad\qquad -$

$(x) \qquad\qquad (x)$

$x = \dfrac{1}{3}$

$K_c = \dfrac{[CH_3COOC_2H_5][H_2O]}{[C_2H_5OH][CH_3COOH]}$

44. (a) $I_2(aq.) \quad I^{\quad}(aq.) \rightleftharpoons I_3(aq.)$

$t \quad 0 \qquad 1$ mole $\quad 0.5$ mole $\qquad -$

$t_{eq.} \qquad (1 \quad x) \quad (0.5 \quad x) \qquad (x)$

$AgNO_3$ reacts with $I^{\quad}$ and forms precipitate.

i.e., $AgNO_3 \quad I^{\quad} \quad AgI(s) \quad NO_3(aq.)$

$(0.5 \quad x) \quad 0.25$

$x \quad 0.25$

$K_c = \dfrac{[I_3]}{[I_2][I^{\quad}]}$

45. (a) $K_c = \dfrac{[H_2S(g)]}{[H_2(g)]}$

$8 \quad 10^{\quad 2} = \dfrac{x}{0.3 \quad x}$

$x \quad 0.022$

$[H_2S(g)] = \dfrac{0.022}{2}$

$0.011M$

46. (a) $K_p = \dfrac{(P_{SO_3})^2}{(P_{SO_2})^2(P_{O_2})^1}$ will be same in both cases

47. (c) $P_{total} \quad 3P$

$P = \dfrac{0.318}{3} \quad 0.106$

$K_p \quad 4P^3 \quad 4.76 \quad 10^{\quad 3}$

48. (d) $K_c \quad [B]^2[C]^3$

and K_c remains constant.

49. (d) $A \qquad B \rightleftharpoons C \quad D$

$t \quad 0 \qquad a \qquad a$

$t_{eq.} \qquad (a \quad x) \quad (a \quad x) \qquad (x) \quad (x)$

$\because \quad [D] \quad 2[A]$

For More Material Join: @JEEAdvanced_2024

$x \quad 2(a \quad x)$

$x \quad 2a/3$

Now $K_c \quad \dfrac{[C][D]}{[A][B]}$

50. (d) $Q \quad 1 \quad K_{eq.}$

reaction moves forward

i.e., $SO_2 \quad NO_2 \rightleftharpoons SO_3 \quad NO$

$t \quad 0 \quad 1 \quad 1 \quad 1 \quad 1$

$t_{eq.} \quad (1 \quad x) \quad (1 \quad x) \quad (1 \quad x) \quad (1 \quad x)$

$K_c \quad \dfrac{[SO_3][NO]}{[SO_2][NO_2]} \quad 16$

$4 \quad \dfrac{1 \quad x}{1 \quad x}$

$x \quad 0.6 \qquad [NO] \quad 1 \quad x \quad 1.6\,M$

53. (a) $2HI(g) \rightleftharpoons H_2(g) \quad I_2(g)$

$t \quad 0 \quad a\,M$

$t_{eq.} \quad (a \quad a) \qquad \dfrac{a}{2} \qquad \dfrac{a}{2}$

Given : 0.5

So $K_{eq} \quad \dfrac{[H_2][I_2]}{[HI]^2}$

54. (b) $H_2(g) \quad CO_2(g) \rightleftharpoons H_2O(g) \quad CO(g)$

moles at eqm $0.8 \quad x \qquad 0.8 \quad x \qquad x \qquad x$

conc. at eqm $\dfrac{0.8 \quad x}{5} \qquad \dfrac{0.8 \quad x}{5} \qquad \dfrac{x}{5} \qquad \dfrac{x}{5}$

$\because \quad n_g \quad 0$

$$K_c \quad \frac{\left(\frac{x}{5}\right)^2}{\frac{0.8 \quad x}{5}}$$

$$2 \quad \frac{x}{0.8 \quad x}$$

$x \quad 0.533$

$[CO_2(g)] \quad \dfrac{0.8 \quad 0.533}{5}$

55. (b) For ideal gas mole % volume %

$N_2O_4(g) \rightleftharpoons 2NO_2(g)$

Initial moles $\quad a \qquad 0$

at eqm $\quad a(1 \quad) \qquad 2a$

As per given original volume

$\dfrac{75}{100}$ Volume at eqm

at constant T and P : $V \quad n$

$a \quad 0.75 \quad a(1 \quad)$

0.33

56. (b) $CO \quad H_2O \rightleftharpoons CO_2 + H_2$

Initial conc. $\quad 3 \qquad 3 \qquad x \qquad 0$

At eqm $\quad 2 \qquad 2 \qquad (x \quad 1) \qquad 1$

$5 \quad \dfrac{(x \quad 1)}{4} \quad 20 \quad x \quad 1$

$x \quad 19$

57. (b) $N_2(g) \quad 3H_2(g) \rightleftharpoons 2NH_3(g)$

$t \quad 0 \quad$ a mole $\qquad 3a$

$t_{eq.} \quad (a \quad x) \qquad (3a \quad 3x) \qquad (2x)$

$x \quad a/4$

$t_{eq.} \quad \dfrac{3a}{4}$ mole $\dfrac{9a}{4}$ mole $\dfrac{a}{2}$ mole

$P_{Gas} \quad X_{Gas} \quad P_{Total}$

58. (b) $2NH_3(g) \rightleftharpoons N_2(g) \quad 3H_2(g)$

$t \quad 0 \qquad a$

$t_{aq.} \quad (a \quad a) \qquad \dfrac{a}{2} \qquad \dfrac{3a}{2}$

$\because \quad PV \quad nRT$

at $t \quad 0\,; 15 \quad V \quad a.R.300$ ...(1)

at $t_{eq.}; 50 \quad V \quad (a \quad a).R.620$ ...(2)

(1) (2); $\dfrac{3}{10} \quad \dfrac{300}{(1 \quad).620}$

59. (c) $N_2O_4(g) \rightleftharpoons 2NO_2(g)$

$t \quad 0 \qquad 0.1$ mole $\qquad -$

$t \quad eq. \qquad (0.1 \quad x) \qquad (2x)$

$K_p \quad \dfrac{(P_{NO_2})^2}{(P_{N_2O_4})^1}$

$$0.14 \quad \frac{\left(\frac{2x}{0.1 \quad x}\right)^2 1}{\left(\frac{0.1 \quad x}{0.1 \quad x}\right)^1 1}$$

60. (c) $2\,SO_2 \qquad O_2 \rightleftharpoons 2\,SO_3$

$t \quad 0 \quad 5$ mole $\qquad 5$ mole

$t_{eq.} \quad 5 \quad 2x \qquad 5 \quad x \qquad 2x$

Given : $(2x) \quad \dfrac{60}{100}(5)$

$x \quad 1.5$

Now $P_{Gas} \quad X_{Gas} \quad P_{Total}$

61. (a) Similar to Q. no. 60.

62. (d) $PCl_5 \rightleftharpoons PCl_3 \quad Cl_2$

$t \quad 0 \qquad 0.4M \qquad (x) \qquad (x)$

$t_{eq.} \quad (0.4 \quad x)$

For More Material Join: @JEEAdvanced_2024

$x \quad \frac{35}{100}(0.4)$

Now $K_c \quad \frac{[PCl_3][Cl_2]}{[PCl_5]}$

63. (a) $2AB_2(g) \rightleftharpoons 2AB(g) \quad B_2(g)$

$t \quad 0 \quad P$

$t_{eq.} \quad P \quad P \quad P \quad \frac{P}{2}$

$K_p \quad \frac{(P_{AB})^2(P_{B_2})^1}{(P_{AB_2})^2}$

64. (a) $H_2(g) \quad CO_2(g) \rightleftharpoons CO(g) \quad H_2O(g)$

$t \quad 0 \quad 1M \quad 1M$

$t_{eq.} \quad (1 \quad x) \quad (1 \quad x) \quad (x) \quad (x)$

$\because \quad n_g \quad 0$

$K_p \quad K_c \quad \frac{x^2}{(1 \quad x)^2}$

65. (c) $A_n(g) \rightleftharpoons n.A(g)$

$t \quad 0 \quad a$ mole

$t_{eq.} \quad (a \quad a) \quad na$

$M_0 \quad \frac{a.M_T}{(a \quad a \quad na)}$

$\frac{M_T \quad M_0}{(n \quad 1)\ M_0}$

$\because \quad D \quad M/2$

$\frac{D_T \quad D_0}{(n \quad 1)\ D_0}$

66. (a) For $PCl_5(g) \rightleftharpoons PCl_3(g) \quad Cl_2(g)$

$n \quad 2$

$\frac{D \quad d}{(2 \quad 1)\ d} \quad \frac{D}{d} \quad 1$

If $\frac{D}{d}$ at x-axis and ( ) at y-axis

then it is a straight line positive slope and negative intercept.

67. (a) $\frac{M_T \quad M_O}{(n \quad 1)\ M_O}$

$0.2 \quad \frac{92 \quad M_O}{M_O}$

$M_O \quad 76.66$

$d_{\text{mixture}} \quad \frac{PM_{\text{mixture}}}{RT}$

$\frac{1 \quad 76.66}{0.821 \quad 300} \quad 3.11$ g/litre

68. (c) $M_O \quad \frac{dRT}{P} \quad 57.47$

$\frac{99 \quad 57.47}{57.47} \quad 0.72$

69. (d) $I_2(g) \rightleftharpoons 2I(g)$

$t \quad 0 \quad a$ mole

$t_{eq.} \quad (a \quad a) \quad (2a)$

$P_{\text{Gas}} \quad x_{\text{Gas}} \quad P_{\text{Total}}$

$K_p \quad \frac{\left(\frac{2}{1} P_{\text{Total}}\right)^2}{\left(\frac{1}{1} P_{\text{Total}}\right)^1}$

K_p will be same in both cases.

70. (a) $A_2(g) \quad B_2(g) \rightleftharpoons 2AB(g)$

Initial conc. $10/2 \quad 15/2 \quad 5/2$

at eqm $5 \quad x \quad 7.5 \quad x \quad 2.5 \quad 2x$

$\because \quad 2.5 \quad 2x \quad 7.5$

$x \quad 2.5$

$K_c \quad \frac{(7.5)^2}{2.5 \quad 5} \quad 4.5$

71. (a) $H_2(g) \quad S(s) \rightleftharpoons H_2S(g)$

Conc. at eqm $0.5 \quad x \quad — \quad x$

$K_c \quad \frac{[H_2S]}{[H_2]}$

$7 \quad 10^{\ 2} \quad \frac{x}{0.5 \quad x}$

$x \quad 0.0327$

$P_{H_2S} \quad \frac{n_{H_2S}}{V} \quad RT$

$0.0327 \quad 0.0821 \quad 360 \quad 0.966$

72. (c) The gaseous mixture contains 40% Cl_2 and 40% PCl_3, since they are produced in 1 : 1 mole ratio. The PCl_5 % is 20.

For ideal gases mole % volume %

$P_{Cl_2} \quad P_{PCl_3}$

$2 \quad 0.40 \quad 0.80$ atm

$P_{PCl_5} \quad 2 \quad 0.2 \quad 0.40$ atm

$K_p \quad \frac{P_{PCl_3} \quad P_{Cl_2}}{P_{PCl_5}}$

$\frac{0.80 \quad 0.80}{0.40}$

$= 1.6$ atm

For More Material Join: @JEEAdvanced_2024

73. (a) For an ideal gas mole % volume %

$$K_p \quad \frac{P^2_{H_2O}}{P^2_{H_2}} \quad \frac{(0.55)^2}{(0.45)^2} \quad 1.49$$

74. (d) Moles of CO_2 present at equilibrium

$$\frac{1.642 \quad 50}{0.0821 \quad 1000} \quad 1$$

Mole % of XCO_3 decomposed

$$\frac{1}{4} \quad 100 \quad 25\%$$

Hence, 75% remains undecomposed.

75. (b) Let initial moles of $H_2(g)$ is 1

$$Fe_2O_3(s) \quad 3H_2(g) \rightleftharpoons 2Fe(s) \quad 3H_2O(g)$$

at eqm — 1 $3x$ — $3x$

$$K_c \quad \frac{\left(\frac{3x}{V}\right)^3}{\left(\frac{1 \quad 3x}{V}\right)^3}$$

$$8 \quad \left(\frac{3x}{1 \quad 3x}\right)^3$$

$$x \quad 0.22$$

% of H_2 unreacted

$$\frac{1 \quad 3 \quad 0.22}{1} \quad 100 \quad 34$$

76. (c) $AB_3(g) \rightleftharpoons AB_2(g) \quad \frac{1}{2}B_2(g)$

At eqm 800 x x $x/2$

$$800 \quad x \quad x \quad \frac{x}{2} \quad 900 \quad x \quad 200$$

$$\% \text{ dissociation} \quad \frac{200}{800} \quad 100 \quad 25$$

77. (b) Let P is initial pressure of NO_2

$$2NO_2(g) \rightleftharpoons 2NO(g) \quad O_2(g)$$

At eqm P $2x$ $2x$ x

as per given x 0.25

$$K_p \quad \frac{(2x)^2(x)}{(P \quad 2x)^2}$$

$$156.25 \quad \frac{(0.5)^2(0.25)}{P^2_{NO_2}}$$

$$P_{NO_2} \quad 0.02$$

78. (b) $2NOCl(g) \rightleftharpoons 2NO(g) \quad Cl_2(g)$

At eqm P $2x$ $2x$ x

$$P \quad x \quad 1; \quad P \quad 2x \quad 0.64, \quad x \quad 0.12$$

$$K_p \quad \frac{(0.24)^2(0.12)}{(0.64)^2}$$

$$= 16.875 \quad 10^{\ 3} \text{ atm}$$

79. (c) $Q_c \quad \frac{1 \quad 1}{1 \quad 1} \quad 1$

$\because \ Q_c \quad K_c$ so reaction will proceed in backward direction

$$A_2(g) \quad B_2(g) \rightleftharpoons C_2(g) \quad D_2(g)$$

Conc eqm $\frac{1 \quad x}{10}$ $\frac{1 \quad x}{10}$ $\frac{1 \quad x}{10}$ $\frac{1 \quad x}{10}$

$$0.25 \quad \frac{\left(\frac{1 \quad x}{10}\right)^2}{\left(\frac{1 \quad x}{10}\right)^2} \qquad x \quad 0.333$$

$$[A_2(g)] \quad \frac{1 \quad x}{10} \quad \frac{1.333}{10} \quad 0.133$$

80. (d) Given :

$$M_{Mix} \quad 62 \quad M_{H_2}$$

$$124$$

For $PCl_5 \rightleftharpoons PCl_3 \quad Cl_2$

$$\frac{(M)_{PCl_5} \quad (M)_{Mix}}{(2 \quad 1)\,(M)_{Mix}}$$

81. (b) Similar to Q. No. 80

82. (b) $K_c \quad \frac{[NO_2]^2}{[N_2O_4]}$

If $[NO_2]$ at y-axis

$[N_2O_4]$ at x-axis

$$y \quad Kc \ x^2$$

Parabolic curve

83. (d) $K_c \quad [Hg(g)]; P \quad CRT$

$$C \quad \frac{0.002}{760} \quad \frac{1}{0.0821 \quad 300}$$

$$= 1.068 \quad 10^{\ 7} \ M$$

84. (b) $K_c \quad \frac{n^2_{AB_2}}{n_{A_2} \ n^2_{B_2}} \quad V \quad \frac{n^2_{AB_2}}{n_{A_2} \ n^2_{B_2}} \quad \frac{n_{total}RT}{P}$

$$\frac{2^2}{5 \quad 3^2} \quad \frac{10 \quad 0.0821 \quad 300}{8.21}$$

85. (b) $A(g) \rightleftharpoons B(g) \quad C(g)$

$t \quad 0 \quad a$ mole

$t_{eq.} \quad (a \quad a) \quad (a) \quad (a)$

if $p_{Total} \quad P_1$

For More Material Join: @JEEAdvanced_2024

then

$$K_{p_1} \quad \frac{\frac{1}{1} p_1 \quad \frac{1}{1} p_1}{\frac{1}{1} p_1}$$

also

$$X(g) \rightleftharpoons 2Y(g)$$

$t \quad 0 \quad b$ mole

$t_{eq.} \quad (b \quad b) \quad (2b)$

if $p_{\text{Total}} \quad p_2$

$$K_{p_2} \quad \frac{\frac{2}{1} p_2^{\,2}}{\frac{1}{1} p_2^{\,1}}$$

Given : $\frac{K_{p_1}}{K_{p_2}} \quad \frac{1}{9}$

86. (b) Addition of inert gas at constant pressure will increase volume so reaction moves in the direction having more gaseous moles.

87. (a) Volume change has no effect for $n_g \quad 0$.

88, 89 (a, b)

Volume increase leads the reaction to move towards more gaseous moles.

90. (c) Addition of inert gas at constant volume has no effect.

91. (a) $Q \quad 1\,\text{atm} \quad K_p$

92. (b) For exothermic reaction yield on temperature

i.e., $T_1 \quad T_2 \quad T_3$

93. (a) $N_2(g) \quad 3H_2(g) \rightleftharpoons 2NH_3(g)$

$(t_{eq})_1 \quad 0.05M \quad 0.3M \quad 0.2M \; ; \; K_c \quad \frac{800}{9}$

$(0.05 \quad 0.1)$

$(t_{eq})_2 \quad (0.15 \quad x) \quad (0.3 \quad 3x) \quad (0.2 \quad 2x)$

$$K_c \quad \frac{[NH_3]^2}{[N_2]^1[H_2]^3}$$

$$\frac{800}{9} \quad \frac{(0.2 \quad 2x)^2}{(0.15 \quad x)(0.3 \quad 3x)^3}$$

94. (c) Similar to Q. No. 90.

95. (b) Le-Chatelier's principle is not applicable in solid phase reaction.

96. (d) When pressure decreases Eq. shift toward more gaseous moles.

97. (d) When temperature decreases Eq. shift toward exothermic direction.

98-99 (b,c)

Forward reaction must be exothermic with more gaseous moles at product side.

101. (d) When volume decreases Eq. shift toward less gaseous moles.

103. (b) For O_3 to be stable the reaction shouldn't be favourable in forward direction.

104. (d) Similar to Q. No. 101.

105. (c) Temperature favours exothermic reaction.

106. (d) Graphite has lesser density

more volume

low pressure

also for exothermic low temperature

107. (b) Molar volume of $H_2O(s)$ is more than $H_2O(l)$ so on increasing pressure formation of $H_2O(l)$.

110. (d) Pressure towards more volume.

111. (c) $\because \quad \ln K_p \quad \frac{H^0}{R} \quad \frac{1}{T} \quad \frac{S^0}{R}$

112. (b) For endothermic; $H \quad 0$, so slope is negative.

113. (a) For exothermic; $H \quad 0$, so slope is positive.

115. (a) For dissociation reaction; $H \quad 0$

116. (c) $\ln \frac{K_{P_2}}{K_{P_1}} \quad \frac{H}{R} \quad \frac{1}{T_1} \quad \frac{1}{T_2}$

$$\ln \frac{10^{\,4}}{10^{\,6}} \quad \frac{H}{8.314} \quad \frac{25}{298 \quad 323}$$

$_rH \quad 147.41$ kJ/mol

117-118 (b, a)

See Q. No. 116

119. (a) The value of K_{eq} is the measure of extent of reaction. For most stable oxide K_{eq} should be minimum.

120 - 125. (c, c, a, b, d, a)

$G \quad RT \ln K_{eq}$

$G \quad (H \quad T S) \quad RT \ln K_{eq}$

123. (b) $_rG \quad _rH \quad T \quad _rS$

$30 \quad 300 \quad 0.1 \quad 0$

$_rG \quad 2.303 RT \log K$

$K \quad 1$

124. (d) $P_{\text{total}} \quad 0.2$ bar

At equilibrium $P_{CO_2} \quad P_{H_2O} \quad 0.1$ bar

$K_p \quad (0.1)^2 \quad 0.01$

$_rG \quad RT \ln K_p$

$8.314 \quad 420 \ln (0.01)$

16083.6 J/mol or 16.083 kJ/mol

For More Material Join: @JEEAdvanced_2024

Level 2

1. (a) $C(s) \quad CO_2(g) \rightleftharpoons 2CO(g); \quad \ldots(1)$

$K_{p_1} \quad 10^{14}$ atm

$2\,CO(g) \quad 2\,Cl_2(g) \rightleftharpoons 2COCl_2(g); \quad \ldots(2)$

$K_{p_2} \quad (6 \quad 10^{-3})^2$ atm $^{-2}$

Add (1) and (2)

$C(s) \quad CO_2(g) \quad 2\,Cl_2(g) \rightleftharpoons 2COCl_2(g);$

$K_p \quad 10^{14} \quad 36 \quad 10^{-6}$

$36 \quad 10^{8}$

For given reaction $n_g \quad 1$

$K_c \quad K_p(RT)$

$36 \quad 10^{8} \quad 0.0821 \quad 1120$

$K_c \quad 3.31 \quad 10^{11}\ M^{-1}$

2. (c) $N_2(g) \quad 3H_2(g) \rightleftharpoons 2NH_3(g)$

Initial moles	1	2	0
at eqm	$1 \quad x$	$2 \quad 3x$	$2x$

where $x \quad 0.5$

$$K_c \quad \frac{(1/4)^2}{\frac{0.5}{4} \quad \frac{0.5}{4}^3} \quad 256$$

equilibrium constant for the reaction

$NH_3(g) \rightleftharpoons \frac{1}{2}N_2(g) \quad \frac{3}{2}H_2(g)$

$K_c^1 \quad \frac{1}{\sqrt{K_c}} \quad \frac{1}{16}$

3. (c) $A_2(g) \quad B_2(g) \rightleftharpoons 2AB(g)$

Moles at eqm $2 \quad x \quad 4 \quad x \quad 2x$

$$K_c \quad \frac{4x^2}{(2 \quad x)(4 \quad x)}$$

$x \quad \frac{32}{24} \quad 1.33$ mole

$[AB(g)] \quad \frac{2 \quad 1.33}{4} \quad 0.66\ M$

4. (d) $P_{total} \quad P_{HNO_3} \quad P_{NO_2} \quad P_{H_2O} \quad P_{O_2}$

$\because \quad P_{NO_2} \quad 4P_{O_2}$ and $P_{H_2O} \quad 2P_{O_2}$

$P_{total} \quad P_{HNO_3} \quad 7P_{O_2}$

$30 \quad 2 \quad P_{O_2} \quad 7$

$P_{O_2} \quad \frac{28}{7} \quad 4$

$$K_p \quad \frac{P_{NO_2}^4 . P_{H_2O} . P_{O_2}}{P_{HNO_3}^4}$$

$$\frac{(4 \quad 4)^4 \quad (2 \quad 4)^2 \quad 4}{2^4} \quad 2^{20}$$

$K_p \quad K_c (RT)^{n_g} \quad K_c (0.08 \quad 400)^3$

$K_c \quad \frac{2^{20}}{(32)^3} \quad 32$

5. (d) $LiCl \quad 3NH_3(s) \rightleftharpoons LiCl \quad NH_3(s) \quad 2NH_3(g);$

$[K_p \quad 9\ atm^2]$

Therefore,

$LiCl \quad NH_3(s) \quad 2NH_3(g) \rightleftharpoons LiCl \quad 3NH_3(s);$

Initial moles	0.1	a	0
Final moles at eqm	0	$(a \quad 0.2)$	0.1

$K_p \quad \frac{1}{9}(atm)^{-2}$

Let initial moles of NH_3 are a

At eqm $K_p \quad \frac{1}{(P_{NH_3})^2}$ or $\frac{1}{9} \quad \frac{1}{(P_{NH_3})^2}$

$P_{NH_3} \quad 3$ atm

$\because \quad PV \quad nRT \quad 3 \quad 5 \quad n \quad 0.0820 \quad 310$

$n \quad 0.59$ *i.e.,* $(a \quad 0.2) \quad 0.59$

initial moles of $NH_3 \quad 0.79$

6. (c) $NH_2COONH_4(s) \rightleftharpoons 2NH_3(g) \quad CO_2(g)$

Let partial pressure at equilibrium of CO_2 be P, then $P_{NH_3} \quad 2P$ and total pressure at equilibrium $3P$

$K_p \quad (2P)^2 \quad P \quad 4P^3 \quad \ldots(1)$

If NH_3 is added and the pressure of NH_3 after addition at equilibrium is $3P$

$K_p \quad 4P^3 \quad (P_{NH_3})^2 \quad P_{CO_2}$

$P_{CO_2} \quad \frac{4}{9}P$

ratio of $\frac{P_{CO_2}}{P_{CO_2}} \quad \frac{4}{9}$

7. (c) $C_2H_6(g) \rightleftharpoons C_2H_4(g) \quad H_2(g)$

At eqm $1 \quad x \quad x \quad x$

$$K_p \quad \frac{P_{C_2H_4} \quad P_{H_2}}{P_{C_2H_6}} \quad \frac{x^2}{1 \quad x} \quad 5 \quad 10^{-2}$$

$x^2 \quad 0.05\ x \quad 0.05 \quad 0$

$$x \quad \frac{0.05 \quad \sqrt{(0.05)^2 \quad 4 \quad 0.05}}{2} \quad 0.20 \text{ atm}$$

Partial pressure of C_2H_6 mole fraction total pressure

0.80 mole fraction 1.2

mole % of $C_2H_6 \quad \frac{0.8}{1.2} \quad 100 \quad 66.66$

For More Material Join: @JEEAdvanced_2024

8. (a) $2NOBr(g) \rightleftharpoons 2NO(g) + Br_2(g)$

Initial pressure	P	0	0
at eqm	$P - 2x$	$2x$	x

where $2x = 0.40\ P$

$x = 0.20\ P$;

$1.20\ P = 0.30$

$P = 0.25$ atm

$$K_p = \frac{P_{NO}^2\ P_{Br_2}}{P_{NOBr}^2} = \frac{(0.4P)^2\ (0.2P)}{(0.6P)^2} = 0.0222$$

K_p for the reaction $2NO(g) + Br_2(g) \rightleftharpoons 2NOBr(g)$ is $\frac{1}{0.0222} = 45$

9. (c) $\because$ Initially only A is present so at eqm B and C should be present in 2 : 1

$$2A(g) \rightleftharpoons 2B(g) + C(g)$$

At eqm 400 mL 200 mL 100 mL

for ideal gases volume % = mole %

$$K_p = \frac{\left(\frac{200}{700}\right)\ 10^{\ 2}\ \left(\frac{100}{700}\right)\ 10}{\left(\frac{400}{700}\right)\ 10^{\ 2}} = \frac{10}{28}$$

10. (c) $SO_2(g) + NO_2(g) \rightleftharpoons SO_3(g) + NO(g)$

Initial moles	2	2	2	2
at eqm	$2 - x$	$2 - x$	$2 + x$	$2 + x$

$(\because Q_p = K_p)$

Total no. of moles of gases at equilibrium $= 8 + 2 = 10$

$$K_p = \frac{P_{SO_3}\ P_{NO}}{P_{SO_2}\ P_{NO_2}}$$

$$25 = \frac{\left(\frac{2 + x}{10}\right)\ P^{\ 2}}{\left(\frac{2 - x}{10}\right)\ P^{\ 2}}$$

$$5 = \frac{2 + x}{2 - x}$$

$x = 1.33$

Partial pressure of $NO_2 = \frac{2 - x}{10}\ P_{total}$

$= \frac{0.666}{10}\ 2$

$= 0.133$ atm

11. (b) $3\,Se_2(g) \rightleftharpoons Se_6(g)$

moles at eqm $a(1 - \quad)$ $\frac{a}{3}$

$$\frac{\text{Observed no. of moles at equilibrium}}{\text{Initial no. of moles of reactant}} = \frac{M_T}{M_O},$$

$$\frac{a(1 - \quad) + \frac{a}{3}}{a}\ \frac{M_T}{M_O} = 0.315$$

where molar mass of Se_2 $(M_T) = 79 \quad 2 = 158$

and molar mass of mixture $(M_O) = \frac{wRT}{PV} = \frac{0.02 \quad 24.63}{1 \quad 2.463 \quad 10^{\ 3}} = 200$

12. (d) $6HCHO \rightleftharpoons C_6H_{12}O_6$

Conc. at eqm $C(1 - \quad)$ $\frac{C}{6}$

$$\frac{\text{Observed molar concentration}}{\text{Initial molar concentration}} = \frac{M_T}{M_O}$$

M_T = Theoretical molar mass of HCHO

$$\frac{C(1 - \quad) + \frac{C}{6}}{C}\ \frac{M_T}{M_O} = \frac{30}{150} = 0.96$$

13. (b) $2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$

$$K_c = \frac{(0.12)^2}{(0.12)^2\ 5} = 0.2$$

Another vessel

$$2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$$

moles at eqm $0.5 - 2x$ $y - x$ $2x$

as per given $2x = \frac{20}{100}\ 0.5 = 0.1$

$$K_c = \frac{(0.1)^2}{(0.4)^2\ (y \quad 0.05)} = 0.20$$

$y = 0.3625$ mole

mass of O_2 added = 11.6 g

14. (b) Let initial mole of $N_2O_4(g)$ is one

$$N_2O_4(g) \rightleftharpoons 2NO_2(g)$$

Initial moles	1	0
At equilibrium	$1 - \alpha$	2α

Total no. of moles at equilibrium $= 1 + \alpha$

$$P_{N_2O_4} = \frac{1 - \alpha}{1 + \alpha}\ P$$

$$P_{NO_2} = \frac{2\alpha}{(1 + \alpha)}\ P$$

For More Material Join: @JEEAdvanced_2024

Hence, $K_p = \frac{P_{NO_2}^2}{P_{N_2O_4}} = \frac{4\ ^2}{(1\ \ ^2)}\ P$

$4.5 = \frac{4\ ^2}{(1\ \ ^2)}\ 2$

0.6

Mole fraction of N_2O_4 :

$X_{N_2O_4} = \frac{1}{1} = 0.25$

$X_{NO_2} = 0.75$

Average molar mass of mixture

$0.25\ \ 92\ \ 0.75\ \ 46\ \ 57.5$

15. (a) $2AB(s) \rightleftharpoons A_2(g)\ \ B_2(g)$

$0.5\ \ x\ \ \ \ x$

$K_p = P_{A_2}\ P_{B_2}$

$0.06 = (0.5\ \ x)\ x$

$x^2 = 0.5x\ \ 0.06$

$x = 0.1$

$P_{total} = P_{A_2}\ \ P_{B_2} = 0.6\ \ 0.1 = 0.70$ atm

16. (a) $Sb_2S_3(s)\ \ 3H_2(g) \rightleftharpoons 2Sb(s)\ \ 3H_2S(g)$

$0.01\ \ x\ \ \ \ 0.01\ \ 3x\ \ \ \ \ \ 2x\ \ \ \ 3x$

where $3x = 0.005$

$H_2S + Pb^{2+} \rightleftharpoons PbS + 2H^+$

no. of moles of PbS formed

$\frac{1.19}{238} = 0.005$ mole

At eqm $[H_2] = \frac{0.005}{250}\ \ \frac{.01\ \ .005}{250}$;

$K_c = \frac{0.005}{0.005}^{\ 3} = 1$

17. (a) $2A(g)\ \ B(g) \rightleftharpoons C(g)\ \ D(g)$

Initial moles	1	1	7	3
at eqm	2 2x	1 x	7 x	3 x

Due to very high value of K_c we can assume that reactant is almost converted into products so

$1\ \ x\ \ y;\ \ 2\ \ 2x\ \ 2y \qquad x \sim 1$

$10^{12} = \frac{8\ \ 4}{(2y)^2 y}$

$y^3 = 8\ \ 10^{\ 12}$

equilibrium concentration of

$A = 2y = 4\ \ 10^{\ 4}$

18. (b) $K_c = \frac{[complex]}{[H_3BO_3]\ [glycerine]} = 0.90$

$\frac{[complex]}{[H_3BO_3]} = \frac{80}{20}$

$[glycerine] = \frac{80}{20\ \ 0.9} = 4.44\ M$

$H_3BO_3 + glycerine \rightleftharpoons (H_3BO_3 - glycerine)$

At eq^m $\quad 0.1\ \ x \qquad a\ \ x \qquad\qquad x$

$x = 0.1\ \ \frac{80}{100} = 0.08;$

$a\ \ x = 4.44$

$a = 4.44\ \ 0.08 = 4.52\ M$

or Initial moles $= 4.52$

19. (c) $\frac{r_{mix}}{r_{O_2}} = \sqrt{\frac{32}{M_{mix}}} = 0.4\sqrt{5}$

$M_{mix} = 40$ g/mol

$O_2(g) \rightleftharpoons \frac{2}{3}O_3(g)$

Initial moles	1	0
at eqm	1	$\frac{2}{3}$

$\frac{32}{40} = \frac{1\ \ \frac{2}{3}}{1}$

0.6

$\% = 60$

20. (c) $2SO_3(g) \rightleftharpoons 2SO_2(g)\ \ O_2(g)$

moles at eqm $\quad 1\ \ 2x \qquad\qquad 2x \qquad\ x$

Only SO_2 (O. No. 4) will oxidized

so equivalent of SO_2 = equivalent of $KMnO_4$

$2x\ \ 2 = 0.2\ \ 5$

$2x = 0.5$

$K_c = \frac{\frac{0.5}{2}^{\ 2}\ \ \frac{0.25}{2}}{\frac{0.5}{2}^{\ 2}} = 0.125$

21. (d) $\frac{r_{mix}}{r_{SO_2}} = \sqrt{\frac{M_{SO_2}}{M_{mix}}}$

$2.56 = \frac{64}{M_{mix}}$

$M_{mixture} = 25$

Let mole fraction of F_2 is x

$25 = \frac{38\ \ x\ \ (1\ \ x)\ \ 19}{1}$

$x = 0.315;$

For More Material Join: @JEEAdvanced_2024

$K_p \quad \frac{P_F^2}{P_{F_2}} \quad \frac{(0.685\ P)^2}{(0.315\ P)} \sim 1.49\text{atm}$

22. (b) $8 \quad 10^{\ 6} \quad \frac{x^2}{0.5}$

$x \quad 2 \quad 10^{\ 3}$

pOH 2.7

So, pH 11.3

23. (c) $_rG \quad 0 \quad 77.1 \quad 2 \quad 154.2$ kJ/mol

$Q \quad \frac{[H\ ]^2}{P_{H_2}\ [Ag\ ]^2}$

$\frac{10^{\ 6}}{0.5 \quad (10^{\ 10})} \quad 2 \quad 10^4$

$G \quad _rG \quad RT \ln Q$

$_rG \quad 154.2 \quad \frac{8.314 \quad 300 \ln (2 \quad 10^4)}{1000}$

129.5 kJ/mol

24. (d) $N_2O_5 \rightleftharpoons N_2O_3 + O_2$

$4 \quad x \qquad x \quad y \qquad x \quad y$

$N_2O_3 \rightleftharpoons N_2O + O_2$

$x \quad y \qquad y \qquad y \quad x$

$\because \ [O_2] \quad x \quad y \quad 2.5$

and $2.5 \quad \frac{(x \quad y)(x \quad y)}{4 \quad x}$

$x \quad 2.166$

$[N_2O_5] \quad 4 \quad x \quad 1.846$

25. (b) Let x is partial pressure of A and y is partial pressure of C when both equilibrium simultaneously established in a vessel

$X(s) \rightleftharpoons A(g) \qquad 2B(g)$

$x \qquad (2x \quad 2y)\ ;$

$K_{P_1} \quad P_A \quad P_{B\,(\text{total})}^2$

$Y(s) \rightleftharpoons C(g) \qquad 2B(g)$

$y \qquad (2y \quad 2x)\ ;$

$K_{p_2} \quad P_C \quad P_{B\,(\text{total})}^2$

$\frac{K_{P_1}}{K_{p_2}} \quad \frac{x}{y} \qquad x \quad 2y$

$K_{P_1} \quad x(2x \quad 2y)^2$

$x \quad 0.1$ atm;

$y \quad 0.05$ atm

Total pressure of gases $P_A \quad P_B \quad P_C$

$3(x \quad y)$

$= 0.45$ atm.

Level 3

Passage-2 :

1,2,3. (a, b, b)

$\ln K_p \quad \frac{H}{R} \quad \frac{1}{T} \quad \frac{S}{R}$

4. (a) $\ln \frac{K_2}{K_1} \quad \frac{H}{R} \quad \frac{1}{T_1} \quad \frac{1}{T_2}$

Passage-3

$N_2O_3(g) \rightleftharpoons NO_2(g) \quad NO(g)\ ; \quad K_{p_1} \quad 2.5$ bar

$2NO_2(g) \rightleftharpoons N_2O_4(g) \qquad ; \quad K_{p_2}$

$x \quad y \qquad y/2$

$P_{NO} \quad x \quad 1.5$ b

$K_{p_1} \quad \frac{P_{NO}\ P_{NO_2}}{P_{N_2O_3}}$

$2.5 \quad \frac{x(x \quad y)}{2 \quad x} \quad \frac{1.5(1.5 \quad y)}{2 \quad 1.5} \quad 4.5 \quad 3y$

$y \quad \frac{2}{3}$ bar

(1) $P_{N_2O_3} \quad 2 \quad x \quad 2 \quad 0.5 \quad 0.5$ b

(2) $P_{NO_2} \quad x \quad y \quad 1.5 \quad \frac{2}{3} \quad \frac{2.5}{3}$ b

(3) $K_{p_2} \quad \frac{y}{2(x \quad y)^2} \quad \frac{2/3}{2 \quad \left(\frac{2.5}{3}\right)^2} \quad 0.48\ \text{bar}^{\ 1}$

Passage-4 :

1. (b) For $X(g) \xrightleftharpoons{H_2O(l)} X_{(aq.)}$; $H \quad 0$

For more solubility forward shift

low temperature (exothermic)

high pressure (less volume)

2. (a) For $Fe(l) \rightleftharpoons Fe(s)$

i.e., $H \quad 0$ low temperature

less volume high pressure

One or More Answers is/are Correct

6. (c, d) Le Chatelier's principle is not quantitative. If both stress would cause the same direction of shift, the shift is determinable. If the two stresses would cause shifts in opposite directions, no deduction is possible.

For More Material Join: @JEEAdvanced_2024

Subjective Problems

1. $\frac{1}{2}C(s) \quad \frac{1}{2}CO_2(g) \rightleftharpoons CO(g)$

$t \quad 0 \qquad P$ atm

$t_{eq.} \qquad (P - P/2) \qquad P$

$P/2 \quad P \quad 12$

$P \quad 8$ atm

$K_p \quad \frac{(p_{CO})^1}{(p_{CO_2})^{1/2}}$

2. $A(s) \rightleftharpoons B(g) \quad 2C(g)$

$t \quad 0 \quad a$

$t_{eq.} \quad a \quad p \qquad p \qquad 2p$

$K_p \quad (P_B)^1(P_C)^2$

$32 \quad (P)^1(2P)^2$

$P \quad 2$ atm

3. $A \qquad 2B \rightleftharpoons 2C \quad D$

$t \quad 0 \quad 2a \qquad 3a$

$t_{eq} \quad (2a \quad x) \; (3a \quad 2x) \quad (2x) \quad (x)$

Given : $(2a \quad x) \quad x$

$x \quad a$

$K_c \quad \frac{[C]^2[D]^2}{[A]^1[B]^2}$

4. $D \rightleftharpoons C; \quad K_C \quad \frac{1}{0.01}$

$B \rightleftharpoons A; \quad K_C \quad \frac{1}{10}$

$C \rightleftharpoons B; \quad K_C \quad \frac{1}{2}$

Addition

$D \rightleftharpoons A; \quad K_C \quad \frac{1}{0.01} \quad \frac{1}{10} \quad \frac{1}{2}$

5. $A(g) \rightleftharpoons B(g)$

$(t_{eq.})_1 \quad 0.4M \qquad 0.4M$

$K_C \quad 1$

$\because \; K_C$ remains constant

6. $K_P \quad k_C(RT)^{n_g}$

7. $SO_2 \qquad NO_2 \rightleftharpoons SO_3 \qquad NO$

$t \quad 0 \quad 3$ mole $\qquad 4 \qquad\qquad 1 \qquad\qquad 4$

$t_{eq.} \qquad (3 \quad x) \qquad (4 \quad x) \qquad (1 \quad x) \qquad (4 \quad x)$

Given : $(3 \quad x) \quad 1$

$x \quad 2$

$K_C \quad \frac{[SO_3][NO]}{[SO_2][NO_2]}$

8. For mixture

$P.M_{Mix} \quad d.RT$ and for $N_2O_4 \rightleftharpoons 2NO_2$

$\frac{(M)_{N_2O_4} \quad (M)_{Mix}}{(2 \quad 1)\;(M)_{Mix}}$

9. $G \quad RT \ln K_{eq}$

At standard states all concentration and partial pressure are unity.

$K_{eq} \quad 1$

10. $G \quad RT \ln K_{eq}$

$830 \quad 2 \quad 300 \quad 2303 \log_{10} K_{eq}$

$K_{eq} \quad 4$

Now

$A \qquad B \rightleftharpoons C \qquad D$

$t \quad 0 \qquad a \qquad\qquad a$

$t_{eq.} \qquad (a \quad x) \quad (a \quad x) \qquad (x) \qquad (x)$

$K_C \quad \frac{[C][D]}{[A][B]}$

$x \quad 2a/3$

11. $NH_4HS(s) \rightleftharpoons NH_3(g) \qquad H_2S(g)$

At eqm. $\qquad - \qquad\qquad (0.1 \quad P) \qquad\qquad P \qquad ;$

As per given $0.1 \quad 2P \quad 1.1; \quad P \quad 0.5$

$K_P \quad P_{NH_3} \; P_{H_2S} \quad (0.1 \quad P) \; P$

$0.6 \quad 0.5 \quad 0.3 \quad 3 \quad 10^{\;1} \quad 3$

12. $A \qquad nB \rightleftharpoons mC$

$t \quad 0 \qquad\qquad 8M \qquad 12\,M$

$t \quad t_{eq.} \qquad\qquad 8 \quad x \qquad 12 \quad nx \qquad mx$

Given: $\qquad [A]_{eq} \quad 6M$

$8 \quad x \quad 6$

$x \quad 2M$

$[B]_{eq} \quad 8M$

$12 \quad nx \quad 8$

$n(2) \quad 4$

$n \quad 2$

$[C]_{eq} \quad 6M$


$mx \quad 6$

$m(2) \quad 6$

$m \quad 3$

$m \quad n \quad 3 \quad 2 \quad 5$

For More Material Join: @JEEAdvanced_2024



6

IONIC EQUILIBRIUM

Arrhenius Theory

When dissolved in water, the substances which releases

(i) H ions are called acids *e.g.* HNO_3, $HClO_4$, CH_3COOH,

(ii) OH ions are called bases, *e.g.* NaOH, KOH, NH_4OH

Bronsted - Lowry Protonic Concept : acid is H donor and base is H acceptor

Conjugate acid and base

To get conjugate acid of a given species add H to it. *e.g.* conjugate acid of N_2H_4 is N_2H_5.

To get conjugate base of any species subtract H from it. *e.g.* Conjugate base of NH_3 is NH_2.

HCl is an acid while its conjugate ion Cl is a base.

HCO_3 can donate proton as well as can accept proton hence amphiprotic.

Lewis Concept : Acids are substances which accept a lone pair of electrons to form a coordinate bond and bases are the substances which donate a lone pair of electrons to form a coordinate bond.

❖ Lewis-acid is lone pair acceptor ; Lewis-base is lone-pair donor.

Ostwald dilution law: Applicable on weak electrolytes (weak acids/bases)

If is negligiable with respect to 1 (0.05), than for mono basic acid and mono acidic base.

K_C C^2 or $\sqrt{\frac{K_C}{C}}$

So $\frac{1}{\sqrt{C}}$ or $\sqrt{V}$ where, V is the volume of solution

at infinite dilution reaches its maximum value, unity.

For More Material Join: @JEEAdvanced_2024

Presence of other solute (Common ion effect)

Degree of dissociation of a weak electrolyte is suppressed by the addition of a substance having an ion common to weak electrolyte is known as common ion effect.

Ionic Product of Water

Pure water is weak electrolyte and dissociated as:

$$H_2O \rightleftharpoons H \quad OH$$

Before dissociation	C	0	0
After dissociation	$C(1 \quad)$	C	C

$$K_{eq.} \quad \frac{[H \ ][OH \ ]}{[H_2O]}$$

where, K_{eq} is dissociation constant or equilibrium constant of water. Since, dissociation of water is too less and undissociated H_2O can be taken as constant, therefore

$$K_{eq}[H_2O] \rightleftharpoons [H \ ]\,[OH \ ]$$

$$K_w \quad [H \ ][OH \ ]$$

The new constant K_w is known as ionic product of water. The numerical value of K_w increases considerably with temperature. It is 1.0 10^{-14} at 25°C. The variation of ionic product of water with temperature is given by

$$\ln \frac{K_{w_2}}{K_{w_1}} \quad \frac{{}_rH}{R} \frac{[T_2 \quad T_1]}{T_1 T_2}$$

pH CONCEPT

S. Sorenson used a new term pH to express the hydrogen ion concentration.

$$pH \quad \log[H \ ] \quad \text{or} \quad pH \quad \log[H_3O \ ] \quad \text{or} \quad [H \ ] \quad 10^{-pH}$$

For simplicity hydrogen ions are simply written as H , with the understanding that H ions in solution are always solvated.

- Ionic product (auto dissociation constant) of water
  $K_w \quad 1 \quad 10^{-14} \quad [H_3O \ ][OH \ ]$ at 298 K
- K_w increases with increase in T
- pH pOH pK_w 14 (at 298 K)
- For weak acid and its conjugate base

$$\text{or} \qquad \frac{K_aK_b \quad K_w}{pK_a \quad pK_b \quad pK_w \quad 14 \quad \text{(at 298 K)}}$$

pH calculation of weak acid (HA)

$$HA(aq) \rightleftharpoons H \ (aq) \quad A \ (aq)$$

Initial conc.	C	–	–
Conc. at eq.	$C(1 \quad)$	C	C

$$K_a \quad \frac{C^{2}}{1}$$

if $\quad 1; \quad K_a \quad C^{2}$

pH calculation of weak base (BOH)

$$BOH(aq) \rightleftharpoons B \ (aq) \quad OH \ (aq)$$

Initial conc.	C		
Conc. at eq.	$C(1 \quad)$	C	C

$$K_b \quad \frac{C^{2}}{1}$$

if $\quad 1; \quad K_b \quad C^{2}$

For More Material Join: @JEEAdvanced_2024

$[H^+] \quad C \quad \sqrt{K_a C}$

$pH \quad \frac{1}{2}(pK_a \quad \log C)$

Degree of dissociation

K_a dissociation constant of weak acid

$[OH^-] \quad C \quad \sqrt{K_b C}$

$pOH \quad \frac{1}{2}(pK_b \quad \log C)$

Degree of dissociation

K_b dissociation constant of weak base

pH Calculation of mixture of two weak acid H*A* and H*B* :

$[H^+] \quad \sqrt{K_{a_1} C_1 \quad K_{a_2} C_2}$; if $\quad_1$ and $\quad_2$ of both acids are negligiable w.r.t. unity.

K_{a_1} and K_{a_2} are dissociation constant of H*A* and H*B* respectively.

C_1 and C_2 are concentration of H*A* and H*B* respectively.

pH calculation of mixture of two weak base *A*OH and *B*OH.

$[OH^-] \quad \sqrt{K_{b_1} C_1 \quad K_{b_2} C_2}$; if $\quad_1$ and $\quad_2$ of both bases are negligiable w.r.t. unity.

K_{b_1} and K_{b_2} are dissociation constant of *A*OH and *B*OH respectively.

C_1 and C_2 are concentration of *A*OH and *B*OH respectively.

Salt Hydrolysis

- Hydrolysis is the interaction of cation/anion/both ions of the salt with H_2O
- Salt of strong acid (SA) and strong base (SB) is not hydrolysed ; solution is neutral, pH 7 at 25°C.
- Solution of salt of SA and weak base (WB) is acidic due to hydrolysis of cation

h degree of hydrolysis $\sqrt{\frac{K_w}{K_b C}} \quad \sqrt{\frac{K_h}{C}}$, K_h (hydrolysis constant) $\frac{Ch^2}{(1 \quad h)}$

$$pH \quad 7 \quad \frac{1}{2}(pK_b \quad \log C) \qquad \text{At } 25°C$$

- Solution of salt of weak acid (WA) and SB is alkaline due to hydrolysis of anion

$$h \quad \sqrt{\frac{K_w}{K_a C}} \quad \sqrt{\frac{K_h}{C}}$$

$$pH \quad 7 \quad \frac{1}{2}(pK_a \quad \log C) \qquad \text{At } 25°C$$

- Solution of salt of WA and WB is neutral ($K_a \quad K_b$), acidic ($K_a \quad K_b$) and basic ($K_b \quad K_a$) due to hydrolysis of both

$$h \quad \sqrt{\frac{K_w}{K_a K_b}} \quad \sqrt{K_h} \quad ; \quad pH \quad 7 \quad \frac{1}{2}(pK_a \quad pK_b) \qquad \text{At } 25°C$$

- For amphiprotic salt like HS^-, HCO_3^-, $pH \quad \frac{pK_1 \quad pK_2}{2}$ or $[H^+] \quad \sqrt{K_1 K_2}$

(K_1 first dissociation constant, K_2 second dissociation constant of acid)

Buffer Solutions are the solutions whose pH does not change significantly on adding a small quantity of strong base or strong acid.

In general buffer solutions are simple (salts of weak acid and weak base), or mixed buffer.

Mixed buffer are of two types :

(A) Acidic buffer solutions : weak acid with its conjugate base :

For More Material Join: @JEEAdvanced_2024

made by mixing a weak acid with its conjugate base. *e.g.*, CH_3COOH with CH_3COONa, $NaHCO_3$ and H_2CO_3, H_3PO_4 and NaH_2PO_4.

(B) Basic buffer solutions : weak base with its conjugate acid :

$NH_3(aq)$ with $NH_4Cl(aq)$, $RNH_2(aq.)$ and $RNH_3Cl(aq.)$ etc,

If K_a for acid (or K_b for base) is not too high, we may write :

Henderson's Equation

For acidic buffer solution : pH pK_a log ([salt]/[acid])

For basic buffer solution : pOH pK_b log ([salt]/[base])

- For good buffer capacity, [salt] : [acid] ratio should be as close to one as possible. In such a case,

  pH pK_a. (This also is the case at midpoint of titration)

- Buffer capacity $\dfrac{\text{No. of moles of acid (or base) added per litre of buffer solution}}{\text{Change in pH}}$

Maximum Buffer Capacity

It can be proved that maximum buffer capacity is achieved when the salt and acid or base concentration is equal.

Indicators

Indicator is a substance which indicates the point of equivalence in a titration by undergoing a change in its colour. They are weak acids or weak bases.

Theory of Indicators : The ionized and unionized forms of indicators have different colours. If 90 % or more of a particular form (ionised or unionised) is present, then its colour can be distinctly seen. In general, for an indicator which is weak acid, HIn $\rightleftharpoons$ H In , the ratio of ionized to unionized form can be determined from

$$\text{pH} \quad pK_a \quad \log \frac{[\text{In}^-]}{[\text{HIn}]}$$

So, for detectable colour change, pH pK_a 1

This roughly gives the range of indicators. Ranges for some popular indicators are

Indicators	pH range	Colour	
		Acid medium	Basic medium
Methyl orange	3.1 4.4	red	yellow
Methyl red	4.2 6.3	red	yellow
Phenolphathlene	8.3 10	colourless	pink

Equivalence point: The point at which exactly equivalent amounts of titrate and titrant have been mixed.

Acid Base Titration : For choosing a suitable indicator titration curves are of great help. In a titration curve, change in pH is plotted against the volume of alkali to a given acid. Four cases arise.

- **Strong acid *vs* strong base :** The curve is almost vertical over the pH range 3.5-10. This abrupt change corresponds to equivalence point. Any indicator suitable.

For More Material Join: @JEEAdvanced_2024

- **Weak acid *vs* strong base :** Final solution is basic at equivalence point. Vertical region (not so sharp) lies in pH range 6.5-10. So, phenolphathlene is suitable.
- **Strong acid *vs* weak base :** Final solution acidic. Vertical point in pH range 3.8-7.2. Methyl red or methyl orange suitable.
- **Weak acid *vs* weak base :** No sharp change in pH. No suitable indicator.

Note : *At midpoint of titration, pH* pK_a, *thus by pH measurements,* K_a *for weak acids (or* K_b *for weak bases) can be determined.*

Polyprotic acids and bases : K_2, K_3 etc. can be safely neglected. If K_1 K_2 K_3 and only K_1 plays a significant role in pH calculation.

Solubility (*S*) and Solubility Product (K_{sp})

This is generally used for sparingly soluble salts,

Let the salt $A_x B_y$ is present in water. Let the solubility of $A_x B_y$, is S

$$A_x B_y(s) \rightleftharpoons xA^y \ (aq) \quad yB^x \ (aq)$$

$$- \qquad xS \qquad yS$$

$$K_{sp} \quad (xS)^x \ (yS)^y \quad x^x y^y \ (S)^{x\ y}$$

e.g., For sparingly soluble salts (*e.g.,* $Ag_2C_2O_4$) an equilibrium which exists as

$$Ag_2C_2O_4(s) \rightleftharpoons 2Ag \ (aq.) \quad C_2O_4^{2} \ (aq.)$$

Then $K_{sp} \quad [Ag\ ]^2 [C_2O_4^{2}\ ]$

Ionic product (*Q*) : It is defined as product of concentration of all the ions of a sparingly soluble salt at time '*t*'.

- **Case-1 :** If Q K_{sp} then solution is unsaturated no precipitation takes place.
- **Case-2 :** If Q K_{sp} then solution is saturated.
- **Case-3 :** If Q K_{sp} solution is super saturated, precipitation takes place.

Common ion effect : Suppression of dissociation by adding an ion common with dissociated products. *e.g.,* Solubility of AgCl decreases in presence of NaCl (Cl ion common) OR in presence of $AgNO_3$(Ag ion common)

$$AgCl(s) \rightleftharpoons Ag \ (aq) \qquad Cl \ (aq)$$

Solubility in water at eq.: – S S

Solubility in 0.01*M* NaCl at eq.: – S' $(S' \ 0.01)$

$$K_{sp} \quad [Ag\ ][Cl\ ]_{Total}$$

$$K_{sp} \quad S'(S' \ 0.01) \qquad \text{where} \quad S' \ S$$

Simultaneous solubility : When two sparingly soluble salts are present in water with atleast a common ion, then solubility of both salts decreases *e.g.,* AgCl and AgBr, $SrSO_4$ and $BaSO_4$

e.g. Calculate simultaneous solubility of AgCl and AgBr in the solution.

If simultaneous solubility of AgCl and AgBr are x and y respectively.

$$AgCl(s) \rightleftharpoons Ag \ (aq.) \quad Cl \ (qq.)$$

At eq. – $(x \ y)$ x

$$K_{sp_1}(AgCl) \quad [Ag\ ]_{Total}[Cl\ ]$$

$$K_{sp_1} \quad (x \ y)\,x$$

$$AgCl(s) \rightleftharpoons Ag \ (aq.) \quad Cl \ (qq.)$$

At eq. – $(x \ y)$ y

$$K_{sp_2}(AgBr) \quad [Ag\ ]_{Total}[Br\ ]$$

$$K_{sp_2} \quad (x \ y)\,y$$

For More Material Join: @JEEAdvanced_2024

Effect of complex formation and hydrolysis on solubility : Solubility increases due to complex formation and hydrolysis of anion (cation/anion)

Solubility of AgBr in aq. sol. of NH_3.

Let solubility of AgBr is x

$$AgBr(s) \rightleftharpoons Ag\ (aq) \quad Br\ (qq) \qquad ; \quad K_{sp}$$

At eq. $\quad - \qquad (x \quad y) \qquad x$

$$Ag\ (aq) \quad 2NH_3(aq) \rightleftharpoons Ag(NH_3)_2(aq) \qquad ; \quad K_f$$

At eq. $\quad (x \quad y) \qquad (z \quad 2y) \qquad y$

Overall $\quad AgBr(s) \quad 2NH_3(aq) \rightleftharpoons Ag(NH_3)_2(aq) \quad Br\ (aq)$

Reaction at eq. $\quad - \qquad (z \quad 2y) \qquad y \qquad x$

Due to very large value of K_f we can assume $y \quad x$

$$\boldsymbol{K_{sp} \quad K_f \quad \frac{x^2}{(z \quad 2x)^2}}$$

For More Material Join: @JEEAdvanced_2024



Level 1

1. Morphine ($C_{17}H_{19}NO_3$), which is used medically to relieve pain is a base. What is its conjugate acid?
(a) $C_{17}H_{18}NO_3^+$ (b) $C_{17}H_{18}NO_3$ (c) $C_{17}H_{20}NO_3$ (d) $C_{17}H_{20}NO_3^+$

2. The conjugate base of $H_2PO_4^-$ is :
(a) H_3PO_4 (b) H_2PO_4 (c) HPO_4^{2} (d) PO_4^{3}

3. The strongest Bronsted base in the following anion is:
(a) CN (b) Cl (c) I (d) Br

4. Which salt can furnish H in its aqueous solution?
(a) NaH_2PO_2 (b) Na_2HPO_3 (c) Na_2HPO_4 (d) All of these

5. Which is the set of amphiprotic species?
(a) H_3O , HPO_4^{2} , HCO_3 (b) H_2O, HPO_3^{2} , H_2PO_2
(c) H_2PO_4, H_2PO_3, H_2O (d) All of these

6. The K_a values for HPO_4^{2} and HSO_3 are 4.8 10 ^{13} and 6.3 10 ^{8} respectively. Therefore, it follows the HPO_4^{2} is a ..... acid than HSO_3 and PO_4^{3} is a ..... base than SO_3^{2} .
(a) weaker, stronger (b) stronger, weaker (c) weaker, weaker (d) stronger, stronger

7. Given the following K_a values, determine which species is the strongest base ?
HSO_4 1.2 10 ^{2}, H_2PO_4 6.3 10 ^{8}, HCO_3 4.7 10 ^{11}
(a) CO_3^{2} (b) H_2SO_4
(c) SO_4^{2} (d) HPO_4^{2}

8. Given that K_w for water is 10 ^{13} M^2 at 62°C, compute the sum of pOH and pH for a neutral aqueous solution at 62°C :
(a) 7.0 (b) 13.30
(c) 14.0 (d) 13.0

9. The value of the ion product constant for water, (K_w) at 60°C is 9.6 10 ^{14} M^2. What is the $[H_3O^+]$ of a neutral aqueous solution at 60°C and an aqueous solution with a pH 7.0 at 60°C are respectively?
(a) 3.1 10 ^{8}, acidic (b) 3.1 10 ^{7}, neutral
(c) 3.1 10 ^{8}, basic (d) 3.1 10 ^{7}, basic

10. For pure water :
(a) pH increases while pOH decreases with rise in temperature
(b) pH decreases while pOH increases with rise in temperature
(c) both pH and pOH decreases with rise in temperature
(d) both pH and pOH increases with rise in temperature

11. A beer has a pH of 4.30. What is the $[H_3O^+]$?
(a) 3.0 10 ^{4} (b) 2.0 10 ^{4} (c) 2.0 10 ^{5} (d) 5.0 10 ^{5}

For More Material Join: @JEEAdvanced_2024

12. The hydrogen ion concentration of the oceans is about 2 10^{-9} M. What is the pH?
(a) 8.85 (b) 9.3 (c) 7.85 (d) 8.7

13. The hydroxide ion concentration of a wine is 8 10^{-11} M. What is the pH of the wine?
(a) 2.10 (b) 2.9 (c) 3.9 (d) 4.9

14. The pH of a solution is 5. To this solution acid was added so that its pH value becomes 2.0. The increase in H concentration is :
(a) 100 times (b) 5 times (c) 2.5 times (d) 1000 times

15. A solution has a pH 9. It is 1000 times more basic than the original solution. What was the pH of the original solution?
(a) 12 (b) 6 (c) 9 (d) 10

16. Equal volumes of two HCl solutions of pH 3 and pH 5 were mixed. What is the pH of the resulting solution?
(a) 3.5 (b) 4.0 (c) 4.5 (d) 3.3

17. pOH of 0.002 M HNO_3 is :
(a) 11 log 2 (b) 11 log 2 (c) 3 log 2 (d) None of these

18. Number of equivalents of HCl present in 100 mL of its solution whose pH is 4 :
(a) 10^{-4} (b) 10^{-3} (c) 10^{-2} (d) 10^{-5}

19. To a 10 mL of 10^{-3} N H_2SO_4 solution water has been added to make the total volume of one litre. Its pOH would be :
(a) 3 (b) 12 (c) 9 (d) 5

20. The pH of a solution of H_2SO_4 is 1. Assuming complete ionisation, find the molarity of H_2SO_4 solution :
(a) 0.1 (b) 0.2 (c) 0.05 (d) 2.0

21. pH of a strong diprotic acid (H_2A) at concentrations :
(i) 10^{-4} M, (ii) 10^{-4} N
are respectively :
(a) 3.7 and 4.0 (b) 4 and 3.7 (c) 4 and 4 (d) 3.7 and 3.7

22. Calcium hydroxide is a strong base. Compute [Ca^{2}] and [OH] for a solution that is prepared by dissolving 0.60 g of $Ca(OH)_2$ in enough water to make a 1500 mL of solution.
[Atomic mass : Ca 40, O 16, H 1]
(a) 5.4 10^{-3}, 9.1 10^{-13} (b) 5.4 10^{-3}, 1.08 10^{-2}
(c) 5.4 10^{-3}, 5.4 10^{-3} (d) 8.1 10^{-3}, 8.1 10^{-3}

23. pH of 10^{-6} M HCl($aq.$) is :
(a) just less than 6 (b) exactly equal to 6
(c) just greater than 6 (d) just less than 7

24. 10^{-5} M NaOH solution at 25°C is diluted 1000 times. The pH of the resultant solution will :
(a) be equal to 8 (b) lie between 7 and 8
(c) lie between 6 and 7 (d) remain unchanged

25. 4.0 g of NaOH and 4.9 g of H_2SO_4 are dissolved in water and volume is made upto 250 mL. The pH of this solution is:
(a) 7.0 (b) 1.0 (c) 2.0 (d) 12.0

For More Material Join: @JEEAdvanced_2024

26. A 25.0 mL sample of 0.10 *M* HCl is titrated with 0.10 *M* NaOH. What is the pH of the solution at the points where 24.9 and 25.1 mL of NaOH have been added?
(a) 3.70, 10.70 (b) 3.30, 10.30 (c) 3.70, 10.30 (d) 3.0, 11.0

27. What is the pH of a solution in which 25.0 mL of 0.1 *M* NaOH is added to 25 mL of 0.08 *M* HCl and final solution is diluted to 500 mL?
(a) 3 (b) 11 (c) 12 (d) 13

28. What is the pH of a solution in which 10.0 mL of 0.010 *M* $Sr(OH)_2$ is added to 10.0 mL of 0.010 *M* HCl?
(a) 2.30 (b) 1.50 (c) 11.70 (d) 7.00

29. At 90°C, pure water has [H] 10 ^{6} *M*. If 100 mL of 0.2 *M* HCl is added to 200 mL of 0.1 *M* KOH at 90°C then pH of the resulting solution will be :
(a) 5 (b) 6 (c) 7 (d) None of these

30. What change will occur for the following reaction if the hypochlorous acid solution is diluted from 0.1 to 0.01 *M*?

$$HOCl(aq.) \quad H_2O(l) \rightleftharpoons OCl^{-}(aq.) \quad H_3O^{+}(aq.)$$

(a) a decrease in the fraction of acid ionized
(b) an increase in the fraction of acid ionized
(c) no change in the fraction of acid ionized
(d) we can not predict

31. Given K_a values of 5.76 10 ^{10} and 4.8 10 ^{10} for NH_4 and HCN respectively. What is the equilibrium constant for the following reaction?

$$NH_4(aq.) \quad CN \ (aq.) \rightleftharpoons NH_3(aq.) \quad HCN(aq.)$$

(a) 0.83 (b) 1.2 (c) 8.0 10 ^{11} (d) 27.6 10 ^{10}

32. Which is the strongest acid (pK_a value is given)?
(a) HCOOH [3.77] (b) C_6H_5COOH [4.22]
(c) CH_3COOH [4.7] (d) CH_3CH_2COOH [4.88]

33. Given : Enthalpy of ionization of two acids :
H (HCN) 45.2 kJ mol ^{1}
H (CH_3COOH) 2.1 kJ mol ^{1}
Which relationship for the two acids is true ?
(a) pK_a(HCN) pK_a(CH_3COOH) (b) pK_a(HCN) pK_a(CH_3COOH)
(c) pK_a(HCN) pK_a(CH_3COOH) (d) pK_a(HCN) $\frac{45.2}{2.1}$ pK_a(CH_3COOH)

34. What is the hydronium ion concentration of a 0.25 *M* H*A* solution? (K_a 4 10 ^{8})
(a) 10 ^{4} (b) 10 ^{5} (c) 10 ^{7} (d) 10 ^{10}

35. What is the percent dissociation ( ) of a 0.01 *M* H*A* solution? (K_a 10 ^{4})
(a) 9.5% (b) 1% (c) 10.5% (d) 17%

36. Given the two concentration of HCN (K_a 10 ^{9}) are 0.1 *M* and 0.001 *M* respectively. What will be the ratio of degree of dissociation?
(a) 1 (b) 0.1 (c) 0.003 (d) 0.01

For More Material Join: @JEEAdvanced_2024

37. A 0.10 *M* solution of HF is 8.0% dissociated What is the K_a?
(a) 6.4 10 ^{4} (b) 8.8 10 ^{4} (c) 6.95 10 ^{4} (d) 7.6 10 ^{4}

38. A weak base *M*OH of 0.1 *N* concentration shows a pH value of 9. What is the percentage degree of dissociation of the base?
(a) 0.01% (b) 0.001% (c) 0.1% (d) 0.02%

39. 0.01 *M* H*A* (*aq.*) is 2% dissociated; [OH] of solution is :
(a) 2 10 ^{4} (b) 10 ^{8} (c) 5 10 ^{11} (d) 5 10 ^{12}

40. If degree of dissociation is 0.01 of decimolar solution of weak acid H*A* then pK_a of acid is :
(a) 2 (b) 3 (c) 5 (d) 7

41. What concentration of HCOO is present in a solution of 0.01 *M* HCOOH (K_a 1.8 10 ^{4}) and 0.01 *M* HCl?
(a) 1.8 10 ^{3} (b) 10 ^{2} (c) 1.8 10 ^{4} (d) 10 ^{4}

42. Chose the correct code

Column-I		Column-II	
(P)	pK_b of X (K_a of HX 10 ^{6})	**(1)**	6.9
(Q)	pH of 10 ^{8} *M* HCl	**(2)**	8
(R)	pH of 10 ^{2} *M* acetic acid solution (Take K_a of acetic acid 1.6 10 ^{5})	**(3)**	10.7
(S)	pOH of a solution obtained by mixing equal volumes of solution with pH 3 and 5.	**(4)**	3.4

Codes :

	P	Q	R	S
(a)	1	2	4	3
(b)	4	3	2	1
(c)	2	1	4	3
(d)	1	2	3	4

43. How much water must be added to 300 mL of 0.2 *M* solution of CH_3COOH (K_a 1.8 10 ^{5}) for the D.O.D. ( ) of the acid to double?
(a) 600 mL (b) 900 mL (c) 1200 mL (d) 1500 mL

44. What is [NH_4] in a solution that contain 0.02 *M* NH_3 (K_b 1.8 10 ^{5}) and 0.01 *M* KOH?
(a) 9 10 ^{6} (b) 1.8 10 ^{5} (c) 3.6 10 ^{5} (d) None of these

45. A hand book states that the solubility of $RNH_2(g)$ in water at 1 atm and 0°C is 22.41 volumes of $RNH_2(g)$ per volume of water. (pK_b of RNH_2 4) Find the max. pOH that can be attained by dissolving RNH_2 in water :
(a) 1 (b) 2 (c) 4 (d) 6

For More Material Join: @JEEAdvanced_2024

46. The [H] of a resulting solution that is 0.01 *M* acetic acid (K_a 1.8 10^{-5}) and 0.01 *M* in benzoic acid (K_a 6.3 10^{-5}) :
(a) 9 10^{-4} (b) 81 10^{-4} (c) 9 10^{-5} (d) 2.8 10^{-3}

47. 6.0 g weak acid H*A* (mol. mass 60 g/mol.) is dissolved in water and formed 10 m^3 solution. If $K_{a(HA)}$ 10^{-9}, then pOH of solution is : [Given : log 4 0.6]
(a) 6.7 (b) greater than 6.7 and less than 7.0
(c) greater than 7.0 and less than 7.3 (d) greater than 7.3

48. Carbonic acid (H_2CO_3), a diprotic acid has K_{a_1} 4.0 10^{-7} and K_{a_2} 7.0 10^{-11}. What is the [HCO_3] of a 0.025 *M* solution of carbonic acid?
(a) 7.8 10^{-3} (b) 6.6 10^{-4} (c) 10^{-10} (d) 1.0 10^{-4}

49. Carbonic acid (H_2CO_3), a diprotic acid has K_{a_1} 4.0 10^{-7} and K_{a_2} 7.0 10^{-11}. What is the [CO_3^{2-}] of a 0.025 *M* solution of carbonic acid?
(a) 5.5 10^{-9} (b) 5.5 10^{-8} (c) 7.0 10^{-9} (d) 7.0 10^{-11}

50. Selenious acid (H_2SeO_3), a diprotic acid has K_{a_1} 3.0 10^{-3} and K_{a_2} 5.0 10^{-8}. What is the [OH] of a 0.30 *M* solution of selenious acid?
(a) 2.85 10^{-3} (b) 5.0 10^{-6} (c) 3.5 10^{-12} (d) 3.5 10^{-13}

51. Which of the hydrated species may exist?
I : $H_5O_2^+$ II : H_3O^+ III : H_3O_2 IV : $H_7O_3^+$
(a) II only (b) I and II (c) I, II and IV (d) I, II, III and IV

52. Consider the following salts. Which one(s) when dissolved in water will produce an acidic solution?
1. NH_4Cl 2. $KHSO_4$ 3. NaCN 4. KNO_3
(a) 2 and 3 (b) 1 and 2 (c) only 3 (d) 2 and 4

53. Consider the following salts. Which one(s) when dissolved in water will produce a basic solution?
1. $RbClO_4$ 2. $NaNO_2$ 3. NH_4Cl 4. NaCl
(a) 1 and 3 (b) only 2 (c) 1 and 2 (d) 3 and 4

54. At 25°C dissociation constants of acid H*A* and base *B*OH in aqueous solution are same. The pH of 0.01 *M* solution of H*A* is 5. The pOH of 10^{-4} *M* solution of *B*OH at the same temperature is:
(a) 3.5 (b) 4 (c) 6 (d) None of these

55. Which of the following solutions has the highest pH?
(a) 0.2 *M* $HClO_4$ (b) 0.20 *M* CH_3COOH
(c) 0.020 *M* HCl (d) 0.2 *M* NaCl

56. pH of solutions of four sodium salts Na*W*, Na*X*, Na*Y* and Na*Z* were found to be 7.0, 9.0, 10.0 and 11.0 respectively. If each solution has concentration 0.1 *M*, the weakest acid is :
(a) H*W* (b) H*X* (c) H*Y* (d) H*Z*

57. The pH values 0.1 *M* solution of HCOONa (I), HCOOH (II), CH_3COONH_4 (III), NaOH (IV), HCl (V), will be in the order :

For More Material Join: @JEEAdvanced_2024

(a) IV > III > I > II > V (b) IV > I > III > II > V
(c) II > III > I > IV > V (d) V > II > III > I > IV

58. pH of an aqueous NaCl solution at 50°C should be :
(a) 7 (b) > 7 (c) < 7 (d) 0

59. Upon hydrolysis of sodium carbonate, the reaction takes place between:
(a) Na and water (b) Na and OH (c) CO_3^{2} and water (d) CO_3^{2} and H

60. The solution of blue vitrol in water is acidic because:
(a) $CuSO_4$ reacts with water (b) Cu^{2} reacts with water
(c) SO_4^{2} reacts with water (d) $CuSO_4$ removes OH ions from water

61. 1 mL of 0.1 *N* HCl is added to 999 mL solution of NaCl. The pH of the resulting solution will be :
(a) 7 (b) 4 (c) 2 (d) 1

62. If a salt of strong acid and weak base hydrolyses appreciably (0.1), which of the following formula is to be used to calculate degree of hydrolysis ' '?
(a) $\sqrt{\frac{K_w}{K_a \ a}}$ (b) $\sqrt{\frac{K_w}{K_b \ a}}$ (c) $\sqrt{\frac{K_w}{K_a \ K_b}}$ (d) None of these

63. The correct formula to calculate the hydroxyl ion concentration of an aqueous solution of NH_4NO_3 is :
(a) $\sqrt{\frac{C \ K_w}{K_b}}$ (b) $\sqrt{\frac{K_w \ K_b}{C}}$ (c) $\sqrt{\frac{C \ K_w}{K_a}}$ (d) $\sqrt{\frac{K_a \ K_w}{C}}$

64. [H] $\sqrt{\frac{K_w K_a}{C}}$ is suitable for
(a) NaCl, NH_4Cl (b) CH_3COONa , NaCN
(c) CH_3COONa, $(NH_4)_2SO_4$ (d) CH_3COONH_4, $(NH_4)_2CO_3$

65. What is the hydrolysis constant of the OCl ion? The ionization constant of HOCl is 3.0 10^{8}.
(a) 3.33 10^{8} (b) 3.33 10^{7} (c) 3.0 10^{7} (d) 3.33 10^{6}

66. What is the pH of a 0.10 *M* C_6H_5O solution? The K_a of C_6H_5OH is 1.0 10^{10}.
(a) 10.51 (b) 11.04 (c) 11.50 (d) 12

67. Calculate the [OH] in 0.01 *M* aqueous solution of NaOCN (K_b for OCN 10^{10}) :
(a) 10^{6} *M* (b) 10^{7} *M* (c) 10^{8} *M* (d) None of these

68. What is the ionization constant of an acid if the hydronium ion concentration of a 0.40 *M* solution is 1.40 10^{4} *M*?
(a) 1.96 10^{8} (b) 1.22 10^{9} (c) 4.90 10^{8} (d) 1.40 10^{6}

69. The degree of hydrolysis of 0.1 *M* RNH_3Cl solution is 1.0%. If the concentration of RNH_3Cl is made 0.4 *M*, what is the new degree of hydrolysis (in percentage)?
(a) 0.01 (b) 0.001 (c) 0.2 (d) 0.5

70. % hydrolysis of 0.1*M* CH_3COONH_4, when $K_a(CH_3COOH)$ $K_b(NH_4OH)$ 1.8 10^{5} is:
(a) 0.55 (b) 7.63 (c) 0.55 10^{2} (d) 7.63 10^{3}

For More Material Join: @JEEAdvanced_2024

71. The enthalpy of neutralisation of four acids H*A*, H*B*, H*C* and H*D* with NaOH are –13, –12, –11, –10 kcal/mol. Which salt has maximum degree of hydrolysis?
(a) 1*M* Na*A* (b) 1*M* Na*B* (c) 1*M* Na*C* (d) 1*M* Na*D*

72. Calculate [H] at equivalent point between titration of 0.1 *M*, 25 mL of weak acid H*A* ($K_{a(HA)}$ 10 ^{5}) with 0.05 *M* NaOH solution :
(a) 3 10 ^{9} (b) 1.732 10 ^{9} (c) 8 (d) 10

73. When a salt of weak acid and weak base is dissolved in water at 25°C, the pH of the resulting solution will always :
(a) be 7 (b) be greater than 7
(c) be less than 7 (d) depend upon K_a and K_b values

74. What will be the pH of an aqueous solution of 1.0 *M* ammonium formate?
Given : pK_a 3.8 and pK_b 4.8
(a) 7.5 (b) 3.4 (c) 6.5 (d) 10.2

75. What will be the pH and % (degree of hydrolysis) respectively for the salt *BA* of 0.1 *M* concentration? Given : K_a for H*A* 10 ^{6} and K_b for *B*OH 10 ^{6}
(a) 5, 1% (b) 7, 10% (c) 9, 0.01% (d) 7, 0.01%

76. The percentage degree of hydrolysis of a salt of weak acid (H*A*) and weak base (*B*OH) in its 0.1 *M* solution is found to be 10%. If the molarity of the solution is 0.05 *M*, the percentage hydrolysis of the salt should be :
(a) 5% (b) 10% (c) 20% (d) None of these

77. What is the hydronium ion concentration of a 0.02 *M* solution of Cu^{2} solution of copper(II) perchlorate? The acidity constant of the following reaction is 5 10 ^{9}.

$$Cu^{2}\ (aq.)\quad 2H_2O(l) \rightleftharpoons Cu(OH)\ (aq.)\quad H_3O^{+}(aq.)$$

(a) 1 10 ^{5} (b) 7 10 ^{4} (c) 5 10 ^{4} (d) 1 10 ^{4}

78. What is the acidity constant for the following reaction given that the hydronium ion concentration of a 0.04 *M* solution of Ni^{2} solution of nickel(II) perchlorate is 4.5 10 ^{6}?

$$Ni^{2}\ (aq.)\quad 2H_2O(l) \rightleftharpoons Ni(OH)^{+}(aq.)\quad H_3O^{+}(aq.)$$

(a) 2 10 ^{12} (b) 4 10 ^{6} (c) 5 10 ^{12} (d) 5 10 ^{10}

79. Calculate the pH at 25°C of a solution that is 0.10 *M* in $Fe(NO_3)_3$. The acid dissociation constant for the reaction given below is 1.0 10 ^{3}.

$$[Fe(H_2O)_6]^{3+} + H_2O(l) \rightleftharpoons H_3O^{+}(aq.)\quad [Fe(H_2O)_5(OH)]^{2+}$$

(a) 2.00 (b) 2.02 (c) 2.30 (d) 2.50

80. Approximate pH of 0.01 *M* NaH*A* is calculated by :
(K_{a_1} 10 ^{6} and K_{a_2} 10 ^{8} are ionization constants of H_2A)
(a) pH 7 $\frac{pK_{a_1}}{2}$ $\frac{\log C}{2}$ (b) pH 7 $\frac{pK_{a_1}}{2}$ $\frac{\log C}{2}$
(c) pH $\frac{pK_{a_1} \quad pK_{a_2}}{2}$ (d) None of these

For More Material Join: @JEEAdvanced_2024

81. H_3PO_4 is a weak triprotic acid; approximate pH of 0.1 *M* $Na_2HPO_4(aq.)$ is calculated by :

(a) $\frac{1}{2}[pK_{a_1} \quad pK_{a_2}]$ (b) $\frac{1}{2}[pK_{a_2} \quad pK_{a_3}]$ (c) $\frac{1}{2}[pK_{a_1} \quad pK_{a_3}]$ (d) $pK_{a_1} \quad pK_{a_2}$

82. Which of the following is a buffer solution?

(a) 500 mL of 0.1 *N* CH_3COOH + 500 mL of 0.1 *N* NaOH

(b) 500 mL of 0.1 *N* CH_3COOH + 500 mL of 0.1 *N* HCl

(c) 500 mL of 0.1 *N* CH_3COOH + 500 mL of 0.2 *N* NaOH

(d) 500 mL of 0.2 *N* CH_3COOH + 500 mL of 0.1 *N* NaOH

83. If 20 mL of 0.1 *M* NaOH is added to 30 mL of 0.2 *M* CH_3COOH (pK_a 4.74), the pH of the resulting solution is :

(a) 4.44 (b) 9.56 (c) 8.96 (d) 9.26

84. H_2CO_3 $NaHCO_3$ found in blood helps in maintaining pH of the blood close to 7.4. An excess of acid entering the blood stream is removed by:

(a) HCO_3 (b) H_2CO_3 (c) H ion (d) CO_3^{2-} ion

85. 100 mL of 0.02 *M* benzoic acid (pK_a 4.20) is titrated using 0.02 *M* NaOH, pH after 50 mL and 100 mL of NaOH have been added are:

(a) 3.50, 7 (b) 4.2, 7 (c) 4.2, 8.1 (d) 4.2, 8.25

86. What is the pH of a solution of 0.28 *M* acid and 0.84 *M* of its conjugate base if the ionization constant of acid is 4 10^{-4}?

(a) 3.88 (b) 3.34 (c) 7 (d) 10.12

87. The toxic compound 2,4-dinitrophenol has K_a 10^{-4}. In an experiment, a buffer solution of 2,4-dinitrophenol was prepared with the pH adjusted to 5. Calculate the ratio of the concentrations of the dissociated ion to the undissociated acid :

(a) 0.01 (b) 0.1 (c) 10 (d) 100

88. Equilibrium constant for the following reaction is 1 10^{-9} :

$$C_5H_5N(aq.) \quad H_2O(l) \rightleftharpoons C_5H_5NH^+(aq.) \quad OH \ (aq.)$$

Determine the moles of pyridinium chloride ($C_5H_5N.HCl$) that should be added to 500 mL solution of 0.4 *M* pyridine (C_5H_5N) to obtain a buffer solution of pH 5 :

(a) 0.1 mole (b) 0.2 mole (c) 0.3 mole (d) 0.4 mole

89. Which one of the following mixture does not act as a buffer solution?

(a) Boric acid and borax

(b) Sodium phosphate & disodium hydrogen phosphate

(c) Sodium propionate and propionic acid

(d) Sod. acetate and sodium propionate

90. The acid dissociation constant of uric acid is K_a 4.0 10^{-6} *M*. The pH of a sample of urine is 6.0. What is the ratio of concentration of urate ion to uric acid in the urine?

(a) 2.0 (b) 4.0 (c) 6.0 (d) 0.25

91. CH_3NH_2 (0.12 mole, pK_b 3.3) is added to 0.08 moles of HCl and the solution is diluted to one litre, resulting pH of solution is :

(a) 10.7 (b) 3.6 (c) 10.4 (d) 11.3

For More Material Join: @JEEAdvanced_2024

92. An aqueous solution at room temperature contains 0.1 M NH_4Cl and 0.01M NH_4OH (pK_b 5), the pH of the solution is :
(a) 7.5 (b) 6.8 (c) 6.5 (d) 8.0

93. Which of the following will decrease with dilution at a given temperature?
(a) pH of 10 ^{3} M acetic acid solution
(b) pH of 10 ^{3} M aniline solution
(c) degree of dissociation of 10 ^{3} M acetic acid
(d) degree of dissociation of 10 ^{3} M aniline solution

94. A 1L solution contains 0.2 M NH_4OH and 0.2 M NH_4Cl. If 1.0 mL of 0.001 M HCl is added to it, what will be the [OH] of the resulting solution (K_b 2 10 ^{5})
(a) 2 10 ^{5} (b) 5 10 ^{10} (c) 2 10 ^{3} (d) None of these

95. 0.1 M formic acid solution is titrated against 0.1 M NaOH solution. What would be the difference in pH between 1/5 and 4/5 stages of neutralization of acid ?
(a) 2 log 3/4 (b) 2 log 1/5 (c) log 1/3 (d) 2 log 4

96. In the neutralization process of H_3PO_4 and NaOH, the number of buffers formed will be:
(a) 3 (b) 1 (c) 2 (d) 4

97. A buffer solution is made up of acetic acid [pK_a 5] having conc. 1.5 M and sodium acetate having conc. 0.15 M. What is the number of OH ions present in 1 litre solution?
(a) 10 ^{10} N_A (b) 10 ^{4} N_A (c) 10 ^{3} N_A (d) 10 ^{6} N_A

98. The pH of a solution of 0.10 M CH_3COOH increases when which of the following substances is added?
(a) $NaHSO_4$ (b) $HClO_4$ (c) KNO_3 (d) K_2CO_3

99. H ion concentration of water does not change by adding :
(a) CH_3COONa (b) $NaNO_3$ (c) NaCN (d) Na_2CO_3

100. pK_a of NH_4 is 9.26. Hence, effective range for NH_4OH – NH_4Cl buffer is about pH :
(a) 8.26 to 10.26 (b) 4.74 to 5.74 (c) 3.74 to 5.74 (d) 8.26 to 9.26


101. 1.0 L solution is prepared by mixing 61 g benzoic acid (pK_a 4.2) with 72 g of sodium benzoate and then 300 mL 1.0 M HBr solution was added. The pH of final solution is:
(a) 3.6 (b) 3.8 (c) 4.2 (d) 4.8


102. The pH of a solution containing 0.4 M HCO_3 and 0.2 M CO_3^{2} is :
[K_{a_1} (H_2CO_3) 4 10 ^{7}; K_{a_2} (HCO_3) 4 10 ^{11}]
(a) 10.4 (b) 10.1 (c) 6.1 (d) 10.7


103. The pH of the resultant solution of 20 mL of 0.1 M H_3PO_4 and 20 mL of 0.1 M Na_3PO_4 is :
(a) pK_{a_1} log 2 (b) pK_{a_1} (c) pK_{a_2} (d) $\frac{pK_{a_1} \quad pK_{a_2}}{2}$


104. Which one of the following curves represents the graph of pH during the titration of NaOH and HCl($aq.$)?

For More Material Join: @JEEAdvanced_2024


(a) 


(b) 


(c) 


(d) 

105. When 100 mL of 0.1 *M* NaCN solution is titrated with 0.1 *M* HCl solution the variation of pH of solution with volume of HCl added will be :

(a) 

(b) 

(c) 

(d) 

106. The best indicator for the detection of end point in titration of a weak acid and a strong base is :

(a) Methyl orange (3.1 to 4.4) (b) Methyl red (4.2 to 6.3)

(c) Bromothymol blue (6 to 7.6) (d) Phenolphthalein (8.2 to 10)

107. Select the best indicator from the given table for titration of 20 mL of 0.02 *M* CH_3COOH with 0.02 *M* NaOH. Given pK_a (CH_3COOH) 4.74

	Indicator	pH range
(I)	Bromothymol blue	6.0–7.6
(II)	Thymolphthalein	9.3–10.5
(III)	Malachite green	11.4–13
(IV)	M-Cresol purple	7.4–9.0

(a) I (b) II (c) III (d) IV

108. Bromothymol blue is an indicator with a K_a value of 6 10^{5}. What % of this indicator is in its basic form at a pH of 5 ?

(a) 40 (b) 85.7 (c) 14.3 (d) 60

For More Material Join: @JEEAdvanced_2024

109. An acid-base indicator has K_{HIn} 3.0 10^{-5}. The acid form of the indicator is red and the basic form is blue. The change in [H] required to change the indicator from 75% red to 75% blue is :

(a) 8 10^{-5} *M* (b) 9 10^{-5} *M* (c) 1 10^{-5} *M* (d) 3 10^{-4} *M*

110. An acid-base indicator which is a weak acid has a pK_a value 5.45. At what concentration ratio of sodium acetate to acetic acid would the indicator show a colour half-way between those of its acid and conjugate base forms ? [pK_a of acetic acid 4.75]

(a) 4 : 1 (b) 6 : 1 (c) 5 : 1 (d) 3 : 1

111. A 20.0 mL sample of a 0.20 *M* solution of the weak diprotic acid H_2A is titrated with 0.250 *M* NaOH. The concentration of solution at the second equivalent point is :

(a) 0.10 *M* Na*HA* (b) 0.153 *M* Na_2A (c) 0.10 *M* Na_2A (d) 0.0769 *M* Na_2A

112. During the titration of a weak diprotic acid (H_2A) against a strong base (NaOH), the pH of the solution half-way to the first equivalent point and that at the first equivalent point are given respectively by :

(a) pK_{a_1} and pK_{a_1} pK_{a_2} (b) $\sqrt{K_{a_1}C}$ and $\frac{pK_{a_1} \quad pK_{a_2}}{2}$

(c) pK_{a_1} and $\frac{pK_{a_1} \quad pK_{a_2}}{2}$ (d) pK_{a_1} and pK_{a_2}

113. In which of the following cases is the solution of AgCl unsaturated?

(a) [Ag][Cl] K_{sp} (b) [Ag][Cl] K_{sp} (c) [Ag][Cl] K_{sp} (d) [Ag][Cl] K_{sp}

114. When equal volumes of the following solutions are mixed, the precipitation of AgCl (K_{sp} 1.8 10^{-10}) will occur with :

(a) 10^{-4} *M* (Ag) and 10^{-4} *M* (Cl) (b) 10^{-5} *M* (Ag) and 10^{-5} *M* (Cl)

(c) 10^{-5} *M* (Ag) and 10^{-6} *M* (Cl) (d) 10^{-10} *M* (Ag) and 10^{-10} *M* (Cl)

115. Choose the correct set of True/False for following statements:

(i) Silver chloride is more soluble in very concentrated sodium chloride solution than in pure water.

(ii) The pH of a buffer solution does not change on addition of small amount of an acid or a base.

(iii) Addition of NH_4Cl does not affect the pH of a solution of NH_4OH

(iv) Degree of hydrolysis of ammonium acetate does not depend upon the concentration of ammonium acetate solution.

(v) A mixture of acetic acid and sodium acetate can act as buffer solution.

(a) TTFTT (b) FTTTF (c) TFTFT (d) FFTTT

116. A 1 litre solution containing NH_4Cl and NH_4OH has hydroxide ion concentration of 10^{-6} mol/litre. Which of the following hydroxides could be precipitated when the solution is added to 1 litre solution of 0.1 *M* metal ions?

(I) $Ba(OH)_2$ (K_{sp} 5 10^{-3}) (II) $Ni(OH)_2$ (K_{sp} 1.6 10^{-16})

(III) $Mn(OH)_2$ (K_{sp} 2 10^{-13}) (IV) $Fe(OH)_2$ (K_{sp} 8 10^{-16})


(a) I, II, IV (b) IV (c) II and IV (d) II, III, IV


For More Material Join: @JEEAdvanced_2024


117. 150 mL of 0.0008 *M* ammonium sulphate is mixed with 50 mL of 0.04 *M* calcium nitrate. The ionic product of $CaSO_4$ will be : (K_{sp} 2.4 10^{5} for $CaSO_4$)


(a) K_{sp} (b) K_{sp} (c) K_{sp} (d) None of these

118. In a saturated solution of AgCl, NaCl is added gradually. The concentration of Ag is plotted against the concentration of Cl . The graph appears as :

(a) 

(b) 

(c) 

(d) 

119. K_{sp} of AgCl is 1 10^{10}. Its solubility in 0.1*M* KNO_3 will be :

(a) 10^{5} moles/litre (b) 10^{5} moles/litre (c) 10^{5} moles/litre (d) None of these

120. 50 mL of a solution containing 10^{3} mole of Ag is mixed with 50 mL of a 0.1 *M* HCl solution. How much Ag remains in solution? (K_{sp} of AgCl 1.0 10^{10})

(a) 2.5 10^{9} (b) 2.5 10^{7} (c) 2.5 10^{8} (d) 2.5 10^{10}

121. The solubility of a sparingly soluble salt A_xB_y in water is '*S*' moles per litre. The solubility product has the value :

(a) S^2 (b) $x^y y^x S^{x\ y}$ (c) $x^x y^y S^{x\ y}$ (d) $S^{x\ y}$

122. What is the molarity of a saturated solution of $CaCO_3$? (K_{sp} 2.8 10^{9})

(a) 2.6 10^{5} (b) 2.8 10^{9} (c) 5.2 10^{5} (d) 5.6 10^{9}

123. K_{sp} of $Zr_3(PO_4)_4$ in terms of solubility (*S*) is :

(a) $108\ S^7$ (b) $4\ S^3$ (c) $6912\ S^7$ (d) None of these

124. The solubility of electrolytes MX_1, MX_2 and MX_3 is 1 10^{3} moles per litre. Hence their respective solubility products are :

(a) $10^{6}, 4\ 10^{9}, 27\ 10^{12}$ (b) $10^{9}, 4\ 10^{9}, 32\ 10^{12}$

(c) $10^{9}, 8\ 10^{8}, 32\ 10^{12}$ (d) None of these

125. A saturated solution of $Ca_3(PO_4)_2$ has $[Ca^{2}]$ 2 10^{8} *M* and $[PO_4^{3}]$ 1.6 10^{5} *M* K_{sp} of $Ca_3(PO_4)_2$ is :

(a) 3.2 10^{13} (b) 3.2 10^{34} (c) 2.048 10^{33} (d) None of these

126. Which of the following is most soluble in water?

(a) $Ba_3(PO_4)_2$ (K_{sp} 6 10^{39}) (b) ZnS (K_{sp} 7 10^{16})

(c) $Fe(OH)_3$ (K_{sp} 6 10^{38}) (d) $Ag_3(PO_4)$ (K_{sp} 1.8 10^{18})

127. Silver ions are added to a solution with [Br] [Cl] $[CO_3^{2}]$ $[AsO_4^{3}]$ 0.1*M*. Which compound will precipitate with lowest [Ag]?

(a) AgBr (K_{sp} 5 10^{13}) (b) AgCl (K_{sp} 1.8 10^{10})

(c) Ag_2CO_3 (K_{sp} 8.1 10^{12}) (d) Ag_3AsO_4 (K_{sp} 1 10^{22})

For More Material Join: @JEEAdvanced_2024

128. The solubility product of different sparingly soluble salts are given as under :

S. No.	Formula Type	Solubility product
1.	AB	4.0 10 ^{20}
2.	A_2B	3.2 10 ^{11}
3.	AB_3	2.7 10 ^{31}

The correct increasing order of solubility is :
(a) 1, 3, 2 (b) 2, 1, 3 (c) 1, 2, 3 (d) 3, 1, 2

129. If K_{sp} for $HgSO_4$ is 6.4 10 ^{5}, then solubility of this substance in mole per m ^{3} is :
(a) 8 10 ^{3} (b) 6.4 10 ^{5} (c) 8 10 ^{6} (d) None of these

130. The solubility of $Ba_3(AsO_4)_2$ (formula mass 690) is 6.9 10 ^{2} g/100 mL. What is the K_{sp} ?
(a) 1.08 10 ^{11} (b) 1.08 10 ^{13} (c) 1.0 10 ^{15} (d) 6.0 10 ^{13}

131. The solubility of $AgBrO_3$ (formula mass 236) is 0.0072 g in 1000 mL. What is the K_{sp} ?
(a) 2.2 10 ^{8} (b) 3.0 10 ^{10} (c) 3.0 10 ^{5} (d) 9.3 10 ^{10}

132. The solubility of PbF_2 (formula mass 245) is 0.46 g/L. What is the solubility product?
(a) 1.1 10 ^{10} (b) 2.6 10 ^{8} (c) 1.1 10 ^{7} (d) 6.8 10 ^{9}

133. How many grams of MgC_2O_4 (formula mass 112) will dissolve in 1.5 L of water? (K_{sp} 8.1 10 ^{5})
(a) 1.0 (b) 1.29 (c) 1.512 (d) 4.65

134. What is the molarity of F ions in a saturated solution of BaF_2? (K_{sp} 1.0 10 ^{6})
(a) 1.0 10 ^{2} (b) 1.0 10 ^{3} (c) 1.26 10 ^{2} (d) 6.3 10 ^{3}

135. What is the molarity of F in a saturated solution of InF_3? (K_{sp} 7.9 10 ^{10})
(a) 2.3 10 ^{3} (b) 8.3 10 ^{3} (c) 1.0 10 ^{3} (d) 7.0 10 ^{3}

136. What is the pH of a saturated solution of $Cu(OH)_2$? (K_{sp} 2.6 10 ^{19})
(a) 6.1 (b) 7.30 (c) 8.42 (d) 7.90

137. The solubility product of AgCl is 10 ^{10} M^2. The minimum volume (in m ^{3}) of water required to dissolve 14.35 mg of AgCl is approximately :
(a) 0.01 (b) 0.1 (c) 100 (d) 10

138. What is the molar solubility of $Fe(OH)_2$ (K_{sp} 8.0 10 ^{16}) at pH 13.0?
(a) 8.0 10 ^{18} (b) 8.0 10 ^{15} (c) 8.0 10 ^{17} (d) 8.0 10 ^{14}

139. What is the minimum pH necessary to cause a precipitate of $Pb(OH)_2$ (K_{sp} 1.2 10 ^{5}) to form in a 0.12 M $PbCl_2$ solution?
(a) 12.4 (b) 10.8 (c) 12.0 (d) 11.1

140. Which of the following would increase the solubility of $Pb(OH)_2$?
(a) Add hydrochloric acid
(b) Add a solution of $Pb(NO_3)_2$
(c) Add a solution of NaOH
(d) None of the above–the solubility of a compound is constant at constant temperature

For More Material Join: @JEEAdvanced_2024

141. What is the molar solubility of Ag_2CO_3 (K_{sp} 4 10 ^{13}) in 0.1 M Na_2CO_3 solution?
(a) 10 ^{6} (b) 10 ^{7} (c) 2 10 ^{6} (d) 2 10 ^{7}

142. What is the concentration of Pb^{2} when $PbSO_4$ (K_{sp} 1.8 10 ^{8}) begins to precipitate from a solution that is 0.0045 M in SO_4^{2} ?
(a) 4.0 10 ^{8} M (b) 1.0 10 ^{6} M (c) 2.0 10 ^{8} M (d) 4.0 10 ^{6} M

143. What is the concentration of Ba^{2} when BaF_2 (K_{sp} 1.0 10 ^{6}) begins to precipitate from a solution that is 0.30 M F ?
(a) 9.0 10 ^{7} (b) 3.3 10 ^{5} (c) 1.1 10 ^{5} (d) 3.0 10 ^{7}

144. Solubility of AgCl in 0.2 M NaCl is x and that in 0.1 M $AgNO_3$ is y then which of the following is correct?
(a) x y (b) x y (c) x y (d) We cannot predict

145. What is the molarity of $Fe(CN)_6^{4-}$ in a saturated solution of $Ag_4[Fe(CN)_6]$? (K_{sp} 1.6 10 ^{41})
(a) 1.6 10 ^{8} (b) 5.2 10 ^{8} (c) 2.0 10 ^{8} (d) 2.3 10 ^{9}

146. At 25°C, K_{sp} for $PbBr_2$ is equal to 8 10 ^{5}. If the salt is 80% dissociated, what is the solubility of $PbBr_2$ in mol/litre?
(a) $\left(\frac{10^{4}}{1.6\ 1.6}\right)^{1/3}$ (b) $\left(\frac{10^{5}}{1.6\ 1.6}\right)^{1/3}$ (c) $\left(\frac{10^{4}}{0.8\ 0.8}\right)^{1/3}$ (d) $\left(\frac{10^{5}}{1.6\ 1.6}\right)^{1/2}$

147. What is the molar solubility of $Mn(OH)_2$(K_{sp} 4.5 10 ^{14}) in a buffer solution containing equal amounts of NH_4 and NH_3 (K_b 1.8 10 ^{5})?
(a) 3.0 10 ^{4} (b) 1.38 10 ^{4} (c) 1.38 10 ^{3} (d) 7.3 10 ^{4}

148. Find moles of NH_4Cl required to prevent $Mg(OH)_2$ from precipitating in a litre of solution which contains 0.02 mole of NH_3 and 0.001 mole of Mg^{2} ions.
Given : K_b (NH_3) 10 ^{5}; K_{sp} [$Mg(OH)_2$] 10 ^{11}.
(a) 10 ^{4} (b) 2 10 ^{3} (c) 0.02 (d) 0.1

149. What mass of AgI will dissolve in 1.0 L of 1.0 M NH_3? Neglect change in conc. of NH_3.
[Given : K_{sp} (AgI) 1.5 10 ^{16}; K_f [$Ag(NH_3)_2$] 1.6 10^{7}]; (At. mass Ag 108; I 127)
(a) 4.9 10 ^{5} g (b) 0.0056 g (c) 0.035 g (d) 0.011 g

150. Consider the following statement and select correct option :
(I) K_{sp} of $Fe(OH)_3$ in aqueous solution is 3.8 10 ^{38} at 298 K. The concentration of Fe^{3} will increase when [H] ion concentration decreases.
(II) In a mixture of NH_4Cl and NH_4OH in water, a further amount of NH_4Cl is added. The pH of the mixture will decreases.
(III) An aqueous solution of each of the following salts (NH_4I, HCOOK) will be basic, acidic respectively.
(a) only I is correct (b) only II is correct
(c) only III is correct (d) II and III are correct

For More Material Join: @JEEAdvanced_2024



Level 2

1. Equilibrium constants of T_2O (T or ^3_1H is an isotope of ^1_1H) and H_2O are different at 298 K. Let at 298 K pure T_2O has pT (like pH) is 7.62. The pT of a solution prepared by adding 10 mL of 0.2 *M* TCl to 15 mL of 0.25 *M* NaOT is :
(a) 2 log 7 (b) 14 log 7 (c) 13.24 log 7 (d) 13.24 log 7

2. Liquid NH_3 dissociation to a slight extent. At a certain temp. its self dissociation constant $K_{SDC\ (NH_3)}$ 10^{-30}. The number of NH_4^+ ions are present per 100 cm^3 of pure liquid are :
(a) 10^{-15} (b) 6.022×10^8 (c) 6.022×10^7 (d) 6.022×10^6

3. To what volume of 10 litre of 0.5 *M* CH_3COOH (K_a 1.8×10^{-5}) be diluted in order to double the hydroxide ion concentration :
(a) 20 L (b) 30 L (c) 40 L (d) None of these

4. 20 mL of 0.1 *M* weak acid HA (K_a 10^{-5}) is mixed with solution of 10 mL of 0.3 *M* HCl and 10 mL of 0.1 *M* NaOH. Find the value of $[A^-]/([HA] + [A^-])$ in the resulting solution :
(a) 2×10^{-4} (b) 2×10^{-5} (c) 2×10^{-3} (d) 0.05

5. What concentration of FCH_2COOH (K_a 2.6×10^{-3}) is needed so that $[H^+]$ 2×10^{-3}?
(a) 2×10^{-3} *M* (b) 2.6×10^{-3} *M* (c) 5.2×10^{-3} *M* (d) 3.53×10^{-3} *M*

6. Calculate the ratio of $[HCOO^-]$ and $[F^-]$ in a mixture of 0.2 *M* HCOOH (K_a 2×10^{-4}) and 0.1 *M* HF (K_a 6.6×10^{-4}) :
(a) 1 : 6.6 (b) 1 : 3.3 (c) 2 : 3.3 (d) 3.3 : 2

7. If first dissociation of $X(OH)_3$ is 100% where as second dissociation is 50% and third dissociation is negligible then the pH of 4×10^{-3} *M* $X(OH)_3$ is :
(a) 11.78 (b) 10.78 (c) 2.5 (d) 2.22

8. H_3A is a weak triprotic acid (K_{a_1} 10^{-5}, K_{a_2} 10^{-9}, K_{a_3} 10^{-13})

What is the value of p*X* of 0.1 *M* H_3A (*aq.*) solution? where p*X* $-\log X$ and X $\dfrac{[A^{3-}]}{[HA^{2-}]}$

(a) 7 (b) 8 (c) 9 (d) 10

9. Calcium lactate is a salt of weak organic acid and strong base represented as $Ca(LaC)_2$. A saturated solution of $Ca(LaC)_2$ contains 0.6 mole in 2 litre solution. pOH of solution is 5.60. If 90% dissociation of the salt takes place then what is pK_a of lactic acid?
(a) $2.8 - \log(0.54)$ (b) $2.8 + \log(0.54)$ (c) $2.8 + \log(0.27)$ (d) None of these

10. What is the concentration of CH_3COOH(*aq.*) in a solution prepared by dissolving 0.01 mole of $NH_4^+CH_3COO^-$ in 1 L H_2O? [$K_{a\,(CH_3COOH)}$ 1.8×10^{-5}; $K_{b\,(NH_4OH)}$ 1.8×10^{-5}]
(a) 5.55×10^{-5} (b) 0.10 (c) 6.4×10^{-4} (d) 5.55×10^{-3}

11. K_a for the reaction; Fe^{3+} (*aq.*) $H_2O(l) \rightleftharpoons Fe(OH)^{2+}(aq.)$ H_3O^+ (*aq.*) is 6.5×10^{-3}. What is the max. pH value which could be used so that at least 80% of the total iron (III) in a dilute solution exists as Fe^{3+} ?
(a) 2 (b) 2.41 (c) 2.79 (d) 1.59

For More Material Join: @JEEAdvanced_2024

12. $Fe(OH)_2$ is diacidic base has K_{b_1} 10 ^{4} and K_{b_2} 2.5 10 ^{6}

What is the concentration of $Fe(OH)_2$ in 0.1 *M* $Fe(NO_3)_2$ solution?

(a) 4 10 ^{9} (b) 2.5 10 ^{6} (c) 10 ^{10} (d) 10 ^{14}

13. How many gm of solid KOH must be added to 100 mL of a buffer solution to make the pH of solution 6.0, if it is 0.1 *M* each w.r.t. acid H*A* and salt K *A*.

[Given : pK_a (H*A*) 5]

(a) 0.458 (b) 0.327 (c) 5.19 (d) None of these

14. Fixed volume of 0.1 *M* benzoic acid (pK_a 4.2) solution is added into 0.2 *M* sodium benzoate solution and formed a 300 mL, resultant acidic buffer solution. If pH of this buffer solution is 4.5 then find added volume of benzoic acid :

(a) 100 mL (b) 150 mL (c) 200 mL (d) None of these

15. A 1.025 g sample containing a weak acid H*X* (mol. mass 82) is dissolved in 60 mL water and titrated with 0.25 *M* NaOH. When half of the acid was neutralised the pH was found to be 5.0 and at the equivalence point the pH is 9.0. Calculate mass percentage of H*X* in sample :

(a) 50% (b) 75% (c) 80% (d) None of these

16. Which of the following expression for % dissociation of a monoacidic base (*B*OH) in aqueous solution at appreciable concentration is not correct?

(a) 100 $\sqrt{\frac{K_b}{c}}$ (b) $\frac{1}{1 \quad 10^{(pK_b \quad pOH)}}$

(c) $\frac{K_w [H \ ]}{K_b \quad K_w}$ (d) $\frac{K_b}{K_b \quad [OH \ ]}$

17. A solution of weak acid H*A* was titrated with base NaOH. The equivalent point was reached when 40 mL of 0.1 *M* NaOH has been added. Now 20 mL of 0.1 *M* HCl were added to titrated solution, the pH was found to be 5.0. What will be the pH of the solution obtained by mixing 20 mL of 0.2 *M* NaOH and 20 mL of 0.2 *M* H*A*?

(a) 7 (b) 9 (c) 11 (d) None of these

18. A buffer solution 0.04 *M* in Na_2HPO_4 and 0.02 *M* in Na_3PO_4 is prepared. The electrolytic oxidation of 1.0 milli-mole of the organic compound *R*NHOH is carried out in 100 mL of the buffer. The reaction is

$$RNHOH + H_2O \qquad RNO_2 + 4H^+ \quad 4e$$

The approximate pH of solution after the oxidation is complete is :

[Given : for H_3PO_4, pK_{a_1} 2.2; pK_{a_2} 7.20; pK_{a_3} 12]

(a) 6.90 (b) 7.20 (c) 7.5 (d) None of these

19. When a 20 mL of 0.08 *M* weak base *B*OH is titrated with 0.08 *M* HCl, the pH of the solution at the end point is 5. What will be the pOH if 10 mL of 0.04 *M* NaOH is added to the resulting solution?

[Given : log 2 0.30 and log 3 0.48]

(a) 5.40 (b) 5.88 (c) 4.92 (d) None of these

For More Material Join: @JEEAdvanced_2024

20. Calculate approximate pH of the resultant solution formed by titration of 25 mL of 0.04 *M* Na_2CO_3 with 50 mL of 0.025 *M* HCl. [Given : pK_{a_1} 6.4 and pK_{a_2} 10.3 for H_2CO_3]

(a) 5.92 (b) 6.88 (c) 6.4 (d) 5.88

21. In the titration of a solution of a weak acid H*A* and NaOH, the pH is 5.0 after 10 mL of NaOH solution has been added and 5.60 after 20 mL NaOH has been added.
What is the value of pK_a for H*A*?

(a) 5.15 (b) 5.3 (c) 5.6 (d) None of these

22. 50 mL of 0.05 *M* Na_2CO_3 is titrated against 0.1 *M* HCl. On adding 40 mL of HCl, pH of the solution will be [Given : For H_2CO_3, pK_{a_1} 6.35, pK_{a_2} 10.33; log 3 0.477, log 2 0.30]

(a) 6.35 (b) 6.526 (c) 8.34 (d) 6.173

23. 10 mL of 0.1 *M* tribasic acid H_3A is titrated with 0.1 *M* NaOH solution. What is the ratio of $\frac{[H_3A]}{[A^{3-}]}$ at 2nd equivalence point? [Given : K_{a_1} 10^{-3}, K_{a_2} 10^{-8}, K_{a_3} 10^{-12}]

(a) $\sim 10^{-4}$ (b) $\sim 10^{-4}$ (c) $\sim 10^{-7}$ (d) $\sim 10^{-6}$

24. A_3B_2 is a sparingly soluble salt of molar mass *M* (g mol $^{-1}$) and solubility *x* g litre $^{-1}$. The ratio of the molar concentration of B^{3-} to the solubility product of the salt is :

(a) $108\frac{x^5}{M^5}$ (b) $\frac{1}{108}\frac{M^4}{x^4}$ (c) $\frac{1}{54}\frac{M^4}{x^4}$ (d) None of these

25. A solution is 0.10 *M* $Ba(NO_3)_2$ and 0.10 *M* $Sr(NO_3)_2$. If solid Na_2CrO_4 is added to the solution, what is $[Ba^{2+}]$ when $SrCrO_4$ begins to precipitate?

[K_{sp} ($BaCrO_4$) 1.2×10^{-10}; K_{sp} ($SrCrO_4$) 3.5×10^{-5}]

(a) 7.4×10^{-7} (b) 2.0×10^{-7} (c) 6.1×10^{-7} (d) 3.4×10^{-7}

26. A solution is 0.01 *M* KI and 0.1 *M* KCl. If solid $AgNO_3$ is added to the solution, what is the $[I^-]$ when AgCl begins to precipitate?

[K_{sp} (AgI) 1.5×10^{-16}; K_{sp} (AgCl) 1.8×10^{-10}]

(a) 3.5×10^{-7} (b) 6.1×10^{-8} (c) 2.2×10^{-7} (d) 8.3×10^{-8}

27. A solution of 0.1 *M* in Cl^- and 10^{-4} *M* CrO_4^{2-}. If solid $AgNO_3$ is gradually added to this solution, what will be the concentration of Cl^- when Ag_2CrO_4 begins to precipitate?

[K_{sp} (AgCl) $10^{-10}\ M^2$; K_{sp} (Ag_2CrO_4) $10^{-12}\ M^3$]

(a) $10^{-6}\ M$ (b) $10^{-4}\ M$ (c) $10^{-5}\ M$ (d) $10^{-9}\ M$

28. If 500 mL of 0.4 *M* $AgNO_3$ is mixed with 500 mL of 2 *M* NH_3 solution then what is the concentration of $Ag(NH_3)^+$ in solution?

Given : $K_{f_1}[Ag(NH_3)]$ 10^3; $K_{f_2}[Ag(NH_3)_2]$ 10^4

(a) $3.33 \times 10^{-7}\ M$ (b) $3\ 33 \times 10^{-5} M$ (c) $3 \times 10^{-4} M$ (d) $10^{-7}\ M$


29. The simultaneous solubility of AgCN (K_{sp} 2.5×10^{-16}) and AgCl (K_{sp} 1.6×10^{-10}) in 1.0 *M* $NH_3(aq.)$ are respectively : [Given : K_f $[Ag(NH_3)_2]$ 10^7]


(a) $0.037, 5.78 \times 10^{-8}$ (b) $5.78 \times 10^{-8}, 0.037$


(c) $0.04, 6.25 \times 10^{-8}$ (d) $1.58 \times 10^{-3}, 1.26 \times 10^{-5}$


For More Material Join: @JEEAdvanced_2024

30. There exist an equilibrium between solid $SrSO_4$ and Sr^{2} and SO_4^{2} ion in aqueous medium. The possible equilibrium states are shown in figure as thick line. Now, if equilibrium is disturbed by addition of (a) $Sr(NO_3)_2$ and (b) K_2SO_4 and dotted line represent approach of system towards equilibrium. Match the columns given below :

(iii) 

(iv) 

(v) 

(vi) 

(I) addition of $Sr(NO_3)_2$ (II) addition of K_2SO_4

(a) (I) (iii), (II) (iv) (b) (I) (iv), (II) (v)

(c) (I) (vi), (II) (v) (d) (I) (iv), (II) (vi)

31. Solubility of AgCN is maximum in :

(a) acidic buffer solution (b) basic buffer solution

(c) in pure water (d) equal in all solution

32. $AgBr(s)$ $2S_2O_3^{2}$ $(aq.)$ $\rightleftharpoons$ $Ag(S_2O_3)_2^{3}$ $(aq.)$ Br $(aq.)$

[Using : K_{sp} (AgBr) 5 10^{13} K_f $(Ag(S_2O_3)_2^{3})$ 5 10^{13}]

What is the molar solubility of AgBr in 0.1 M $Na_2S_2O_3$?

(a) 0.5 M (b) 0.45 M (c) 0.045 M (d) None of these

33. What is [Ag] in a solution made by dissolving both Ag_2CrO_4 and $Ag_2C_2O_4$ until saturation is reached with respect to both salts?

[K_{sp} $(Ag_2C_2O_4)$ 2 10^{11}, K_{sp} (Ag_2CrO_4) 2 10^{12}]

(a) 2.80 10^{4} (b) 7.6 10^{5} (c) 6.63 10^{6} (d) 3.52 10^{4}

34. What is the minimum pH required to prevent the precipitation of ZnS in a solution that is 0.01 M $ZnCl_2$ and saturated with 0.10 M H_2S?

[Given : K_{sp} 10^{21}, K_{a_1} K_{a_2} 10^{20}]

(a) 0 (b) 1 (c) 2 (d) 4

35. The $Al(OH)_3$ is involved in the following two equilibria,

$$Al(OH)_3(s) \rightleftharpoons Al^{3}\ (aq.)\ \ 3OH\ (aq.);\ K_{sp}$$

$$Al(OH)_3(s)\ \ OH\ (aq.) \rightleftharpoons Al(OH)_4(aq.);\ K_c$$

Which of the following relationship is correct at which solubility is minimum?

(a) [OH] $\left(\frac{K_{sp}}{K_c}\right)^{1/3}$ (b) [OH] $\left(\frac{K_c}{K_{sp}}\right)^{1/4}$

(c) [OH] $\sqrt{\left(\frac{K_{sp}}{K_c}\right)^{1/4}}$ (d) None of these

For More Material Join: @JEEAdvanced_2024

Level 3

Passage 1

Solution of a weak acid and its anion (that is, its conjugate base) or of a base and its common cation are buffered. When we add a small amount of acid or base to any one of them, the pH of solution changes very little. pH of buffer solution can be computed as

for acidic buffer : pH pK_a $\log \frac{[\text{Conjugate base}]}{[\text{Acid}]}$

for basic buffer : pOH pK_b $\log \frac{[\text{Conjugate acid}]}{[\text{Base}]}$

It is generally accepted that a solution has useful buffer capacity (pH change resisting power) provided that the value of [salt or conjugate base]/[acid] for acidic buffer lies within the range of 1 : 10 to 10 : 1. Buffer capacity is maximum when [conjugate base] [acid]

1. One litre of an aqueous solution contains 0.15 mole of CH_3COOH (pK_a 4.8) and 0.15 mole of CH_3COONa. After the addition of 0.05 mole of solid NaOH to this solution, the pH will be :
(a) 4.5 (b) 4.8 (c) 5.1 (d) 5.4

2. Calculate the pH of a solution made by adding 0.01 mole of HCl in 100 mL of a solution which is 0.2 *M* in NH_3 (pK_b 4.74) and 0.3 *M* in NH_4 :
(Assuming no change in volume)
(a) 5.34 (b) 8.66 (c) 7.46 (d) None of these

3. Useful buffer range of weak acid H*A* (K_a 10^{-5}) is :
(a) 5 to 7 (b) 4 to 6 (c) 3 to 5 (d) None of these

4. Select correct statement :
(a) When we add small amount of NaOH in acidic buffer solution, pOH of solution is increases
(b) When we add small amount of NaOH in basic buffer solution, pH of solution is increases
(c) When we add small amount of water in acidic buffer solution, pH of solution is decreases
(d) When 100 mL of 0.2 *M* CH_3COOH react with 200 mL of 0.1 *M* NaOH buffer solution is formed

Passage 2

Hydrolysis is an acid-base reaction of a cation or anion or both ions of a salt with water. Resultant solution of hydrolysis may be acidic, basic or neutral. The anion *A* which is a weaker base than OH and which has its conjugate acid H*A* stronger than water but weaker than H_3O^+ shows the phenomenon of hydrolysis Ex. : CH_3COO , CN , NO_2 , etc.

The cation *B* which is a weaker acid than H_3O^+ and which has its conjugate base *B*OH stronger than water but weaker than OH shows the phenomenon of hydrolysis Ex. : NH_4 , $C_6H_5NH^+$, $N_2H_5^+$ etc.

For More Material Join: @JEEAdvanced_2024

The hydrolysis constant of anion and cation are given by

$A^-(aq.) + H_2O(l) \rightleftharpoons HA(aq.) + OH^-(aq.)$

$$K_h = \frac{K_w}{K_a} = \frac{[HA(aq.)][OH^-(aq.)]}{[A^-(aq.)]}$$

$B^+(aq.) + H_2O(l) \rightleftharpoons BOH(aq.) + H^+(aq.)$

$$K_h = \frac{K_w}{K_b} = \frac{[BOH(aq.)][H^+(aq.)]}{[B^+(aq.)]}$$

1. Which of the following statement is true?
(a) Weaker the acid, greater will be hydrolysis of its anion
(b) Weaker the base, greater will be hydrolysis of its cation
(c) Both (a) and (b)
(d) None of these

2. Select the correct statement :
(a) KCl undergoes hydrolysis
(b) $K_h = K_b(A^-)$ and $K_h = K_a(B^+)$
(c) 0.1 M solution of NaCN is acidic
(d) resultant solution of equal volume of 0.1 M NH_3 and 0.1 M HCl is basic

3. When pure ammonium chloride is dissolved in pure water, the pH of the resulting solution is not 7. This is because :
(a) ammonium ions accept protons from water molecules leaving free OH^- ions in solution
(b) ammonium ions donate protons to water molecules forming H_3O^+ ions in solution
(c) ammonium ions combine with water molecule to give the weak base, ammonium hydroxide
(d) chloride ion made the solution acidic

4. Calculate percentage degree of hydrolysis in a 0.1 M solution of CH_3COONa.
(K_a of $CH_3COOH = 10^{-5}$)
(a) 0.1 (b) 0.01 (c) 10^{-4} (d) None of these



Passage 3

Acid-base indicators are either weak organic acids or weak organic bases. Indicator change colour in dilute solution when the hydronium ion concentration reaches a particular value. For example, phenolphthalein is a colourless substance in any aqueous solution with a pH less than 8.3. In between the pH range 8.3 to 10, transition of colour (colourless to pink) takes place and if pH of solution is greater than 10 solution is dark pink. Considering an acid indicator HIn, the equilibrium involving it and its conjugate base In^- can be represented as :

$$\underset{\text{acidic form}}{HIn} \rightleftharpoons H^+ + \underset{\text{basic form}}{In^-}$$

pH of solution can be computed as :

$$pH = pK_{In} + \log \frac{[In^-]}{[HIn]}$$

In general, transition of colour takes place in between the pH range $pK_{In} \pm 1$.

For More Material Join: @JEEAdvanced_2024

1. An indicator is a weak acid and pH range is 4.0 to 6.0. If indicator in 50% ionized in a given solution then what is the ionization constant of the acid?

(a) 10^{-4} (b) 10^{-5} (c) 10^{-6} (d) None of these

2. Select the correct statement(s) :


(a) At midway in the transition of an acidic indicator, pH pK_{In}

(b) Methyl orange (3.1 to 4.4) is a suitable indicator for titration of weak acid and strong base

(c) Bromothymol blue (6.0 to 7.6) is a good indicator for titration of HCl and NaOH

(d) Thymol blue (1.2 to 2.8) is a very good indicator for titration of 100 mL of 0.1 *M* NH_4OH (pK_b 4.74) and 0.1 *M* HCl

3. Following is the titration curve of two acids H*A* and H*B* (5 milli-moles each) titrated against strong base NaOH (0.1 *M*)



What is pK_a for H*B* acid ?

(a) 3 (b) 4 (c) 5 (d) 6

4. What is the equilibrium constant for the reaction :

$$HB(aq.) \quad NaA(aq.) \rightleftharpoons HA(aq.) \quad NaB(aq.)?$$

(a) 10 (b) 0.1 (c) 10^{-7} (d) 10^{-11}

5. Calculate the pH at equivalent point when H*B* is titrated with NaOH.

(a) 8.75 (b) 8.85 (c) 9.0 (d) None of these

6. Which of the following indicator is most suitable for titration of H*B* with strong base :

(a) Phenolphthalein (8.3 10) (b) Bromothymol blue (6 7.6)

(c) Methyl red (4.2 6.3) (d) Malachite green (11.4 13)

For More Material Join: @JEEAdvanced_2024



Passage 4

Consider a saturated solution of silver chloride that is in contact with solid silver chloride. The solubility equilibrium can be represented as

$$AgCl(s) \rightleftharpoons Ag\ (aq.)\quad Cl\ (aq.);\quad K_{sp}\quad [Ag\ (aq.)][Cl\ (aq.)]$$

Where K_{sp} is called the solubility product constant or simply the solubility product. In general, the solubility product of a compound is the product of the molar concentrations of the constituent ions, each raised to the power of its stoichiometric coefficient in the equilibrium equation.

For concentrations of ions that do not necessarily correspond to equilibrium conditions we use the reaction quotient (Q) which is called the ion or ionic product (Q), to predict whether a precipitate will form. Note that Q has the same form as K_{sp}.

The possible relationships between Q and K_{sp} are

Q K_{sp} Unsaturated solution
Q K_{sp} Saturated solution
Q K_{sp} Supersaturated solution; precipitate will form

1. Will a precipitate form if 50 cm^3 of 0.01 M $AgNO_3$ and 50 cm^3 of 2 10^{5} M NaCl are mixed? [Given : K_{sp} (AgCl) 10^{10} M^2]

(a) Yes
(b) No
(c) Ionic product is less than solubility product, hence precipitate will form
(d) Data insufficient

2. Will a precipitate form if 1 volume of 0.1 M Pb^2 ion solution is mixed with 3 volume of 0.3 M Cl ion solution? [Given : K_{sp} ($PbCl_2$) 1.7 10^{5} M^3]

(a) Yes
(b) No
(c) Ionic product is less than solubility product, hence precipitate will form
(d) Data insufficient

3. At 25°C, will a precipitate of $Mg(OH)_2$ form when a 0.0001 M solution of $Mg(NO_3)_2$ is adjusted to a pH of 9.0? At what minimum value of pH will precipitation start? [Given : K_{sp} ($Mg(OH)_2$) 10^{11} M^3]

(a) No, pH 3.5 (b) No, pH 10.5 (c) No, pH 6.0 (d) Yes, pH 8.5

4. Determine the molar solubility of MgF_2 from its solubility product K_{sp} 4 10^{9} :

(a) 10^{3} (b) 6.32 10^{5} (c) 2 10^{5} (d) None of these

5. The solubility product of ferric hydroxide in aqueous solution is 6 10^{38} at 298 K. The solubility of Fe^3 ion will increase when the :

(a) pH is increased
(b) pH is 7.0
(c) pH is decreased
(d) saturated solution is exposed to the atmosphere

For More Material Join: @JEEAdvanced_2024

One or More Answers is/are Correct



1. Which is/are wrong statement(s)?
(a) Arrhenius acids are also Bronsted acids but all Arrhenius bases are not Bronsted bases
(b) All Lewis bases are Bronsted bases
(c) All Bronsted acids are Lewis acids
(d) Conjugate base of a strong acid is weak

2. Which of the following are conjugate acid-base pairs ?
(a) HCO_3, CO_3^{2} (b) $C_6H_5\overset{+}{N}H_3, C_6H_5NH_2$
(c) $H_2C_2O_4, HC_2O_4$ (d) OH , H

3. Which are the set of amphiprotic species?
(a) H_2O, H_2PO_4, HPO_4^{2} (b) HPO_4^{2} , HCO_3
(c) $H_2PO_2, H_2PO_3, HC_2O_4$ (d) HPO_3^{2} , H_2O, CO_3^{2}

4. Which of the following statements is/are not correct?
(a) A substance which can provide OH in aqueous medium is a base
(b) A substance which can accept a pair of electron is a base
(c) A substance which can accept a proton in aqueous medium is a base
(d) A substance which can donate a pair of electron is a base

5. If degree of ionization ( ) of a weak electrolyte *AB* is very less then is :
(a) directly proportional to the square root of volume of solution
(b) inversely proportional to the dilution
(c) inversely proportional to the square root of concentration
(d) directly proportional to concentration

6. Factor influencing the degree of ionization of a weak electrolyte is :
(a) dilution (b) temperature
(c) presence of other ions (d) nature of solvent

7. Which of the following statement(s) is/are correct about the ionic product of water?
(a) K_i (ionization constant of water) $< K_w$ (ionic product of water)
(b) pK_i pK_w
(c) At 25°C, K_i 1.8 10 ^{14}
(d) Ionic product of water at 10°C is 10 ^{14}

8. Which among the following statement is/are correct?
(a) pH $\log_{10}[H_3O^+]$ for dilute solution
(b) pH of H_2O decreases with increase of temperature
(c) pH can not more than 14
(d) If a solution is diluted ten times, its pH always increases by 1

9. If concentration of two weak acids are different and D.O.D. ( ) are very less then their relative strength can be compared by :
(a) $\frac{[H\ ]_1}{[H\ ]_2}$ (b) $\frac{1}{2}$ (c) $\frac{C_1\ \ 1}{C_2\ \ 2}$ (d) $\frac{K_{a_1}C_1}{K_{a_2}C_2}$

For More Material Join: @JEEAdvanced_2024

10. If concentration of two weak bases are same and D.O.D. ( ) are very less then their relative strength can be compared by :

(a) $\frac{[OH^-]_1}{[OH^-]_2}$ (b) $\frac{K_{b_1}}{K_{b_2}}$ (c) $\frac{1}{2}$ (d) $\frac{\sqrt{K_{b_1}}}{\sqrt{K_{b_2}}}$

11. Which of the following expressions is/are true?

(a) $[H^+]\ [OH^-] \quad \sqrt{K_w}$ for a neutral solution

(b) $[OH^-] \quad \sqrt{K_w}$ for an acidic solution

(c) pH + pOH 14 at all temperature

(d) $[OH^-] \quad 10^{-7}$ *M* for a neutral solution at 25°C

12. If K_{a_1}, K_{a_2} and K_{a_3} be the first, second and third dissociation constant of H_3PO_4 and $K_{a_1} \quad K_{a_2} \quad K_{a_3}$ which is/are correct :

(a) $[H^+] \sim \sqrt{K_{a_1}[H_3PO_4]}$ (b) $[H^+] \sim [HPO_4^{2-}]$

(c) $K_{a_2} \sim [HPO_4^{2-}]$ (d) $[HPO_4^{2-}] \quad [PO_4^{3-}]$

13. Which of the following mixtures constitute a buffer?

(a) HCOOH + HCOONa (b) $Na_2CO_3 + NaHCO_3$

(c) NaCl + HCl (d) $NH_4Cl + (NH_4)_2SO_4$

14. Which of the following mixtures can act as a buffer?

(a) NaOH + HCOONa (1 : 1 molar ratio) (b) HCOOH + NaOH (2 : 1 molar ratio)

(c) NH_4Cl + NaOH (2 : 1 molar ratio) (d) HCOOH + NaOH (1 : 1 molar ratio)

15. Which of the following will function as buffer?

(a) NaCl + NaOH (b) Borax + boric acid

(c) $NaH_2PO_4 + Na_2HPO_4$ (d) $NH_4Cl + NH_4OH$

16. Which of the following statements is/are correct?

(a) The conjugate acid of NH_2^- is NH_3

(b) Solubility product constant increases with increase in concentration of ions

(c) On diluting a buffer solution pH change is negligible

(d) In alkaline buffer solution, if some HCl is added, its $[OH^-]$ will increase

17. Degree of hydrolysis ( ) for a salt of strong acid and weak base is :

(a) independent of dilution (b) increases with dilution

(c) increases with decrease in K_b (d) increases with increase in temperature

18. The compound whose 0.1 *M* solution is acidic :

[pK_a of HCOOH 3.75, pK_b of NH_4OH = 4.74]

(a) Ammonium formate (b) Ammonium sulphate

(c) Ammonium chloride (d) Sodium formate

19. Formic acid is a weak acid and hydrochloric acid is a strong acid. It shows that the :

(a) $[OH^-]$ of 0.01 *M* HCl (*aq.*) will be less than that of 0.01 *M* HCOOH (*aq.*)

(b) solution containing 0.1 *M* NaOH (*aq.*) and 0.1 *M* HCOONa (*aq.*) is a buffer solution

(c) pH of 10^{-9} *M* HCl (*aq.*) will be approximately 7 at 25°C

(d) pH of a solution formed by mixing equimolar quantities of HCOOH and HCl will be less than that of a similar solution formed from HCOOH and HCOONa

For More Material Join: @JEEAdvanced_2024

20. If you have a saturated solution of CaF_2, then :

(a) $[Ca^{2}\ ]\ \ (K_{sp}/4)^{1/3}$ (b) $2\ \ [Ca^{2}\ ]\ \ [F\ ]$

(c) $[Ca^{2}\ ]\ \ 2\,[F\ ]$ (d) $[Ca^{2}\ ]\ \ \sqrt{K_{sp}}$

21. Which is/are correct statement(s) about the solubility of AgCl(*s*).?

Given : $K_{sp(AgCl)}\ \ 10^{\ 10}$; $K_f[Ag(NH_3)_2]\ \ 10^{8}$.

(a) Solubility of AgCl in pure water is $10^{\ 5}$ gm/litre

(b) Solubility of AgCl in 2 *M* KBr is $10^{\ 5}$ mol/litre

(c) Solubility of AgCl in 2 *M* $AgNO_3$ is $5\ \ 10^{\ 11}\ M$

(d) Solubility of AgCl in 2*M* NH_3 is 0.166 *M*

22. H_2A is a weak diprotic acid. If the pH of 0.1*M* H_2A solution is 3 and concentration of A^{2} is $10^{\ 12}$ at 25°C.

Select correct statement(s)

(a) $[H\ ]_{total}\ \ [H\ ]$ from first step of ionization of acid H_2A

(b) Concentration of OH in solution is $10^{\ 3}\ M$

(c) The value of K_{a_1} is nearly $10^{\ 5}$

(d) $pK_{a_2}\ \ pK_{a_1}\ \ 9$

23. Which is/are correct statement(s)?

(a) CH_3COONH_4 have greater degree of hydrolysis in 0.2 *M* solution in comparison of 0.4 *M* solution.

(b) Anions which are weaker base than OH , do not hydrolyse

(c) The CH_3COO have greater degree of hydrolysis in comparision of HCOO when their salt solution have equal conc.

(d) SO_4^{2} hydrolyses but HSO_4 does not undergo hydrolysis

24. 0.01*M* NH_4Cl (*aq*) solution at 25°C has:

(a) $[Cl\ (aq)]\ \ 10^{\ 2}M$ (b) $[NH_4(aq)]\ \ 10^{\ 2}M$

(c) pOH > 7 (d) $[H\ ]\ \ 10^{\ 7}\ M$

25. In an acidic indicator HIn has ionization constant is $10^{\ 8}$. The acid form of indicator is yellow and alkaline form is red. Which is correct statement ?

(Given : log2 = 0.3, log3 = 0.48)

(a) The pH range of indicator is 7 to 9

(b) Change in pH is 0.96 when 75% yellow colour change to 75% red colour

(c) This indicator is suitable for the titration of strong acid *vs.* strong base

(d) pH of indicator is 8.3 when ratio of acid form to alkaline form is 2.

For More Material Join: @JEEAdvanced_2024

Match the Column



Column-I and **Column-II** contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

1.

Column-I	Column-II
(A) HCl	(P) Bronsted base
(B) NH_3	(Q) Bronsted acid
(C) H_2O	(R) Arrhenius acid
(D) CN	(S) Lewis base in adduct displacement reaction

2.

Column-I	Column-II
(A) Conjugate acid-base pair	(P) Bronsted-Lowry concept
(B) Acid-base adduct	(Q) Lewis concept
(C) An acid-base reaction	(R) Arrhenius concept
(D) Proton donation	(S) K_a K_b K_w

3.

Column-I	Column-II
(A) $Fe(NO_3)_2(aq.)$	(P) Only cationic hydrolysis
(B) $KClO_4(aq.)$	(Q) Only anionic hydrolysis
(C) HCOONa$(aq.)$	(R) Both cationic as well as anionic hydrolysis
(D) $NH_4CN(aq.)$	(S) No hydrolysis

4.

Column-I	Column-II
(A) Salt of weak acid and weak base	(P) pH 1/2 [pK_w pK_a log C]
(B) Salt of weak acid and strong base	(Q) pH 1/2 [pK_w pK_a pK_b]
(C) Salt of strong acid and strong base	(R) pH 1/2 [pK_w pK_b log C]
(D) Salt of strong acid and weak base	(S) pH 1/2 [pK_w]

For More Material Join: @JEEAdvanced_2024

5.

Column-I	Column-II
(A) Salt of weak acid and weak base (pK_a pK_b)	(P) pH of solution at 25°C less than 7
(B) Salt of weak acid and strong base	(Q) pH of solution at 25°C greater than 7
(C) Salt of strong acid and strong base	(R) pH of solution at 25°C equal to 7
(D) Salt of strong acid and weak base	(S) pH cannot be find until the value of K_a/K_b is given

6.

Column-I	Column-II
(A) pH of 0.1 M HA (pK_a 5) and 0.01 M NaA	(P) 4
(B) pH of 0.1 M BOH (pK_b 6) and 0.1 M BCl	(Q) 7
(C) pH of 0.1 M salt of weak acid (pK_a 5) and weak base (pK_b 7)	(R) 6
(D) pH of 500 litre of 0.02 M HNO_3 and 500 litre 0.01 M $Sr(OH)_2$	(S) 8

7.

Column-I	Column-II
(A) CH_3COOH (pK_a 4.74, 0.1 M) CH_3COONa (0.1 M)	(P) Acidic buffer at its maximum capacity
(B) CH_3COOH (0.1 M) + HCl (0.1 M)	(Q) Buffer solution
(C) CH_3COOH (pK_a 4.74, 0.1 M) + NH_4OH (pK_b 4.74, 0.1 M)	(R) pH < 7 at 25°C
(D) CH_3COONa (300 mL of 0.1 M) + HCl (100 mL of 0.1 M)	(S) pH 7 at 25°C

8.

Column-I	Column-II
(A) Titration of a strong acid with strong base	(P) Methyl orange (3.1 4.4)
(B) Titration of weak acid with strong base	(Q) Methyl red (4.2 6.3)
(C) Titration of strong acid with weak base	(R) Phenolphthalein (8.3 10)
(D) Titration of weak acid with weak base	(S) No general indicator is suitable

For More Material Join: @JEEAdvanced_2024

9. When we titrate sodium carbonate solution (in beaker) with hydrochloric acid

Column-I	Column-II
(A) At the start of titration	(P) Buffer solution of HCO_3 and CO_3^{2}
(B) Before the first equivalent point	(Q) Buffer solution of H_2CO_3 and HCO_3
(C) At the first equivalent point	(R) Amphiprotic anion, pH 1/2 (pK_{a_1} pK_{a_2})
(D) Between the first and second equivalent points	(S) Hydrolysis of CO_3^{2}

10.

Column-I	Column-II
(A) Mercurous iodide	(P) $108\ S^5$
(B) Aluminium phosphate	(Q) $4\ S^3$
(C) Calcium phosphate	(R) S^2
(D) Zirconium phosphate	(S) $6912\ S^7$

Assertion-Reason Type Questions

Each question contains STATEMENT-1 (Assertion) and STATEMENT-2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below :

(A) If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1

(B) If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1

(C) If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

(D) If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

1. STATEMENT-1 : All strong monoprotic acid with same concentration in dilute solution show same pH.

STATEMENT-2 : Water shows levelling effect.

2. STATEMENT-1 : If water is heated to 50°C then pH will increase.

STATEMENT-2 : K_w increases with increase in temperature.

3. STATEMENT-1 : Addition of HCl(*aq.*) to HCOOH(*aq.*) decrease the dissociation of HCOOH(*aq.*)

STATEMENT-2 : Due to common ion effect of H , dissociation of HCOOH decrease.

4. STATEMENT-1 : pH of 10^{7} *M* HCl is less than 7 at 25°C.

STATEMENT-2 : At very low concentration of HCl, contribution of H from water is considerable.

For More Material Join: @JEEAdvanced_2024

5. STATEMENT-1 : The dissociation constants of weak diprotic acid are in the order of K_{a_1} K_{a_2}.

STATEMENT-2 : Removal of H from anion is difficult as compared to neutral molecule.

6. STATEMENT-1 : When 0.1 *M* weak diprotic acid H_2A dissociates with its dissociation constants K_{a_1} 10^{-3} and K_{a_2} 10^{-8}, then $[A^{2-}]$ is almost equal to 10^{-3} *M*.

STATEMENT-2 : Since K_{a_2} K_{a_1} for 0.1 *M* H_2A, so $[A^{2-}]$ is negligible w.r.t. [H*A*].

7. STATEMENT-1 : pH value of acidic buffer solution change, if buffer solution is diluted upto very larger extent.

STATEMENT-2 : [H] decrease due to change in concentration as well as and decrease in concentration is more as compared to increase in .

8. STATEMENT-1 : In a titration of weak acid with strong base, the pH at the half equivalence point is pK_a.

STATEMENT-2 : At half equivalence point, it will form acidic buffer at its maximum capacity where [acid] = [salt].

9. STATEMENT-1 : In the titration of Na_2CO_3 with HCl using methyl orange indicator, the volume of the acid required at the equivalence point is twice that of the acid required using phenolphthalein as indicator.

STATEMENT-2 : Two moles of HCl are required for the complete neutralisation of one mole of Na_2CO_3.

10. STATEMENT-1 : In the acid-base titration involving strong base and weak acid, methyl red can be used as an indicator.

STATEMENT-2 : Methyl red changes its colour in the pH range 4.2 to 6.3.

11. STATEMENT-1 : Sparingly soluble salts *AB* and XY_2 with the same solubility product, will have different solubility.

STATEMENT-2 : Solubility of sparingly soluble salt depend upon solubility product.

12. STATEMENT-1 : Solubility product of BaF_2 will increase on dilution.

STATEMENT-2 : Solubility of BaF_2 will change on changing temperature.

13. STATEMENT-1 : Solubility of sparingly soluble salt decreases due to common ion effect.

STATEMENT-2 : Solubility product constant does not depend on common ion effect.

14. STATEMENT-1 : Solubility of AgCl in $NH_3(aq.)$ is greater than in pure water.

STATEMENT-2 : When AgCl dissolve in $NH_3(aq.)$, complex ion formation $[Ag(NH_3)_2]$ takes place and solubility equilibria of AgCl shifted in forward direction.

15. STATEMENT-1 : Solubility of AgCN in acidic solution is greater than that in pure water.

STATEMENT-2 : Solubility equilibria of AgCN in acidic solution is shifted in forward direction due to formation of HCN.

For More Material Join: @JEEAdvanced_2024

Subjective Problems

1. Calculate pOH of 0.1M aq. solution of weak base BOH (K_b 10 ^{7}) at 25 C.

2. pH of 0.01M aq. solution of HA is 4. Find the value of pK_a of HA at 25 C.

3. Calculate approximate pH of 10 ^{10}M NaOH at 25 C.

4. Calculate pH of a resultant solution of 25 mL of 0.1M HCl, 50 mL of 0.02 M HNO_3 and 25 mL of 0.1M NaOH.

5. Calculate pH of a resultant solution of 0.1M HA (K_a 10 ^{6}) and 0.45M HB (K_a 2 10 ^{6}) at 25 C.

6. 0.16 g N_2H_4 (K_b 4 10 ^{6}) are dissolved in water and the total volume made up to 500 mL. Calculate the percentage of N_2H_4 that has reacted with water in this solution

7. Calculate pH of a buffer solution that contains 0.1M NH_4OH (K_b 10 ^{5}) and 0.1M NH_4Cl.

8. Calculate the ratio of sodium formate and formic acid (K_a 2 10 ^{4}) in a buffer solution of pH 4.3.

9. What is the pOH of 0.1M KB (salt of weak acid and strong base) at 25 C ? (Given : pK_b of B 7)


10. A certain weak acid has K_a 10 ^{5}. If the equilibrium constant for its reaction with a strong base is represented as 1 10^{y} then find the value of y.

11. If solubility of AgCl in 0.2 M solution of $AgNO_3$ is represented as y 10 ^{10} then find the value of y.
(Given : $K_{sp(AgCl)}$ 10 ^{10})

12. When one litre of a saturated solution of $PbCl_2$ (mol. mass 278) is evaporated, the residue is found to weight 2.78 g. If K_{sp} of $PbCl_2$ is represented as y 10 ^{6} then find the value of y.

13. A solution is saturated in $SrCO_3$ and SrF_2. The CO_3^{2} was found to be 10 ^{3} mol/L.
If the concentration of F in solution is represented as y 10 ^{2}M then what is the value of y ?
[Given : $K_{sp}(SrCO_3)$ 2.5 10 ^{10}; $K_{sp}(SrF_2)$ 10 ^{10}]

14. 10 mL of H_2A (weak diprotic acid) solution is titrated against 0.1M NaOH. pH of the solution is plotted against volume of strong base added and following observation is made.



If pH of the solution at first equivalence point is pH_1 and at second equivalence point is pH_2.
Calculate the value of (pH_2 pH_1) at 25°C
Given: For H_2A, pK_{a_1} 4.6 and pK_{a_2} 8, log 25 1.4

For More Material Join: @JEEAdvanced_2024

15. Amongst the following, the total number of compounds whose aqueous solution turns red litmus paper blue is:

NaCN	KCl	CH_3COONH_4	NaH_2PO_4	$ZnCl_2$	Na_3PO_4
$Fe(NO_3)_3$	Na_2CO_3	NH_4Cl	$NaHCO_3$	$Na_2C_2O_4$	Na_2HPO_4

Given:

Acid	Ka_1	Ka_2	Ka_3
H_3PO_4	10^{-3}	10^{-8}	10^{-12}
H_2CO_3	10^{-6}	10^{-11}	–
$H_2C_2O_4$	10^{-2}	10^{-5}	–

For More Material Join: @JEEAdvanced_2024

Answers

Level 1

1. (d)	2. (c)	3. (a)	4. (c)	5. (c)	6. (a)	7. (a)	8. (d)	9. (d)	10. (c)
11. (d)	12. (d)	13. (c)	14. (d)	15. (b)	16. (d)	17. (a)	18. (d)	19. (c)	20. (c)
21. (a)	22. (b)	23. (a)	24. (b)	25. (a)	26. (c)	27. (b)	28. (c)	29. (b)	30. (b)
31. (b)	32. (a)	33. (b)	34. (a)	35. (a)	36. (b)	37. (c)	38. (a)	39. (c)	40. (c)
41. (c)	42. (c)	43. (b)	44. (c)	45. (b)	46. (a)	47. (c)	48. (d)	49. (d)	50. (d)
51. (d)	52. (b)	53. (b)	54. (c)	55. (d)	56. (d)	57. (b)	58. (c)	59. (c)	60. (b)
61. (b)	62. (d)	63. (b)	64. (b)	65. (b)	66. (c)	67. (a)	68. (c)	69. (d)	70. (a)
71. (d)	72. (b)	73. (d)	74. (c)	75. (b)	76. (b)	77. (a)	78. (d)	79. (b)	80. (c)
81. (b)	82. (d)	83. (a)	84. (a)	85. (c)	86. (a)	87. (c)	88. (b)	89. (d)	90. (b)
91. (c)	92. (d)	93. (b)	94. (a)	95. (d)	96. (a)	97. (a)	98. (d)	99. (b)	100. (a)
101. (a)	102. (b)	103. (c)	104. (b)	105. (c)	106. (d)	107. (d)	108. (b)	109. (a)	110. (c)
111. (d)	112. (c)	113. (a)	114. (a)	115. (a)	116. (c)	117. (a)	118. (c)	119. (a)	120. (a)
121. (c)	122. (c)	123. (c)	124. (a)	125. (c)	126. (d)	127. (a)	128. (a)	129. (d)	130. (b)
131. (d)	132. (b)	133. (c)	134. (c)	135. (d)	136. (d)	137. (a)	138. (d)	139. (c)	140. (a)
141. (a)	142. (d)	143. (c)	144. (c)	145. (d)	146. (a)	147. (b)	148. (b)	149. (d)	150. (b)

Level 2

1. (d)	2. (c)	3. (c)	4. (a)	5. (d)	6. (c)	7. (a)	8. (d)	9. (a)	10. (a)
11. (d)	12. (c)	13. (a)	14. (b)	15. (c)	16. (c)	17. (b)	18. (c)	19. (b)	20. (b)
21. (b)	22. (d)	23. (c)	24. (c)	25. (d)	26. (d)	27. (a)	28. (b)	29. (a)	30. (b)
31. (a)	32. (c)	33. (d)	34. (b)	35. (d)					

For More Material Join: @JEEAdvanced_2024



Level 3

Passage–1:	**1.**	(c)	**2.**	(b)	**3.**	(b)	**4.**	(b)			
Passage–2:	**1.**	(c)	**2.**	(b)	**3.**	(b)	**4.**	(b)			
Passage–3:	**1.**	(b)	**2.**	(a,c)	**3.**	(c)	**4.**	(a)	**5.**	(b)	**6.** (a)
Passage–4:	**1.**	(a)	**2.**	(a)	**3.**	(b)	**4.**	(a)	**5.**	(c)	

One or More Answer is/are Correct

1. (b,c) **2.** (a,b,c) **3.** (a,b) **4.** (b) **5.** (a,c) **6.** (a,b,c,d) **7.** (a,b) **8.** (a,b)
9. (a,c) **10.** (a,c,d) **11.** (a,b,d) **12.** (a,c) **13.** (a,b) **14.** (b, c) **15.** (b, c, d) **16.** (a, c)
17. (b, c, d) **18.** (a, b, c) **19.** (a, c, d) **20.** (a, b) **21.** (b,c,d) **22.** (a,c) **23.** (c,d) **24.** (b,c,d)
25. (a,b,c)

Match the Column

1.	A	Q, R;	B	P, S;	C	P Q, S;	D	P, S
2.	A	P, S;	B	Q;	C	P, Q, R;	D	P, R
3.	A	P;	B	S;	C	Q;	D	R
4.	A	Q;	B	P;	C	S;	D	R
5.	A	R;	B	Q, S;	C	R;	D	P, S
6.	A	P;	B	S;	C	R;	D	Q
7.	A	P, Q, R;	B	R;	C	Q, S;	D	Q, R
8.	A	P, Q, R;	B	R;	C	P, Q;	D	S
9.	A	S;	B	P;	C	R;	D	Q
10.	A	Q;	B	R;	C	P;	D	S

Assertion-Reason Type Questions

1. (A) **2.** (D) **3.** (A) **4.** (A) **5.** (A) **6.** (D) **7.** (A) **8.** (A) **9.** (B) **10.** (D)
11. (B) **12.** (D) **13.** (B) **14.** (A) **15.** (A)

Subjective Problems

1. 4	**2.** 6	**3.** 7	**4.** 2	**5.** 3	**6.** 2	**7.** 9	**8.** 4	**9.** 4	**10.** 9
11. 5	**12.** 4	**13.** 2	**14.** 4	**15.** 6					

For More Material Join: @JEEAdvanced_2024

Hints and Solutions

Level 1

1,2 (d, c)

$HX \rightleftharpoons H \quad X$

If, HX acid

then, X conjugate base

3. (a) Among HF, HCl, HBr and HI,

HF is weakest acid

So, F is strongest base.

4. (c) H_3PO_2 and H_3PO_3 are mono and dibasic acids. Hence, NaH_2PO_2 and Na_2HPO_3 are normal salts, while Na_2HPO_4 is acid-salt as one proton is there in the compound.

7. (a) For a conjugate acid-base pair,

$K_a \quad K_b \quad K_w$

8. (d) pH pOH pK_w

9. (d) In neutral solution

$[H \ ] \quad \sqrt{K_w}$

10. (c) As temperature , [H] and [OH] both

11-12-14. (d, d, d)

pH $\log_{10}[H \ ]$

13. (c) $[H \ ] \quad \dfrac{K_w}{[OH \ ]}$ and pH $\log_{10}[H \ ]$

15. (b) $[H \ ] \quad 10^{-9}$

If it is 1000 times more basic then,

$[H \ ]_{Original} \quad 10^{-6} M$

16. (d) When equal volumes are taken, the concentration becomes half.

$$[H \ ] \quad \frac{1}{2}(10^{-3} \quad 10^{-5})$$

$$505 \quad 10^{-6}$$

pH 3.3

17. (a) $[H \ ] \quad 0.002 \ M$

pH $\log 2 \quad 10^{-3}$

$3 \quad \log 2$

pOH $11 \quad \log 2$

18. (d) pH 4, $[H \ ] \quad 10^{-4}$ mol/litre

Equivalent of HCl in 100 mL solution

$$\frac{10^{-4} \quad 100}{1000} \quad 10^{-5}$$

19. (c) $N_1V_1 \quad N_2V_2$

$10^{-3} \quad 10 \quad N_2 \quad 1000$

$N_2 \quad 10^{-5}$

pH 5

and pOH $14 \quad 5 \quad 9$

20. (c) pH 1 $[H \ ] \quad 0.1 \ M$

For $H_2SO_4(aq.)$

$[H \ ] \quad 2$ [Acid]

21. (a) $H_2A(aq) \xrightarrow{100\%} 2H \ (aq) \quad A^{2-}(aq)$

pH $\log [H \ ]$

22. (b) $[Ca(OH)_2(aq.)] \quad \dfrac{0.60}{74} \quad \dfrac{1000}{1500}$

$5.40 \quad 10^{-3}$

$Ca(OH)_2(aq.) \qquad Ca^{2+}(aq.) \quad 2OH \ (aq.)$

$[Ca^{2+} \ ] \quad 5.40 \quad 10^{-3}$

$[OH \ ] \quad 2 \quad 5.40 \quad 10^{-3} \quad 1.08 \quad 10^{-2}$

23-24. (a, b)

At these concentration consider ionization of water also.

25. (a) $H_2SO_4 \quad 2\,NaOH \qquad Na_2SO_4 \quad 2H_2O$

0.05 mole 0.1 mole

– – 0.05 0.1

Aqueous solution of *SASB* type salt

i.e., pH 7

26. (c) HCl NaOH NaCl H_2O

(i) 2.5 m mole 2.49 mole

$$[H \ ] \quad \frac{0.01}{50} \quad 2 \quad 10^{-4} \ M$$

pH $4 \quad \log 2 \quad 3.7$

(ii) 2.5 m mole 2.51 M mole

$$[OH \ ] \quad \frac{0.01}{50} \quad 2 \quad 10^{-4} \ M$$

$pOH \quad 3.7$

27. (b) Milli-equivalents of HCl $25 \quad 0.08 \quad 2$

Milli-equivalents of NaOH 2.5

remaining $[OH \ ] \quad \dfrac{2.5 \quad 2}{500} \quad 10^{-3}$

$[H \ ] \quad 10^{-11}$ or pH 11

For More Material Join: @JEEAdvanced_2024

28. (c) $\underset{0.1\,m\text{ mole}}{H} \quad \underset{0.2\,m\text{ mole}}{OH} \quad H_2O$

LR

$[OH^-]_{Rem.} \quad \frac{(0.2 \quad 0.1)}{10 \quad 10} M$

29. (b) 20 meq. of HCl reacts completely with 20 meq. of KOH. Hence, the final solution is neutral. Thus, at 90°C, pH 6.

30. (b) In aqueous reaction, upon dilution, the reaction moves towards more equation moles.

31. (b) $NH_4(aq) \rightleftharpoons NH_3(aq) \quad H^+(aq) : K_1$

$HCN(aq) \rightleftharpoons H^+(aq) \quad CN^-(aq) ; K_2$

$NH_4(aq) \quad CN^-(aq) \rightleftharpoons NH_3(aq) \quad HCN(aq)$

$K_{eq} \quad \frac{K_1}{K_2}$

32. (a) $\because$ Strength of acid $\quad \frac{1}{pK_a}$

Formic acid will be the strongest acid.

33. (b) More H^+_{ion} weaker is the acid.

34. (a) $\sqrt{\frac{K_a}{C}} \quad 0.05$

$i.e., [H^+] \quad \sqrt{K_a \; C}$

35. (a) $\underset{\text{at equilibrium} \quad C(1 \quad)}{HA(aq.)} \rightleftharpoons \underset{C}{H^+(aq.)} \quad \underset{C}{A^-(aq.)}$

$K_a \quad \frac{C^{\;2}}{1}$

$10^{\;4} \quad \frac{(0.01)^{\;2}}{1}$

0.095

$\% \quad 9.5$

36. (b) $\sqrt{\frac{K_a}{C}}$

37. (c) $\because \quad 0.05$

$K_a \quad \frac{C^{\;2}}{1}$

38. (a) pH 9; $[OH^-] \quad 10^{\;5}$; $C \quad 10^{\;5}$

$10^{\;4}$

% dissociation $\quad 10^{\;4} \quad 100 \quad 0.01\%$

39. (c) $[H^+] \quad C$ and $[OH^-] \quad \frac{K_w}{[H^+]}$

40. (c) is negligible w.r.t. 1 so $K_a \quad C^{\;2}$

$0.1 \quad (0.01)^2 \quad 10^{\;5}$

$pK_a \quad 5$

41. (c) $\underset{C \quad x}{HCOOH} \rightleftharpoons \underset{x \quad 0.01}{H^+} \quad \underset{x}{HCOO^-}$

$K_a \quad \frac{(x \quad 0.01)\,x}{0.01 \quad x}$

neglect x w.r.t. 0.01 due to common ion effect

$K_a \quad x$

$x \sim [HCOO^-] \quad 1.8 \quad 10^{\;4}$

42. (c) (P) $(pK_a)_{HX} \quad (pK_b)_X \quad pK_w$

(Q) Very dilute solution

(R) pH $\quad \frac{1}{2} pK_a \quad \frac{1}{2} \log_{10} C$

(S) $[H^+]_1 \quad 10^{\;3} M$, $[H]_2 \quad 10^{\;5} M$

$[H^+]_{Final} \quad \frac{M_1V_1 \quad M_2V_2}{V_1 \quad V_2}$

43. (b) $\because$ is negligible w.r.t. 1

$K_a \quad C_1^{\;2}{}_1 \quad C_2^{\;2}{}_2$

$C_2 \quad C_1 \quad \frac{1}{2}$

$0.2 \quad \frac{1}{4} \quad 0.05$

$C_1V_1 \quad C_2V_2$

$300 \quad 0.2 \quad 0.05 \quad V_2$

$V_2 \quad 1200$ mL

Volume of H_2O added $\quad 1200 \quad 300$

$= 900$ mL

44. (c) Due to low value of K_b and common ion effect we can neglect x w.r.t. 0.01 when $x \quad c$

$K_b \quad \frac{x \quad 0.01}{0.02}$

$x \quad [NH_4^+] \quad 3.6 \quad 10^{\;5}$

45. (b) $n_{RNH_2(g)} \quad \frac{1 \quad 22.41}{0.0821 \quad 273} \quad 1$

$[RNH_2(aq.)] = \frac{n_{RNH_2}}{\text{Volume of solution}} \quad 1$

is negligible w.r.t. 1 so

$pOH \quad \frac{1}{2}[pK_b \quad \log C] \quad 2$

46. (a) is negligible w.r.t. 1 for both acid so

$[H^+] \quad \sqrt{K_{a_1}C_1 \quad K_{a_2}C_2}$

$\sqrt{1.8 \quad 10^{\;5} \quad 0.01 \quad 6.3 \quad 10^{\;5} \quad 0.01}$

$[H^+] \quad \sqrt{81 \quad 10^{\;8}} \quad 9 \quad 10^{\;4}$

For More Material Join: @JEEAdvanced_2024

47. (c) $[HA]$ $\frac{\frac{6}{60}}{10000}$ $10^{-5}M$

$\sqrt{\frac{10^{-9}}{10^{-5}}}$ 0.01 (negligible w.r.t.1)

$[H^+]$ $\sqrt{10^{-9} \quad 10^{-5}}$ $\sqrt{10^{-14}}$

$10^{-7}M$, $\because$ $[H^+]$ $10^{-6}M$

So we should consider $[H^+]$ from H_2O

$[H^+]_{total}$ 2 10^{-7} without considering common ion effect pH = 6.7, so pOH = 7.3

but due to C.I. effect H from H_2O will be less

so pH > 6.7

pOH of solution is exist between 7.0 and 7.3.

48. (d) and 49. (d)

K_{a_1} $\frac{[H^+]_{total}\,[HCO_3^-]}{[H_2CO_3]}$

$\because$ K_{a_1} K_{a_2} and concentration is appreciable so we can neglect w.r.t. 1

$[H^+]_{total} \sim [HCO_3^-]$;

$[HCO_3^-]$ $\sqrt{4 \quad 10^{-7} \quad 0.025}$ 10^{-4}

and $[CO_3^{2-}] \sim K_{a_2}$

50. (d) $[H^+]_{total} \sim [H^+]$ from first step

K_{a_1} $\frac{C^{\ 2}}{(1 \quad)}$; is not negligible w.r.t. 1 so

after solving quadratic equation 0.095

$[OH^-]$ $\frac{10^{-14}}{0.095 \quad 0.3}$

$3.5 \quad 10^{-13}$

52. (b) Salt of *SAWB* type.

53. (b) Salt of *WASB* type.

54. (c) pH $\frac{1}{2}pK_a$ $\frac{1}{2}\log C$

5 $\frac{1}{2}pK_a$ $\frac{1}{2}$ 2

pK_a 8 pK_b 8;

pOH $\frac{1}{2}pK_b$ $\frac{1}{2}\log C$ 4 2 6

56. (d) All are *WASB* salts,

So, $[OH^-]$ $\sqrt{\frac{K_w \ C}{K_a}}$

For weakest acid, K_a should be minimum.

$[OH^-]$ minimum

$[H^+]$ maximum

pH minimum

57. (b) I *WASB* salt

II *WA*

III *WAWB* salt

IV *SB*

V SA

58. (c) As temperature pKw

pH also

61. (b) $[H^+]_{final}$ $\frac{1 \quad 0.1}{(1 \quad 999)}$ $10^{-4}M$

pH 4

65. (b) Similar to Q. No.7

66. (c) $(K_b)_{C_6H_5O^-}$ $\frac{K_w}{10^{-10}}$

$[OH^-]$ $\sqrt{K_b \ c}$

67. (a) $K_{a(HOCN)}$ $\frac{10^{-14}}{10^{-10}}$ 10^{-4}

K_h $K_{b(OCN^-)}$ $\frac{10^{-10}}{\sqrt{\frac{K_h}{C}}}$ $\sqrt{\frac{10^{-10}}{0.01}}$ 10^{-4}

$\because$ 0.1; K_h $c^{\ 2}$

$[OH^-]$ c ; $[OH^-]$ $0.01 \quad 10^{-4}$ $10^{-6}M$

68. (c) $[H^+]$ C

0.05

K_a $\frac{C^{\ 2}}{1}$

69. (d) $\frac{1}{2}$ $\sqrt{\frac{C_2}{C_1}}$ $\frac{0.01}{2}$ $\sqrt{\frac{0.4}{0.1}}$

$_2$ 0.005 or % $_2$ 0.5

70. (a) For *WAWB* salt, $\sqrt{\frac{K_w}{K_a \ K_b}}$ $\frac{}{1 \quad C}$ $\sqrt{K_h}$

71. (d) $\sqrt{\frac{K_w}{K_a C}}$

|Enthalpy of neutralisation| K_a h

72. (b) [Na*A*] $\frac{25 \quad 0.1}{25 \quad 50}$ $\frac{2.5}{75}$

$[H^+]$ $\sqrt{\frac{K_w \ K_a}{C}}$ $1.732 \quad 10^{-9}$

74. (c) pH $\frac{1}{2}(pK_w \quad pK_a \quad pK_b)$ 6.5

75. (b) $\because$ pH $\frac{1}{2}pK_w$ $\frac{1}{2}pK_a$ $\frac{1}{2}pK_b$ 7

For More Material Join: @JEEAdvanced_2024

and $\sqrt{\frac{K_w}{K_a K_b}}$ $\sqrt{\frac{10^{-14}}{10^{-12}}}$ 0.1

or % 10

76. (b) does not depend on concentration in this case.

77. (a) $\sqrt{\frac{K_h}{C}}$; $[H_3O^+]$ C

$\sqrt{K_h C}$ $\sqrt{0.02 \times 5 \times 10^{-9}}$ 10^{-5}

78. (d)

$Ni^{2+}(aq.)$ $H_2O(l) \rightleftharpoons Ni(OH)^+(aq.)$ $H^+(aq.)$

t 0 0.04M

$t_{eq.}$ (0.04 x) (x) (x)

x 4.5×10^{-6}

K_a $\frac{x^2}{(0.04 - x)}$

79. (b) is not negligible w.r.t. 1 so after solving quadratic equation 0.095

$[H^+]$ C 0.0095; pH 2.02

80. (c) For amphiprotic salt,

81. (b) pH $\frac{pK_{a_2} + pK_{a_3}}{2}$

82. (d) Similar to Q. No. 83

83. (a)

	CH_3COOH + NaOH		CH_3COONa + H_2O	
initial milli-moles	6	2	—	—
after reaction	4	—	2	—

pH pK_a $\log \frac{[CH_3COO^-]}{[CH_3COOH]}$ 4.44

85. (c)

	HX	NaOH	NaX	H_2O
(I)	2m mole	1m mole	–	–
	(2 1)	–	1	

ABS

pH pK_a $\log \frac{[Salt]}{[Acid]}$

	HX	NaOH	NaX	H_2O
(II)	2m mole	2m mole	–	–
	2	2	2	

$[NaX]$ 0.01M

pH $7 - \frac{1}{2}[pK_a - \log C]$

86, 87, 90 (a, c, b)

For *ABS*

pH pK_a $\log \frac{[Salt]}{[Acid]}$

88. (b) For *BBS*

pOH pK_b $\log \frac{[Salt]}{[Base]}$

91. (c)

	CH_3NH_2 + HCl		CH_3NH_3Cl
initial moles	0.12	0.08	
after reaction	0.04	—	0.08

pOH pK_b $\log \frac{[CH_3NH_3^+]}{[CH_3NH_2]}$

3.3 $\log \frac{0.08}{0.04}$ 3.6

pH 10.4

92. (d) pOH pK_b $\log \frac{[NH_4^+]}{[NH_4OH]}$ 6

pH 14 6 8.0

94. (a) *BBS*

$NH_4OH \rightleftharpoons NH_4^+ + OH^-$

Addition of HCl takes equilibrium to right

NH_4OH	H^+	NH_4^+	H_2O
0.2 mole	10^{-6}	0.2	
$(0.2 - 10^{-6})$	–	$(0.2 + 10^{-6})$	

$[OH^-]$ $K_b \frac{[Base]}{[Salt]}$

95. (d) HCOOH NaOH HCOONa H_2O

(I) a $a/5$

$(a - a/5)$ – $(a/5)$

pH_1 pK_a $\log \frac{[HCOOH]}{[HCOONa]}$

(II) a $4a/5$

$(a - 4a/5)$ – $4a/5$

pH_2 pKa $\log \frac{[HCOOH]}{[HCOONa]}$

pH pH_2 pH_1

97. (a) Henderson equation is

pH pK_a $\log \frac{[salt]}{[acid]}$ 4

pOH 10, $[OH^-]$ 10^{-10}

Number of OH^- ions $10^{-10} N_A$;

where N_A Avogadro's number.

98. (d) pH upon addition of base.

99. (b) SASB salt.

100. (a) $pK_b(NH_4OH)$ 14 $pK_a(NH_4^+)$ 4.74

So, buffer range,

pK_b 1

101 (a) Mol of C_6H_5COOH $\frac{61}{122}$ 0.5;

Mol of C_6H_5COONa $\frac{72}{144}$ 0.5

For More Material Join: @JEEAdvanced_2024

pH pK_a $\log \frac{(C_6H_5COONa)}{(C_6H_5COOH)}$;

pH 4.2 $\log \frac{0.2}{0.8}$ 3.6

102. (b) pH pK_{a_2} $\log \frac{[CO_3^{2-}]}{[HCO_3^-]}$

(11 log 4) $\log \frac{0.2}{0.4}$ 10.1

103. (c) H_3PO_4 PO_4^{3-} $H_2PO_4^-$ HPO_4^{2-}

initial milli moles	20 0.1	20 0.1		
after reaction	—	—	2	2

Buffer solution of $H_2PO_4^-$ (acid) and HPO_4^{2-} (conjugate base) is formed;

pH pK_{a_2} $\log \frac{[HPO_4^{2-}]}{[H_2PO_4^-]}$ pK_{a_2}

106. (d) At eq. point, *WASB* salt forms means base solution

So, *CTR* 7

107. (d) At equivalent point

pH $\frac{1}{2}[pK_w \quad pK_a \quad \log C]$

$\frac{1}{2}[14 \quad 4.74 \quad 2]$ 8.37

Where [Salt] $\frac{\text{milli-moles of acid}}{\text{Total volume}}$

$\frac{20 \quad 0.02}{20 \quad 20}$ 0.01

For best indicator pH at equivalent point should lie between colour transition range of indicator.

108. (b) $HIn \rightleftharpoons H \quad In$

Acid Basic

$Ka \quad \frac{[H^+][In^-]}{[HIn]}$

$6 \quad 10^{-5} \quad \frac{10^{-5}(x)}{(100 \quad x)}$

109. (a) $HIn \rightleftharpoons In^- \quad H^+$; $K_{In} \quad \frac{[H^+][In^-]}{[HIn]}$

red blue

Case I : $3 \quad 10^{-5} \quad \frac{[H^+]_1 \quad 0.25}{0.75}$

Case II : $3 \quad 10^{-5} \quad \frac{[H^+]_2 \quad 0.75}{0.25}$

$[H^+]_1 \quad 9 \quad 10^{-5} \; [H^+]_2 \quad 10^{-5}$

Change in $[H^+]$ 8 10^{-5}

110. (c) pH pK_{In} $\log \frac{[In^-]}{[HIn]}$ pK_a $\log \frac{[Salt]}{[Acid]}$

111. (d) H_2A 2NaOH Na_2A H_2O

For second eq. pt

$(n)_{NaOH} \quad 2(n)_{H_2A} \quad [Na_2A] \quad \frac{0.4}{(32 \quad 250)}$

$0.25 \quad V \quad 2(0.2 \quad 20)$

V 32 mL

112. (c) H_2A NaOH NaHA H_2O

(I) a $a/2$

$(a \quad a/2)$ $a/2$

WA its conjugate ABS

pH pKa_1 $\log \frac{[NaHA]}{[H_2A]}$ $\frac{[NaHA]}{[H_2A]}$ pKa_1

(II) a a

– – (a)

aq. solution of amphiprotic salt

pH $\frac{pK_{a_1} \quad pK_{a_2}}{2}$

114. (a) When equal volume mixed concentration become half also, for ppt,

$[Ag^+][Cl^-] \quad K_{sp}$

116. (c) When 1 litre each solution are mixed

$[OH^-] \quad 10^{-6} \; M$ (Buffer solution)

M^{n+} 0.05 M

for $Q \quad [0.5][10^{-6}]^2 \quad 5 \quad 10^{-14}$

$Q \quad K_{sp}$ for Fe^{2+} and Ni^{2+}

So $Fe(OH)_2$ and $Ni(OH)_2$ are ppt.

117. (a) For $[Ca^{2+}]$, $M_1V_1 \quad M_2V_2$

$0.04 \quad 50 \quad M_2 \quad 200 \quad M_2 \quad 0.01$

Similarly, for $[SO_4^{2-}]$

$0.0008 \quad 150 \quad M_2 \quad 200$

M_2 0.0006

ionic product (Q) $[Ca^{2+}] \quad [SO_4^{2-}]$

$6 \quad 10^{-8}$ So $Q \quad K_{sp}$

118. (c) For AgCl

$K_{sp} \quad [Ag^+][Cl^-]$

$y \quad x$

xy constant

i.e., Rectangular hyperbolic

119. (a) KNO_3 does not effect the equilibrium.

120. (a) $[Ag^+] \quad \frac{10^{-3}}{50 \quad 50}$ 1000 0.01

For More Material Join: @JEEAdvanced_2024

$[Cl^-] = \frac{0.1 \times 50}{50 + 50} = 0.05$

$Q = [Ag^+][Cl^-] = 5 \times 10^{-4}$

$\because Q > K_{sp}$

$Ag^+(aq.) + Cl^-(aq.) \rightarrow AgCl(s)$

	Ag^+	Cl^-
at given condition	0.01	0.05
after precipitation	x	~ 0.04

$K_{sp} = [Ag^+][Cl^-]$; $10^{-10} = x \times 0.04$

$x = [Ag^+] = 2.5 \times 10^{-9}$

121. (c) $A_xB_y(s) \rightleftharpoons x\ A^{y+}_{(aq.)} + y\ B^{x-}_{(aq.)}$

$t = 0 \quad a$

$t_{eq.} \quad (a - s) \quad (x \times s) \quad (x \times y)$

$k_{sp} = [A^{y+}]^x[B^{x-}]^y$

$= (x \times s)^x(y \times s)^y$

122. (c) Molarity of saturated solution = Solubility

123. (c) $Zr_3(PO_4)_4(s) \rightleftharpoons 3\ Zr^{4+}_{(aq.)} + 4PO^{3-}_{4(aq.)}$

$t = 0 \quad a$

$t_{eq.} \quad (a - s) \quad 3s \quad 4s$

$K_{sp} = [Zr^{4+}]^3[PO_4^{3-}]^4$

124. (a) The corresponding values of solubility products of the given electrolytes are :

$MX_1 \quad MX_2 \quad MX_3$

$S^2 = 10^{-6}, \quad 4S^3 = 4 \times 10^{-9}, \quad 27S^4 = 27 \times 10^{-12}$

125. (c) $K_{sp} = [Ca^{2+}]^3[PO_4^{3-}]^2$

126. (d) For most soluble salt, solubility should be maximum.

127. (a) (A) $[Ag^+] = \frac{K_{sp}}{[Br^-]} = 5 \times 10^{-12} M$

(B) $[Ag^+] = \frac{K_{sp}}{[Cl^-]} = 1.8 \times 10^{-7} M$

(C) $[Ag^+] = \sqrt{\frac{K_{sp}}{[CO_3^{2-}]}} = 9 \times 10^{-6} M$

(D) $[Ag^+] = \left(\frac{K_{sp}}{[ASO_4^{3-}]}\right)^{1/3} = 10^{-7} M$

128. (a) Solubility of $AB = \sqrt{K_{sp}} = 2 \times 10^{-10}$

Solubility of $A_2B = \sqrt[3]{\frac{K_{sp}}{4}} = 2 \times 10^{-4}$

Solubility of $AB_3 = \left(\frac{K_{sp}}{27}\right)^{1/4} = 10^{-8}$

129. (d) $HgSO_4(s) \rightleftharpoons Hg^{2+}(aq.) + SO_4^{2-}(aq.)$

$S = \sqrt{6.4 \times 10^{-5}} = 8 \times 10^{-3}$ mol/L

$= 8$ mol/m^3

130. (b) $s = \frac{6.9 \times 10^{-2}}{690}$ mol/100 mL $= 10^{-3}$ mol/L

For $Ba_3(AsO_4)_2$, $K_{sp} = 108\, s^5$

131-132-133. (d, b, c)

Similar to Q.No. 130

134. (c) $BaF_2(s) \rightleftharpoons Ba^{2+}(aq.) + 2F^-(aq.)$

$S \quad 2S$

$K_{sp} = 4S^3$; $S = \left(\frac{10^{-6}}{4}\right)^{1/3} = 0.63 \times 10^{-2}$

Molarity of F$^-$ $= 2S = 2 \times 0.63 \times 10^{-2} = 1.26 \times 10^{-2}$

135. (d) $InF_3(s) \rightleftharpoons In^{3+}(aq) + 3F^-(aq)$

$t = eq \quad s \quad 3s$

$K_{sp} = (s)^1(3s)^3$

136. (d) $Cu(OH)_2(s) \rightleftharpoons Cu^{2+}(aq.) + 2OH^-(aq.)$

$S \quad 2S$

$K_{sp} = 4S^3$

$S = 4 \times 10^{-7}$

$[OH^-] = 2S = 8 \times 10^{-7}$

pOH $= 6.1$ pH $= 7.90$

137. (a) $[Ag^+]$ or $S = \sqrt{K_{sp}} = 10^{-5} M$

$10^{-5} = \frac{\frac{14.35 \times 10^{-3}}{143.5}}{V \text{ (in litre)}}$

$V = 10$ litre or 0.01 m^3

138. (d) $Fe(OH)_2(s) \rightleftharpoons Fe^{2+}(aq.) + 2OH^-(aq.)$

$t_{eq.} \quad s \quad 10^{-1} M$

$K_{sp} = [Fe^{2+}][OH^-]^2$

$8 \times 10^{-16} = (s)^1(10^{-1})^2$

139. (c) To precipitate

$[Pb^{2+}][OH^-]^2 > K_{sp}$

$[OH^-]_{Min} = \sqrt{\frac{K_{sp}}{[Pb^{2+}]}}$

141. (a) Let solubility of Ag_2CO_3 in presence of Na_2CO_3 is x,

$Ag_2CO_3(s) \rightleftharpoons 2Ag^+(aq.) + CO_3^{2-}(aq.)$

$2x \quad (x + 0.1) \sim 0.1$

$K_{sp} = [Ag^+]^2[CO_3^{2-}]$

$4 \times 10^{-13} = (2x)^2 \times 0.1$

$x = 10^{-6}$

142-143. (d, c)

Similar to Q.No. 139

144. (c) More common ion more in solubility.

For More Material Join: @JEEAdvanced_2024

145. (d) K_{sp} 256 s^5

146. (a) $PbBr_2(s) \rightleftharpoons Pb^{2}(aq.)$ $2Br^{-}(aq.)$

$0.8\,S$ $2\ 0.8\ S$

K_{sp} $[Pb^{2}][Br^{-}]^2$

$8\ 10^{-5}$ $(0.8\,S)(1.6\,S)^2$

S $\left(\frac{10^{-4}}{1.6\ 1.6}\right)^{1/3}$

147. (b) For basic buffer solution

pOH pK_b $\log\frac{[NH_4]}{[NH_3]}$ pK_b

$[OH^-]$ $1.8\ 10^{-5}$

K_{sp} $[Mn^{2}][OH^-]^2$

$[Mn^{2}]$ $1.38\ 10^{-4}\ M$

148. (b) $[OH^-]$ $\sqrt{\frac{K_{sp}}{[Mg^{2}]}}$ $\sqrt{\frac{10^{-11}}{10^{-3}}}$ 10^{-4}

pOH pK_b $\log\frac{[NH_4Cl]}{[NH_3]}$

4 5 $\log\frac{[NH_4Cl]}{[NH_3]}$

0.10 $\frac{[NH_4Cl]}{[NH_3]}$

moles of NH_4Cl required 0.1 0.02

$2\ 10^{-3}$

149. (d) Due to higher value of K_f, mostly Ag converted into complex.

$AgI(s) \rightleftharpoons Ag^{+}(aq.)$ $I^-(aq.)$; K_{sp}

$Ag^{+}(aq.) + 2NH_3(aq.) \rightleftharpoons Ag(NH_3)_2^{+}(aq.)$; K_f

overall reaction is

$AgI(s)$ $2NH_3(aq.) \rightleftharpoons Ag(NH_3)_2(aq.)$ $I^-(aq.)$

K_{eq} K_{sp} K_f

K_{sp} K_f $\frac{x^2}{1^2}$

x $4.9\ 10^{-5}$ mol/litre

So mass of AgI required is

$4.9\ 10^{-5}$ 235 0.011 g

Level 2

1. (d) TCl + $NaOT$ $\rightarrow$ $NaCl$ + T_2O

Initial milli-moles 2 3.75

milli-moles of remaining NaOT 1.75

$[OT^-]$ $\frac{1.75}{25}$ $7\ 10^{-2}$ moles

pOT 2 $\log 7$

$pT + pOT = 7.62$ 2

pT 15.24 2 $\log 7$

13.24 $\log 7$

2. (c) $2NH_3 \rightleftharpoons NH_4^{+}$ NH_2^-; $[NH_4^+]$ $\sqrt{K_{SIC}}$

K_{SIC} $[NH_4^+][NH_2^-]$

$[NH_4^+]$ $10^{-15}\ M$

$\because$ Number of NH_4^+ ions in $1000\ cm^3$

10^{-15} 6.022 10^{23}

Number of NH_4^+ ions in $100\ cm^3$

10^{-15} 6.022 10^{23} $\frac{100}{1000}$

6.022 10^{7}

3. (c) is very less so $[H^+]$ $\sqrt{K_a\ C}$ $3\ 10^{-3}$

Now $[H^+]$ $\frac{3\ 10^{-3}}{2}$ $1.5\ 10^{-3}$

$(1.5\ 10^{-3})$ $\sqrt{K_a\ C_2}$

C_2 0.125

milli-moles of acid remain constant

C_1V_1 C_2V_2

10 0.5 0.125 V_2

V_2 40 litre

4. (a) $[H^+]$ $\frac{10\ 0.3\ 10\ 0.1}{10\ 10\ 20}$ 0.05

$[HA]$ $\frac{20\ 0.1}{20\ 20}$ 0.05

HA $\rightleftharpoons$ H^+ A^-

$0.05\ x$ $0.05\ x$ x

Due to common ion effect neglect x w.r.t. 0.05

K_a $\frac{(0.05\ x)\ x}{(0.05\ x)} \sim x$

x 10^{-5}

$\frac{[A^-]}{[HA]}$ $[A^-]$ $\frac{x}{x\ 0.05}$ $2\ 10^{-4}$

5. (d) $FCH_2COOH(aq.) \rightleftharpoons FCH_2COO^-(aq.)$ $H^+(aq.)$

at eqm. $C(1\ \ )$ C C

K_a $\frac{[H^+][FCH_2COO^-]}{[FCH_2COOH]}$

$2.6\ 10^{-3}$ $\frac{(2\ 10^{-3})^2}{[FCH_2COOH]}$

at eqm. $[FCH_2COOH]$ $1.53\ 10^{-3}$

Total concentration $2\ 10^{-3}$ $1.53\ 10^{-3}$

$3.53\ 10^{-3}$

For More Material Join: @JEEAdvanced_2024

6. (c) $HCOOH(aq.)$ $H_2O(l) \rightleftharpoons HCOO^-(aq.)$ $H_3O^+(aq.)$
C_1 x x x
$HF(aq.) + H_2O(l) \rightleftharpoons F^-(aq.)$ $H_3O^+(aq.)$
C_2 y y x y
$\because$ is very less for both acid so
$K_a(HCOOH) \frac{x(x \quad y)}{C_1}$;
$K_a(HF) \frac{y(x \quad y)}{C_2}$
$\frac{x}{y} \frac{K_a(HCOOH)C_1}{K_a(HF)C_2}$
$\frac{[HCOO^-]}{[F^-]} \frac{x}{y}$
$\frac{2 \quad 10^{-4} \quad 0.2}{6.6 \quad 10^{-4} \quad 0.1} \frac{2}{3.3}$

7. (a) First dissociation
$X(OH)_3$ $X(OH)_2^+$ OH^-
Second dissociation :
$X(OH)_2^+$ $X(OH)^{2+}$ OH^-
Total $[OH^-]$ $4 \quad 10^{-3}$ $2 \quad 10^{-3}$ $6 \quad 10^{-3}$
pOH $3 - \log 6 \sim 2.22$
pH 11.78

8. (d) is negligible w.r.t. 1
$[H^+]$ mainly from first step
$[H^+] \sqrt{K_{a_1} \quad C} \sqrt{10^{-5} \quad 0.1} \quad 10^{-3}$
$K_{a_3} \frac{[H^+][A^{3-}]}{[HA^{2-}]} \quad 10^{-13} \quad \frac{10^{-3}[A^{3-}]}{[HA^{2-}]}$
$X \frac{[A^{3-}]}{[HA^{2-}]} \quad 10^{-10}$;
$pX \quad 10$

9. (a) $Ca(LaC)_2$ Ca^{2+} $2LaC^-$
Initial concentration $0.6/2 \quad 0.3\ M$
After dissociation $[LaC^-] \quad 2 \quad 0.3 \quad 0.9$
$0.54\ M$
$pH \frac{1}{2}[pK_w \quad pK_a \quad \log(LaC^-)]$
$14 \quad 5.6 \frac{1}{2}[14 \quad pK_a \quad \log(0.54)]$
$pK_a \quad 2.8 \quad \log(0.54)$

10. (a) $CH_3COO^-(aq.)$ $NH_4(aq.)$ $H_2O(l) \rightleftharpoons$
at equilibrium $C(1 \quad)$ $C(1 \quad)$
$CH_3COOH(aq.) + NH_4OH(aq.)$
C C
$K_h \frac{K_w}{K_a K_b} \frac{\quad^2}{(1 \quad)^2}$
$\frac{10^{-14}}{(1.8 \quad 10^{-5})^2} \frac{\quad^2}{(1 \quad)^2}$
$\frac{10^{-2}}{1.8} \frac{\quad}{1}$; $5.55 \quad 10^{-3}$
$[CH_3COOH] \quad C$
$0.01 \quad 0.0055 \quad 5.55 \quad 10^{-5}$

11. (d) $Fe^{3+}(aq.)$ $H_2O(l) \rightleftharpoons Fe(OH)^{2+}(aq.)$
$+ H_3O^+(aq.)$
$K_a \frac{[Fe(OH)^{2+}][H_3O^+]}{[Fe^{3+}]}$
$6.5 \quad 10^{-3} \frac{0.20}{0.80} [H_3O^+]$
$[H_3O^+] \quad 2.6 \quad 10^{-2}$; $pH \quad 1.59$

12. (c) Due to larger difference between K_{h_1} and K_{h_2} and common ion effect we can assume that y is negligible w.r.t. x
$Fe^{2+} + H_2O \rightleftharpoons Fe(OH)^+$ H^+
at equilibrium $C \quad x$ $x \quad y$ $x \quad y$
$Fe(OH)^+ + H_2O \rightleftharpoons Fe(OH)_2 + H^+$
at equilibrium $x \quad y$ y $y \quad x$
$K_{h_2} \frac{K_w}{K_{b_1}} \frac{10^{-14}}{10^{-4}} \quad 10^{-10}$
$\because$ $x \quad y \quad x \quad y$
$K_{h_2} \quad y \quad Fe(OH)_2 \quad 10^{-10}$

13. (a) Let x milli-moles of KOH is added
$6 \quad 5 \log \frac{s \quad x}{a \quad x}$; $\frac{s \quad x}{a \quad x} \quad 10$
or $\frac{10 \quad x}{10 \quad x} \quad 10 \qquad x \quad 8.18$
Mass $8.18 \quad 10^{-3} \quad 56 \quad 0.458$ gm

14. (b) $pH \quad pK_a \quad \log \frac{[C_6H_5COO^-]}{[C_6H_5COOH]}$
$\frac{[C_6H_5COO^-]}{[C_6H_5COOH]} \quad 2$
Let volume of acid is V mL
$\frac{0.2 \quad (300 \quad V)}{0.1 \quad V} \quad 2 \qquad V \quad 150$ mL

For More Material Join: @JEEAdvanced_2024

15. (c) When half acid is neutralized pH pK_a and at the equivalent point

$$pH \quad \frac{1}{2}[pK_w \quad pK_a \quad \log C]$$

$$9 \quad \frac{1}{2}[14 \quad 5 \quad \log C] = C \quad 0.1$$

Let V mL of NaOH is used in titration milli eq. of NaOH $0.25 \quad V$ milli eq. of salt formed

$$\frac{0.25 \quad V}{V \quad 60} \quad 0.1 = V \quad 40 \text{ mL}$$

milli equivalent or milli-moles of acid 10

Mass of acid $10 \quad 10^{-3} \quad 82 \quad 0.82$ gm

Mass % of acid $\frac{0.82}{1.025} \quad 100 \quad 80\%$

16. (c) $BOH(aq.) \rightleftharpoons B^+(aq.) \quad OH^-(aq.)$

At equilibrium : $c(1 \quad) \qquad c \qquad c$

$$K_b \quad \frac{[B^+][OH^-]}{[BOH]} \qquad \frac{c^2}{(1 \quad)};$$

neglect w.r.t. 1 as concentration is appreciable, % $100\sqrt{\frac{K_b}{c}}$

Total dissolved base present in solution as BOH and B^+

So $\frac{[B^+]}{[B^+] \quad [BOH]}$

$$\frac{1}{1 \quad \frac{[BOH]}{[B^+]}} \qquad \frac{1}{1 \quad \frac{[OH^-]}{K_b}}$$

$$\frac{K_b}{K_b \quad [OH^-]} \qquad \frac{K_b[H^+]}{K_b[H^+] \quad K_w}$$

also $pOH = \quad \log[OH^-]; [OH^-] = 10^{-pOH};$

$$K_b \quad 10^{-pK_b}$$

$$\frac{1}{1 \quad \frac{10^{-pOH}}{10^{-pK_b}}} \qquad \frac{1}{1 \quad 10^{(pK_b \quad pOH)}}$$

17. (b) $HA + NaOH \qquad NaA + H_2O$

milli-moles of salt NaA or $A^- \quad 40 \quad 0.1 \quad 4$

Now

	A^-	H^+	HA
Initial milli-moles	4	2	
Final milli-moles	2	—	2

Acidic buffer solution is formed and $[A^- \quad] \quad [HA]$

$$pH \quad pK_a \quad \log \frac{[A^-]}{[HA]} \qquad pK_a \quad 5$$

Now $HA + NaOH \qquad NaA + H_2O$

hydrolysis of A^- will takes place

For More Material Join: @JEEAdvanced_2024

$$[NaA] \quad \frac{\text{milli moles of acid}}{\text{Total volume}} \quad \frac{20 \quad 0.2}{20 \quad 20} \quad 0.1$$

$$pH \quad \frac{1}{2}(pK_w \quad pK_a \quad \log C) \quad \frac{1}{2}[14 \quad 5 \quad 1] \quad 9$$

18. (c) Moles of H produced $4 \quad 10^{\ 3}$

$$[H^{\ }] \quad \frac{4 \quad 10^{\ 3}}{0.1} \quad 0.04\ M$$

	$PO_4^{3}\ (aq.)$	$H^{\ }\ (aq.)$	$HPO_4^{2}\ (aq.)$
Initial milli-moles	0.02	0.04	
Final milli-moles	—	0.02	0.02

	$HPO_4^{2}\ (aq.)$	$H^{\ }\ (aq.)$	$H_2PO_4(aq.)$
Initial milli-moles	0.06	0.02	—
Final milli-moles	0.04	—	0.02

Finally buffer solution of H_2PO_4 (acid) and HPO_4^{2} (conjugate base) is formed

$$pH \quad pK_{a_2} \quad \log \frac{[HPO_4^{2}\ ]}{[H_2PO_4]} \quad 7.2 \quad \log \frac{0.04}{0.02} \quad 7.5$$

19. (b) $BOH + HCl \qquad BCl + H_2O$

at equilibrium point $N_1V_1 \quad N_2V_2$; $V_2 \quad 20$ mL

$$[BCl] \quad \frac{20 \quad 0.08}{20 \quad 20} \quad 0.04$$

$$pH \quad \frac{1}{2}[pK_w \quad pK_b \quad \log (C)]$$

$$pK_b \quad 5.4$$

$B \quad OH \qquad BOH$; Basic Buffer is formed

	B	OH	BOH
Initial milli-moles	1.6	0.4	
Final milli-moles	1.2	—	0.4

$$pOH \quad pK_b \quad \log \frac{[B^{\ }]}{[BOH]}$$

$$5.4 \quad \log \frac{1.2}{0.4} \quad 5.4 \quad 0.48$$

$$pOH \quad 5.88$$

20. (b) $CO_3^{2} \qquad H^{\ } \qquad HCO_3$ acidic buffer is formed.

	CO_3^{2}		$H^{\ }$		HCO_3
Initial milli-moles	25	0.04	50	0.025	1
Final milli-moles	—		0.25		

	HCO_3	$H^{\ }$	H_2CO_3
Initial milli-moles	1	0.25	
Final milli-moles	0.75	—	0.25

$$pH \quad pK_{a_1} \quad \log \frac{[HCO_3]}{[H_2CO_3]} \quad 6.4 \quad \log \frac{0.75}{0.25} \quad 6.88$$

21. (b) Let a is initial milli-moles of HA and molarity of NaOH solution is x

$$HA + NaOH \qquad NaA + H_2O$$

In both case solution is acidic, so acidic buffer solution is formed.

$$5 \quad pK_a \quad \log \frac{10x}{a \quad 10x}$$

For More Material Join: @JEEAdvanced_2024

$$5.60 \quad pK_a \quad \log \frac{20x}{a \quad 20x}$$

$$0.60 \quad \log \frac{20x}{(a \quad 20x)} \quad \frac{(a \quad 10x)}{10x}$$

$$4 \quad \frac{2\,(a \quad 10x)}{(a \quad 20x)} \quad a \quad 30x$$

$$5 \quad pK_a \quad \log \frac{10x}{20x}$$

or $\quad pK_a \quad 5.3$

22. (d)

	CO_3^{2}		H		HCO_3
Initial milli-moles	50	0.05	40	0.1	—
Final milli-moles	—		1.5		2.5

	HCO_3	H	H_2CO_3
Initial milli-moles	2.5	1.5	—
Final milli-moles	1	—	1.5

$$pH \quad pK_{a_1} \quad \log \frac{[HCO_3]}{[H_2CO_3]} \quad 6.173$$

23. (c) At second equivalence point the only species present at appreciable concentration is HA^{2}

So, $\quad pH \quad \frac{pK_{a_2} \quad pK_{a_3}}{2} \quad \frac{8 \quad 12}{2} \quad 10$

$$[H \ ] \quad 10^{\ 10}$$

and $\quad \frac{[H_3A]}{[A^{3} \ ]} \quad \frac{[H \ ]^3}{K_{a_1}K_{a_2}K_{a_3}} \quad 10^{\ 7}$

24. (c) Solubility $\quad S \quad \frac{x}{M}$

$$A_3B_2(s) \rightleftharpoons 3A^{2}\,(aq.) \quad 2B^{3}\,(aq.)$$

$$3S \qquad 2S$$

$$K_{sp} \quad (3\,S)^3\,(2\,S)^2 \quad 108S^5$$

$$\frac{[B^{3} \ ]}{K_{sp}} \quad \frac{2\,S}{108S^5} \quad \frac{1}{54}\,\frac{M^4}{x^4}$$

25. (d) $\quad K_{sp}\,(SrCrO_4) \quad [Sr^{2} \ ][\ CrO_4^{2} \ ]$

$$[CrO_4^{2} \ ] \quad \frac{3.5 \quad 10^{\ 5}}{0.1} \quad 3.5 \quad 10^{\ 4}$$

$K_{sp}\,(BaCrO_4) \quad [Ba^{2}]\,[CrO_4^{2} \ ]$

$[CrO_4^{2} \ ]_{total} \sim [CrO_4^{2} \ ]$ from $SrCrO_4$

$$[Ba^{2} \ ] \quad \frac{1.2 \quad 10^{\ 10}}{3.5 \quad 10^{\ 4}} \quad 3.4 \quad 10^{\ 7}$$

27. (a) $[Ag \ ]$ required for precipitation of AgCl

$$[Ag \ ] \quad \frac{K_{sp}\,(AgCl)}{[Cl \ ]} \quad 10^{\ 9}$$

$[Ag \ ]$ required for precipitation of Ag_2CrO_4

$$\sqrt{\frac{K_{sp}\,(Ag_2CrO_4)}{[CrO_4^{2} \ ]}} \quad \sqrt{\frac{10^{\ 12}}{10^{\ 4}}} \quad 10^{\ 4}$$

For More Material Join: @JEEAdvanced_2024

At this point, the concentration of Cl ion in the solution can be calculated from K_{sp} (AgCl)

$$[\text{Cl}^-] = \frac{K_{sp}(\text{AgCl})}{[\text{Ag}^+]} = \frac{10^{-10}}{10^{-4}} = 10^{-6}\ M$$

28. (b) After mixing $[\text{Ag}^+] = 0.2\ M$; $[\text{NH}_3] = 1\ M$

Due to very high value of K_f; Ag mainly converted into complex

$$\text{Ag}^+(aq.) + 2\text{NH}_3(aq.) \rightleftharpoons \text{Ag}(\text{NH}_3)_2^+(aq.)$$

Initial concentration	0.2	1	
At equilibrium	x	0.6	~ 0.2

$$\text{Ag}(\text{NH}_3)_2^+(aq.) \rightleftharpoons \text{Ag}(\text{NH}_3)^+(aq.) + \text{NH}_3(aq.)$$

At equilibrium	$0.2 - y$	y	$0.6 + y$
	~ 0.2		~ 0.6

$$\frac{1}{K_{f_2}} = \frac{y \times 0.6}{0.2} = \frac{0.1}{10^4};\ y = [\text{Ag}(\text{NH}_3)^+] = 3.33 \times 10^{-5} M$$

29. (a)

$$\text{AgCl}(s) + 2\text{NH}_3(aq.) \rightleftharpoons \text{Ag}(\text{NH}_3)_2^+(aq.) + \text{Cl}^-(aq.)$$

$$K_1 = 1.6 \times 10^{-10} \times 10^7 = 1.6 \times 10^{-3}$$

$$= \frac{[\text{Ag}(\text{NH}_3)_2][\text{Cl}^-]}{[\text{NH}_3]^2}$$

$$\text{AgCN}(s) + 2\text{NH}_3(aq.) \rightleftharpoons \text{Ag}(\text{NH}_3)_2^+(aq.) + \text{CN}^-(aq.)$$

$$K_2 = 2.5 \times 10^{-16} \times 10^7 = 2.5 \times 10^{-9}$$

$$= \frac{[\text{Ag}(\text{NH}_3)_2][\text{CN}^-]}{[\text{NH}_3]^2}$$

$$\frac{[\text{Cl}^-]}{[\text{CN}^-]} = \frac{1.6 \times 10^{-3}}{2.5 \times 10^{-9}} = 6.4 \times 10^5$$

$$K_1 = \frac{x^2}{(1 - 2x)^2} \approx \frac{x}{1 - 2x} = 0.04$$

$$x = 0.037$$

$$[\text{CN}^-] = \frac{0.037}{6.4 \times 10^5} = 5.78 \times 10^{-8}$$

32. (c)

$$\text{AgBr}(s) \rightleftharpoons \text{Ag}^+(aq.) + \text{Br}^-(aq.);\ K_{sp}$$

$$\text{Ag}^+(aq.) + 2\text{S}_2\text{O}_3^{2-}(aq.) \rightleftharpoons \text{Ag}(\text{S}_2\text{O}_3)_2^{3-}(aq.);\ K_f$$

$$\text{AgBr}(s) + 2\text{S}_2\text{O}_3^{2-}(aq.) \rightleftharpoons \text{Ag}(\text{S}_2\text{O}_3)_2^{3-}(aq.) + \text{Br}^-(aq.);\ K_{sp} \times K_f = 25$$

$$0.1 \quad 2x \qquad\qquad x \qquad\qquad x$$

Value K_f is very higher so we can assume almost Ag converts into complex

For More Material Join: @JEEAdvanced_2024

33. (d)

$Ag_2CrO_4(s) \rightleftharpoons 2Ag^+(aq.) + CrO_4^{2-}(aq.)$

$\overset{2x \quad 2y}{Ag_2C_2O_4(s) \rightleftharpoons 2Ag^+(aq.)} \quad \overset{x}{C_2O_4^{2-}(aq.)}$

$\overset{2y \quad 2x}{\qquad\qquad} \qquad \overset{y}{\qquad}$

$\frac{K_{sp_1}}{K_{sp_2}} \quad \frac{x}{y} \quad \frac{2 \quad 10^{-12}}{2 \quad 10^{-11}} \quad \frac{x}{y} \quad 0.1$

$2 \quad 10^{-11} \quad (2x \quad 2y)^2 \quad y$

$2 \quad 10^{-11} \quad 4.84y^3$

$y \quad 1.6 \quad 10^{-4};$

$x \quad 0.16 \quad 10^{-4}$

Total $[Ag^+] \quad 2x \quad 2y$

$(2 \quad 0.16 \quad 2 \quad 1.6) \quad 10^{-4}$

$3.52 \quad 10^{-4}$

34. (b) $K_{sp} \quad [Zn^{2+}][S^{2-}]$

$[S^{2-}] \quad \frac{10^{-21}}{0.01} \quad 10^{-19}$

for $K_{a_1} \quad K_{a_2} \quad \frac{[H^+]^2[S^{2-}]}{[H_2S]}$

$10^{-20} \quad \frac{[H^+]^2 \quad 10^{-19}}{0.1} \qquad [H^+] \quad 0.1$

or $\qquad$ pH $\quad$ 1

35. (d) Dissolved $Al(OH)_3$ present in solution as $Al^{3+}(aq.)$ as well as $[Al(OH)_4^-](aq.)$

$S \quad [Al^{3+}(aq.)] \quad [Al(OH)_4(aq.)]$

$S \quad \frac{K_{sp}}{[OH^-]^3} \quad K_c[OH^-]$

for minimum solubility

$\frac{dS}{d(OH^-)} \quad 0$

or $\qquad \frac{K_{sp} \quad 3}{[OH^-]^4} \quad K_c \quad 0$

$[OH^-] \quad \left(\frac{3K_{sp}}{K_c}\right)^{1/4}$

Level 3

Passage-1

2. (b) $[HCl] \quad \frac{0.01}{0.1} \quad 0.1\,M$

pOH $\quad 4.74 \quad \log \frac{0.3 \quad 0.1}{0.2 \quad 0.1} \quad 5.34$

pH $\quad 14 \quad 5.34 \quad 8.66$

Passage-3

3. (c) When acid is half neutralized pH $\quad pK_a$

$pK_a \quad 5$

4. (a) $K_{eq} \quad \frac{K_a(HB)}{K_a(HA)} \quad \frac{10^{-5}}{10^{-6}} \quad 10$

5. (b) When $V_{NaOH} \quad 0$ mL; pH $\quad 3$

or $\qquad [H^+] \quad 0.001$

$K_{a(HB)} \quad \frac{C \quad ^2}{(1 \quad)} \quad \frac{(0.001)}{1} \quad 10^{-5}$

0.01

$C \quad \frac{0.001}{0.01} \quad 0.1$

$C \quad \frac{n_{HB}}{V}$

$0.1 \quad \frac{5}{V}; V \quad 50$ mL

$[NaB] \quad \frac{\text{milli-moles of acid}}{\text{total volume}}$

$\frac{5}{50 \quad 50} \quad 0.05$

pH $\quad \frac{1}{2}[pK_w \quad pK_a \quad \log C]$

$\frac{1}{2}[14 \quad 5 \quad \log(0.05)]$

8.85

6. (a) For best indicator, pH $\sim pK_{In}$ of an indicator as well as pH transition range of the indicator must coincide with the steep portion of the titration curve.

One or More Answers is/are Correct

21. (b,c,d)

(a) $\quad S \quad \sqrt{K_{sp}} \quad 10^{-5}$ mol lit^{-1}

(b) $\quad s \quad 10^{-5}$ mol lit^{-1}

(c) $\quad AgCl(s) \rightleftharpoons Ag \quad Cl$

$S \quad 2 \quad S$

$10^{-10} \quad (S \quad 2)S \qquad S \quad \frac{10^{-10}}{2} \quad 5 \quad 10^{-11}$

For More Material Join: @JEEAdvanced_2024

(d) $AgCl(s) \rightleftharpoons [Ag(NH_3)_2] \quad Cl$

$Ag \quad 2NH_3 \rightleftharpoons [Ag(NH_3)_2]^2 \quad Cl$

$AgCl(s) \quad 2NH_3 \rightleftharpoons [Ag(NH_3)_2] \quad Cl$

$K \quad \frac{[Ag(NH_3)_2]\ [Cl\ ]}{[NH_3]^2} \quad K_{sp} \quad K_f$

$\frac{S^2}{(2 \quad 2S)^2} \quad 10^{\ 8} \quad 10^{\ 10} \quad 10^{\ 2}$

$S \quad 0.166\,M$

22. (a,c) $H_2A \rightleftharpoons H \quad HA$; $\because$ y is very less so $x \quad y \quad x$

$x \quad y \quad x$

At equilibrium $c \quad x \quad x \quad y \quad x \quad y$ $K_{a_2} \quad y \quad 10^{\ 12}$; $pK_{a_2} \quad 12$

$HA \rightleftharpoons H \quad A^2$ $K_{a_1} \quad \frac{x^2}{c \quad x}$

At equilibrium $x \quad y \quad x \quad y \quad y$

$pK_{a_2} \quad pK_{a_1} \quad 12 \quad 5 \quad 7$ $pH \quad \frac{1}{2}[pK_{a_1} \quad \log c]$

$3 \quad \frac{1}{2}[pK_{a_1} \quad 1] \quad pK_{a_1} \quad 5$

25. (a,b,c)

$pH_1 \quad 8 \log \frac{25}{75} \quad 8 \quad \log \frac{1}{3}$

$pH_2 \quad 8 \quad \log \frac{75}{25} \quad 8 \quad \log 3$

Change in pH $\quad \log 3 \quad \log \frac{1}{3} \quad 2 \log 3$

Subjective Problems

1. $[OH\ ] \quad \sqrt{K_b \ c}$

2. $[H\ ] \quad 10^{\ 4}M \quad C$

$\because \quad 0.05 \quad K_a \quad \frac{C^{\ 2}}{1}$

3. Consider ionisation of water.

4. $H \quad OH \quad H_2O$

(2.5 1) mole (2.5) m mole

L.R.

$[H\ ]_{Final} \quad \frac{(3.5 \quad 2.5)}{25 \quad 50 \quad 25} \quad 10^{\ 3}M$

5. $[H\ ]_{Final} \quad \sqrt{K_{a_1} \ c_1 \quad K_{a_2} \ c_2}$

6. $C \quad \frac{0.16/32}{500/1000}M$

$\sqrt{K_b \ c}$

For More Material Join: @JEEAdvanced_2024

7. pOH pK_b $\log \frac{[NH_4Cl]}{[NH_4OH]}$

8. pH pK_a $\log \frac{[Salt]}{[Acid]}$

9. For *WASB* salt

$[OH^-] \sqrt{\frac{K_w \ c}{ka}}$

10. $HX(aq.) \rightleftharpoons H^+(aq.) \ X^-(aq.); K_a \ 10^{-5}$

$H^+(aq.) \ OH^-(aq.) \rightleftharpoons H_2O(l); Kw \ 10^{14}$

Addition

$HX(aq.) \ OH^-(aq.) \rightleftharpoons H_2O(l) \ X^-(aq.)$

$K_{eq.} \ Ka \ Kw$

11. $AgCl(s) \rightleftharpoons \underset{0.2M}{Ag^+(aq)} \ \underset{s}{Cl^-(aq)}$

$K_{sp} \ [Ag^+][Cl^-]$

12. Upon evaporation of saturated solution, all water will be escaped and only dissolved $PbCl_2$ will be left.

i.e., Solubility $\frac{2.78/278}{1} \ 10^{-2}$ mol/L

for $PbCl_2 \ K_{sp} \ 4s^2$

$y \ 10^{-6} \ 4(10^{-2})^3$

$y \ 4$

13. Let's assume simultaneous solubility of $SrCO_3$ as $x\,M$ while SrF_2 as $y\,M$

$SrCO_3(s) \rightleftharpoons Sr^{2+}(aq) \ CO_3^{2-}(aq)$

At eq. $(x \ y)M \ M \quad ; K_{sp_1} \ (x \ y)x$

$SrF_2(s) \rightleftharpoons Sr^{2+}(aq) \ 2F^-(aq) \quad ; 2.5 \ 10^{-10} \ (x \ y)x$ ...(1)

At eq. $(y \ x)M \ 2y\,M \quad ; K_{sp_2} \ (x \ y)(2y)^2$

$10^{-10} \ 4y^2(x \ y)$ ...(2)

$2.5 \ \frac{x}{4y^2}$

$10y^2 \ x \ 10^{-3}$

$y^2 \ 10^{-4}$

$y \ 10^{-2}M \qquad [F^-] \ 2y \ 2 \ 10^{-2}M$

14. $H_2CO_3 \ NaOH \ NaHCO_3 \ H_2O$

$pH_1 \ \frac{1}{2}(pK_{a_1} \ pK_{a_2}) \ \frac{1}{2}(4.6 \ 8) \ 6.3$

$N_1V_1 \ N_2V_2$

$N_1 \ 10 \ 0.1 \ 20$

$N_1 \ 0.2$


$NaHCO_3 \ NaOH \ Na_2CO_3 \ H_2O$

$c \ \frac{2m \text{ mole}}{50 \text{ mL}} \ \frac{1}{25}M$

$pH_2 \ pH_1 \ 10.3 \ 6.3 \ 4$

15. $NaCN; Na_3PO_4; Na_2CO_3; NaHCO_3; Na_2C_2O_4; Na_2HPO_4$

For More Material Join: @JEEAdvanced_2024



7

CHEMICAL KINETICS & NUCLEAR CHEMISTRY

Rate of Appearance or Disappearance of a Substance *C*

(i) Average $\frac{[c]}{t}$ (ii) Instantaneous $\frac{d[c]}{dt}$

❖ **Expressions of the rate:**

For a general reaction: $aA \quad bB \quad cC \quad dD$,

Rate of disappearance of A $\frac{d[A]}{dt}$; Rate of disappearance of B $\frac{d[B]}{dt}$

Rate of appearance of C $\frac{d[C]}{dt}$ & Rate of appearance of D $\frac{d[D]}{dt}$

The positive sign shows that concentrations of C and D increases with time and the negative sign is indicating that concentrations of A and B decrease with time.

Instantaneous rate of reaction : $\frac{1}{a}\frac{d[A]}{dt} \quad \frac{1}{b}\frac{d[B]}{dt} \quad \frac{1}{c}\frac{d[C]}{dt} \quad \frac{1}{d}\frac{d[D]}{dt}$

Average rate of reaction: $\frac{1}{a}\frac{[A]}{t} \quad \frac{1}{b}\frac{[B]}{t} \quad \frac{1}{c}\frac{[C]}{t} \quad \frac{1}{d}\frac{[D]}{t}$

Rate of reaction (ROR) $\frac{\text{Rate of disappearance of reactant}}{\text{Stoichiometry coefficient of reactant}}$

$\frac{\text{Rate of appearance of product}}{\text{Stoichiometry coefficient of product}}$

❖ **Units of Rate of Reaction:** $\text{mol L}^{-1}\text{s}^{-1}$ or $\text{mol L}^{-1}\text{min}^{-1}$ (concentration time $^{-1}$).

For More Material Join: @JEEAdvanced_2024

Rate Law and Rate Constant

Rate $[A]^a.[B]^b$ or Rate $k[A]^a[B]^b$

The constant of proportionality, k is known as the rate constant (specific reaction rate) and may be defined as the rate at unit concentrations of the reactants.

- k depends on the temperature and is independent of the concentration of the reactants.
- At a fixed temperature, k is a constant and is a characteristic of the reaction. Larger value of k indicates fast reaction and small k indicates slow reactions.

Molecularity

It is always a whole number (not zero) and never a fraction. Its value does not exceed 3 & it has no meaning for a complex reaction.

Order of Reaction

$aA \quad bB \qquad cC$










Rate of reaction $= k[A]^m[B]^n$

Order of reaction $m \quad n$ & the order w.r.t. A, B are m, n respectively.

- The order of reaction is obtained from the experimentally determined rate and may be zero, integral or a fraction. In a multi-step complex reaction, the order of the reaction can be determined with the help of slowest step, which is called rate determining step.
- In elementry reaction stoichiometric coefficient of reactants is equal to order of reaction : $m \quad n \quad a \quad b$.

Order	Differential rate law	Integrated rate law	Straight line plot	Half life	Units of k	Example
0	$d[R]/dt \quad k$	$kt \quad [R]_0 \quad [R]$	$[R]_0$ vs t	$[R]_0/2k$	conc time ^{1} or mole L ^{1}s ^{1}	$H_2(g) \quad Cl_2(g) \xrightarrow{h\nu} 2HCl(g)$ $2HI(g) \xrightarrow[\text{surface}]{Au} H_2(g) \quad I_2(g)$
1	$d[R]/dt \quad k[R]$ $kt \quad \ln\{[R]_0/[R]\}$	$[R] \quad [R]_0 e^{\ kt}$	$\ln[R]$ vs t	$(\ln 2)/k$ or $\dfrac{0.693}{k}$	time ^{1} or s ^{1}	Decomposition of H_2O_2 in aqueous solution Radioactive disintegration
2	$d[R]/dt \quad k[R]^2$	$kt \quad \dfrac{1}{[R]} \quad \dfrac{1}{[R]_0}$	$\dfrac{1}{[R]}$ vs t	$\dfrac{1}{k[R]_0}$	mol ^{1}L s ^{1}	$CH_3COOC_2H_5 + NaOH \downarrow CH_3COONa + C_2H_5OH$ $C_2H_4 + H_2 \xrightarrow{100^\circ C} C_2H_6$

For More Material Join: @JEEAdvanced_2024

Important Graphical Representation		
		
		
		

where a_0 Initial Concentration of reactant, r rate of reaction

- Some useful relationships between times for different fractions of reaction of first order to complete

$$t_{3/4} \text{ or } t_{75\%} \quad 2t_{1/2}$$
$$t_{87.5\%} \quad 3t_{1/2}$$
$$t_{99.9\%} \quad 10t_{1/2}$$

- Amount of the substance left after n half-lives $\frac{A_0}{2^n}$

n^{th} Order Reaction

A Product

$$k \; t \quad \frac{1}{n \quad 1} \; \frac{1}{(a \quad x)^{n \; 1}} \quad \frac{1}{a^{n \; 1}} \qquad [n \quad 1, n \quad \text{order}]$$


$$t_{1/2} \quad \frac{1}{k(n \quad 1)} \; \frac{2^{n \; 1} \quad 1}{a^{n \; 1}}$$

- **Reversible reaction** $A \underset{k_b}{\overset{k_f}{\rightleftharpoons}} B$

$$(k_f \quad k_b) \quad \frac{1}{t} \ln \frac{x_{eq}}{x_{eq} \quad x} \quad ; \quad \text{At equilibrium } k_{eq} \quad \frac{k_f}{k_b} \quad \frac{[B]}{[A]} \quad \frac{x_{eq}}{a \quad x_{eq}}$$

For More Material Join: @JEEAdvanced_2024

Parallel/Side Reactions



; $(k_1 \quad k_2)t \quad \ln\frac{[A]_0}{[A]_t}$; $\frac{[B]}{[C]} \quad \frac{k_1}{k_2}$

% Yield of B $\frac{k_1}{k_1 \quad k_2}$ 100 and % Yield of C $\frac{k_2}{k_1 \quad k_2}$ 100

Series/Consecutive Reactions

$A \qquad B \qquad C$

$t_{max} \quad \frac{1}{k_1 \quad k_2}\ln\frac{k_1}{k_2}$

Temperature coefficient ( ) $\frac{k_{T\ 10}}{k_T}$

- For general chemical reactions with rise in temperature by 10°C, the rate constant is nearly doubled.

Arrhenius Equation


$$k \quad Ae^{\ Ea/RT}$$

where k Rate constant, A Pre-exponential factor, T Temperature in kelvin, E_a Activation Energy

- The factor $e^{\ Ea/RT}$ represents fraction of molecules that have kinetic energy greater than E_a
- **Logarithmic expression**

$$\log_{10}\frac{k_2}{k_1} \quad \frac{E_a}{2.303R}\ \frac{1}{T_1} \quad \frac{1}{T_2}$$

- Energy profile for exothermic process & endothermic process:




- Threshold Energy Activation Energy Energy possessed by reactant molecules.

For More Material Join: @JEEAdvanced_2024

❖ **Activation energy diagram for two step endothermic reaction:**

(A) (i) I step-Slow (ii) II step-Fast

(B) (i) Rapid equilibrium (ii) Slow



Factors affecting reaction rates : Concentration of reactants and Reaction temperature. Besides these, presence of catalyst and surface area (if a reactant or a catalyst is a solid) exposure to radiation also affect the reaction rates.

Nuclear Chemistry

Nuclear chemistry is the study of the properties and reactions of atomic nuclei. In nuclear reaction only the nuclides (nuclide : the nucleus of a specific isotope) participates and overall (mass + energy) remains conserved.

Radioactivity

The property of a nucleus emitting radiations like , and is known as radioactivity and the substance possessing the property is called a radioactive substance. The emission of these particles takes place because of the instability of the nucleus. Radioactivity is a property of nucleus.

Properties of , -particles and -rays.

Properties	**Alpha**	**Beta**	**Gamma**
Nature	Fast moving He nuclei (He^{2})	Fast moving electrons	High energy electromagnetic
Representation	$^{4}_{2}He$ or	$^{0}_{1}e$ or $^{0}_{1}$	or $^{0}_{0}$
Charge	2 unit (+ve)	1 unit (–ve)	No charge
Mass	6.65 10 ^{24} g/particle	9.11 10 ^{28} g/particle	0
Velocity	1/10 of light (min.)	nearly velocity of light	Same as light waves (max.)
Relative penetrating power	1	100	10000
Relative ionising power	10000	100	1
Effect of electromagnetic field	Deflected toward –ve electrode	Deflected toward +ve electrode	No effect
Kinetic energy	High	Low	–
Effect on ZnS Screen	Maximum effect	Little effect	–

Types of Radioactive Decay

1. -decay causes decrease of atomic number of 2 units and mass number by 4 units, *e. g.*,

$^{215}_{84}Po$ $^{211}_{82}Pb$ $^{4}_{2}He$; $^{226}_{88}Ra$ $^{226}_{86}Rn$ $^{226}_{86}Rn$ $^{4}_{2}He$

All nucleides with atomic number greater than 83 are beyond the band of stability and are radioactive.

For More Material Join: @JEEAdvanced_2024

2. **-decay** causes increase of atomic number by 1 unit and no change in mass number, *e. g.*,

$^{228}_{88}\text{Ra} \quad ^{228}_{89}\text{Ac} \quad ^{0}_{1}e$; $^{14}_{6}\text{C} \quad ^{14}_{7}\text{N} \quad ^{0}_{1}e$

A neutron is converted to proton in this process.

$^{1}_{0}n \quad ^{1}_{1}p \quad ^{0}_{1}e$

Such emission occurs for the nuclei lying above the stability belt.

3. **-ray** emission ($^{0}_{0}$) causes no change in atomic number and mass number, since it represents the energy lost, *e. g.*,

$^{60m}_{28}\text{Ni} \quad ^{60}_{28}\text{Ni} \quad ^{0}_{0}$

4. **Positron Emission :** Causes decrease the atomic number by 1 unit, *e. g.*,

$^{15}_{8}\text{O} \quad ^{15}_{7}\text{N} \quad ^{0}_{1}e$; $^{118}_{54}\text{Xe} \quad ^{0}_{1}e \quad ^{118}_{53}\text{I}$

A proton is converted to neutron and positron in this process.

$^{1}_{1}p \quad ^{1}_{0}n \quad ^{0}_{1}e$

Such emission occurs for the nuclei lying below the stability belt.

❖ Positron ($^{0}_{1}e$) is a particle having the same mass as an electron, but positively charged.

5. **Electron-capture :**

Causes decrease the atomic number by 1 unit, *e. g.*,

$^{204}_{84}\text{Po} \quad ^{0}_{1}e \quad ^{204}_{83}\text{Bi}$; $^{40}_{19}\text{K} \quad ^{0}_{1}e \quad ^{40}_{18}\text{Ar}$

A proton is converted to neutron in this process.

$^{1}_{1}p \quad ^{0}_{1}e \quad ^{1}_{0}n$

Electron capture occurs with the nuclei lying below the stability belt, in which an electron from the K-shell is captured by the nucleus.

Group Displacement Law (Soddy-Fajans)

In an -particle emission, the resulting element has a mass number less by four units and atomic number less by two units and so lies two places to the left in the periodic table.


In a -particle emission the resulting element has the same mass number but has an atomic number greater by one unit and so lies one place to the right in the periodic table.

For More Material Join: @JEEAdvanced_2024

Neutron/proton ratio and stability zone

- For atomic number < 20, most stable nuclei have *n: p* ratio nearly 1 : 1 (except H & Ar)
- For *n/p* ratio > 1.5, nucleus is unstable. Largest stable nucleus is $^{208}_{82}Pb$ for which *n/p* ratio is 1.53.
- For atomic number > 82, there are no stable nuclei.

Note: Heaviest stable nuclide is $^{208}_{82}Pb$ $t_{1/2}$ for $_{83}Bi$ 1.9 10^{19} yr., which is -emitter reference. Pierre De Marcillae exp det.....
nature, 422, 876-878, 2003



Magic numbers and nuclear stability

Nuclei with 2, 8, 20, 28, 50, 82 or 126 protons or neutrons are unusually stable and have a larger number of stable isotopes than neighboring nuclei in the periodic table. These numbers are called magic numbers. They are supposed to represent completely filled nuclear shells of energy levels.

- Nuclei with magic number of protons as well as neutrons have notably high stabilities. [*e.g.*, $^{4}_{2}He$, $^{16}_{8}O$, $^{40}_{20}Ca$ and $^{208}_{82}Pb$].

Even Odd Theory

Most naturally occurring nuclides have even numbers of neutrons and even numbers of protons. 165 such stable nuclei are known. There exist 55 known nuclei with even number of protons and odd number of neutrons, and 50 known stable nuclei with odd number of protons and even number of neutrons. On the other hand, the number of known stable nuclei having odd numbers of both neutrons and protons is only 5.

Artificial nuclear reactions

The first artificial transmutation was carried out by Rutherford in 1919 who bombarded nitrogen gas with alpha particles and obtained hydrogen and oxygen.

$$^{14}_{7}N \quad ^{4}_{2}He \qquad ^{17}_{8}O \quad ^{1}_{1}H$$

1. Alpha particle induced or (,*n*) type reactions :

$$^{9}_{4}Be \quad ^{4}_{2}He \qquad ^{12}_{6}C \quad ^{1}_{0}n$$

Since -particle is used and a neutron is produced, the reaction may be termed as (,*n*) reaction. In another -bombardment nuclear reaction, the isotope ($^{30}_{15}P$) produced is itself radioactive.

$$^{27}_{13}Al \quad ^{4}_{2}He \qquad ^{30}_{15}P \quad ^{1}_{0}n$$

2. Deuteron-induced or (*D*,) type reaction :

$$^{16}_{8}O \quad ^{2}_{1}H \qquad ^{14}_{7}N \quad ^{4}_{2}He$$

For More Material Join: @JEEAdvanced_2024

3. Proton-induced or (p,) reaction :

$^{14}_{7}\mathrm{N}$ $^{1}_{1}\mathrm{H}$ $^{15}_{8}\mathrm{O}$

4. Neutron-induced or (n,) reaction :

$^{31}_{15}\mathrm{P}$ $^{1}_{0}n$ $^{32}_{15}\mathrm{P}$

Radioactive decay

- Radioactive decay is a first order process. The rate of nuclear decay is determined from measurements of the activity (A) of a sample.

Hence $\frac{dN}{dt}$ N or N $N_0 e^{t}$

where N number of radioactive nuclei at any time t ; N_0 number of radioactive nuclei at t 0; decay constant.

S.I. units : The SI unit is becquerel (Bq).

1 disintegration per second (dps) $1Bq$.

Other units : 1 curie (Ci) 3.7 10^{10} dps, 1 Rutherford (Rd) 10^{6} dps.

- **Specific activity :** Activity per unit mass of radioactive sample (dps/g)
- **Half life ($t_{1/2}$) :** The time taken by half the nuclei (originally present) to decay. $t_{1/2}$ $0.693/$

Note : After n half-lives have passed, activity is reduced to $\frac{1}{2^n}$ of its initial value.

- Average life (t_{av}): t_{av} $1/$ $1.44\,t_{1/2}$

Radioactive equilibrium

Among the members of a decay chain, the state which prevails when the ratios between the activities of successive members remain constant. (This is not an equilibrium in the strict sense since radioactive decay is an irreversible process).

Secular equilibrium

Radioactive equilibrium where the half life of the intermediate isotope is so long that the change of its activity can be ignored during the period of interest and all activities remain constant.

A ^{A} B ^{B} C

Number of nuclei of B is max. at $t_{\max}$; $t_{\max}$ $\frac{1}{(\ _1\ \ _2)}\ln\ \frac{1}{2}$

$\frac{dN_B}{dt}$ $_A N_A$ $_B N_B$.

Secular equilibrium occurs when $\frac{dN_B}{dt}$ 0 or $\frac{N_B}{N_A}$ $\frac{A}{B}$

- Parallel decay A ^{1} B, A ^{2} C

% of B $\frac{1}{1\ \ 2}$ 100

% of C $\frac{2}{1\ \ 2}$ 100

For More Material Join: @JEEAdvanced_2024

Applications of Radioactivity :

- **Age of Minerals and Rocks :** We can determine the age of the rock by measuring the ratio of $^{238}_{92}U$ and $^{206}_{82}Pb$.

$$^{238}_{92}U \qquad ^{206}_{82}Pb \quad 8\,^{4}_{2}He \quad 6\,^{0}_{1}e$$

At time t w gram y gram

Age of Rock, $t = \frac{1}{\lambda}\ln\frac{w_0}{w}$ where $w_0 = w + x$,

w_0 Initial mass of $^{238}_{92}U$; x mass of $^{238}_{92}U$ disintegrated. $x = \frac{238}{206}y$

- **Radiocarbon Dating :** $^{14}_{6}C$ is used to date historical artifacts.

Age of wood artifacts , $t = \frac{1}{\lambda}\ln\frac{A_0}{A}$ (where $t_{1/2}$ of $^{14}_{6}C$ = 5770 years)

A_0 Activity of fresh wood of plant/tree. ; A Activity of old wood.

- Uses in Medicines and other Areas

Nuclear Fission

In a nuclear fission reaction, a heavy nucleus splits up into two main fragments of lighter nuclei and several neutrons.

Of the three natural isotopes of uranium ($^{238}_{92}U$, $^{235}_{92}U$ and $^{234}_{92}U$) the $^{235}_{92}U$ nucleus undergoes nuclear fission when bombarded with slow neutrons. The $^{236}_{92}U$ formed breaks up in several different ways, for example :

$$^{235}_{92}U + {}^{1}_{0}n \rightarrow {}^{236}_{92}U \rightarrow {}^{140}_{56}Ba + {}^{93}_{36}Kr + 3\,{}^{1}_{0}n$$
$$^{235}_{92}U + {}^{1}_{0}n \rightarrow {}^{236}_{92}U \rightarrow {}^{144}_{54}Xe + {}^{90}_{38}Sr + 2\,{}^{1}_{0}n$$
$$^{235}_{92}U + {}^{1}_{0}n \rightarrow {}^{236}_{92}U \rightarrow {}^{144}_{54}Cs + {}^{90}_{37}Rb + 2\,{}^{1}_{0}n$$

- The key to the liberation of energy in the nuclear fission reaction is the production of two or more neutrons per reaction initiated by one neutron.
- Atom bomb is based on nuclear fission.

A nuclear fission chain reaction has three general steps :

1. Initiation : The reaction of a single atom is needed to start the chain reaction Fission of $^{235}_{92}U$ is initiated by the absorption of a neutron.

2. Propagation : This part of the process repeats itself over and over, with each step yielding more product. The fission of $^{236}_{92}U$ releases neutrons that initiate the fission of other uranium atoms.

3. Termination : Eventually, the chain will end. Termination could occur if the reactant ($^{235}_{92}U$) is used up, or if the neutrons that continue the chain escape from the sample without being captured by $^{235}_{92}U$.

Nuclear Fusion

Just as the fission of heavy nuclei is accompanied by mass losses resulting into the liberation of large amounts of energy, the fusion of light nuclei is accompanied by mass losses and the evolution of large quantities of energy. Some such reactions and energy release in each process are shown below,

For More Material Join: @JEEAdvanced_2024

Fusion reaction	Mass loss	Energy released (kJ mol ^{1})
$^{2}_{1}H$ $^{2}_{1}H$ $^{4}_{2}He$	0.026	2.3 10^{9}
$^{2}_{1}H$ $^{3}_{1}H$ $^{4}_{2}He$ $^{1}_{0}n$	0.018	1.79 10^{9}

Compared with fission reactions, fusion reactions have the advantage that large amounts of highly radioactive nuclides are not obtained as by-products which may pose problem of safe storage.

- Very high temperatures is required for a fusion reaction.
- Hydrogen bomb is based on nuclear fusion.
- **Einstein's mass-energy equation :** Energy changes in nuclear reactions can determined by using Einstein's mass-energy equation E mc^{2}

Where E is the energy equivalent of mass m and c is the velocity of light.

For a change of 1 amu (atomic mass unit), the corresponding energy change, E 931 10^{6} eV 931MeV, *i.e.*, a mass of 1 amu is equivalent to 931 MeV of energy.

Mass Defect

It has invariably been found that the actual mass of an isotope of an element is less than the sum of masses of the protons, neutrons and electrons present in it. This difference is called mass defect. The mass defect is nothing, but the loss of mass during the formation of the nucleus of the isotope.

- **Mass defect in nuclear reaction :**

m mass of nuclei of reactants – mass of nuclei of products.

Energy liberated in nuclear reaction : E $m.c^{2}$.

- **Mass defect in an isotope formation :**

Let m_{p}, m_{n} and m_{e} are the respective masses of proton, neutron and electron. Then, the calculated mass of this isotope,

M $Z m_{p}$ Zm_{e} $(A \quad Z)m_{n}$ $Z m_{\mathrm{H}}$ $(A \quad Z)m_{n}$

($\because m_{p}$ m_{e} Mass of hydrogen atom m_{n})

Let M Actual atomic mass as determined experimentally.

then, m Mass defect M M

Binding Energy

Loss of mass during the formation of the nucleus from nucleons is converted into energy. The release of energy imparts stability to the nucleus. The energy released when constituent nucleons combine to form a nucleus, is called binding energy of the nucleus. In other words, energy equal to binding energy will be needed to break up the nucleus into its constituent nucleons. Consequently, the greater the binding energy, the more stable is the nucleus.

Binding Energy and Nuclear Stability

$$\text{B.E. per nucleon} \quad \frac{\text{B.E.}}{\text{No. of nucleons}}$$

Binding energy per nucleon is a direct indicator of its nuclear stability. Higher the binding energy per nucleon of an isotope, greater is its nuclear stability.

For More Material Join: @JEEAdvanced_2024



Level 1

1. The differential rate law equation for the elementary reaction $A \quad 2B \xrightarrow{k} 3C$, is :

(a) $\frac{d[A]}{dt} \quad \frac{d[B]}{dt} \quad \frac{d[C]}{dt} \quad k[A][B]^2$

(b) $\frac{d[A]}{dt} \quad \frac{1}{2}\frac{d[B]}{dt} \quad \frac{1}{3}\frac{d[C]}{dt} \quad k[A]^2[B]$

(c) $\frac{d[A]}{dt} \quad \frac{1}{2}\frac{d[B]}{dt} \quad \frac{1}{3}\frac{d[C]}{dt} \quad k[A][B]^2$

(d) None of these

2. The rate of a reaction is expressed in different ways as follows:

$\frac{1}{2}\frac{d[C]}{dt} \quad \frac{1}{3}\frac{d[D]}{dt} \quad \frac{1}{4}\frac{d[A]}{dt} \quad \frac{d[B]}{dt}$

The reaction is:

(a) $4A \quad B \quad 2C \quad 3D$ (b) $B \quad 3D \quad 4A \quad 2C$

(c) $A \quad B \quad C \quad D$ (d) $B \quad D \quad A \quad C$

3. In the reaction, $A \quad 2B \quad 6C \quad 2D$, if the initial rate $\frac{d[A]}{dt}$ at $t \quad 0$ is $2.6 \quad 10^{\ 2}$M sec $^{\ 1}$, what will be the value of $\frac{d[B]}{dt}$ at $t \quad 0$?

(a) $8.5 \quad 10^{\ 2} M$ sec $^{\ 1}$ (b) $2.5 \quad 10^{\ 2} M$ sec $^{\ 1}$

(c) $5.2 \quad 10^{\ 2} M$ sec $^{\ 1}$ (d) $7.5 \quad 10^{\ 2} M$ sec $^{\ 1}$

4. For the reaction $2A \quad B \quad 3C$; if $\frac{d[A]}{dt} \quad k_1[A]^2$; $\frac{d[B]}{dt} \quad k_2[A]^2$; $\frac{d[C]}{dt} \quad k_3[A]^2$

the correct relation between k_1, k_2 and k_3 is :

(a) $k_1 \quad k_2 \quad k_3$ (b) $2k_1 \quad k_2 \quad 3k_2$ (c) $4k_1 \quad k_2 \quad 3k_2$ (d) $\frac{k_1}{2} \quad k_2 \quad \frac{k_3}{3}$

5. The rate constant of nth order has units :

(a) litre$^{\ 1\ n}$ mol$^{\ 1\ n}$ sec $^{\ 1}$ (b) Mol$^{\ 1\ n}$ litre$^{\ 1\ n}$ sec

(c) Mol$^{\ 1\ n^2}$ litre$^{n^2}$ sec $^{\ 1}$ (d) Mole$^{\ 1\ n}$ litre$^{n\ 1}$ sec $^{\ 1}$

6. Which of the following statement is incorrect?

(a) Unit of rate of disappearance is $Ms^{\ 1}$

(b) Unit of rate of reaction is $Ms^{\ 1}$

(c) Unit of rate constant k depends upon order

(d) Unit of k for first order reaction is $Ms^{\ 1}$

7. Which of the following relation is correct for k_f and k_b in an equilibrium process that contains equal moles of reactants and products.

(a) $k_f \quad k_b$ (b) $k_f \quad k_b$ (c) $k_f \quad k_b$ (d) we cannot predict

8. Listed in the table are forward and reverse rate constants for the reaction

$$2NO(g) \rightleftharpoons N_2(g) + O_2(g)$$

For More Material Join: @JEEAdvanced_2024

Temperature (K)	k_f (M ^{1}s ^{1})	k_b ($M^{-1}s^{-1}$)
1400	0.29	1.1 10 ^{6}
1500	1.3	1.4 10 ^{5}

Select the correct statement :

(a) Reaction is exothermic and value of equilibrium constant (K_{eq}) at 1400 K is 3.79 10 ^{6}

(b) Reaction is endothermic and value of K_{eq} at 1400 K is 2.63 10^{5}

(c) Reaction is exothermic and value of K_{eq} at 1400 K is 2.63 10^{5}

(d) Reaction is endothermic and value of K_{eq} at 1500 K is 9.28 10^{4}

9. The rate constant of a reaction depends on

(a) temperature

(b) pressure

(c) extent of reaction

(d) initial concentration of the reactant

10. In the following reaction, how is the rate of appearance of the underlined product related to the rate of disappearance of the underlined reactant

$$\underline{BrO_3(aq)} \quad 5Br\ (aq) \quad 6H\ (aq) \quad \underline{3Br_2(l)} \quad 3H_2O(aq)$$


(a) $\frac{d[BrO_3^-]}{dt}$ $\frac{d[Br_2]}{dt}$


(b) $\frac{1}{3}\frac{d[BrO_3]}{dt}$ $\frac{d[Br_2]}{dt}$


(c) $\frac{d[BrO_3]}{dt}$ $\frac{1}{3}\frac{d[Br_2]}{dt}$


(d) None of these

11. Consider a reaction $A(g)$ $^{k\ 0.1\,M\ \min\ 1}$ $2B(g)$. If initial concentration of A is 0.5M then select correct graph.

(a)


(b)


(c)


(d)


For More Material Join: @JEEAdvanced_2024

12. Which of the following statements is incorrect?

(a) A second order reaction must be a bimolecular elementary reaction

(b) A bimolecular elementary reaction must be a second order reaction

(c) Zero order reaction must be a complex reaction


(d) First order reaction may be complex or elementary reaction


13. The molecularity of a complex reaction given below is :


$$2N_2O_5(g) \quad 4NO_2(g) \quad O_2(g)$$


(a) 1 (b) 2 (c) 3 (d) has no meaning

14. Decomposition of NH_4NO_2 (aq) into N_2 (g) and $2H_2O$ (l) is first order reaction. Which of the following graph is correct?


(a) 


(b) 


(c) 


(d) 

15. Decomposition of HI (g) on Gold surface is zero order reaction. Initially few moles of H_2 are present in container then which of the following graph is correct?

(a) 


(b) 

(c) 

(d) 

16. Consider the plots for the types of reaction


$$nA \quad B \quad C$$




These plots respectively correspond to the reaction orders :

(a) 0, 2, 1 (b) 0, 1, 2 (c) 1, 1, 2 (d) 1,0, 2


17. If decomposition reaction $A(g)$ $B(g)$ follows first order kinetics then the graph of rate of formation (R) of B against time t will be:

(a) 

(b) 


For More Material Join: @JEEAdvanced_2024

(c) 

(d) 

18. Consider the plots, given below, for the types of reaction

$nA \quad B \quad C$



These plots respectively correspond to the reaction orders :


(a) 0, 1, 2 (b) 1, 2, 0 (c) 1, 0, 2 (d) None of these

19. For a zero order reaction, the plot of conc. ($a - x$) vs time is linear with

(a) ve slope and zero intercept (b) ve slope and zero intercept

(c) ve slope and non-zero intercept (d) – ve slope and non-zero intercept

20. What will be the order of reaction for a chemical change having $\log t_{1/2}$ *vs* $\log a$? (where a = initial concentration of reactant; $t_{1/2}$ half-life)




(a) Zero order

(b) First order

(c) Second order

(d) None of these


21. A graph between $\log t_{1/2}$ and log a (abscissa), a being the initial concentration of A in the reaction For reaction A Product, the rate law is :



(a) $\frac{d[A]}{dt} \quad K$ (b) $\frac{d[A]}{dt} \quad K[A]$


(c) $\frac{d[A]}{dt} \quad K[A]^2$ (d) $\frac{d[A]}{dt} \quad K[A]^3$


22. For the reaction A B, for which graph between half life ($t_{1/2}$) and initial concentration (a) of the reactant is as given below





Hence graph between $\frac{d[A]}{dt}$ and time will be:

For More Material Join: @JEEAdvanced_2024

(a) 

(b) 

(c) 

(d) 

23. For the ideal gaseous reaction, the rate is generally expressed in terms of $\frac{dP}{dt}$ instead of $\frac{dC}{dt}$ or $\frac{dn}{dt}$ (where C $\frac{n}{V}$ is concentration and n the no. of moles). What is the relation among these three expressions if T and V are constant?

(a) $\frac{dC}{dt}$ $\frac{dn}{dt}$ $\frac{dP}{dt}$ (b) $\frac{dC}{dt}$ $\frac{1}{V}\frac{dn}{dt}$ $\frac{1}{RT}\frac{dP}{dt}$

(c) $RT\frac{dC}{dt}$ $\frac{dn}{dt}$ $\frac{dP}{dt}$ (d) None of these

24. A_2 B_2 $2AB$; R.O.R. $k[A_2]^a[B_2]^b$

Initial $[A_2]$	Initial $[B_2]$	R.O.R. (r) Ms ^{1}
0.2	0.2	0.04
0.1	0.4	0.04
0.2	0.4	0.08

Order of reaction with respect to A_2 and B_2 are respectively :

(a) a $1, b$ 1 (b) a $2, b$ 0 (c) a $2, b$ 1 (d) None

25. For a reaction the initial rate is given as: R_0 $k[A_0]^2[B_0]$. By what factor, the initial rate of reaction will increase if initial concentration of A is taken 1.5 times and of B is tripled?

(a) 4.5 (b) 2.25 (c) 6.75 (d) None of these

26. For $A_{(s)}$ $B_{(s)}$ $C_{(s)}$; rate $k[A]^{1/2}[B]^2$, if initial concentration of A and B are increased by factors 4 and 2 respectively, then the initial rate is changed by the factor:

(a) 4 (b) 6 (c) 8 (d) None of these

27. Reaction A B follows second order kinetics. Doubling the concentration of A will increase the rate of formation of B by a factor of :

(a) 1/4 (b) 1/2 (c) 2 (d) 4

For More Material Join: @JEEAdvanced_2024

28. The reaction of A_2 and B_2 follows the equation

$$A_2(g) \quad B_2(g) \qquad 2AB(g)$$

The following data were observed

$[A_2]_0$	$[B_2]_0$	Initial rate of appearance of $AB(g)$ (in Ms^{-1})
0.10	0.10	2.5 10^{-4}
0.20	0.10	5 10^{-4}
0.20	0.20	10 10^{-4}

The value of rate constant for the above reaction is:

(a) 2.5 10^{-4} (b) 2.5 10^{-2} (c) 1.25 10^{-2} (d) None of these

29. Above given question, the value of rate constant for appearance of $AB(g)$ is :

(a) 2.5 10^{-4} (b) 2.5 10^{-2} (c) 1.25 10^{-2} (d) None of these

30. The following data pertain to reaction between A and B

S. No	[A] mol L^{-1}	[B] mol L^{-1}	Rate mol L^{-1} sec^{-1}
I	1 10^{-2}	2 10^{-2}	2 10^{-4}
II	2 10^{-2}	2 10^{-2}	4 10^{-4}
III	2 10^{-2}	4 10^{-2}	8 10^{-4}

Which of the following inference(s) can be drawn from the above data

(a) Rate constant of the reaction 10^{-4}

(b) Rate law of the reaction is $k[A][B]$

(c) Rate of reaction increases four times on doubling the concentration of both the reactant, Select the correct answer

codes

(a) a, b and c (b) a and b (c) b and c (d) c alone

31. The unit of rate constant of elementary reaction depends upon the :

(a) temperature of the reaction (b) concentration of reactants

(c) activation energy of the reaction (d) molecularity of the reaction

32. Select the rate law that corresponds to the data shown for the reaction $A \quad B \qquad C$

Exp.	[A]	[B]	Rate
1.	0.012	0.035	0.10
2.	0.024	0.070	0.80
3.	0.024	0.035	0.10
4.	0.012	0.070	0.80

(a) Rate $k[B]^3$ (b) Rate $k[B]^4$ (c) Rate $k \quad [A][B]^3$ (d) Rate $k[A]^2[B]^2$

33. An elementary reaction between A and B is a second order reaction. Which of the following rate equations must be correct?

(a) $r \quad k[A]^2[B]^0$ (b) $r \quad k[A]^{3/2}[B]^{1/2}$ (c) $r \quad k[A]^0[B]^2$ (d) $r \quad k[A][B]$

For More Material Join: @JEEAdvanced_2024

34. If 'a' is the initial concentration of the reactant, the half-life period of the reaction of nth order is inversely proportional to :
(a) a^{n-1} (b) a^{n} (c) a^{1-n} (d) a^{n-1}

35. Which of the following expressions is correct for zero order and first order respectively [where a is initial concentration]?
(a) $t_{1/2} \quad a; t_{1/2} \quad \frac{1}{a}$ (b) $t_{1/2} \quad a; t_{1/2} \quad a^{0}$
(c) $t_{1/2} \quad a^{0}; t_{1/2} \quad a$ (d) $t_{1/2} \quad a; t_{1/2} \quad \frac{1}{a^{2}}$

36. The unit of rate constant of zero order and first order chemical reactions are respectively :
(a) $\text{mol L}^{-1}\text{ s}^{-1}, \text{mol L}^{-1}\text{ s}^{-1}$ (b) $\text{s}^{-1}, \text{mol L}^{-1}\text{ s}^{-1}$
(c) $\text{mol L}^{-1}\text{ s}^{-1}, \text{s}^{-1}$ (d) None of these

37. The units of rate of reaction and rate constant are same for a :
(a) zero order reaction (b) first order reaction
(c) second order reaction (d) third order reaction

38. $CH_3COOC_2H_5(aq) \quad H_2O(l) \xrightarrow{H^{+}(aq)} CH_3COOH(aq) \quad C_2H_5OH(aq)$. What type of reaction is this?
(a) Unimolecular elementary (b) Pseudo first order
(c) Zero order (d) Second order

39. When ethyl acetate was hydrolysed in presence of 0.1 M HCl, the rate constant was found to be $5.4 \quad 10^{-5}\text{ s}^{-1}$. But in presence of 0.1 M H_2SO_4 the rate constant was found to be $6.25 \quad 10^{-5}\text{ s}^{-1}$. Thus it may be concluded that :
(a) H_2SO_4 furnishes more H than HCl
(b) H_2SO_4 furnishes less H than HCl
(c) both have the same strength
(d) will depend on concentration of ethyl acetate

40. For an elementary reaction $2A \quad B \quad A_2B$ if the volume of vessel is quickly reduced to half of its original volume then rate or reaction will
(a) remain unchanged (b) increase four times
(c) increase eight times (d) decrease eight times

41. In the reaction $A \quad B \quad C$, rate constant is $0.001\ M\text{s}^{-1}$. If we start with 1 M of A then conc. of A and B after 10 minutes are respectively :
(a) 0.5 M, 0.5 M (b) 0.6 M, 0.4 M (c) 0.4 M, 0.6 M (d) none of these

42. For a reaction $A \xrightarrow{k_r \ 0.6M\text{ min}^{-1}} 2B$
starting with 1 M of 'A' only, concentration of B (in M) after 100 sec. and 200 sec. is respectively ?
(a) 2 and 4 (b) 1 and 2 (c) 2 and 3 (d) None of these

43. Half-life ($t_{1/2}$) and completion time (T) of the above reaction ($A \quad 2B$) are :
(a) 500 min, 750 min (b) 500 sec, 750 sec (c) 50 sec, 100 sec (d) None of these

For More Material Join: @JEEAdvanced_2024

44.

Column I	Column II
P. Zero order reaction	**1.** $t_{1/2}$ $\frac{1}{[A]_0}$
Q. First order reaction	**2.** $t_{100\%}$ $[A]_0 / k$
R. Second order reactions	**3.** Involves at least two reactants
S. Pseudo unimolecular reaction	**4.** $[A]$ $[A]_0 e^{kt}$

Code:

	P	Q	R	S
(a)	2	1	4	2
(b)	2	4	1	3
(c)	2	1	3	4
(d)	3	2	1	4

45. For the zero order reaction A B C; initial concentration of A is 0.1 M. If $[A]$ 0.08 M after 10 minutes, then its half-life and completion time are respectively :

(a) 10 min; 20 min (b) 2 10^{3} min; 4 10^{3} min

(c) 25 min, 50 min (d) 250 min, 500 min

46. For an elementary reaction, $X(g)$ $Y(g)$ $Z(g)$

the half life period is 10 min. In what period of time would the concentration of X be reduced to 10% of original concentration ?

(a) 20 Min. (b) 33 Min (c) 15 Min (d) 25 Min

47. In the presence of an acid, the initial concentration of cane sugar was reduced from 0.20 to 0.10 molar in 5 hours and from 0.2 to 0.05 molar in 10 hours. The reaction is of :

(a) Zero order (b) First order (c) Second order (d) Third order

48. A first order reaction is 75% completed in 100 minutes. How long time will it take for its 87.5% completion?

(a) 125 min (b) 150 min (c) 175 min (d) 200 min

49. The rate constant for a first order reaction whose half life is 480 sec, is :

(a) 1.44 10^{3} sec^{1} (b) 1.44 sec^{1} (c) 0.72 10^{3} sec^{1} (d) 2.88 10^{3} sec^{1}

50. Rate constant k 2.303 min^{1} for a particular reaction. The initial concentration of the reactant is 1 mol/litre then rate of reaction after 1 minute is :

(a) 2.303 M min^{1} (b) 0.2303 M min^{1} (c) 0.1 M min^{1} (d) none of these

51. For the reaction $3A(g)$ $\overset{k}{}$ $B(g)$ $C(g)$, k is 10^{4} L/mol.min.

If $[A]$ 0.5M then the value of $\frac{d[A]}{dt}$ (in Ms^{1}) is:

(a) 7.5 10^{5} (b) 3 10^{4} (c) 2.5 10^{5} (d) none of these

52. 99% of a first order reaction was completed in 32 minutes when 99.9% of the reaction will complete :

(a) 50 min (b) 46 min (c) 48 min (d) 49 min

For More Material Join: @JEEAdvanced_2024

53. Which of the following represents the expression for $\frac{3}{4}$th life of first order reaction

(a) $\frac{k}{2.303}\log 4/3$ (b) $\frac{2.303}{k}\log 3/4$ (c) $\frac{2.303}{k}\log 4$ (d) $\frac{2.303}{k}\log 3$

54. Consider the following first order competing reactions:

$X \xrightarrow{k_1} A \quad B$ and $Y \xrightarrow{k_2} C \quad D$


If 50% of the reaction of X was completed when 96% of the reaction of Y was completed, the ratio of their rate constants (k_2 / k_1) is:

(a) 4.06 (b) 0.215 (c) 1.1 (d) 4.65

55. For the reactions

(i) $A \xrightarrow{k_I} P$

(ii) $B \xrightarrow{K_{II}} Q$, following observation is made.



Calculate $\frac{k_I}{K_{II}}$, where k_I and k_{II} and rate constant for the respective reaction.

(a) 2.303 (b) 1 (c) 0.36 (d) 0.693

56. The decomposition of N_2O_5 in carbon tetrachloride was followed by measuring the volume of O_2 gas evolved : $2N_2O_5\,(CCl_4) \quad 2N_2O_4\,(CCl_4) + O_2(g)$. The maximum volume of O_2 gas obtained was 100 cm^3. In 500 minutes, 90 cm^3 of O_2 were evolved. The first order rate constant (in min^{-1}) for the disappearance of N_2O_5 is :

(a) $\frac{2.303}{500}$ (b) $\frac{2.303}{500}\log\frac{100}{90}$ (c) $\frac{2.303}{500}\log\frac{90}{100}$ (d) $\frac{100}{10 \quad 500}$

57. For a first order homogeneous gaseous reaction $A \quad 3B$, if pressure after time t was P_T and after completion of reaction, pressure was P then select correct relation

(a) $k \quad \frac{1}{t}\ln\frac{P}{3(P \quad P_t)}$ (b) $k \quad \frac{1}{t}\ln\frac{2P}{(P \quad P_T)}$

(c) $k \quad \frac{1}{t}\ln\frac{3P}{2P \quad P_t}$ (d) $k \quad \frac{1}{t}\ln\frac{2P}{3(P \quad P_T)}$

58. The half-life of first order decomposition of NH_4NO_3 is 2.10 hr at 288 K temperature

$NH_4NO_3(aq) \quad N_2O(g) \quad 2H_2O(l)$. If 6.2 g of NH_4NO_3 is allowed to decompose, the time required for NH_4NO_3 to decompose 90% is :

(a) 6.978 hr (b) 0.319 (c) 0.319 hr (d) None of these

For More Material Join: @JEEAdvanced_2024

59. For a first order homogeneous gaseous reaction, $A \quad 2B \quad C$ then initial pressure was P_i while total pressure after time 't' was P_t. The right expression for the rate constants k in terms of P_i, P_t and t is :

(a) $k \quad \frac{2.303}{t} \log \quad \frac{2P_i}{3P_i \quad P_t}$ (b) $k \quad \frac{2.303}{t} \log \quad \frac{2P_i}{2P_t \quad P_i}$

(c) $k \quad \frac{2.303}{t} \log \quad \frac{P_i}{P_i \quad P_t}$ (d) none of these

60. The decomposition of azo methane, at certain temperature according to the equation $(CH_3)_2N_2 \quad C_2H_6 + N_2$ is a first order reaction.
After 40 minutes from the start, the total pressure developed is found to be 350 mm Hg in place of initial pressure 200 mm Hg of azo methane. The value of rate constant k is :

(a) $2.88 \quad 10^{\ 4}\text{sec}^{\ 1}$ (b) $1.25 \quad 10^{\ 4}\text{sec}^{\ 1}$ (c) $5.77 \quad 10^{\ 4}\text{sec}^{\ 1}$ (d) None of these

61. The hydrolysis of sucrose was studied with the help of polarimeter and following data were collected time (min.) : 0 70
observed rotation (degrees) : 44 16.5 –11
The time taken when the reaction mixture will be optically inactive ? (Given: ln 2=0.7, ln 3=1.1, ln 5=1.6)

(a) 16 min. (b) 69.47 min. (c) 160 min. (d) none of these

62. For a particular reaction with initial conc. of the reactants as a_1 and a_2, the half-life period are t_1 and t_2 respectively. The order of the reaction (n) is given by :

(a) $n \quad 1 \quad \frac{\log (t_2/t_1)}{\log (a_2/a_1)}$ (b) $n \quad \frac{\log (t_1/t_2)}{\log (a_2/a_1)}$

(c) $n \quad 1 \quad \frac{\log (t_1/t_2)}{\log (a_2/a_1)}$ (d) none of these

63. The value of $\frac{t_{0.875}}{t_{0.50}}$ for n^{th} order reaction is

(a) $2^{(2n \ 2)}$ (b) $2^{(2n \ 2) \ 1}$ (c) $\frac{8^{n \ 1} \quad 1}{2^{n \ 1} \quad 1}$ (d) None of these

64. $A \quad B$ first order reaction A is optical active and B is optically inactive. A series of experiment were conducted on a solution of A

Time	0	60 min
optical rotation	82	77 2

Assume some impurity is present calculate the optical rotation after 5 hours.
(Given ln 1.066 0.064, $e^{0.16}$ 1.17)

(a) 60 (b) 30 (c) 20 (d) 120

65. At 300 K the half-life of a sample of a gaseous compound initially at 1 atm is 100 sec. When the pressure is 0.5 atm the half-life is 50 sec. The order of reaction is :

(a) 0 (b) 1 (c) 2 (d) 3

For More Material Join: @JEEAdvanced_2024

66. A decomposes as $A(g) \xrightarrow{k_1 \quad 2 \quad 10^{-3}\ s^{-1}} 2B(g)$ and $A(g) \xrightarrow{k_2 \quad 1 \quad 10^{-3}\ s^{-1}} C(g)$. The rate of appearance of B, taking 2 M concentration of A, is equal to :

(a) $2 \quad 10^{-3}\ M\ s^{-1}$ (b) $4 \quad 10^{-3}\ M\ s^{-1}$ (c) $8 \quad 10^{-3}\ M\ s^{-1}$ (d) None of these

67. For an endothermic reaction where H represents the enthalpy of reaction in kJ/mol, the minimum value for the energy of activation will be :

(a) less than H (b) more than H (c) equal to H (d) zero

68. The activation energy of the reaction, $A \quad B \quad C \quad D$ 38 kcal is 20 kcal. What would be the activation energy of the following reaction.

$C \quad D \quad A \quad B$

(a) 20 kcal (b) –20 kcal (c) 18 kcal (d) 58 kcal


69. When the activation energies of the forward and backward reactions are equal, then:


(a) $U \quad 0, \ S \quad 0$ (b) $U \quad 0; \ G \quad 0$ (c) $S \quad 0; \ G \quad 0$ (d) only $U \quad 0$


70. For an exothermic chemical process occurring in two steps as follows


(i) $A \quad B \quad X$ (slow) (ii) $X \quad AB$ (fast)

The progress of reaction can be best described by:


(a)



(b)



(c)



(d)


71. Select the correct diagram for an endothermic reaction that proceeds through two steps, with the second step is rate determining :

(a)


(b)


(c)


(d)


For More Material Join: @JEEAdvanced_2024

72. $\frac{k_{35}}{k_{34}}$ 1, this means that

(a) rate increases with the rise in temperature
(b) rate decreases with rise in temperature
(c) rate does not change with rise in temperature
(d) None of the above

73. The plot of ln k *versus* $1/T$ is linear with slope of:

(a) E_a/R (b) E_a/R (c) $E_a/2.303\ R$ (d) $E_a/2.303\ R$

74. Rate constant for a chemical reaction taking place at 500K is expressed as k $A.\ e^{1000}$ The activation energy of the reaction is:

(a) 100 cal/mol (b) 1000 kcal/mol (c) 10^4 kcal/mol (d) 10^6 kcal/mol

75. For a complex reaction $A \xrightarrow{k}$ products

E_{a_1} 180 kJ/mol; E_{a_2} 80 kJ/mol; E_{a_3} 50 kJ/mol

Overall rate constant k is related to individual rate constant by the equation k $\left(\frac{k_1 . k_2}{k_3}\right)^{2/3}$.

Activation energy (kJ/mol) for the overall reaction is :

(a) 100 (b) 43.44 (c) 150 (d) 140

76. For reaction A B, the rate constant k_1 $A_1(e^{E_{a_1}/RT})$ and for the reaction X Y, the rate constant k_2 $A_2(e^{E_{a_2}/RT})$. If A_1 10^9, A_2 10^{10} and E_{a_1} 1200 cal/mol and E_{a_2} 1800 cal/mol, then the temperature at which k_1 k_2 is: (Given; R 2 cal/K-mol)

(a) 300K (b) 300×2.303K (c) $\frac{300}{2.303}K$ (d) None of these

77. The activation energies of the forward and backward reactions in the case of a chemical reaction are 30.5 and 45.4 kJ/mol respectively. The reaction is :

(a) exothermic (b) endothermic
(c) neither exothermic nor endothermic (d) independent of temperature

78. A reaction rate constant is given by : k 1.2 $10^{14} e^{\frac{25000}{RT}}$ sec ^{1}. It means :


(a) log k *versus* log T will give a straight line with a slope as 25000
(b) log k *versus* log T will give a straight line with a slope as –25000
(c) log k *versus* T will give a straight line with a slope as –25000
(d) log k *versus* $1/T$ will give a straight line


79. The temperature coefficient of a reaction is :


(a) the rate constant
(b) the rate constant at a fixed temperature
(c) the ratio of rate constant at two temperature
(d) the ratio of rate constant differing by 10°C preferably k_{308}/k_{298}


For More Material Join: @JEEAdvanced_2024

80. Which graph shows zero activation energy?


(a) 

(b) 

(c) 

(d) 

81. A homogeneous catalytic reaction takes place through the three alternative plots *A, B* and *C* shown in the given figure. Which one of the following indicates the relative ease with which the reaction can take place ?



(a) *A B C* (b) *C B A* (c) *A C B* (d) *A B C*

82. A first order reaction is 50% completed in 20 minutes at 27°C and in 5 minutes at 47°C. The energy of activation of the reaction is :

(a) 43.85 kJ/mol (b) 55.14 kJ/mol (c) 11.97 kJ/mol (d) 6.65 kJ/mol

83. The rate of a reaction gets doubled when temperature increases by 27 C to 37°C. By what factor will it change for the temperature range 17 C to 27 C.

(a) 1.81 (b) 1.71 (c) 2.1 (d) 2.41

84. Which of the following explains the increase of the reaction rate by catalyst ?

(a) Catalyst decreases the rate of backward reaction so that the rate of forward reaction increases

(b) Catalyst provides extra energy to reacting molecules so that they may reduce effective collisions

(c) Catalyst provides an alternative path of lower activation energy to the reactants

(d) Catalyst increases the number of collisions between the reacting molecules.

85. Collision theory is satisfactory for:

(a) First order reactions (b) Zero order reactions

(c) Bimolecular elementary reactions (d) Any order reactions

86. For the first order reaction *A* *B* *C*, carried out at 27°C. If 3.8×10^{16}% of the reactant molecules exists in the activated state, the E_a (activation energy) of the reaction is :

(a) 12 kJ/mol (b) 831.4 kJ/mol (c) 100 kJ/mol (d) 88.57 kJ/mol

87. A catalyst lowers the activation energy for a certain reaction from 83.314 to 75 kJ mol^{1} at 500 K. What will be the rate of reaction as compared to uncatalysed reaction? Assume other things being equal.

For More Material Join: @JEEAdvanced_2024

(a) Double (b) 28 times (c) 7.38 times (d) 7.38 10^3 times

88. A following mechanism has been proposed for a reaction

$2A \quad B \quad D \quad E$

$A \quad B \quad C \quad D$ (slow)

$A \quad C \quad E$ (fast)

The rate law expression for the reaction by RDS method is:

(a) $r \quad k[A]^2[B]$ (b) $r \quad k[A][B]$ (c) $r \quad k[A]^2$ (d) $r \quad k[A][C]$

89. A hypothetical reaction $A_2 \quad B_2 \quad 2AB$ follows the mechanism as given below

$A_2 \rightleftharpoons A \quad A$ (fast)

$A \quad B_2 \quad AB \quad B$ (slow)

$A \quad B \quad AB$ (fast)

The order of the over all reaction by RDS method is

(a) 2 (b) 1 (c) $\frac{3}{2}$ (d) 0

90. Chemical reaction occurs as a result of collisions between reacting molecules. Therefore, the reaction rate is given by

(a) total number of collisions occurring in a unit volume per second

(b) fraction of molecules which possess energy less than the threshold energy

(c) total number of effective collisions which have enough activation energy

(d) none of the above

91. Radioactivity is affected by :

(a) temperature (b) pressure

(c) electric and magnetic field (d) none of these

92. The radiation from naturally occurring radioactive substance as seen after deflection by a magnetic field in one direction are :

(a) -rays (b) -rays

(c) both and rays (d) either or -rays

93. In the radioactive decay

$_ZX^A \quad _{Z\ 1}Y^A \quad \underset{\text{high energy}}{_{Z\ 1}Z^{A\ 4}} \quad \underset{\text{low energy}}{_{Z\ 1}Z^{A\ 4}}$ the sequence of the radiation emitted is :

(a) , , (b) , ,

(c) , , (d) , ,

94. A radioactive nuclide emitts -rays due to the :

(a) emission of an electron from its orbital

(b) nuclear energy transition from a higher state to a lower state

(c) presence of less neutrons than protons

(d) presence of more neutrons than protons

95. Consider the following decay $^A_ZX \quad _Z{}^A_1Y \quad {}^0_1e, X$ is unstable because :

(a) its nucleus has excess energy (b) $\frac{n}{p}$ ratio is high

(c) $\frac{n}{p}$ ratio is low (d) none of these

For More Material Join: @JEEAdvanced_2024

96. Consider the following decay ${}^{A}_{Z}X \quad {}_{Z}\ {}^{A}_{1}Y \quad {}^{0}_{1}e(\quad)\ X$ is unstable because :

(a) its nucleus has excess energy (b) $\frac{n}{p}$ ratio is high

(c) $\frac{n}{p}$ ratio is low (d) none of these

97. During -decay :

(a) $\frac{n}{p}$ ratio decreases (b) $\frac{n}{p}$ ratio increases

(c) $\frac{n}{p}$ ratio remains constant (d) $\frac{n}{p}$ ratio may increase or decrease

98. Which of the following processes causes the emission of X-ray?

(a) -emission (b) -emission

(c) (Positron) emission (d) electron capture

99. Which of the following processes result in an increase in the atomic number of a nuclide?

(a) -emission (b) electron capture (c) -emission (d) -(Beta)emission

100. ..... is produced when a positron and an electron collide.

(a) X-ray (b) Neutron (c) -radiation (d) Neutrino

101. ${}^{165}_{67}$Ho is stable isotope. ${}^{150}_{67}$Ho is expected to disintegrate by:

(a) -emission (b) -emission (c) positron emission (d) -emission

102. ${}^{1}_{1}$H is a stable isotope. ${}^{3}_{1}$H is expected to disintegrated by :

(a) -emission (b) -emission (c) positron emission (d) proton emission

103. Emission of -particle is equivalent to :

(a) increase of one proton only (b) decrease of one neutron only

(c) both (a) and (b) (d) none of these

104. Atoms ${}^{A}_{7}X$, ${}^{B}_{8}Y$ and ${}^{17}_{9}Z$ are such that ${}_{8}Y$ is an isobar of ${}_{7}X$ and atom ${}^{17}_{9}Z$ is isotone of ${}_{8}Y$. Mass no. of X and no. of neutrons in Y are respectively :

(a) 8, 8 (b) 17, 7 (c) 9, 8 (d) 16, 8

105. ${}^{234}_{90}$Th disintegrates to give ${}^{206}_{82}$Pb as the final product. Total no. of and particles emitted out during this process are :

(a) 6 (b) 7 (c) 8 (d) 13

106. An isotone of ${}^{76}_{32}$Ge is :

(a) ${}^{77}_{32}$Ge (b) ${}^{77}_{33}$As (c) ${}^{77}_{34}$Se (d) ${}^{77}_{36}$Se

107. Pair of isobar is :

(a) ${}^{13}_{6}$C, ${}^{13}_{7}$N (b) ${}^{13}_{6}$C, ${}^{14}_{7}$N (c) ${}^{14}_{6}$C; ${}^{15}_{8}$N (d) none of these

108. Isodiaphers are atoms having:

(a) n/p same (b) p/n same (c) (n p) same (d) (n p) different

109. The 'Group displacement law' was given by:

(a) Bacqueral (b) Rutherford (c) Madam Curie (d) Soddy and Fajan

110. ${}^{7}_{3}$Li ${}^{1}_{1}p$ X; Identify X if reaction is (p,) type.

(a) ${}^{8}_{4}$Be (b) ${}^{4}_{2}$He (c) ${}^{0}_{0}$ (d) none of these

For More Material Join: @JEEAdvanced_2024

111. Identify reaction type:

$$^{27}_{13}\text{Al} \quad ^{2}_{1}\text{H} \qquad ^{28}_{13}\text{Al} \quad ^{1}_{1}\text{H}$$

(a) (d, p) (b) (p, p) (c) (p, d) (d) none of these

112. $^{27}_{13}\text{Al} \quad ^{1}_{1}p \qquad X \quad ^{0}_{0}$; Identify X if reaction is $(p, \quad)$ type artificial radioactive reaction.

(a) $^{28}_{13}\text{Al}$ (b) $^{27}_{14}\text{Si}$ (c) $^{28}_{14}\text{Si}$ (d) none of these

113. The number of neutrons accompanying in the formation of $^{139}_{54}\text{Xe}$ and $^{94}_{38}\text{Sr}$ from the absorption of slow neutron by $^{235}_{92}\text{U}$ followed by nuclear fission is :

(a) 0 (b) 1 (c) 2 (d) 3

114. What will be the product of reaction $^{255}_{101}\text{Md}(\quad, 2n)$?

(a) $^{256}_{103}\text{Lr}$ (b) $^{257}_{102}\text{No}$ (c) $^{257}_{103}\text{Lr}$ (d) $^{205}_{82}\text{Pb}$

115. Complete the following nuclear equation by supplying the symbol for the other product of the fission :

$$^{235}_{92}\text{U} \quad ^{1}_{0}n \qquad ^{94}_{38}\text{Sr} \quad \ldots\ldots\ldots \quad 2\,^{1}_{0}n$$

(a) $^{139}_{54}\text{Xe}$ (b) $^{140}_{54}\text{Xe}$ (c) $^{104}_{64}\text{Gd}$ (d) none of these

116. $^{235}_{92}\text{U} \quad ^{1}_{0}n \qquad ^{139}_{56}\text{Ba} \quad ^{94}_{36}\text{Kr} \quad 3\,^{1}_{0}n \quad 200\text{MeV}$

Total energy released (in MeV) after 5th stage of fission is :

(a) 48600 (b) 16200 (c) 24200 (d) None of these

117. Proton bombardment on Th^{230} followed by emission of two alpha particles will produce :

(a) Rn^{232} (b) Ra^{233} (c) Fr^{223} (d) Fr^{222}

118. $X \qquad ^{206}_{82}\text{Pb} \quad ^{4}_{2}\text{He}$. In this reaction predict the position of group of X :

(a) II B (b) IV B (c) VI A (d) VI B

119. $_{90}\text{Th}$ is a member of III group. After losing -particle it forms a new element belonging to :

(a) I group (b) II group (c) III group (d) IV group

120. Alpha decay of $^{238}_{92}\text{U}$ forms $^{234}_{90}\text{Th}$. What kind of decay from $^{234}_{90}\text{Th}$ produces $^{234}_{89}\text{Ac}$?

(a) (b) (c) (positron) (d) -emission

121. $^{214}_{83}\text{Bi}$ decays to A by -emission. A then decays to B by beta emission, which further decays to C by another beta emission. Element C decays to D by still another beta emission, and D decays by -emission to form a stable isotope E. What is element E?

(a) $^{207}_{81}\text{Tl}$ (b) $^{206}_{80}\text{Hg}$ (c) $^{206}_{79}\text{Au}$ (d) $^{206}_{82}\text{Pb}$

122. The activity of a radioactive nuclide (X^{100}) is 6.023 curie at a certain time 't'. If its disintegration constant is 3.7 10^{4} s ^{1} the mass of X after t sec is :

(a) 6.022 10^{6} g (b) 10 ^{13} g (c) 10 ^{15} g (d) 10 ^{17} g

123. Activity of a radioactive substance is A_1 at time t_1 and A_2 at time $t_2 (t_2 \quad t_1)$, then the ratio of $\frac{A_2}{A_1}$ is :

(a) $e^{(t_2 \quad t_1)}$ (b) $e^{(t_1 \quad t_2)}$ (c) $e^{(t_1 \quad t_2)}$ (d) $\frac{t_2}{t_1}$

For More Material Join: @JEEAdvanced_2024

124. The half-life of $^{14}_{6}C$ is 5730 year. What fraction of its original C^{14} would left after 22920 year of storage?
(a) 0.50 (b) 0.25 (c) 0.125 (d) 0.0625

125. A radioactive sample had an initial activity of 56 dpm. After 69.3 minutes, it was found to have an activity of 28 dpm. Find the number of atoms in a sample having an activity of 100 dpm.
(a) 693 (b) 100 (c) 1000 (d) 10,000

126. A radioactive sample has initial activity of 28 dpm 30 minutes later its activity 14 dpm. How many atoms of nuclide were present initially?
(a) 2800 (b) 1212 (c) 528 (d) 2802

127. The value of decay constant of Co^{60} is 2.5 10^{7} min ^{1}. The activity of 2.0 g of the sample is nearly :
(a) 5 10^{5} dpm (b) 2.5 10^{10} dpm (c) 5 10^{15} dpm (d) 10^{10} dpm

128. Half-life ($t_{1/2}$) for a radioactive decay is 6930 sec. The time required to fall the rate of decay to $\frac{1}{100}^{\text{th}}$ of its initial value is :
(a) 69.3 sec (b) 20,000 sec (c) 46060 sec (d) none of these

129. A sample of radioactive substance is found 90% of its initial amount after one day. What % of the original sample can be found after 3 days?
(a) 81 (b) 72.9 (c) 25 (d) 65.61

130. If time t is required for a radioactive substance to become one third of its initial amount, what fraction would be left after 0.5 t?
(a) $\frac{1}{2}$ (b) $\frac{1}{\sqrt{3}}$ (c) $\frac{1}{3}$ (d) $\sqrt{\frac{2}{3}}$

131. The present activity of the hair of Egyption mummy is 1.75 dpm. $t_{1/2}$ of $^{14}_{6}C$ is 5770 year and disintegration rate of fresh sample of C^{14} is 14 dpm. Find out age of mummy.
(a) 23080 year (b) 138480 year (c) 11998.3 year (d) 17310 year

132. The amount of $^{14}_{6}C$ isotope in a piece of wood is found to one fourth (1/4) of that present in a fresh piece of wood. Calculate the age of the piece of wood (t_{12} of $^{14}_{6}C$ 5770 years)
(a) 7999 year (b) 11540 year (c) 16320 year (d) 23080 year

133. A radioactive element undergoing decay is left 20% of its initial weight after certain period of time t. How many such periods should elapse from the start for the 50% of the element to be left over?
(a) 3 (b) 4 (c) 5 (d) None of these

134. In a sample of wood, the reading of a counter is 32 dpm and in a fresh sample of tree it is 122dpm. Due to error counter gives the reading 2 dpm in absence of ^{14}C. Half life of ^{14}C is 5770 years.
The approximate age (in years) of wood sample is:
(a) 7997.2 (b) 57570 (c) 11,540 (d) 15140

135. A 0.50g sample of rock was found to have 2.5 10^{6} mol of $^{40}_{19}K$ ($t_{1/2}$ 1.3 10^{9} yr) and 7.5 10^{6} mol of $^{40}_{20}Ca$. How old is the rock ?
(a) 6.5 10^{8} yr (b) 1.3 10^{9} yr (c) 2.6 10^{9} yr (d) 5.2 10^{9}yr

For More Material Join: @JEEAdvanced_2024

136. Indium-112 is radioactive and has a very short half-life ($t_{1/2}$ 14 min). Its decay constant and average life are respectively :
(a) 0.0495 min ^{1}, 9.7 min (b) 0.495 min ^{1}, 20.2 min
(c) 9.7 min ^{1}, 20.2 min (d) 0.0495 min ^{1}, 20.2 min

137. The half-life of a radioactive element is 100 minutes. The time interval between the stages to 50% and 87.5% decay will be :
(a) 100 min (b) 50 min (c) 200 min (d) 25 min

138. The half-life of Tc^{99} is 6.0 hr. The total residual activity in a patient 30 hr after receiving an injection containing Tc^{99} must be more than 0.01 C_i. What is the maximum activity (in C_i) that the sample injected can have?
(a) 0.16 (b) 0.32 (c) 0.64 (d) 0.08

139. A pure radio-chemical preparation was observed to disintegrate at the rate of 2140 counts/minutes at 12.35 P.M. At 3.55 P.M. of the same day, the disintegration rate of the sample was only 535 count/minutes. What is the half-life of the material?
(a) 50 min (b) 100 min (c) 200 min (d) None of these

140. A certain radioactive isotope $^{A}_{Z}X$ ($t_{1/2}$ 100 days) decays to $^{A}_{Z}{}^{8}_{2}Y$. If 1 mole of $^{A}_{Z}X$ is kept in sealed container, how much He gas will accumulate at 1 atm and 273 K in 200 days?
(a) 11.2 litre (b) 33.6 litre (c) 22.4 litre (d) 44.8 litre

141. A radioactive substance decay 25% in 10 minute. If at start there are 4 10^{20} atoms present. After how much time will the number of atoms be reduced to 10^{20} atoms? (given ln 3 1.098)
(a) 10.98 min (b) 21.97 min (c) 48.19 min (d) None of these

142. The time of decay for a nuclear reaction is given by t 4 $t_{1/2}$. The relation between the mean life (T) and time of decay (t) is given by :
(a) 2 T ln 2 (b) 4 T ln 2 (c) 2 T^{4} ln 2 (d) $\frac{1}{T^{2}}$ ln 2

143. Two radio isotopes A and B of atomic mass X and Y are mixed in equal amount by mass. After 20 days, their mass ratio is found to be 1 : 4. Isotope A has a half-life of 1 day. The half-life of isotope B is :
(a) 1.11 $\frac{Y}{X}$ day (b) 0.11 $\frac{X}{Y}$ day (c) 0.6237 day (d) 1.10 day

144. Two radioactive nuclides A and B have half-lives 50 min and 10 min respectively. A fresh sample contains the nuclide of B to be eight times that of A. How much time should elapse so that the number of nuclides of A becomes double of B?
(a) 30 (b) 40 (c) 50 (d) 100

145. A radioactive nuclide is produced at a constant rate of per second. It's decay constant is . If N_0 be the no. of nuclei at time t 0, then max. no. of nuclei possible are :
(a) N_0 (b) (c) N_0 — (d) — N_0s

146. An analysis of the rock shows that the relative number of Sr^{87} and Rb^{87} ($t_{1/2}$ 4.7 10^{10} year) atoms is 0.05. What is the age of the rock? Assume all the Sr^{87} have been formed from Rb^{87} only.
(a) 7.62 10^{9} year (b) 1.43 10^{9} year
(c) 3.28 10^{9} year (d) 4.32 10^{8} year

For More Material Join: @JEEAdvanced_2024

147. There are two radio nuclei A and B. A is a -emitter and B is -emitter. Their disintegration constant are in the ratio of 1 : 2. What should be the number of atoms ratio of A and B at time t 0, so that initially probability of getting of and -particles are same.


(a) 2 : 1 (b) 4 : 1 (c) 1 : 2 (d) 1 : 4

148. A radioactive substance (parent) decays to its daughter element. The age of radioactive substance (t) is related to the daughter (d)/parent (p) ratio by the equation :

(a) $t \quad \frac{1}{}\ln \ 1 \quad \frac{p}{d}$ (b) $t \quad \frac{1}{}\ln \ 1 \quad \frac{d}{p}$

(c) $t \quad \frac{1}{}\ln \ \frac{d}{p}$ (d) $t \quad \frac{1}{}\ln \ \frac{p}{d}$

149. Ac^{227} has a half-life of 22 years. The decays follows two parallel paths



What are the decay constants ( ) for Th and Fr respectively?

(a) 0.03087, 0.00063 (b) 0.00063, 0.03087 (c) 0.02, 0.98 (d) None of these

150. $^{218}_{84}Po\,(t_{1/2} \quad 183\ \text{sec})$ decay to $_{82}Pb\,(t_{1/2} \quad 161\ \text{sec})$ by -emission, while Pb^{214} is a -emitter. In an experiment starting with 1 mole of pure Po^{218}, how much time would be required for the number of nuclei of $^{214}_{82}Pb$ to reach maximum?

(a) 147.5 (b) 247.5 (c) 182 (d) 304

For More Material Join: @JEEAdvanced_2024



Level 2

1. The forward rate constant for the elementary reversible gaseous reaction

$$C_2H_6 \rightleftharpoons 2CH_3 \text{ is } 1.57 \quad 10^{\ 3} \text{ s}^{\ 1} \text{ at 100 K}$$

What is the rate constant for the backward reaction at this temperature if $10^{\ 4}$ moles of CH_3 and 10 moles of C_2H_6 are present in a 10 litre vessel at equilibrium.

(a) $1.57 \quad 10^{9}$ L mol ^{1} s ^{1} (b) $1.57 \quad 10^{10}$ L mol ^{1} s ^{1}

(c) $1.57 \quad 10^{11}$ L mol ^{1} s ^{1} (d) $1.57 \quad 10^{7}$ L mol ^{1} s ^{1}

2. For a hypothetical reaction,

$A \quad 3B \quad P \quad H \quad 2x$ kJ/mole of A

& $M \quad 2Q \quad R \quad H \quad x$ kJ/mole of M

These reactions are carried simultaneously in a reactor such that temperature is not changing. If rate of disappearance of B is $y\,M$ sec ^{1} then rate of formation (in M sec ^{1}) of Q is:

(a) $\frac{2}{3}y$ (b) $\frac{3}{2}y$ (c) $\frac{4}{3}y$ (d) $\frac{3}{4}y$

3. The kinetic data for the given reaction $A(g) + 2B(g) \overset{k}{\longrightarrow} C(g)$ is provided in the following table for three experiments at 300 K

Ex. No.	[A/M]	[B/M]	Initial rate (M sec ^{1})
1.	0.01	0.01	$6.930 \quad 10^{\ 6}$
2.	0.02	0.01	$1.386 \quad 10^{\ 5}$
3.	0.02	0.02	$1.386 \quad 10^{\ 5}$

In another experiment starting with initial concentration of 0.5 and 1 M respectively for A and B at 300 K, find the rate of reaction after 50 minutes from the starting of experiment (in M/sec).

(a) $6.93 \quad 10^{\ 4}$ (b) $0.25 \quad 10^{\ 7}$ (c) $4.33 \quad 10^{\ 5}$ (d) $3.46 \quad 10^{\ 4}$

4. The reaction $A(g) + 2B(g) \longrightarrow C(g)$ is an elementary reaction. In an experiment involving this reaction, the initial partial pressures of A and B are $P_A \quad 0.40$ atm and $P_B \quad 1.0$ atm respectively. When $P_C \quad 0.3$ atm, the rate of the reaction relative to the initial rate is :

(a) $\frac{1}{12}$ (b) $\frac{1}{50}$ (c) $\frac{1}{25}$ (d) none of these

5. Which of the following is incorrect statement ?

(a) Stoichiometry of a reaction tells about the order of the elementary reaction

(b) For a zero order reaction, rate and the rate constant are identical.

(c) A zero order reaction is controlled by factors other than concentration of reactants

(d) A zero order reaction is an elementary reaction

For More Material Join: @JEEAdvanced_2024


6. Two first order reaction have half-lives in the ratio 8 : 1. Calculate the ratio of time intervals $t_1 : t_2$. The time t_1 and t_2 are the time period for $\frac{1}{4}^{th}$ and $\frac{3}{4}^{th}$ completion.

(a) 1 : 0.301 (b) 0.125 : 0.602 (c) 1 : 0.602 (d) none of these

7. Reaction A B C D follows rate law, r $k[A]^{1/2}[B]^{1/2}$. Starting with 1 M of A and B each, what is the time taken for concentration of A become 0.1 M?
[Given k 2.303 10^{-2} sec $^{-1}$].

(a) 10 sec (b) 100 sec (c) 1000 sec (d) 434 sec

8. For the two reactions I: A B ; II: C D following graph is obtained.



Which of the following is true:

(a) If $[B]$ $[A]$ then at that time $[D]$ 0.75 M (b) If $[C]$ $[A]$ then at that time $[B]$ $[D]$

(c) $(t_{100\%})_{\text{Reaction I}}$ $(t_{100\%})_{\text{Reaction II}}$ (d) $[A]$ $[C]$ at t $\frac{\sqrt{3}}{2}$ min.

9. The reaction $A(g)$ $B(g)$ $2C(g)$ is a first order reaction with rate constant 2.772 10^{-3} s $^{-1}$. Starting with 0.1 mole of A in 2 litre vessel, find the concentration of A after 250 sec when the reaction is allowed to take place at constant pressure and at 300 K ?

(a) 0.0125 M (b) 0.025 M (c) 0.05 M (d) none of these

10. For a first order homogeneous gaseous reaction, A $2B$ C
if the total pressure after time t was P_t and after long time (t) was P then k in terms of P_t, P and t is :

(a) k $\frac{2.303}{t}\log\frac{P}{P \quad P_t}$ (b) k $\frac{2.303}{t}\log\frac{2P}{P \quad P_t}$

(c) k $\frac{2.303}{t}\log\frac{2P}{3(P \quad P_t)}$ (d) none of these

11. $A(aq)$ $B(aq)$ $C(aq)$ is a first order reaction.
Time t
moles of reagent n_1 n_2
Reaction progress is measured with the help of titration of reagent 'R'. If all A, B and C react with reagent and have 'n' factors [n factor; eq. mass $\frac{\text{mol. mass}}{n}$] in the ratio of 1 : 2 : 3 with the reagent, the k in terms of t, n_1 and n_2 is :

(a) k $\frac{1}{t}\ln\frac{n_2}{n_2 \quad n_1}$ (b) k $\frac{1}{t}\ln\frac{2n_2}{n_2 \quad n_1}$


For More Material Join: @JEEAdvanced_2024

(c) $k \quad \frac{1}{t}\ln \frac{4n_2}{n_2 \quad n_1}$ (d) $k \quad \frac{1}{t}\ln \frac{4n_2}{5(n_2 \quad n_1)}$

12. The gaseous decomposition reaction, $A(g) \quad 2B(g) \quad C(g)$ is observed to first order over the excess of liquid water at 25°C. It is found that after 10 minutes the total pressure of system is 188 torr and after very long time it is 388 torr. The rate constant of the reaction (in hr^{-1}) is : [Given : vapour pressure of H_2O at 25° is 28 torr ($\ln 2 \quad 0.7, \ln 3 \quad 1.1, \ln 10 \quad 2.3$)]
(a) 0.02 (b) 1.2 (c) 0.2 (d) none of these

13. The reaction, Sucrose $\xrightarrow{H}$ Glucose + Fructose, takes place at certain temperature while the volume of solution is maintained at 1 litre. At time zero the initial rotation of the mixture is 34°. After 30 minutes the total rotation of solution is 19° and after a very long time, the total rotation is –11°C. Find the time when solution was optically inactive.
(a) 135 min (b) 103.7 min (c) 38.7 min (d) 45 min

14. A gaseous compound *A* reacts by three independent first order processes (as shown in figure) with rate constant $2 \quad 10^{-3}$, $3 \quad 10^{-3}$ and $1.93 \quad 10^{-3} sec^{-1}$ for products *B*, *C* and *D* respectively. If initially pure *A* was taken in a closed container with $P \quad 8$ atm, then the partial pressure of *B* (in atm) after 100 sec from starting the experiment.



(a) 0.288 (b) 0.577 (c) 1.154 (d) none of these


15. A compound *A* dissociates by two parallel first order paths at certain temperature

$A(g) \xrightarrow{k_1(min^{-1})} 2B(g) \qquad k_1 \quad 6.93 \quad 10^{-3} min^{-1}$

$A(g) \xrightarrow{k_2(min^{-1})} C(g) \qquad k_2 \quad 6.93 \quad 10^{-3} min^{-1}$

The reaction is started with 1 mole of pure '*A*' in 1 litre closed container with initial pressure 2 atm. What is the pressure (in atm) developed in container after 50 minutes from start of experiment?
(a) 1.25 (b) 0.75 (c) 1.50 (d) 2.50

16. For given hypothetical elementary parallel reaction,



Initially only 2 moles of *A* are present. The total no. of moles of *A, B* and *C* at the end of 75% reaction are:
(a) 2 (b) 3 (c) 4 (d) 3.5

For More Material Join: @JEEAdvanced_2024


17. The reaction *cis*-$X \underset{k_b}{\overset{k_f}{\rightleftharpoons}}$ *trans*-X is first order in both directions. At 25°C, the equilibrium constant is 0.10 and the rate constant k_f 3 10 ^{4} s ^{1}. In an experiment starting with the pure *cis*-form, how long would it take for half of the equilibrium amount of the *trans*-isomer to be formed?

(a) 150 sec (b) 200 sec (c) 240 sec (d) 210 sec

18. A hydrogenation reaction is carried out at 500 K. If the same reaction is carried out in the presence of a catalyst at the same rate with same frequency factor, the temperature required is 400 K. What is the activation energy of the reaction, if the catalyst lowers the activation energy barrier by 16 kJ/mol?

(a) 100 kJ/mol (b) 80 kJ/mol (c) 60 kJ/mol (d) None of these

19. Consider the reaction.



The rate constant for two parallel reactions were found to be 10 ^{2} dm ^{3} mol ^{1} s ^{1} and 4 10 ^{2} dm ^{3} mol ^{1} s ^{1}. If the corresponding energies of activation of the parallel reaction are 100 and 120 kJ/mol respectively, what is the net energy of activation (E_a) of A?

(a) 100 kJ/mol (b) 120 kJ/mol (c) 116 kJ/mol (d) 220 kJ/mol

20. A reaction takes place in various steps. The rate constant for first, second, third and fifth steps are k_1, k_2, k_3 and k_5 respectively. The overall rate constant is given by

$$k \quad \frac{k_2}{k_3} \quad \frac{k_1}{k_5}^{1/2}$$

If activation energy are 40, 60, 50 and 10 kJ/mol respectively, the overall energy of activation (kJ/mol) is :


(a) 10 (b) 20 (c) 25 (d) none of these

21. For reaction A B, the rate constant k_1 $A_1 e^{-Ea_1/(RT)}$ and for the reaction X Y, the rate constant k_2 $A_2 e^{-Ea_2/(RT)}$. If A_1 10^8, A_2 10^{10} and E_{a_1} 600 cal/mol, E_{a_2} 1800 cal/mol, then the temperature at which k_1 k_2 is (Given : R 2 cal/K-mol)

(a) 1200 K (b) 1200 4.606 K (c) $\frac{1200}{4.606}$ K (d) $\frac{600}{4.606}$ K

22. For first order parallel reactions k_1 and k_2 are 4 and 2 min ^{1} respectively at 300 K. If the activation energies for the formation of B and C are respectively 30,000 and 38,314 joule/mol respectively, the temperature at which B and C will be obtained in equimolar ratio is :

For More Material Join: @JEEAdvanced_2024




(a) 757.48 K (b) 378.74 K
(c) 600 K (d) none of these


23. In the series reaction
$A \xrightarrow{k_1} B \xrightarrow{k_2} C \xrightarrow{k_3} D$, if $k_1 \quad k_2 \quad k_3$ then the rate determing step of the reaction is
(a) $A \quad B$ (b) $C \quad D$
(c) $B \quad C$ (d) Any step


24. The following mechanism has been proposed for the exothermic catalyzed complex reaction


$$A \quad B \underset{\text{fast}}{\overset{\text{slow}}{\rightleftharpoons}} IAB \xrightarrow{k_1} AB \quad I \xrightarrow{k_2} P \quad A$$

If k_1 is much smaller than k_2, the most suitable qualitative plot of potential energy (P. E.) *versus* reaction co-ordinate (R.C.) for the above reaction

(a)



(b)


(c)


(d)



For More Material Join: @JEEAdvanced_2024

25. The mechanism of esterification in presence of acid catalyst (H_2SO_4) is proposed as follows :




Which of the following potential energy *Vs* reaction co-ordinate diagram is consistent with given mechanism?


(a)




(b)



(c)



(d)



26. For the first order reaction $A \quad B \quad C$, carried out at 27°C. If $3.8 \quad 10^{16}$% of the reactant molecules exists in the activated state, the E_a (activation energy) of the reaction is :

(a) 12 kJ/mol (b) 831.4 kJ/mol (c) 100 kJ/mol (d) 88.57 kJ/mol

27. Upon irradiating californium with neutrons, a scientist discovered a new nuclide having mass number of 250 and a half-life of 30 min. After 90 min. of irradiation, the observed radioactivity due to nuclide was 100 dis/min. How many atoms of the nucleide were prepared initially?

(a) $2.4 \quad 10^4$ (b) $3.46 \quad 10^4$ (c) 1900 (d) 800

For More Material Join: @JEEAdvanced_2024

28. A radioactive isotope X with half-life of 6.93 10^9 years decay to Y which is stable. A sample of rock from the moon was found to contain both the elements X and Y in the mole ratio 1 : 7. What is the age of the rock?

(a) 2.079 10^{10} years (b) 1.94 10^{10} years

(c) 1.33 10^9 years (d) 10^{10} years

29. The average (mean) life of a radio nuclide which decays by parallel path is

A B; $_1$ 1.8 10 ^{2} sec ^{1}

$2A$ ^{2} B; $_2$ 10 ^{3} sec ^{1}

(a) 52.63 sec (b) 500 sec (c) 50 sec (d) none of these

30. The radioactive decay $^{211}_{83}$Bi $^{207}_{81}$ Tl, takes place in 100 L closed vessel at 27°C. Starting with 2 moles of $^{211}_{83}$Bi ($t_{1/2}$ 130 sec), the pressure development in the vessel after 520 sec will be :

(a) 1.875 atm (b) 0.2155 atm (c) 0.4618 atm (d) 4.618 atm

31. A fresh radioactive mixture contains short lived species A and B. Both emitting -particles initially of 8000 -particles per minute. 20 minutes later, they emits at the rate of 3500 -particles per minute. If the half-lives of the species A and B are 10 minutes and 500 hours respectively, then the ratio of activities of A : B in the initial mixture was :

(a) 4 : 6 (b) 6 : 4 (c) 3 : 4 (d) 3 : 1

32. In order to determine the volume of blood in an animal, a 1.0 mL sample of solution of 10^3 dpm of $^3_1$H is injected into the animal blood stream. After sufficient time for circulatory equilibrium to be established, 2 mL of blood is found to have activity to 10 dpm. The volume of blood in animal is :

(a) 199 mL (b) 198 mL (c) 200 mL (d) 20 mL

33. The ratio of activities of two radio nuclides X and Y in a mixture at time t 0 was found to be 4 : 1. After two hours, the ratio of activities become 1 : 1. If the $t_{1/2}$ of radio nuclide X is 20 min then $t_{1/2}$ [in minutes] of radio nuclide Y is :

(a) 10 (b) 20 (c) 30 (d) 40

34. Find the age of an ancient Egyptian wooden article (in years) from the given information.

(i) Activity of 1 g of carbon obtained from ancient wooden article 7 counts/min/g

(ii) Activity of 1 g carbon obtained from fresh wooden sample 15.4 counts per min/g

(iii) Percentage increase in level of C^{14} due to nuclear explosions in past 100 years is 10%

(iv) $t_{1/2}$ of $^{14}_6$C 5770 years

(a) 5.770 10^3 (b) 16.87 10^3 (c) 2488 (d) none of these

35. The isotopes ^{238}U and ^{235}U occur in nature in the mass ratio 140 : 1. It is assumed that initially they were found in equal mass. If half life ($t_{1/2}$) of ^{238}U 4.5 10^9 and $t_{1/2}$ of ^{235}U 5 10^8 year respectively then the age of earth is (log 7 0.846; log 2 0.3)

(a) 4.02 10^9 year (b) 2.01 10^9 year (c) 8.72 10^9 year (d) None of these

For More Material Join: @JEEAdvanced_2024

Level 3

Passage 1

Two consecutive irreversible first order reactions can be represented by

$$A \xrightarrow{k_1} B \xrightarrow{k_2} C$$

The rate equation for A is readily integrated to obtain


$$[A]_t = [A]_0 . e^{-k_1 t}; \text{ and } [B]_t = \frac{k_1 [A]_0}{k_2 - k_1} [e^{-k_1 t} - e^{-k_2 t}]$$

1. At what time will B be present in maximum concentration?


(a) $\frac{k_1}{k_2 - k_1}$ (b) $\frac{1}{k_1 - k_2} \ln \frac{k_1}{k_2}$ (c) $\frac{1}{k_2 - k_1} \ln \frac{k_1}{k_2}$ (d) None of these

2. When $k_1 = 1\,s^{-1}$ and $k_2 = 500\,s^{-1}$; select most appropriate graph


(a)




(b)



(c)



(d)



3. Select the correct statement for given reaction :

(a) A decreases linearly

(b) B rises to a max. and then constant

(c) B rises to a max. and then falls

(d) The slowest rate of increases of C occuring where B is max.

For More Material Join: @JEEAdvanced_2024



Passage 2

Arrhenius studied the effect of temperature on the rate of a reaction and postulated that rate constant varies with temperature exponentially as $k \quad Ae^{\ E_a/RT}$. This method is generally used for finding the activation energy of a reaction. Keeping temperature constant, the effect of catalyst on the activation energy has also been studied.

1. The pre-exponential factor in the Arrhenius equation of a first order reaction has the unit :
(a) mol L ^{1}s ^{1} (b) L mol ^{1}s ^{1} (c) s ^{1} (d) dimensionless

2. If x is the fraction of molecules having energy greater than E_a it will be given by :
(a) $x \quad \frac{E_a}{RT}$ (b) $\ln x \quad \frac{E_a}{RT}$ (c) $x \quad e^{E_a/RT}$ (d) Any of these

3. If the rate of reaction doubles for 10°C rise of temperature form 290K to 300K, the activation energy of the reaction will be approximately:
(a) 40 kcal mol ^{1} (b) 12 kcal mol ^{1} (c) 60 kcal mol ^{1} (d) 70 kcal mol ^{1}



Passage 3

An important parameter of a photochemical reaction is the quantum efficiency or quantum yield ( ) which is defined as

$$\frac{\text{moles of the substance reacted}}{\text{moles of photons absorbed}}$$

Absorption of UV radiation decompose acetone according to the reaction

$$(CH_3)_2CO \quad ^{h\nu} \quad C_2H_6 + CO$$

1. The quantum yield of a reaction at 330 nm is 0.4. A sample of acetone absorbs monochromatic radiation at 330 nm at the rate of 7.2 10 ^{3} Js ^{1} (Given : N_A 6 10^{23}; h 6.6 10 ^{34} in S.I. unit). The rate of formation of CO(mol/s) is :
(a) 2 10 ^{8} (b) 8 10 ^{8} (c) 8 10 ^{9} (d) none of these

2. If quantum yield in 0.8 then rate of formation of C_2H_6 (mol/s) is :
(a) 2 10 ^{8} (b) 1.6 10 ^{9} (c) 16 10 ^{9} (d) 8 10 ^{9}



Passage 4

Radioactive disintegration is a first order reaction and its rate depends only upon the nature of nucleus and does not depend upon external factors like temperature and pressure. The rate of radioactive disintegration (Activity) is represented as

$$\frac{dN}{dt} \quad N$$

Where decay constant; N number of nuclei at time t; N_0 initial no. of nuclei

For More Material Join: @JEEAdvanced_2024

The above equation after integration can be represented as

$$\frac{2.303}{t}\log\frac{N_0}{N}$$

1. Half-life period of U^{237} is 2.5 10^5 years. In how much time will the amount of U^{237} remaining be only 25% of the original amount?
(a) 2.5 10^5 years (b) 1.25 10^5 years (c) 5 10^5 years (d) 10^6 years

2. Calculate the half-life period of a radioactive element which remains only 1/16 of its original amount in 4740 years :
(a) 1185 years (b) 2370 years (c) 52.5 years (d) none of these

3. What is the activity in Ci (curie) of 1.0 mole of Plutonium–239? ($t_{1/2}$ 24,000 years)
(a) 1.49 Ci (b) 14.9 Ci (c) 5.513 10^{11} Ci (d) None of these



Passage 5

Size of nucleus was obtained by the equation r $R_0A^{1/3}$, where r is the radius of nucleus of mass no. A and R_0 is a constant whose value is equal to 1.5 10^{-15} metre.
(Given : 1 amu 1.66 10^{-24} g)

1. What is the density of a nucleus of mass number A?
(a) $\frac{4}{3}$ $(1.5\ 10^{-15})^3 A$ (b) 1.17 10^{17} kg/cm^3 (c) 1.17 10^{-17} kg/m^3 (d) None of these

2. Nucleus radius of $_6C^{12}$ is 3 10^{-15} metre. What is density ratio of d_C/d_{H_2O}?
(a) 1.76 10^{17} (b) 1.76 10^{14} (c) 17.6 10^7 (d) 17.6 10^{17}

One or More Answers is/are Correct



1. Select the correct statement(s):
(a) Rate constants are never negative
(b) Partial orders are never negative
(c) Molecularity and order of reaction both are equal for elementary reactions
(d) Order of reaction may be change with change in practical conditions (temp. and pressure)

2. Select the correct statement(s) :
(a) The rate of reaction decreases with decrease in temperature
(b) The rate of reaction is uniform in zero order reaction
(c) The rate of reaction depends upon the surface area of the solid reactants
(d) Average and instantaneous rate of reaction defined for micro and macro-scopic time interval respectively

For More Material Join: @JEEAdvanced_2024

3. Select the correct statement(s) :

(a) The rate law of the elementary reaction; $2A \quad B \quad C$, must be $r \quad k[A]^2$

(b) The rate law for the complex reaction

$$A \quad B \qquad C, \text{ might not be } r \quad k[A][B]$$

(c) If the partial orders differ from the stoichiometric coefficients in the balanced reaction, the reaction must be complex

(d) If the partial orders are equal to corresponding coefficients in the balanced reaction, the reaction must be elementary

4. Select the correct statement(s):

(a) Every substance that appears in the rate law of reaction must be a reactant or product in that reaction

(b) If we know the rate law of a reaction; we can deduce its mechanism

(c) If the reaction has rate $r \quad k[A][B]^{3/2}$ then reaction may be elementary

(d) A zero order reaction must be a complex reaction

5. Select the correct statement(s) :

(a) When $T \qquad$ or $E_a \quad 0$ then $k \quad A$

(b) A positive catalyst can change $\quad H$ of the reaction

(c) A mixture of reactants may be thermodynamically unstable but kinetically stable

(d) A negative catalyst increases the activation energy of the reaction

6. Consider a reaction $A \quad B \quad C$, in which both reactants are in the same phase, may be

(a) unimolecular elementary reaction

(b) Exothermic

(c) Heterogeneous

(d) Photochemical

7. In the following gaseous phase first order reaction

$$A(g) \qquad 2B(g) \quad C(g)$$

initial pressure was found to be 400 mm of Hg and it changed to 1000 mm of Hg after 20 min. Then

(a) half life for A is 10 min

(b) rate constant is 0.0693 min ^{1}

(c) partial pressure of C at 30 min is 350 mm of Hg

(d) total pressure after 30 min is 1100 mm of Hg

8. Identify the true statement(s)

(a) A catalyst is chemically unchanged at the end of a reaction

(b) A catalyst may appear in the kinetic rate equation of the reaction


(c) A catalyst will not affect the composition of an equilibrium mixture

(d) A catalyst cannot cause a non-spontaneous ($G \quad 0$) reaction to proceed

9. For the reaction $A \underset{k_2 sec^{-1}}{\overset{k_1 sec^{-1}}{\rightleftharpoons}} B$ following graph is given,

$k_1 \quad 4 \quad 10^{\ 2}$ sec ^{1}. Which is/are correct statement (s) (ln 2 $\quad$ 0.7, ln 8/7=0.14)

For More Material Join: @JEEAdvanced_2024



(a) Equilibrium constant is 4.0
(b) Time taken for the completion of 50% of equilibrium conc. of B is 14 sec.
(c) Time taken for the completion of 10% of initial conc. of A is 2.8 sec.
(d) Rate constant of backward reaction is 10^{-2} sec^{-1}

10. Select the correct statement(s) :
(a) -particles are simply helium atoms
(b) -rays travel with higher speed as compared to -particle and have higher ionization power as compare to -particle
(c) Loss of -particles results in the production of isobars
(d) -particles are considered as the best bombarding particles

11. Select the correct statement(s) :
(a) In the reaction $^{235}_{92}U + {}^{1}_{0}n$ $^{140}_{56}Ba + 2\ {}^{1}_{0}n + x$, x is $^{94}_{36}Kr$
(b) In the reaction $^{23}_{11}Na + z$ $^{23}_{12}Mg + {}^{1}_{0}n$, the bombarding particle z is deuteron
(c) Very large amount of energy is produced during nuclear fission and nuclear fussion
(d) In a fission reaction, a loss in mass occurs releasing a vast amount of energy

12. Select the correct statement(s):
(a) SI unit of radioactivity is becquerel (Bq)
(b) 1 Ci 3.7 10^{7} Bq
(c) $^{7}_{3}Li + {}^{1}_{1}H$ $^{4}_{2}He$ is (P,) type reaction
(d) The half-life of a particular radioactive isotope is a characteristics constant of that isotope

13. Select the correct statement(s) :
(a) On bombarding $^{14}_{7}N$ nuclei with -particle, the nuclei of the product formed after release of proton would be $^{17}_{8}O$
(b) Decay constant does not depend upon temperature
(c) Nuclide and its decay product after -emission are called isodiaphers
(d) Half-life of radium is 1580 years. Its average life will be 1097.22 years

14. In the decay process

$$A \quad B \quad C \quad D$$

(a) A and B are isobars
(b) A and D are isotopes

For More Material Join: @JEEAdvanced_2024

(c) B, C and D are isobars

(d) A and C are isotones

15. In electron capture (radioactive process) :

(a) a neutron is formed (b) a proton is consumed

(c) -ray emission takes place (d) X-ray emission takes place

16. Select the correct statement (s) for positron emission by unstable nucleus.

(a) X-ray emission takes place (b) A neutron is formed

(c) $\frac{n}{p}$ of daughter nucleus increases (d) A neutron is consumed

17. Select the correct statement (s) :

(a) Mass number remains constant when positron emission takes place

(b) One neutron converts into proton in $({}^{0}_{1}e)$ emission process

(c) Activity of a radioactive substance double when temp. increases from 300 K to 310 K

(d) Isodiaphers formed when one alpha particle emitted and isotopes formed when 2 beta particles emitted

For More Material Join: @JEEAdvanced_2024

Match the Column



Column-I and Column-II contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

1.

Column-I	Column-II
(A) Unit of k is always equals to	(P) 1/time
(B) Unit of k in zero order	(Q) M/time
(C) Unit of k in first order	(R) Time ^{1} M ^{1}
(D) Unit of k in second order	(S) Unit of A (pre-exponential factor)

2.

Column-I	Column-II
(A) Molecularity of a reaction	(P) 0, 1 Possible
(B) Order of reaction	(Q) 1, 2 Possible
(C) The dissociation of H_2O_2 (aq) is	(R) First order reaction
(D) $H_2(g)$ $Cl_2(g)$ $\overset{h\nu}{}$ 2HCl is	(S) Zero order reaction

3.

Column-I (Curve)	Column-II (Slope)
(A) C vs t (abscissa) for zero order	(P) unity
(B) log C vs t (abscissa) for first order	(Q) zero
(C) $\frac{dc}{dt}$ vs C for zero order	(R) k
(D) ln $\frac{dc}{dt}$ vs ln C for first order	(S) $\frac{k}{2.303}$




4.

Column-I Linear plots (with non zero slope)	Column-II (Order)
(A) ln $\frac{d[A]}{dt}$ vs. ln [A]	(P) 2
(B) $\log_e k$ vs. $\frac{1}{T}$	(Q) $\frac{1}{2}$
(C) $\log t_{1/2}$ vs. $\log [A]_0$	(R) 0
(D) $\frac{d[A]}{dt}$ vs. $[A]^2$	(S) 1

$[A]$ = Reactant concentration at time 't'
k rate constant
T Absolute temperature

For More Material Join: @JEEAdvanced_2024

5.

Column-I (First order of reaction *A* → *B*)	Column-II
(A) $[B]$ vs t	(P) 
(B) $[A]$ vs t	(Q) 
(C) $\frac{1}{[A]}$ vs t	(R) 
(D) $\ln k$ vs $\frac{1}{T}$	(S) 

6.

Column-I	Column-II
(A) Isotones	(P) $^{234}_{91}Pa$ and $^{234}_{90}Th$
(B) Isobars	(Q) $^{12}_{6}C$ and $^{14}_{6}C$
(C) Isotopes	(R) $^{39}_{19}K$ and $^{19}_{9}F$
(D) Isodiaphers	(S) $^{39}_{18}Ar$ and $^{40}_{19}K$

7.

Column-I	Column-II
(A) -emission	(P) Change in mass no.
(B) -emission	(Q) No change in atomic no. and mass no.
(C) -emission	(R) Atomic no. decreases
(D) (Positron) emission	(S) Atomic no. increases

For More Material Join: @JEEAdvanced_2024

Assertion-Reason Type Questions



Each question contains STATEMENT-1 (Assertion) and STATEMENT-2 (Reason).
Examine the statements carefully and mark the correct answer according to the instructions given below :
(A) If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
(B) If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
(C) If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
(D) If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

1. **STATEMENT-1 :** Molecularity has no meaning for a complex reaction
   STATEMENT-2 : Molecularity defined only for RDS
2. **STATEMENT-1 :** An elementary reaction cannot have fractional order.
   STATEMENT-2 : Stoichiometric coefficients in an elementary reaction can be fractional.
3. **STATEMENT-1 :** Concentration of reactant in zero order reaction is constant.
   STATEMENT-2 : For zero order reaction A B, successive half life of reaction decreases with the progress of the reaction.
4. **STATEMENT-1 :** Acid catalysed hydrolysis of esters is pseudo first order reaction.
   STATEMENT-2 : Water is present in excess in given reaction.
5. **STATEMENT-1 :** The order of reaction can have fractional value.
   STATEMENT-2 : For an elementary reaction, the partial orders are determined by the reaction stoichiometry.
6. **STATEMENT-1 :** Catalyst may increase the rate constant to a large extent.
   STATEMENT-2 : By using suitable catalyst, we can significantly increase yield.
7. **STATEMENT-1 :** For each 10°C rise of temperature the k is nearly double.
   STATEMENT-2 : Energy wise distribution of molecules in a gas sample is an exponential function of temperature so $e^{E_a/RT}$ is doubled.
8. **STATEMENT-1 :** Product is formed only when the required orientation and energy conditions are met.
   STATEMENT-2 : All collisions between reactants yield the desired product.
9. **STATEMENT-1 :** The plot of k *versus* $1/T$ is linear.
   STATEMENT-2 : $k \quad A \; e^{E_a/(RT)}$
10. **STATEMENT-1 :** For exothermic reaction equilibrium constant decreases with increase in temperature.
    STATEMENT-2 : For exothermic reaction rate constant decreases with decrease in temperature.
11. **STATEMENT-1 :** If the activation energy of reaction is zero, temperature will have no effect on the rate constant.
    STATEMENT-2 : Lower the activation energy faster is the reaction.
12. **STATEMENT-1 :** Active complex is an intermediate product.
    STATEMENT-2 : Active complex is unstable with high vibrational energy.

For More Material Join: @JEEAdvanced_2024

13. STATEMENT-1 : The pre-exponential factor A has the same units for all reactions.
STATEMENT-2 : $e^{E_a/RT}$ has no unit.

14. STATEMENT-1 : -rays have very high penetrating power.
STATEMENT-2 : -rays are electromagnetic radiations of high energy.

15. STATEMENT-1 : Nuclide $^{30}_{13}$Al is less stable than $^{40}_{20}$Ca.
STATEMENT-2 : Nuclide having odd number of protons and neutrons are generally unstable.

16. STATEMENT-1 : Disintegration of $^{3}_{1}$H (tritium) is accompanied by -emission.
STATEMENT-2 : Tritium has high n/p ratio.

17. STATEMENT-1 : The life of radioactive object (organic origin) can found with the help of carbon dating.
STATEMENT-2 : $^{14}_{6}$C is a and -emitter.

18. STATEMENT-1 : Neutrons are the best bombarding particles.
STATEMENT-2 : Neutrons are neutral particles.

19. STATEMENT-1 : Nucleus does not contain free electrons, yet it emit beta-particles
STATEMENT-2 : At high n/p ratio, one neutron supposed to give 1 proton and 1 e ( ).

20. STATEMENT-1 : Rate of disintegration of thorium increases with the increase in moles of thorium.
STATEMENT-2 : Rate of disintegration does not depend upon temperature, pressure

Subjective Problems

1. The rate of decomposition of $NH_3(g)$ at 10 atm on platinum surface is zero order. What is rate of formation (in M min ^{1}) of $H_2(g)$, if rate constant of reaction $2NH_{3(g)}$ $N_2(g)$ $3H_2(g)$ is $2.0\,M$ min ^{1} ?

2. $5A$ Product

In above reaction, half-life period is directly proportional to initial concentration of reactant. The initial rate of reaction is 400 mol lit ^{1} min ^{1}.

Calculate the half-life period (in sec) when initial concentration of reactant is 200 mol lit ^{1}.

3. In an elementary reaction $A(g)$ $2B(g)$ $C(g)$ the initial pressure of A and B are P_A 0.40 atm and P_B 0.60 atm respectively. After time T, if pressure of C is observed 0.1 atm, then find the value of $\dfrac{r_i \text{ (initial rate of reaction)}}{r_t \text{ (rate of reaction after time } t)}$.

4. Carbon monoxide reacts with O_2 to form CO_2: $2CO(g)$ $O_2(g)$ $2CO_2(g)$
Informations about this reaction are given in the table below.

For More Material Join: @JEEAdvanced_2024

[CO] mol/L	**[O_2]** mol/L	**Rate of reaction** (mol/L.min)
0.02	0.02	4 10^{5}
0.04	0.02	1.6 10^{4}
0.02	0.04	8 10^{5}

What is the value for the rate constant for the reaction in properly related unit ?


5. Half-life for the zero order reaction $A(g)$ $B(g)$ $C(g)$ and half-life for the first order reaction $X(g)$ $Y(g)$ $Z(g)$ are equal. If completion time for the zero order reaction is 13.86 min, then calculate the rate constant (in hr^{1}) for the reaction $X(g)$ $Y(g)$ $Z(g)$.

6. For any acid catalysed reaction, $A \xrightarrow{H} B$

half-life period is independent of concentration of A at given pH. At difinite concentration of A half -life time is 10 min at pH 2 and half-life time is 100 min at pH 3. If the rate law expression of reaction is r $k[A]^{x}[H]^{y}$ then calculate the value of $(x \quad y)$.

7. For a reaction, $A \rightleftharpoons B$ equilibrium constant is 1.66 and $k_{forward}$ 0.166 hr^{1}.

Calculate the time (in hours) when concentration of B is 80% of its equilibrium concentration.
(Given : ln 25 3.20)

8. 

; At time t 0, initial mole of A is 1.

Overall half life of the reaction is 15 days. Then calculate the number of mole of C after 45 days if the ratio of $k_1 : k_2 : k_3$ is 4 : 2 : 1

9. Iodine-131 is a radioactive isotope. If 1.0 mg of ^{131}I has an activity of 4.6 10^{12} Bq, what is the half -life of ^{131}I (in days) ?

10. The average life of a radioactive element is 7.2 min. Calculate the time interval (in min.) between the stages of 33.33% and 66.66% decay

11. A, B and C are isodiaphers while C, D and E are isobars. Calculate the difference of protons between A and E $^{206}_{82}A$ B C D E

Given : Isodiaphers and isobars are formed in successive and -emission respectively.

12. In the given radioactive disintegration series,

$^{232}_{90}Th$ $^{208}_{82}Pb$

Calculate value of $(n \quad 2)$.
Where value of n is number of isobars formed in this series, suppose there is successive emission of -particles.

13. In the given radioactive disintegration series

$^{235}_{92}U$ $^{207}_{82}Pb$

Calculate difference between number of and number of particles emitted in this series.

For More Material Join: @JEEAdvanced_2024

Answers

Level 1

1. (c)	2. (b)	3. (c)	4. (d)	5. (d)	6. (d)	7. (d)	8. (c)	9. (a)	10. (c)
11. (c)	12. (a)	13. (d)	14. (d)	15. (b)	16. (d)	17. (c)	18. (c)	19. (d)	20. (a)
21. (c)	22. (c)	23. (b)	24. (a)	25. (c)	26. (c)	27. (d)	28. (c)	29. (b)	30. (c)
31. (d)	32. (a)	33. (d)	34. (a)	35. (b)	36. (c)	37. (a)	38. (b)	39. (a)	40. (c)
41. (c)	42. (d)	43. (c)	44. (b)	45. (c)	46. (b)	47. (b)	48. (b)	49. (a)	50. (b)
51. (d)	52. (c)	53. (c)	54. (d)	55. (d)	56. (a)	57. (d)	58. (a)	59. (a)	60. (c)
61. (c)	62. (c)	63. (c)	64. (a)	65. (a)	66. (c)	67. (b)	68. (d)	69. (d)	70. (c)
71. (d)	72. (a)	73. (a)	74. (b)	75. (d)	76. (c)	77. (a)	78. (d)	79. (d)	80. (c)
81. (b)	82. (b)	83. (c)	84. (c)	85. (c)	86. (c)	87. (c)	88. (b)	89. (c)	90. (c)
91. (d)	92. (d)	93. (d)	94. (b)	95. (b)	96. (c)	97. (b)	98. (d)	99. (d)	100. (c)
101. (c)	102. (b)	103. (c)	104. (d)	105. (d)	106. (b)	107. (a)	108. (c)	109. (d)	110. (b)
111. (a)	112. (c)	113. (d)	114. (c)	115. (b)	116. (c)	117. (c)	118. (c)	119. (b)	120. (c)
121. (d)	122. (c)	123. (b)	124. (d)	125. (d)	126. (b)	127. (c)	128. (c)	129. (b)	130. (b)
131. (d)	132. (b)	133. (d)	134. (c)	135. (c)	136. (d)	137. (c)	138. (b)	139. (b)	140. (b)
141. (c)	142. (b)	143. (d)	144. (c)	145. (b)	146. (c)	147. (a)	148. (b)	149. (b)	150. (b)

Level 2

1. (d)	2. (c)	3. (c)	4. (c)	5. (d)	6. (c)	7. (b)	8. (a)	9. (a)	10. (c)
11. (d)	12. (b)	13. (b)	14. (c)	15. (d)	16. (d)	17. (d)	18. (b)	19. (c)	20. (c)
21. (d)	22. (b)	23. (b)	24. (b)	25. (a)	26. (c)	27. (b)	28. (a)	29. (c)	30. (c)
31. (d)	32. (a)	33. (c)	34. (a)	35. (a)					

For More Material Join: @JEEAdvanced_2024



Level 3

Passage–1	**1.**	(b)	**2.**	(b)	**3.**	(c)		
Passage–2	**1.**	(c)	**2.**	(b)	**3.**	(b)		
Passage–3	**1.**	(c)	**2.**	(c)				
Passage–4	**1.**	(c)	**2.**	(a)	**3.**	(b)		
Passage–5	**1.**	(b)	**2.**	(b)				

One or More Answers is/are Correct

1. (a,c,d) **2.** (a,b,c) **3.** (a,b,c) **4.** (d) **5.** (a,c,d) **6.** (b,c,d) **7.** (a,b,c,d) **8.** (a,b,c,d)

9. (a,b,c,d) **10.** (c) **11.** (a,c,d) **12.** (a,c,d) **13.** (a,b,c) **14.** (b,c) **15.** (a,b,d) **16.** (b,c)

17. (a,b)

Match the Column

1.	A	S;	B	Q;	C	P;	D	R
2.	A	Q;	B	P, Q;	C	R;	D	S
3.	A	R;	B	S;	C	Q;	D	P
4.	A	P, Q, S;	B	P, Q, R, S;	C	P, Q, R;	D	P
5.	A	P;	B	R;	C	Q;	D	S
6.	A	S;	B	P;	C	Q;	D	R
7.	A	P, R;	B	S;	C	Q;	D	R

Assertion-Reason Type Questions

1. (C) **2.** (C) **3.** (D) **4.** (A) **5.** (B) **6.** (C) **7.** (A) **8.** (C) **9.** (D) **10.** (B)

11. (B) **12.** (D) **13.** (D) **14.** (A) **15.** (B) **16.** (A) **17.** (C) **18.** (A) **19.** (A) **20.** (B)

Subjective Problems

1. 6	**2.** 3	**3.** 3	**4.** 5	**5.** 6	**6.** 2	**7.** 6	**8.** 2	**9.** 8	**10.** 5
11. 2	**12.** 7	**13.** 3							

For More Material Join: @JEEAdvanced_2024

Hints and Solutions

Level 1

3. (c)

Rate of reaction $\frac{\text{Rate of disappearance/appearance}}{\text{Stoichiometric coefficient}}$

$\frac{d[A]}{dt}$ $\frac{1}{2}\frac{d[B]}{dt}$ 2.6 10^{-2}

$\frac{1}{2}\frac{d[B]}{dt}$

5.2 10^{-2} $\frac{d[B]}{dt}$

4. (d) Rate of reaction

$\frac{1}{2}\frac{d[A]}{dt}$ $\frac{d[B]}{dt}$ $\frac{1}{3}\frac{d[C]}{dt}$

or $\frac{k_1[A]^2}{2}$ $\frac{k_2[A]^2}{1}$ $\frac{k_3[A]^3}{3}$

$\frac{k_1}{2}$ k_2 $\frac{k_3}{3}$

5. (d) Rate of reaction $K[\text{conc.}]^n$

K $\frac{\text{Rate}}{[\text{conc.}]^n}$ $\frac{Ms^{-1}}{M^n}$ $\frac{\text{mol}\,L^{-1}\,s^{-1}}{\text{mol}^n L^{-n}}$

$\text{mol}^{1-n}L^{n-1}s^{-1}$

6. (d) As explain in Q. No. 5

7. (d) K_{eq} $\frac{k_f}{k_b}$ $\frac{\text{Conc. of products as per rate law}}{\text{Conc. of reactants as per rate law}}$, stoichiometry of reaction is required.

8. (c) K_{eq} at $1400\,K$ $\frac{0.29}{1.1 \quad 10^{-6}}$ 2.63 10^5; K_{eq} at $1500\,K$ 9.28 10^4; Value of K_{eq} is reduces with rise in temperature so reaction is exothermic.

9. (a) Arrhenious eq^n. Rate constant K Ae $^{-Eq/RT}$

10. (c) As explain in Q. 3

Rate of disappearance $\frac{d[BrO_3^-]}{dt}$

Rate of appearance $\frac{d[Br_2]}{dt}$

Rate of reaction $\frac{d[BrO_3^-]}{dt}$ $\frac{1}{3}\frac{d[Br_2]}{dt}$

11. (c) x kt

$t_{\text{completion}}$ $\frac{a}{k}$ $\frac{0.5}{0.1}$ 5 min

13. (d) Reaction mechanism required.

14. (d) $(a-x)$ ae^{-kt} or A_t A_0e^{-kt}

where A_t $(a-x)$ conc. of reactant remained after time t

A_0 a Initial concentration of reactant when t 0


17. (c) Rate of reaction decreases with respect to times in 1st order Rx^n.

18. (c) $t_{1/2}$ $\frac{1}{a^{n-1}}$;

where, a : initial concentration of reactant

n : order of reaction ($\because$ n 1)

19. (d) $a-x$ $a-kt$



where ,

a : initial conc. of reactant

x : conc. of reactant convert into product.

20. (a) $t_{1/2}$ $\frac{a}{2k}$ $k\,.\,a$; $\log(t_{1/2})$ $\log k$ $\log a$

21. (c) $t_{1/2}$ $\frac{1}{K\ a}$ for 2nd order ;

where a : initial conc. of reactant

22. (c) $t_{1/2}$ a ; zero order

23. (b) C $\frac{n}{V}$ $\frac{P}{RT}$ $\frac{dC}{dt}$ $\frac{1}{V}\frac{dn}{dt}$ $\frac{1}{RT}\frac{dP}{dt}$

24. (a) Rate of reaction $K[A_2]^a[B_2]^b$

0.04 $K(0.2)^a(0.2)^b$ ...(i)

0.04 $K(0.1)^a(0.4)^b$ ...(ii)

0.08 $K(0.2)^a(0.4)^b$ ...(iii)

$\frac{(i)}{(iii)}$ $\frac{1}{2}$ $\frac{1}{2}^b$; b 1

$\frac{(i)}{(ii)}$ 1 2^a $\frac{1}{2}^b$; a 1

For More Material Join: @JEEAdvanced_2024

25. (c) $R_0 \quad K[A_0]^2[B_0]$ ...(i)

$R_0 \quad K[1.5\,A_0]^2[3B_0]$

$R_0 \quad (1.5)^2 \quad 3[A_0]^2[B_0]$ ...(ii)

From eq. (i) and (ii)

$R_0 \quad 6.75\,R_0$

26. (c) As explain in Q. 25

27. (d) Rate of reaction $k[A]^2$

Rate of disappearance of $A \quad \frac{d[A]}{dt}$

Rate of appearance of $B \quad \frac{d[B]}{dt}$

Rate of reaction $\frac{d[A]}{dt} \quad \frac{d[B]}{dt} \quad k[A]^2$

Rate of appearance of B after double the conc. of A.

$\frac{d[B]}{dt} \quad k[2\,A]^2$

28. (c) Order w.r.t. $A \quad 1$; order w.r.t. $B \quad 1$

R.O.R. $\frac{1}{2}\frac{d(AB)}{dt} \quad k_r[A][B]$;

$\frac{1}{2} \quad (2.5 \quad 10^{-4}) \quad k_r\,(0.1)(0.1)$

$k_r \quad 1.25 \quad 10^{-2}$

29. (b) Rate of appearance of $AB(g)$,

$\frac{d(AB)}{dt} \quad k_b[A][B]$

$k_b \quad 2.5 \quad 10^{-2}$

30. (c) As explain in Q. 24

31. (d) For elementary reaction molecularity of reaction = order of reaction.

32. (a) As explain in Q. 24

33. (d) $r \quad k[A]^a[B]^b$

Order of reaction $\quad n \quad a \quad b \quad 2$

In elementary reaction, partial order should be integer.

So, $a \quad 1$ and $b \quad 1$

34. (a) $t_{1/2} \quad \frac{1}{(n \quad 1)k} \quad \frac{2^{n\ 1} \quad 1}{a^{n\ 1}}$

35. (b) As explain in Q. 18

36. (c) As explain in Q. 5

37. (a) Rate $\quad k[\text{conc.}]^0 \quad k$;

for zero order reaction.

38. (b) Conc. of water (solvent (l)) remains constant so rate is independent from conc. of H_2O.

39. (a) $CH_3COOC_2H_5(aq) \quad H_2O(l)$

$\xrightarrow{H} \quad CH_3COOH(aq) \quad C_2H_5OH(aq)$

$r \quad \frac{(k_{net})H_2SO_4}{(k_{net})HCl}$ R.O.R. in presence of H_2SO_4 > R.O.R. in presence of HCl

40. (c) $r_1 \quad k[A]^2[B]; r_2 \quad k[2A]^2[2B] \quad 8\,r_1$

41. (c) $[A]_t \quad [A] \quad kt \quad 1 \quad 0.001 \quad 10 \quad 60$

$0.4\,M$

$[B]_t \quad 0.001 \quad 10 \quad 60 \quad 0.6\,M$

42. (d) $t_{100\%} \quad \frac{a}{k} \quad \frac{1}{0.6}\,\text{min} \quad 100\,\text{sec.}$

Reaction will be finish in 100 sec., so concentration of B will reach maximum value 2 M after 100 sec.

43. (c) $x \quad kt$ so $t_{1/2} \quad \frac{a}{2k}$ and $T \quad \frac{a}{k}$;

$t_{1/2} \quad \frac{1}{2 \quad 0.001} \quad 50\,\text{sec}$ and $T \quad 100\,\text{sec}$

45. (c) $x \quad 0.1 \quad 0.08 \quad 0.02\,M$;

$k \quad \frac{x}{t} \quad \frac{0.02}{10} \quad 2 \quad 10^{-3}\,M\,\text{min}^{-1}$

$t_{1/2} \quad \frac{[A]_0}{2K} \quad \frac{0.1}{2 \quad 2 \quad 10^{-3}} \quad 25\,\text{min}$

$t_{\text{completion}} \quad 2 \quad t_{1/2} \quad 50\,\text{min.}$

46. (b) Elementary reaction,

$x(g) \quad y(g) \quad z(g)$

Rate $\quad k[x]$, first order

$k \quad \frac{1}{t}\ln\frac{[A_0]}{[A_t]}$

$\frac{\ln 2}{10\,\text{min}} \quad \frac{1}{t}\ln\frac{100}{10}$

$\frac{0.3}{10} \quad \frac{1}{t} \qquad t \quad 33\,\text{min.}$

47. (b) $K_1 \quad \frac{1}{5}\ln\frac{0.2}{0.1}$; $K_2 \quad \frac{1}{10}\ln\frac{0.2}{0.05}$; $K_1 \quad K_2$.

48. (b) $K \quad \frac{1}{100}\ln\frac{100}{25} \quad \frac{1}{t}\ln\frac{100}{12.5}$

49. (a) $K \quad \frac{0.693}{t_{1/2}} \quad \frac{0.693}{480} \quad 1.44 \quad 10^{-3}\,\text{s}^{-1}$

50. (b) $k \quad \frac{2.303}{t}\log\frac{C_{A_0}}{C_A}$;

$2.303 \quad 1 \quad 2.303\log\frac{C_{A_0}}{C_A}$

For More Material Join: @JEEAdvanced_2024

$\frac{C_{A_0}}{C_A} = 10$

$C_A = \frac{1}{10} = 0.1$

rate after 1 min $r_1 = k\,C_A$

$2.303 \quad 0.1 \quad 0.2303\ M\ \text{min}^{-1}$

51. (d) Reaction is second order; $k = 10^{-4}$ L/mol. min.

$$\frac{1}{3}\frac{d[A]}{dt} = k[A]^2$$

$$\frac{d[A]}{dt} = 10^{-4} \times 3 \times (0.5)^2$$

$$7.5 \times 10^{-5} M\ \text{min}^{-1}$$

$$\frac{7.5 \times 10^{-5}}{60} = 1.25 \times 10^{-6} \text{Ms}^{-1}$$

52. (c) $k = \frac{2.303}{t}\log\frac{a}{a-x}$

$$k = \frac{2.303}{32}\log\frac{100}{1} \qquad \frac{2.303}{16}$$

$$t = \frac{2.303 \times 16}{2.303}\log\frac{100}{0.1} \quad 48\ \text{min}$$

53. (c) $t_{3/4} = \frac{1}{K}\ln\frac{100}{\frac{1}{4}} \quad 100$

54. (d) $K_2 = \frac{1}{t}\ln\frac{100}{4}$ ...(i)

$K_1 = \frac{1}{t}\ln\frac{100}{50}$ ...(ii)

$$\frac{K_2}{K_1} = \frac{\ln 25}{\ln 2} \quad 4.65$$

55. (d) $k_I = \frac{0.693}{30}$

$$k_{II} = \frac{a}{2t_{1/2}} \quad \frac{2}{2\ t_{1/2}} \quad \frac{1}{t_{1/2}} \quad \frac{1}{30}$$

$$\frac{k_I}{k_{II}} = \frac{0.693}{30} \times \frac{30}{1} \quad 0.693$$

56. (a) $2N_2O_5(CCl_4) \rightarrow 2N_2O_4(CCl_4) + O_2(g)$

$t = 0$	a mol		0
$t = t$	$a - x$		$\frac{x}{2}$ mol
$t = \infty$	0		$\frac{a}{2}$ mol

$a = 100$

$x = 90$

$K = \frac{1}{t}\ln\frac{a}{a-x} \quad \frac{2.303}{500}$

57. (d) $A \rightarrow 3B$; $P_T = P_0 + 2x$

After time t, $P_0 - x \quad 3x$

$x = \frac{P_T - P_0}{2} \qquad k = \frac{1}{t}\ln\frac{P_0}{P_0 - x}$

After long time, $0 \quad 3P_0$

$$x = \frac{P_T - \frac{P}{3}}{2} \quad k = \frac{1}{t}\ln\frac{\frac{P}{3}}{\frac{P}{3} - \frac{3P_T - P}{6}}$$

$P = 3P_O \quad x = \frac{3P_T - P}{6} \quad k = \frac{1}{t}\ln\frac{P/3}{\frac{P}{2} - \frac{P_T}{2}}$

58. (a) $k = \frac{0.693}{2.1} = 0.33\ \text{hr}^{-1}$

Let t be the time for 90% decomposition,

so $a = 100\%$ $x = 90$ or $(a - x) = 10$

$$k = \frac{2.303}{t}\log\frac{a}{a-x}$$

$$t = \frac{2.303}{0.33}\log\frac{100}{10}; \quad t = 6.978\ \text{hr}$$

$n_{NH_4NO_3}$ taken $\frac{6.2}{62} = 0.1$;

as per given no. of moles of N_2O produced

$$0.1 \quad \frac{90}{100}$$

Vol. of N_2O produced at STP

$$0.09 \quad 22.4 \quad 2.016\ \text{L}$$

59. (a)

	A		$2B$		C
$t = 0$	P_i		0		0
$t = t$	$P_i - y$		$2y$		y

Total pressure $P_t = P_i - y + 2y + y = P_i + 2y$

$$\frac{P_t - P_i}{2} = y$$

$$K = \frac{2.303}{t}\log\frac{2P_i}{3P_i - P_t}$$

60. (c)

$k = \frac{2.303}{t}\log\frac{P_i}{P_t}$; $(CH_3)_2N_2 \rightarrow C_2H_6 + N_2$

$t = 0 \quad 200$; $t = 200 - x \quad x \quad x$

as per given $200 - x = 350 \quad x = 150$

For More Material Join: @JEEAdvanced_2024

$k \quad \frac{2.303}{t} \log \quad \frac{200}{200 \quad 150}$

$k \quad 5.77 \quad 10^{\ 4} \text{ sec}^{\ 1}$

61. (c) $k \quad \frac{1}{t} \ln \frac{r \quad r_0}{r \quad r_t} \quad \frac{1}{t} \ln \frac{r_0 \quad r}{r_t \quad t}$

$k \quad \frac{1}{70} \ln \frac{44 \quad 11}{16.5 \quad 11}$

$k \quad \frac{\ln 2}{70}$ when solution is optically inactive

$r_t \quad 0$

$k \quad \frac{1}{t} \ln \frac{r_o \quad r}{0 \quad r}$

$\frac{\ln 2}{70} \quad \frac{1}{t} \ln \frac{44 \quad 11}{11}$;

$\frac{0.7}{70} \quad \frac{1}{t} \quad 1.6 ; t \quad 160 \text{ min}$

62. (c) $\frac{(t_{1/2})_1}{(t_{1/2})_2} \quad \frac{a_2}{a_1}^{\ n\ 1}$; $n \quad 1 \quad \frac{\log \ t_{1/2\ 1}}{\log \ t_{1/2\ 2}}$

63. (c) $kt \quad \frac{1}{n \quad 1} \frac{1}{A_t^{\ n\ 1}} \quad \frac{1}{A_0^{\ n\ 1}}$

$kt_{0.5} \quad \frac{1}{n \quad 1} \frac{2^{n\ 1} \quad 1}{A_0^{\ n\ 1}}$ ...(1)

$kt_{0.875} \quad \frac{1}{n \quad 1} \frac{8^{n\ 1}}{A_0^{\ n\ 1}}$

$\frac{t_{0.875}}{t_{0.5}} \quad \frac{8^{n\ 1} \quad 1}{2^{n\ 1} \quad 1}$

64. (a) $kt \quad \ln \frac{\ 0}{\ t}$; $k \quad 60 \quad \ln \frac{80}{75}$

$k \quad \frac{1}{60} \ln \frac{80}{75}$

$k \quad 5 \quad 60 \quad \ln \frac{80}{r_t} ; 300 \quad \frac{1}{60}$.

$\ln \frac{80}{75} \quad \ln \frac{80}{r_t}$; $\ln \frac{80}{r_t} \quad 0.32$

$r_t \quad 58$

optical rotation observed after 5 hrs


$58 \quad 2 \quad 60$

65. (a) For zero order $t_{1/2} \quad [\text{conc.}]^0$

66. (c) $\frac{1}{2} \frac{dC_B}{dt} \quad k_1 . C_A ; \quad \frac{dC_B}{2\,t} \quad 2k_1 C_A$


$2 \quad 2 \quad 10^{\ 3} \quad 2 \quad 8 \quad 10^{\ 3} \, M \, s^{\ 1}$

67. (b)



E_a is always greater than H

68. (d) $Ea_2 \quad 58$



70. (c) First step is slow (require more activation energy) second step is fast (less activation energy) and overall reaction exothermic, so product energy level should be less as compared to reactants.

71. (d) Slow step require larger activation energy and product energy level is always more than reactant energy level in endothermic reaction.

72. (a) Rate of $Rx^n \quad K\,[\text{Conc.}]^n$

73. (a) $K \quad A \ e^{\ E_a/RT}$

$\ln K \quad \ln A \quad \frac{E_a}{RT}$

74. (b) $K \quad A \ e^{\ E_a/RT}$

Given $K \quad A \ e^{\ 1000}$

$\frac{E_a}{RT} \quad 1000$

75. (d) $k \quad \frac{k_1 . k_2}{k_3}^{2/3}$; $E \quad \frac{2}{3}[E_{a_1} \quad E_{a_2} \quad E_{a_3}]$

$\frac{2}{3}[180 \quad 80 \quad 50] \quad 140 \text{ kJ/mol}$

76. (c) $A_1 e^{\ E_{a_1}/RT} \quad A_2 e^{\ E_{a_2}/RT}$

$\frac{A_2}{A_1} \quad e^{(E_{a_2} \ E_{a_1})/RT}$

$10 \quad \text{Exp} \ \frac{600}{RT}$, $R \quad 2 \text{ cal/K-mol}$

$\ln 10 \quad \frac{600}{2T}$

$T \quad \frac{300}{2.303} \text{K}$

For More Material Join: @JEEAdvanced_2024

77. (a) Exothermic because of activation energy $E_b \quad E_f$

78. (d) $K \quad 1.2 \quad 10^{14} e^{\frac{25000}{RT}}$

$\ln K \quad \ln 1.2 \quad 10^{14} \quad \frac{25000}{RT}$

81. (b) More activation energy, slow reaction rate.

82. (b) $k_{1(300)} \quad \frac{0.693}{20}; \quad k_{2(320)} \quad \frac{0.693}{5}$

$\ln \frac{k_{2(320)}}{k_{1(300)}} \quad \frac{E_a}{R} \frac{1}{T_1} \quad \frac{1}{T_2}$

$E_a \quad \frac{2.303\ RT_1T_2}{(T_2 \quad T_{1)}} \log \frac{k_2}{k_1}$

$\frac{2.303 \quad 8.314}{20 \quad 1000} \quad 300 \quad 320 \log 4$

$= 55.14$ kJ/mol

83. (c) $\ln \frac{K_2}{K_{1\ 1}} \quad \ln 2 \quad \frac{E_a}{R} \frac{1}{280} \quad \frac{1}{290}$

$\ln \frac{K_2}{K_{1\ 2}} \quad \frac{E_a}{R} \frac{1}{290} \quad \frac{1}{300}$

$\frac{\ln 2}{\ln \frac{K_2}{K_{1\ 2}}} \quad \frac{10}{\frac{280 \quad 290}{10}}$

$\frac{290 \quad 300}{0.693} \quad \frac{300}{280}$

$\frac{K_2}{K_1} \quad 1.9$

84. (c) As explain in Q. 81

86. (c) $e^{\frac{E_a}{RT}} \quad 3.8 \quad \frac{10^{\ 16}}{100}$

$\frac{E_a}{RT} \quad \ln 3.8 \quad 10^{\ 18}$

$E_a \quad 100$ kJ/mol

87. (c) $k_1 \quad A.e^{\frac{83.314}{R \quad 500}}; \quad k_2 \quad A.e^{\frac{75}{R \quad 500}};$

$\frac{k_2}{k_1} \quad \exp \frac{8.314 \quad 1000}{8.314 \quad 500}$

$\exp (2) \quad 7.38$

88. (b) From slow reaction

Rate $K[A][B]$

89. (c) Rate of reaction $K_1[A][B_2]$ ...(i)

$K_2 \quad \frac{[A]^2}{[A_2]}$ ...(ii)

From Eq. (i) and (ii)

$R \quad K_1 \quad \sqrt{K_2[A_2]}[B_2] \quad K_1 \quad K_2^{1/2}[A_2]^{1/2}[B]$

$K[A_2]^{1/2}[B_2]$

$(\because K \quad K_1K_2^{1/2})$

91. (d) No significance of arrhenius equation in radioactivity.

92. (d) Due to charge on the and . Rays.

93. (d) ${}^{A}_{Z}X \quad {}^{A}_{Z\ 1}Y \quad {}^{0}_{1}$

${}^{A}_{Z\ 1}Y \quad {}^{A\ 4}_{Z\ 1}Z \quad {}^{4}_{2}$

${}^{A\ 4}_{Z\ 1}Z \quad {}^{A\ 4}_{Z\ 1}Z \quad {}^{0}_{0}$

High energy nucleous — Low energy nucleous — Energy

95. (b) ${}^{A}_{Z}x \quad {}^{A}_{Z\ 1}y \quad {}^{0}_{1}e$

Neutron $A \quad Z \quad A \quad (Z \quad 1)$

Proton $Z \quad Z \quad 1$

$\frac{n}{p} \quad \frac{A \quad Z}{Z} \quad \frac{A \quad Z \quad 1}{Z \quad 1}$

96. (c) As explain in Q. 95

104. (d) Isotones have same no. of neutron;

$17 \quad 9 \quad B \quad 8$

$B \quad 16$

Isobars have same mass no.

$A \quad B \quad 16$

no. of neutrons $16 \quad 8 = 8$

105. (d) ${}^{234}_{90}\text{Th} \quad {}^{206}_{82}\text{Pb} \quad x^4_2 \quad {}^{2} \quad y^0_1$

$234 \quad 206 \quad 4x \quad 0$

$x \quad 7$ -particle

$90 \quad 82 \quad 2x \quad y$

$y \quad 6$

106. (b) Same number of neutron.

107. (a) Same mass number

108. (c) Same difference between n and p

110. (b) ${}^{7}_{3}\text{Li} \quad {}^{1}_{1}\text{P} \quad {}^{4}_{2}x \quad {}^{4}_{2} \quad {}^{2}$

112. (c) As explain Q. No. 110

113. (d) ${}_{92}\text{U}^{235} \quad {}_{0}n^{1} \quad {}_{54}X^{139} \quad {}_{38}\text{Sr}^{194} \quad 3_0n^1$

114. (c) ${}^{255}_{101}\text{Md} \quad {}^{4}_{2} \quad 2^1_0n \quad {}^{257}_{103}\text{Lr}$

115. (b) ${}^{235}_{92}\text{U} \quad {}^{1}_{0}n \quad {}^{94}_{38}\text{Sr} \quad {}^{140}_{54}\text{Xe} \quad 2^1_0n$

116. (c) $E_{\text{total}} \quad 200 \frac{(3^5 \quad 1)}{3 \quad 1}$

$100 \quad 242 \quad 24200$ MeV

117. (c) ${}^{230}_{90}\text{Th} \quad {}^{1}_{1}\text{P} \quad 2^4_2\text{He} \quad {}^{223}_{87}\text{Fr}$

118. (c) On emission of one particle, formed nuclei occupied two position left in periodic table.

For More Material Join: @JEEAdvanced_2024

120.(c) On emission of positron atomic no. increases mass no. remain constant.

121.(d) $^{214}_{83}Bi$ $^{4}_{2}$ $^{210}_{81}A$ $^{0}_{1}$ $^{210}_{82}B$ $^{0}_{1}$ $^{210}_{83}C$ $^{0}_{1}$ $^{210}_{84}D$ $^{4}_{2}$ $^{206}_{82}E$

122. (c) A N

6.023 3.7 $10^{10} = 3.7$ $10^{4}N$;

N 6.023 10^{6} atoms

1 mole or 6.023 10^{23}atoms 100 g of X

6.023 10^{6} atoms

$$\frac{100}{6.023 \quad 10^{23}} \quad 6.023 \quad 10^{6} \quad 10^{15} \text{ g}$$

123.(b) A_1 $A_0 e^{\quad t_1}$ ...(i)

A_2 $A_0 e^{\quad t_2}$ ...(ii)

$$\frac{A_2}{A_1} \quad e^{(t_1 \quad t_2)} \qquad \text{...(iii)}$$

124. (d) Four half-lives (Total time n half-life so, n 4), hence 0.0625

125. (d) $\frac{0.693}{69.3}$ $10^{\ 2}$ min ^{1};

$$N \quad \frac{\frac{dN}{dt}}{} \quad \frac{100}{10^{\ 2}} \quad 10{,}000$$

126.(b) A_0 N_0

28 $\frac{\ln 2}{30} N_0$

N_0 1212

129. (b) Equal fraction decay in equal periods of time, fraction of sample remaining after

3 days $(0.9)^3$ 0.729

131. (d) $\frac{N_0}{N}$ $\frac{A_0}{A}$ 8;

t $\frac{1}{}\ln \frac{A_0}{A}$ t $\frac{5770}{0.693}\ln 8$

17310 year

133. (d) t for 20% left

$$t_1 \quad \frac{2.303}{} \log \frac{1}{1 \quad 0.8}$$

$$\frac{\ln 5}{}$$

t for 50% left

t_2 $\frac{1}{}\ln 2$

$$\frac{t_2}{t_1} \quad \frac{\frac{1}{}\ln 2}{\frac{1}{}\ln 5} \quad 0.43$$

t_2 0.43 t_1

134. (c) t $\frac{t0.5}{0.693}.\ln \frac{r_0}{r}$ $\frac{5770}{0.693}$ $\ln \frac{120}{30}$;

t $\frac{5770}{0.693}$ 2 0.693 11,540 years

135. (c) $^{40}_{19}K$ $^{40}_{20}Ca$ $^{0}_{1}e$

At present 2.5 $10^{\ 6}$ 7.5 $10^{\ 6}$ mole

moles of $^{40}_{19}K$ consumed 7.5 $10^{\ 6}$ mole

initial mole of ^{40}K was (7.5 2.5) $10^{\ 6}$ mole

$$t \quad \frac{1}{}\ln \frac{10 \quad 10^{\ 6}}{2.5 \quad 10^{\ 6}} \quad \frac{1.3 \quad 10^{9}}{\ln 2} \quad \ln(4)$$

1.3 10^{9} 2 2.6 10^{9} yr

136.(d) t_{avg} $\frac{1}{}$ $\frac{14}{0.693}$

137.(c) t $|t_1$ $3t_{1/2}|$ 200

138. (b) Total time $nt_{1/2}$; n 5; $\frac{\text{Initial activity}}{2^n}$

Initial activity 0.01 $2^5 = 0.32$ C_i

139. (b) t $\ln \frac{A_0}{A}$

$$\frac{0.693}{t_{1/2}} \quad 200 \quad \ln \frac{2140}{535}$$

$t_{1/2}$ 100 min.

141. (c) $\frac{1}{10}\ln \frac{100}{100 \quad 25}$

t $\frac{1}{}\ln \frac{N_0}{N}$

$$t \quad \frac{10}{\ln \frac{4}{3}} \quad \ln \frac{4 \quad 10^{20}}{10^{20}}$$

t 48.19

142. (b) $t_{1/2}$ $\frac{t}{4}$; $t_{1/2}$ $T \ln 2$

so $\frac{t}{4}$ $T \ln 2$; t $4t \ln 2$

143. (d) W_0 initial mass, W mass. after 20 days

$$A \quad \frac{2.303}{t} \log \frac{W_0}{W_A} \ ;$$

For More Material Join: @JEEAdvanced_2024

$_B \quad \frac{2.303}{t} \log \frac{W_0}{W_B}$

$_B \quad \frac{2.303}{t} \log \frac{W_B}{W_A}$

$(t_{1/2})_B \quad 1.10$ day

144. (c) Given $(n_0)_B \quad 8 \quad (n_0)_A; (n)_A \quad 2 \quad (n)_B$

$_{A \quad B} \quad \frac{2.303}{t} \log \frac{(N_A)_0}{(N_A)} \quad \frac{(N_B)}{(N_B)_0}$;

$$t \quad \frac{\ln \frac{1}{16}}{\frac{0.693}{50} \quad \frac{0.693}{10}}$$

$= 50$ min

145. (b) as per given $\frac{dN}{dt}$ N for max. no. of nuclei;

$\frac{dN}{dt} \quad 0; \qquad N, \; N \quad —$

146. (c) $\frac{\text{Atoms of Sr}}{\text{Atoms of Rb}} \quad 0.05$

or $\frac{\text{Atoms of (Sr + Rb)}}{\text{atoms of Rb}} \quad 1.05$

so, $\frac{\text{Initial no. of atoms of Rb}}{\text{Present no. of atoms of Rb}} \quad 1.05$

$t \quad \frac{2.303}{} \log (n_0/n)$

$\frac{2.303}{0.693} \quad 4.7 \quad 10^{10} \log (1.05)$

$= 3.28 \quad 10^9$ year

147. (a) $_A N_A \quad _B N_B$

($\because$ rate of disintegration are same)

$\frac{N_A}{N_B} \quad \frac{_B}{_A}$

$\frac{2}{1}$

148. (b) $N \quad N_0 . e^{\;t}$; where N Parent remaining (p)

and N_0 Initial parent Parent remaining (p) + daughter formed (d)

$p \quad (p \quad d) . e^{\;t}$ or $\ln \frac{(p \quad d)}{p} \quad t$

$t \quad \frac{1}{} \ln \; 1 \quad \frac{d}{p}$

149. (b) $_1 \quad _2; \quad \frac{0.693}{22}$ and $\frac{_1}{_2} \quad \frac{2}{98}$

$_1 \quad 0.00063$ year $^{1}; \quad _2 \quad 0.03087$ year ^{1}

150. (b) $^{218}_{84}Po \quad ^{214}_{82}Pb + ^{4}_{2}He$

$^{214}_{83}Bi + ^{0}_{-1}e$

Pb^{214} to reach max. no. of nuclei

$t_{max.} \quad \frac{1}{1 \quad 2} \ln \frac{_1}{_2}$

$= 247.5$ sec

where $_1 \quad \frac{0.693}{183}; \quad _2 \quad \frac{0.693}{161}$

Level 2

1. (d) $K_{eq} \quad \frac{k_f}{k_b} \quad \frac{[CH_3]^2}{[C_2H_6]}$

$[CH_3] \quad \frac{10^{\;4}}{10} \quad 10^{\;5} \; M$

$\frac{1.57 \quad 10^{\;3}}{k_b} \quad \frac{(10^{\;5})^2}{1}$

$k_b \quad 1.57 \quad 10^7$ L mol ^{1} s ^{1}

3. (c) $r_1 \quad k[0.01]^a [0.01]^b \quad 6.93 \quad 10^{\;6}$ …(i)

$r_2 \quad k[0.02]^a [0.01]^b \quad 1.386 \quad 10^{\;5}$ …(ii)

$r_3 \quad k[00.2]^a [0.02]^b \quad 1.386 \quad 10^{\;5}$ …(iii)

From data $a \quad 1; \; b \quad 0;$

overall order $1; k \quad 6.93 \quad 10^{\;4}$ sec ^{1}

$6.93 \quad 10^{\;4} \quad \frac{1}{50 \quad 60} \ln \frac{A_o}{A_t};$

when $[A]_0 \quad 0.5$

$A_t \quad 0.0625$

rate of reaction $6.93 \quad 10^{\;4} \quad 0.0625$

$4.33 \quad 10^{\;5}$ Ms ^{1}

4. (c) $\frac{r_2}{r_1} \quad \frac{P_{A_2} . P_{B_2}^2}{P_{A_1} . P_{B_1}^2} \quad \frac{0.1 \quad (0.4)^2}{0.4 \quad 1^2}$

$\frac{1}{25}$

6. (c) $t_1 \quad \frac{(t_{1/2})_1}{0.693} \ln \frac{1}{(1 \quad (1/4)}$

$t_2 \quad \frac{(t_{1/2})_2}{0.693} \ln \frac{1}{1 \quad 3/4}$

$\frac{t_1}{t_2} \quad \frac{8}{1} \quad \frac{\ln (4/3)}{\ln (4)}$

$1 : 0.602$

7. (b) $A \qquad B \qquad C \qquad D$

$t \quad 0 \quad 1 \qquad 1$

$t \quad 1 \quad x \quad 1 \quad x \qquad x \qquad x$

For More Material Join: @JEEAdvanced_2024

$$r \quad k[A]^{1/2}[B]^{1/2}$$

$$\frac{dx}{dt} \quad k(1 \quad x)^{1/2}(1 \quad x)^{1/2}$$

or $$\frac{dx}{dt} \quad k(1 \quad x)$$

$$t \quad \frac{1}{k}\ln \quad \frac{1}{1 \quad x} ;$$

$$t \quad \frac{2.303}{2.303 \quad 10^{\ 2}}\log \quad \frac{1}{0.1}$$

$$=100 \text{ sec}$$

8. (a)

(a) True $[B]_t \quad k_{\mathrm{I}}\, t ; 0.25 \quad \frac{1}{\sqrt{3}} t ; t \quad 0.25\sqrt{3}$

$[D]_t \quad k_{\mathrm{II}} t \quad \sqrt{3} \quad 0.25 \quad \sqrt{3} \quad 0.75\, M$

(b) False if $[C] \quad [A]$ then at that time $[B] \quad [D]$

(c) False $t_{100\%} \quad \frac{a}{k}$ (for zero order)

$$\frac{(t_{100\%})_{\mathrm{I}}}{(t_{100\%})} \quad \frac{a_{\mathrm{I}}}{a_{\mathrm{II}}} \quad \frac{k_{\mathrm{II}}}{k_{\mathrm{I}}} \quad \frac{0.5}{1} \quad \frac{\sqrt{3}}{1/\sqrt{3}} \quad \frac{3}{2}$$

(d) $[A]_t \quad [A]_0 \quad k_{\mathrm{I}} t$ or $[A]_t \quad 0.5 \quad \frac{1}{\sqrt{3}} t$

$[C]_t \quad [C]_0 \quad k_{\mathrm{II}} t$ or $[C]_t \quad 1 \quad \sqrt{3} t$

if $[A] \quad [C]_t$

i.e. $0.5 \quad \frac{1}{\sqrt{3}} t \quad 1 \quad \sqrt{3} t$ or $(\sqrt{3} \quad \frac{1}{\sqrt{3}}) t \quad 0.5$

$$t \quad \frac{\sqrt{3}}{4} \text{ min.}$$

9. (a) $$t_{1/2} \quad \frac{0.693}{k}$$

$$\frac{0.693}{2.772 \quad 10^{\ 3}}$$

$$= 250 \text{ sec}$$

Since volume is changing therefore half-life should be defined on the basis of moles, so moles of A remaining after 250 sec. Also final volume of the container

$$V_f \quad \frac{n_f}{n_i} \quad V_i \quad \frac{0.05 + 0.15}{0.1} \quad 2$$

$$= 4 \text{ litre}$$

Final concentration of $A \quad \frac{0.05}{4}$

$$= 0.0125\, M$$

10. (c)

	A	$2B$	C
$t \quad 0$	P_i	0	0
t	$P_i \quad x$	$2x$	x
t	0	$2P_i$	P_i

$P \quad 3P_i$ or $P_i \quad \frac{P}{3} ; P_i \quad 2x \quad P_t$

$$x \quad \frac{P_t \quad P_i}{2}$$

As we know $k \quad \frac{2.303}{t}\log \quad \frac{P_i}{P_i \quad x}$

so $k \quad \frac{2.303}{t}\log \quad \frac{2P}{3(P \quad P_t)}$

11. (d) Let n is the moles of reagent 'R' when R is reacted with A at time $t \quad 0$

	A	B	C
$t \quad 0$	n	0	0
at t	$n \quad x$	$2x$	$3x$
at t	0	$2n$	$3n$

$$5n \quad n_2 \quad n \quad \frac{n_2}{5}$$

$$n \quad 4x \quad n_1 \quad x \quad \frac{n_1 \quad n}{4}$$

$$k \quad \frac{2.303}{t}\log \quad \frac{n}{n \quad x}$$

so $$k \quad \frac{1}{t}\ln \quad \frac{4n_2}{5(n_2 \quad n_1)}$$

12. (b)

	$A(g)$	$2B(g)$	$C(g)$
Let initial pressure	P_0	0	0
After 10 min.	$(P_0 \quad x)$	$2x$	x
After long time (t)	0	$2P_0$	P_0

as per given $(P_0 \quad x) \quad 2x \quad x$ vapour pressure of $H_2O \quad 188$

$$P_0 \quad 2x \quad 160 \text{ and } 3P_0 \quad 28 \quad 388$$

so, $$P_0 \quad 120 \text{ and } x \quad 20 \text{ torr}$$

$$k \quad \frac{1}{t}\ln \quad \frac{P_0}{P_0 \quad x}$$

$$\frac{1}{10}\ln \quad \frac{120}{100} \quad \frac{1}{10} \quad (\ln 4 \quad \ln 3 \quad \ln 10)$$

$$0.02 \text{ min}^{\ 1} \quad 1.2 \text{ hr}^{\ 1}$$

13. (b) $$k \quad \frac{2.303}{t}\log \quad \frac{r \quad r_0}{r \quad r_t}$$

$$\frac{2.303}{30}\log \quad \frac{11 \quad 34}{11 \quad 19}$$

$$= 1.35 \quad 10^{\ 2} \text{ min}^{\ 1}$$

when solution is optically inactive $r_t \quad 0$;

For More Material Join: @JEEAdvanced_2024

$$t \quad \frac{1}{k}\ln \frac{45}{11} \quad 103.7 \text{ min}$$

14. (c) Overall rate constant

$$k \quad k_1 \quad k_2 \quad k_3 \quad 6.93 \quad 10^{-3}$$

$$t_{1/2} \quad \frac{0.693}{6.93 \quad 10^{-3}} \quad 100 \text{ sec};$$

After half-life, $P_B \quad P_C \quad P_D \quad 4$ atm

$$\frac{P_B}{P_B \quad P_C \quad P_D} \quad \frac{k_1}{k_1 \quad k_2 \quad k_3} \quad \frac{200}{693}$$

$$P_B \quad 4 \quad \frac{200}{693}$$

$$= 1.154 \text{ atm}$$

15. (d) $A \xrightarrow{k_1} 2B$; $A \xrightarrow{k_2} C$

$a_0 \quad x \quad y \qquad 2x \qquad a_0 \quad x \quad y \qquad y$

$$\frac{d[A]}{dt} \quad (k_1 \quad k_2)[A]; \quad \frac{[B]}{[C]} \quad \frac{2k_1}{k_2} \quad \frac{2x}{y}$$

$$\frac{k_1}{k_2} \quad \frac{x}{y}$$

$\because \quad k_1 \quad k_2 \quad x \quad y$

$$(k_1 \quad k_2)t \quad \ln \frac{A_0}{A_t}$$

$$2 \quad 6.93 \quad 10^{-3} \quad 50 \quad \ln \frac{A_0}{A_t}$$

$$[A]_t \quad \frac{[A_0]}{2} \quad a_0 \quad x \quad y \quad a_0/2 \quad (\because x \quad y)$$

$x \quad a_0/4$. Total moles in container

$a_0 \quad x \quad y \quad 2x \quad y \quad a_0 \quad x \quad 1.25\, a_0$

% increase in moles 25, So final pressure

$2 \quad 1.25 = 2.5$ atm

16. (d) 1 mole of A will form 2 moles of B and C. After completion of reaction when 75% A converted into B and C then total no. of moles $0.5 + 2 \quad 1.5 = 3.5$ moles

17. (d) $cis\text{-}X \underset{k_b}{\overset{k_f}{\rightleftharpoons}} trans\text{-}X$

Initial $a \qquad 0$; $K_{(aq)} \quad \frac{k_f}{k_b}$; $k_b \quad \frac{3 \quad 10^{-4}}{0.1} \quad 3 \quad 10^{-3}$

at time $t \quad a \quad x \qquad x$

at eqm $a \quad x_e \qquad x_e$

As we know $(k_f \quad k_b) \quad \frac{1}{t}\ln \frac{x_e}{x_e \quad x}$

given $x \quad \frac{x_e}{2}$

$$(k_f \quad k_b) \quad \frac{1}{2}\ln 2$$

or $(3 \quad 10^{-3} \quad 3 \quad 10^{-4}) \quad \frac{0.693}{t}$

$$t \quad 210 \text{ sec}$$

18. (b) $k \quad A\, e^{-E_a/RT}$; $A\, e^{-\frac{E_a}{RT_1}} \quad A\, e^{-\frac{(E_a \quad x)}{RT_2}}$

$$\frac{E_a}{500} \quad \frac{(E_a \quad 16)}{400} \quad E_a \quad 80 \text{ kJ/mol}$$

19. (c) For parallel reactions

$$E_a \quad \frac{k_1 E_{a_1} \quad k_2 E_{a_2}}{k_1 \quad k_2}$$

$$\frac{10^{-2} \quad 100 \quad 4 \quad 10^{-2} \quad 120}{10^{-2} \quad 4 \quad 10^{-2}}$$

$$E_a \quad 116 \text{ kJ/mol}$$

20. (c) $k \quad A\, e^{-E_a/(RT)}$

Effective overall energy of activation

$$E_a \quad E_{a(2)} \quad E_{a(3)} \quad \frac{1}{2}E_{a(1)} \quad \frac{1}{2}E_{a(5)}$$

$$60 \quad 50 \quad \frac{1}{2} \quad 40 \quad \frac{1}{2} \quad 10$$

$$25 \text{ kJ/mol}$$

21. (d) $A_1\, e^{-E_{a_1}/RT} \quad A_2\, e^{-E_{a_2}/RT}$;

$$\frac{A_2}{A_1} \quad e^{(Ea_2 \quad Ea_1)/RT}$$

$$10^2 \quad \text{Exp} \quad \frac{1200}{RT}; 2\ln 10 \quad \frac{1200}{2T}$$

$$T \quad \frac{600}{4.606} \text{ K}$$

22. (b) $\ln \frac{k_1}{k_1} \quad \frac{E_1}{R} \quad \frac{1}{T_1} \quad \frac{1}{T_2}$ …(i)

$$\ln \frac{k_2}{k_2} \quad \frac{E_2}{R} \quad \frac{1}{T_1} \quad \frac{1}{T_2} \qquad \ldots(ii)$$

Solving we get (ii)–(i),

$$\ln \frac{k_2}{k_2} \quad \ln \frac{k_1}{k_1} \quad \frac{E_2 \quad E_1}{R} \quad \frac{1}{T_1} \quad \frac{1}{T_2}$$

(for equimolar formation of B and C, $k_2 \quad k_1$)

$$\ln \frac{k_1}{k_2} \quad \frac{8314}{8.314} \quad \frac{(T_2 \quad 300)}{300 \quad T_2}$$

$$\ln 2 \quad (1000) \quad \frac{T_2 \quad 300}{300 \quad T_2}$$

$$T_2 \quad 378.74 \text{ K}$$

For More Material Join: @JEEAdvanced_2024

24. (b) $A + B \underset{\text{fast}}{\overset{\text{slow}}{\rightleftharpoons}} IAB$; So $E_{a(f)}$ is high and $E_{a(b)}$ is low. $k_1 \quad k_2$; So, E_a for this step is very high and for next step is low and overall reaction is exothermic.

25. (a) Activation energy

$E_I \quad E_{II} \quad E_V \quad E_{VI}$

$E_{1(f)} \quad E_{1(b)} \quad E_{2(b)} \quad E_{3(f)} \quad E_{3(b)}$

$E_{4(f)} \quad E_{5(f)} \quad E_{6(b)}$

$E_{2(f)} \quad E_{4(b)}$ (slow steps so larger)

26. (c) $e^{\frac{E_a}{RT}} \quad 100 \quad 3.8 \quad 10^{16}$; $e^{\frac{E_a}{RT}} \quad 3.8 \quad 10^{18}$

$\frac{E_a}{RT} \quad \ln (3.8 \quad 10^{18})$;

$R \quad 8.314$ and $T \quad 300$

$E_a \quad 100$ kJ/mol

27. (b) $t \quad \ln \frac{A_0}{A}$

$\frac{0.693}{30} \quad 90 \quad \ln \frac{A_0}{A} \quad A_0 \quad A \quad 8$

$N_0 \quad \frac{A_0}{} \quad \frac{800}{0.693} \quad 30 \quad 3.46 \quad 10^4$ atoms

28. (a) at time $X \quad Y$; $\frac{n_0 \quad x}{x} \quad \frac{1}{7}$; $n_0 \quad \frac{8x}{7}$;

$\frac{0.693}{6.93 \quad 10^9} \quad 10^{-10}$ years^{-1}

$t \quad \frac{2.303}{} \log \frac{w_0}{w_0 \quad x}$

$\frac{2.303}{10^{-10}} \log (8)$

$= 2.079 \quad 10^{10}$ years

29. (c) $_{net}$ or $\quad _1 \quad 2 \quad _2 \quad 1.8 \quad 10^{-3} \quad 2 \quad 10^{-3}$

$2 \quad 10^{-2}$

Average life (T_{av}) $\quad \frac{1}{-}$

$\frac{1}{2 \quad 10^{-2}} \quad 50$ sec

30. (c) $_{83}Bi^{211} \quad _{81}Tl^{207} + {}_2He^4$; total time n half-life

moles of substance left after n halves

$\frac{\text{initial moles}}{2^n} \quad \frac{2}{2^4} \quad 0.125$

moles of He produced

$2 \quad 0.125 \quad 1.875$

Pressure developed due to

He $\quad \frac{1.875 \quad 0.0821 \quad 300}{100}$

$= 0.4618$ atm

31. (d) Let initial activities of A and B are A_0 and B_0

($\because$ after 2 half-lives of activity of A will remain $\frac{A_0}{4}$)

$A_0 \quad B_0 \quad 8000$ and also $\frac{A_0}{4} \quad B_0 \quad 3500$

(we can assume that activity of B remains constant due to larger half-life)

so $\frac{3A_0}{4} \quad 4500$; $A_0 \quad 6000$; $B_0 \quad 2000$;

$\frac{A_0}{B_0} \quad \frac{6000}{2000} \quad \frac{3}{1}$

32. (a) No change in activity of sample during establishment of circulatory equilibrium.

Let volume of blood is V mL, so total vol. $(V \quad 1)$ mL after injection of sample.

2 mL sample has activity of 10 dpm, so $(V \quad 1)$ mL sample has activity of $\frac{10}{2} \quad (V \quad 1)$

Since rate is constant so

$\frac{10}{2} \quad (V \quad 1) \quad 1000$; $V \quad 199$ mL

33. (c) $\frac{A_{0(X)}}{A_{0(Y)}} \quad \frac{4}{1}$; $\frac{A_x}{A_y} \quad 1$,

$_y \quad _x \quad \frac{1}{t} \ln \frac{(A_0)_y}{(A_0)_x} \quad \frac{A_x}{A_y}$

$(_y \quad _x)t \quad \ln \frac{1}{4}$; $(t_{1/2})_y \quad 30$ min.

34. (a) Corrected C^{14} content :

Let initial C^{14} count $\quad A_0$;

$A_0 \quad A_0/10 \quad 15.4$; $A_0 \quad 14$

$t \quad \frac{1}{-} \log \frac{14}{7} \quad 5.770 \quad 10^3$ years; $t \quad \frac{\ln 2}{}$

$t_{1/2}$; $t \quad 5770$ years

35. (a) For U^{238}; $_1t \quad \ln \frac{n_0}{n_1}$;

For U^{235}; $_2t \quad \ln \frac{n_0}{n_2}$

$(_1 \quad _2)t \quad \ln \frac{n_0}{n_1} \quad \frac{n_2}{n_0}$;

For More Material Join: @JEEAdvanced_2024

$\ln 2 \quad \frac{1}{0.5 \quad 10^9} \quad \frac{1}{4.5 \quad 10^9} \quad t \quad \ln(140)$

$t \quad 4.02 \quad 10^9$ year

Level 3

Passage-1

1. (b) For max. con. of B;

$\frac{d[B]}{dt} \quad 0$; so $t_{max} \quad \frac{1}{k_1 \quad k_2} \ln \frac{k_1}{k_2}$

2. (b) $k_1 \quad k_2$

So rate of appearance of B is much lesser than rate of disappearance of B.

Passage-2

1. (c) $e^{-E_a/RT}$ is fraction of molecules having energy greater than activation energy (E_a), it is unit less. So A has same unit of K.

for 1st order reaction unit of K is s^{-1}.

2. (b) Answer is given in (1)

$x \quad e^{-E_a/RT}$

$\ln x \quad \frac{E_a}{RT}$

3. (b) $\ln \frac{K_2}{K_1} \quad \frac{E_a}{R} \quad \frac{1}{T_1} \quad \frac{1}{T_2}$

$\ln 2 \quad \frac{E_a}{2 \text{ cal}} \quad \frac{1}{290} \quad \frac{1}{300}$

$E_a \quad \frac{2.303 \quad 0.3 \quad 2 \quad 290 \quad 300}{10}$ cal

12 K Cal

Passage-3

1. (c) $E \quad \frac{nhC}{}$

$7.2 \quad 10^{-3}$ J / s $\quad \frac{n \quad 6.6 \quad 10^{-34} \quad 3 \quad 10^8}{330 \quad 10^{-9}}$

No. of photon

$\frac{7.2 \quad 10^{-3} \quad 330 \quad 10^{-9}}{6.6 \quad 10^{-34} \quad 3 \quad 10^8} \quad 120 \quad 10^{14}$

Mole of photon $\quad \frac{120 \quad 10^{14}}{6 \quad 10^{23}} \quad 20 \quad 10^{-9}$

Mole of CO formed

$0.4 \quad 20 \quad 10^{-9} \quad 8 \quad 10^{-9}$ mol/s

2. (c) Rate of formation of $C_2H_6 \quad 0.8 \quad 20 \quad 10^{-9}$

$16 \quad 10^{-9}$ mol/s

Passage-4

1. (c) $\quad \frac{1}{t} \ln \frac{N_0}{N}$

$\frac{\ln 2}{t_{1/2}} \quad \frac{1}{t} \ln \frac{N_0}{\frac{N_0}{4}}$

$\frac{\ln 2}{2.5 \quad 10^5} \quad \frac{1}{t} \ln 4$

$t \quad 2 \quad 2.5 \quad 10^5 \quad 5 \quad 10^5$ years.

2. (a) $\quad \frac{N}{N_0} \quad \frac{1}{x}^n$; where n no. of halves;

$\frac{N}{N_0} \quad \frac{1}{10} \quad \frac{1}{x}^n$

$n \quad 4$

Total time $\quad n \quad t_{1/2}$

$4740 \quad 4 \quad t_{1/2}$;

$t_{1/2} \quad 1185$ years

3. (b) $\quad \frac{dN}{dt} \quad N \quad \frac{0.693}{t_{1/2}} \quad n \quad N_A$

$\frac{dN}{dt} \quad \frac{0.693}{24{,}000 \quad 365 \quad 24 \quad 60 \quad 60}$

$1 \quad 6.02 \quad 10^{23}$ dps or 14.9 Ci

Passage-5

1. (b) $r \quad R_0 A^{1/3}$

Volume of nucles $(4/3) \quad r^3$

$\frac{4}{3} \quad (1.5 \quad 10^{-15})^3 \quad A \text{ m}^3$

density $d \quad \frac{m}{V} \quad \frac{A \quad 1.66 \quad 10^{-27} \text{ kg}}{4/3 \quad (1.5 \quad 10^{-15})^3 \quad A}$

$1.17 \quad 10^{17}$ kg/m^3

2. (b) $\quad d_C \quad \frac{\text{Mass}}{\text{Volume}} \quad \frac{12 \quad 1.66 \quad 10^{-27}}{(4/3) \quad 3.14 \quad (3 \quad 10^{-15})^3}$

$1.76 \quad 10^{17}$ kg/m^3

$d_{H_2O} \quad 1000$ kg/m^3

$\frac{d_C}{d_{H_2O}} \quad \frac{1.76 \quad 10^{17}}{1{,}000} \quad 1.76 \quad 10^{14}$

One or More Answers is/are Correct

2. (A) $K \quad Ae^{-E_a/RT}$ as T , K rate of reaction

For More Material Join: @JEEAdvanced_2024

(B) Rate of reaction $k[A]^0$ K constant

(C) As surface area No. of molecules will be more to react per unit time, more surface site available for the reaction.

(D) Average rate defined for macro-scopic time and instantaneous rate defined for micro-scopic time.

3. For an elementary reaction aA bB cC dD rate law is always r $K[A]^a[B]^b$ but not *vice versa.*

For a complex reaction aA bB cC dD rate law may or may not be r $K[A]^a[B]^b$.

4. (A) In a rate law reactant, product or catalyst may be appeared.

(B) We can not

(C) In an elementary reaction only positive integer stoichiometric coefficient appears.

(D) A zero order reaction is always complex

5. (A) K $Ae^{-E_a/RT}$

As T , K so if T , then K A

or as Ea , K so if E_a 0, then K A

(B) Catalyst does not change H of reaction.

(C) A negative catalyst decrease rate of reaction by increasing activation energy.

6. A B C

(A) If it is an elementary reaction then it will be bimolecular reaction.

(B) It may be exothermic eq. $NH_3(g)$ $HCl(g)$ $NH_4Cl(s)$

(C) It may be heterogeneous

(D) It may be photochemical reaction

7. $A(g)$ $2B(g)$ $C(g)$

t 0 400

t 20 400 x $2x$ x

So 400 x $2x$ x 1000

x 300 mmHg

K $\frac{1}{20}\ln\frac{400}{100}$ $\frac{2\ln 2}{20}$ 0.0693 min $^{-1}$

$t_{1/2}$ $\frac{\ln 2}{K}$ $\frac{0.693}{0.0693}$ 10 min

$\frac{30}{10}$ $\ln 2$ $\ln\frac{400}{400\ x}$

8 $\frac{400}{400\ x}$ $400\ x$ 50

x 350 mmHg

Partial pressure of C after 30 min 350 mmHg

Total pressure after 30 min

400 $2x$ 400 700 1100 mmHg

9. (a,b,c,d)

k $\frac{Be}{Ae}$ $\frac{1.6}{0.4}$ 4 ; k $4\frac{k_1}{k_2}$ $\frac{4\ 10^{-2}}{k_2}$

k_2 10^{-2}

$t_{0.5x_e}$ $\frac{1}{k\ k_2}\ln\frac{x_e}{x_e\ x}$

$\frac{1}{5\ 10^{-2}}\ln 2$ $\frac{70}{2}$; $t_{0.5x_e}$ 14 min ;

$t_{0.1a}$ $\frac{1}{5\ 10^{-2}}\ln\frac{x_e}{x_e\ x}$ $\frac{1}{5\ 10^{-2}}\ln\frac{1.6}{1.6\ 0.2}$

$\frac{100}{5}\ln\frac{16}{14}$ 20 0.14 2.8

10. (A) -particles are just nucleus of helium ($_2^4He^{2-}$)

(C) $_Z^AX$ $_{Z\ 1}^AY$ $_1e^0$

so X and Y are isobars.

11. (A) $_{92}U^{235}$ $_0n^1$ $_{56}Ba^{140}$ $2\ _0n^1$ $_Z^AX$

Atomic no. 92 0 56 (2 0) Z

Z 36

Mass no. 235 1 140 (2 1) A

A 236 172 94

So X is $_{36}^{94}Kr$

(B) $_{11}Na^{23}$ $_Z^AX$ $_{12}Mg^{23}$ $_0n^1$

Atomic no. 11 Z 12 0 Z 1

Mass no. 23 A 23 1 A 1

X $_1^1H$ Proton

(C) and (D) both are correct.

12. (A) SI unit of radioactivity is Bq

1 Bq 1 dps

(B) 1 Ci 3.7 10^{10} dps or Bq

13. Similar as question 11.

14. $_Z^MA$ $_{Z\ 2}^{M\ 4}B$ $_{Z\ 1}^{M\ 4}C$ $_Z^{M\ 4}D$

15. (a, b)

$_1^1P$ $_1e^0$ $_0^1n$ X-rays

17. (a,b)

$_1^1P$ $_0^1n$ $_1^0e$ (positron) ; $_0^1n$ $_1^1P$ $_1^0e$ (Beta)

Activity N, Decay constant does not depend upon temperature.

By the emission of -particle, isodiaphers are formed.

By the emission of -particle, isobars are formed.

For More Material Join: @JEEAdvanced_2024

Match the Column

1. (A) K $Ae^{-E_a/RT}$, unit of K unit of A

(B) K $\dfrac{\text{Rate}}{(\text{Concentration})^n}$ $\dfrac{M/s}{(M)^n}$ $M^{1-n}s^{-1}$

n 0 Unit of K Ms^{-1}

(C) n 1 Unit of K s^{-1}

(D) n 2 Unit of K $M^{-1}s^{-1}$

2. (A) Molecularity of a reaction never be zero.

(B) Order of reaction may be 0, 1, 2, 3 fraction

(C) H_2O_2 H_2O $1/2\,O_2$ if its an elementary then it is 1st order reaction.

(D) It is zero order reaction because reaction rate is depend on due to absorption of photon.

3. (A) C_t C_0 Kt slope K

(B) $\log C_t$ $\log C_0$ $\dfrac{K\ t}{2.303}$; slope $K/2.303$

(C) $\dfrac{dC}{dt}$ $K[C]$; slope zero

(D) $\ln\dfrac{dc}{dt}$ $\ln K$ $\ln C$; slope 1

4. (A) $\dfrac{d[A]}{dt}$ *vs* $[A]$, Let the order or reaction w.r.t. A is x then for non zero slope $[A]^x$, and x 0

(B) K $Ae^{-E_a/RT}$

$\ln K$ $\ln A$ $\dfrac{E_a}{RT}$

It is applicable for all possible order in this slope 0

(C) $t_{1/2}$ $[A]_0$ it is not possible for 1st order.

(D) $\dfrac{d[A]}{dt}$ *vs* $[A]^2$, $[A]^2$ second order.

5. (A) Concentration of product increases w.r.t time.

(B) Concentration of reactant decreases w.r.t time.

(D) K $Ae^{-E_a/RT}$

$\ln K$ $\ln A$ $\dfrac{E_a}{RT}$ slope $\dfrac{E_a}{R}$

6. (A) Isotone atoms having same number of neutrons (*i.e.* value of mass number (A) – atomic number (Z).

(B) Isobare Atoms has same mass no. and different atomic no.

(C) Isotope Atoms has same atomic no. and difference mass no.

(D) Isodiaphers Same difference between neutrons and proton.

7. (A) ${}^{A}_{Z}X$ ${}^{A}_{Z-2}Y$

(B) ${}^{A}_{Z}X$ ${}^{0}_{-1}$ ${}^{A}_{Z+1}Y$, atomic no. increase.

(C) ${}^{A}_{Z}X$ ${}^{0}_{0}$ ${}^{A}_{Z}Y$, no. change in mass no. and atomic no.

(D) ${}^{A}_{Z}X$ ${}^{0}_{-1}$ ${}^{A}_{Z-1}Y$, Atomic no. decrease.

Subjective Problems

1. $2NH_3(g) \rightleftharpoons N_2(g)\ 3H_2(g)$

Rate $K[NH_3]^0$

Rate 2

Rate $\dfrac{1}{3}\ \dfrac{H_2]}{t}$ $\dfrac{H_2]}{t}$ 6

2. $t_{1/2}$ $\dfrac{[A_0]}{2K_A}$ $\dfrac{[A_0]}{2\ 5K}$ $\dfrac{200\text{ min.}}{2\ 5\ 400}$

$\dfrac{60\text{ sec.}}{20}$ 3

3. $A(g)$ $2B(g)$ $C(g)$

t 0 0.4 0.6

t t 0.3 0.4 0.1

r_i $KP_A^1\ P_B^2$

r_i K 0.4 $(0.6)^2$

r_t $K(0.3)(0.4)^2$

$\dfrac{r_i}{r_t}$ 3

4. Rate of reaction $K[CO]^x[O_2]^y$

4 10^{-5} $K(0.02)^x(0.02)^y$ ...(i)

1.6 10^{-4} $K(0.04)^x(0.02)^y$ ...(ii)

8 10^{-5} $K(0.02)^x(0.04)^y$ ...(iii)

$\dfrac{(i)}{(ii)}$ $\dfrac{1}{4}$ $\dfrac{1}{2}^{x}$

x 2

$\dfrac{(i)}{(iii)}$ $\dfrac{1}{2}$ $\dfrac{1}{2}^{y}$

y 1

From eq. (1)

4 10^{-5} $K(0.02)^2(0.02)^1$

K $\dfrac{4\ 10^{-5}}{4\ 10^{-4}\ 2\ 10^{-2}}$ $\dfrac{10}{2}$ 5

5. For zero order $2(t_{1/2})_0$ t_{Complete};

$(t_{1/2})_0$ $\dfrac{1.386}{2}$ ...(i)

For More Material Join: @JEEAdvanced_2024

For first order $(t_{1/2})_1 \quad \frac{0.693}{k_1}$ ...(ii)

Given $(t_{1/2})_0 \quad (t_{1/2})_1$

$\frac{13.86}{2} \quad \frac{0.693}{k_1}$

$k_1 \quad 0.1\,\text{min}^{-1} \quad 0.1 \quad 60\,\text{hr}^{-1} \quad 6\,\text{hr}^{-1}$

6. $r \quad k[A]^x[\text{H}^+]^y$

Order of reaction w.r.t. $A \quad 1$

$(x \quad 1)$; $r \quad k[\text{H}^+]^y \quad (\because k \quad k[A]^1)$

$\frac{a/2}{10} \quad k[10^{-2}]^y$ ...(i)

$\frac{a/2}{100} \quad k[10^{-3}]^y$ ...(ii)

(i)/(ii) ; $\quad y \quad 1$;


where a initial concentration.

7. $(k_f \quad k_b)t \quad \ln\frac{x_e}{x_e \quad x} \quad \ln\frac{x_e}{x_e \quad 0.8x_e}$;

$(k_f \quad k_b)t \quad \ln\frac{x_e}{0.2x_e} \quad \ln 5 \quad 1.6$;

$\frac{k_f}{k_b} \quad K_c$, $k_b \quad 0.1t \quad 6$ hrs.

8.



$K_1 : K_2 : K_3 = 4 : 2 : 1$

mole of A remain after 45 days

$\frac{N_0}{2^n} \quad \frac{N_0}{2^{\frac{T}{t_{1/2}}}} \quad \frac{1}{2^{45/15}} \quad \frac{1}{2^3} \quad \frac{1}{8}$

moles of A convert into product $\frac{7}{8}$ mol

moles of $[C] \quad \frac{K_2}{K_1 \quad K_2 \quad K_3} \quad \frac{7}{8} \quad \frac{8}{1} \quad 2$

9. $A \quad N$

$4.6 \quad 10^{12}\,\text{Bq} \quad \frac{\ln 2}{t_{1/2}} \quad \frac{1 \quad 10^{-3}}{131} \quad 6 \quad 10^{23}$

$t_{1/2} \quad \frac{0.693 \quad 6 \quad 10^{20}}{4.6 \quad 10^{12} \quad 131 \quad 3600 \quad 24} \quad 7.98\,\text{day}$

$t_{1/2} \quad 8$

10. $\frac{1}{t_{avg}} \quad \frac{1}{t}\ln\frac{[A_0]}{[A_t]}$; $\frac{1}{7.2} \quad \frac{1}{t_1}\ln\frac{100}{100 \quad \frac{100}{3}}$;

$\frac{1}{7.2} \quad \frac{1}{t_2}\ln\frac{100}{100 \quad \frac{200}{3}}$;

$t_2 \quad t_1 \quad 7.2\ln 3 \quad 7.2\ln\frac{3}{2} \quad 5$

11. ${}^{206}_{82}A \quad {}^{202}_{80}B \quad {}^{298}_{78}C \quad {}^{198}_{79}D \quad {}^{198}_{80}E$.

Proton difference in A and $E \quad 2$

12. Number of -particle $\frac{232 \quad 208}{4} \quad 6$

$Z_2 \quad Z_1 \quad 2x \quad y$

(where x number of and y number of)

$82 \quad 90 \quad 12 \quad y$

$y \quad 4$

Series

${}^{A}_{Z}\text{P} \quad {}^{A}_{Z\ 1}Q \quad {}^{A}_{Z\ 2}R \quad {}^{A}_{Z\ 3}S \quad {}^{A}_{Z\ 4}T$

Total number of isobars $\quad 5$

$n \quad 5$, so $n \quad 2 \quad 7$

13. Number of -particle $\frac{235 \quad 207}{4} \quad 7$

$Z_2 \quad Z_1 \quad 2x \quad y$

(where x number of and y number of)

$82 \quad 92 \quad 14 \quad y$

$y \quad \mathbf{4}$

Number of particle-number of particle) **3**

For More Material Join: @JEEAdvanced_2024



8

ELECTROCHEMISTRY

Electrolytic Cell

- **Electrolysis :** Chemical changes in electrolyte solution by passage of electric current, resulting into deposition of metals or liberation of gases at electrodes is known as electrolysis.
- **Electrolytic Cell :** This cell converts electrical energy into chemical energy. The entire assembly except that of the external battery is known as the electrolytic cell.

Quantitative Aspect of Electrolysis

Faraday's laws of electrolysis :

- **First law of electrolysis :** Amount of substance deposited or liberated at an electrode is directly proportional to amount of charge passed (utilized) through the solution.

$$W \quad Q$$

$$W \quad ZQ \quad Z \quad i \quad t$$

$$\text{Current efficiency} \quad \frac{\text{Actual charge utilised in process}}{\text{Charge passed through battery}} \quad 100$$

W mass liberated/deposited, Q charge in coulomb, i current, t time

Z electrochemical equivalent, 1 Faraday 96500 coulomb

- **Second law of electrolysis :** When same amount of charge is passed through different electrolytic solutions connected in series then mass of substances deposited or dissolved at anode or cathode are in ratio of their equivalent mass. *i.e.*, $W_1/W_2 \quad E_1/E_2$

$$\frac{W_1}{W_2} \quad \frac{E_1}{E_2}$$


For More Material Join: @JEEAdvanced_2024

Products of Electrolysis

S.No.	Electrolyte	Anode Product	Cathode Product
1.	NaCl (Molten) with Pt electrode	$Cl_2(g)$	$Na(l)$
2.	NaCl (aq) with Pt electrode	$Cl_2(g)$	$H_2(g)$
3.	$Na_2SO_4(aq)$ with Pt electrode	$O_2(g)$	$H_2(g)$
4.	$NaNO_3(aq)$ with Pt electrode	$O_2(g)$	$H_2(g)$
5.	$AgNO_3(aq)$ with Pt electrode	$O_2(g)$	$Ag(s)$
6.	$CuSO_4(aq)$ with inert electrode	$O_2(g)$	$Cu(s)$
7.	$CuSO_4(aq)$ with copper electrode	Cu dissolves	$Cu(s)$

Galvanic Cell

This cell converts chemical energy into electrical energy.



Galvanic cell is made up of two half cells *i.e.*, anodic and cathodic. Oxidation takes place at anode and reduction at cathode. It is also known as **voltaic cell.** It may be represented as shown in Fig. Zinc rod immersed in $ZnSO_4$ behaves as anode and copper rod immersed in $CuSO_4$ behaves as cathode.

- **Anode half cell reaction :**

$$Zn(s) \quad Zn^{2}\ (aq) \quad 2e \quad \text{(oxidation)}$$

- **Cathode half cell reaction :**

$$Cu^{2}\ (aq) \quad 2e \quad Cu(s)\ \text{(reduction)}$$

- **Over all process :**

$$Zn(s) \quad Cu^{2}\ (aq) \quad Cu(s) \quad Zn^{2}\ (aq)$$

In galvanic cell like Daniel cell; electrons flow from anode (zinc rod) to the cathode (copper rod) through external circuit; zinc dissolves as Zn^{2} ; Cu^{2} ion in the cathode cell picks up two electron and become deposited at cathode.

Salt Bridge

It is U-shaped tube contains saturated solution of inert electrolyte like KCI, KNO_3, NH_4Cl and NH_4NO_3 etc. in agar-agar gel.

- Cation and anion of inert electrolyte have same mobility.
- Ions of inert electrolyte do not mix with electrolytic solution present in the half cell.
- Ions of inert electrolyte do not participate in electrochemical change.

For More Material Join: @JEEAdvanced_2024

Function of Salt Bridge

- ❖ It complete the electrical circuit.
- ❖ It maintained the two half cell electrically neutral by the flow of ions.

Representation of a cell (IUPAC conventions) : Let us explain the convention taking the example of Daniel cell.

(i) Anodic half cell is written on left and cathodic half cell on right hand side.

$$Zn(s)|ZnSO_4(aq)||CuSO_4(aq)|Cu(s)$$

(ii) Two half cells are separated by double vertical lines and it indicate salt bridge or any type of porous partition.

(iii) EMF (electromotive force) may be written on the right hand side of the cell.

(iv) Single vertical lines indicate the phase separation between electrode and electrolyte solution.

$$\underset{\text{Anode}}{Zn(s) \mid Zn^{2}(aq)} \,||\, \underset{\text{Catode}}{Cu^{2}(aq) \mid Cu(s)}$$

E°_{cell} $E^{\circ}_{OP(LHS)}$ $E^{\circ}_{RP(RHS)}$; E_{cell} $E_{OP(LHS)}$ $E_{RP(RHS)}$; E_{cell} $E_{RP(RHS)}$ $E_{RP(LHS)}$

Nernst's Equation

$$E_{cell} = E^{\circ}_{cell} - \frac{2.303RT}{nF}\log Q$$

At 298 K temp. $E_{cell} = E^{\circ}_{cell} - \frac{0.0591}{n}\log Q$; where Q is reaction quotient.

Cell Thermodynamics

(i) $G = nFE_{cell}$

(ii) $G^{\circ} = nFE^{\circ}_{cell}$

(iii) Equilibrium constant (K), $\log K = \frac{n E^{\circ}_{cell}}{0.0591}$

(iv) Temp. coefficient of cell $\left(\frac{E}{T}\right)_P$

(v) Enthalpy of reaction inside the cell, $H = nFE - nFT\left(\frac{E}{T}\right)_P$

(vi) Entropy change inside the cell, $S = nF\left(\frac{E}{T}\right)_P$

Concentration Cell

Anode and cathode both are formed by same substance so $E_{cell} = 0$, such type of cell is known as concentration cell. Concentration gradient may arise either in electrode material or in electrolyte. Thus there are two types of concentration cells.

- ❖ **Electrode gas conc. cell :**

$$\underset{P_1 \text{ atm}}{Pt, H_2(g)|H^{+}(C)|H_2(g), Pt}\underset{P_2 \text{ atm}}{}$$

$$E_{cell} = \frac{0.0591}{2}\log\frac{P_1}{P_2}$$

For More Material Join: @JEEAdvanced_2024

- **Electrolyte concentration cell :**

$$Zn(s)|ZnSO_4(C_1)||ZnSO_4(C_2)|Zn(s)$$

$$E_{cell} \quad \frac{0.0591}{2}\log\frac{C_2}{C_1}$$

Different Type of Electrodes

S.No.	Name of Electrode	Anode	Cathode
1.	Hydrogen electrode	$Pt(s)\|H_2(g)\|H^+(aq)$ $E_{OP} \quad E^\circ_{OP} \quad \frac{0.0591}{2}\log\frac{[H^+]^2}{P_{H_2}}$	$H^+(aq)\|H_2(g)\|Pt(s)$ $E_{RP} \quad E^\circ_{RP} \quad \frac{0.0591}{2}\log\frac{P_{H_2}}{[H^+]^2}$
2.	Metal-metal ion electrode	$M(s)\|M^{n+}(aq)$ $E_{OP} \quad E^\circ_{OP} \quad \frac{0.0591}{n}\log[M^{n+}]$	$M^{n+}(aq)\|M(s)$ $E_{RP} \quad E^\circ_{RP} \quad \frac{0.0591}{n}\log\frac{1}{[M^{n+}]}$
3.	Calomel electrode	$Hg(l), Hg_2Cl_2(s)\|Cl^-(aq)$ $E_{OP} \quad E^\circ_{OP} \quad \frac{0.0591}{2}\log\frac{1}{[Cl^-]^2}$	$Cl^-(aq)\|Hg_2Cl_2(s), Hg(l)$ $E_{RP} \quad E^\circ_{RP} \quad \frac{0.0591}{2}\log[Cl^-]^2$
4.	Redox electrode	$Pt(s)\|Fe^{2+}(aq), Fe^{3+}(aq)$ $E_{OP} \quad E^\circ_{OP} \quad \frac{0.0591}{1}\log\frac{[Fe^{3+}]}{[Fe^{2+}]}$	$Fe^{3+}(aq), Fe^{2+}(aq)\|Pt(s)$ $E_{RP} \quad E^\circ_{RP} \quad \frac{0.0591}{1}\log\frac{[Fe^{2+}]}{[Fe^{3+}]}$
5.	Metal insoluble salt anion electrode	$Ag(s)\|AgCl(s)\|Cl^-(aq)$ $E_{OP} \quad E^\circ_{OP} \quad \frac{0.0591}{1}\log\frac{1}{[Cl^-]}$	$Cl^-(aq)\|AgCl(s)\|Ag(s)$ $E_{RP} \quad E^\circ_{RP} \quad \frac{0.0591}{1}\log[Cl^-]$

Batteries

Electrochemical cells can be used as **batteries.** Batteries are of two kinds :

- **Primary battery :** Where the reaction occurs only once and can not be reused once it becomes dead over the course of time. For examples, dry cell (Leclanche cell), Mercury cell.
- **Secondary battery :** Which can be recharged by passing current through it in the opposite direction so that it can be used again. For example, lead-acid battery, Nickel-cadmium cell.

Fuel Cells

Fuel cells are the galvanic cells that convert the energy of combustion of fuels (*e.g.*, hydrogen, methane, methanol, etc.) directly into electrical energy.

Fuel cells are more efficient thermodynamically, and more of the energy of the reaction can be made available for useful work provided that the supply of reactants is maintained.

Methanol, ethanol, hydrazine, formaldehyde, carbon monoxide can be used as fuels in fuel cells apart from hydrogen.

For More Material Join: @JEEAdvanced_2024

Corrosion

When the metal exposed to some environment it gets converted to its oxides. The oxidative deterioration of metal is known as corrosion.

Example : Rusting of iron, tarnishing of silver, development of green coating on copper and bronze etc.

Oxidation $Fe(s) \quad Fe^{2}(aq) \quad 2e$

Reduction $O_2(g) \quad 4H \ (aq) \quad 4e \quad 2H_2O(l)$

Atmospheric oxygen oxidised Fe^{2}

$$2Fe^{2}(aq) \quad 2H_2O(l) \quad \frac{1}{2}O_2(g) \quad Fe_2O_3(s) \quad 4H \ (aq)$$

Hydrated feric oxide ($Fe_2O_3 \ xH_2O(s)$) is known as rust.

❖ **Factors which enhance corrosion :**

(i) Presence of impurities in the metal.
(ii) Presence of moisture.
(iii) Presence of electrolyte.

❖ **Prevention of corrosion :**

(i) Barrier protection by oil/grease layer, paints or electroplating.
(ii) Sacrificial protection by mixing the metal with more electropositive metal.

Conductance

❖ **Conductance (*G*) :** It is defined as the reciprocal of the electrical resistance *i.e.*, $G \quad 1/R$. It measures the ease with which the current flows through a conductor.

Unit : Siemens, S or ^{1}.

❖ **Specific resistance or resistivity ( ) :** The resistivity or specific resistance is defined as the resistance in ohm of a conductor having length equal to 1 cm and area of cross-section equal to 1 cm^{2}.

$R \quad \frac{l}{A}$ or $R \quad \frac{l}{A}$ where specific resistance, l length of conductor

A area of cross-section of the conductor.

Unit : ohm cm.

❖ **Specific conductance or conductivity (*k*) :** It is define as the reciprocal of specific resistance.

i.e., $k \quad \frac{1}{} \quad R \quad \frac{l}{A} \quad k \quad G \quad \frac{l}{A}$

$k \quad G \quad G^{*}$

$G^{*} \quad \frac{l}{A}$, where G^{*} cell constant

When $l \quad 1$ cm and $A \quad 1\,cm^{2}$; $k \quad G$

Thus conductivity is the conductance of one centimeter cube of the solution of an electrolyte.

Unit : $^{1}cm^{-1}$, Scm^{-1}.

For More Material Join: @JEEAdvanced_2024


- **Molar conductivity ($_M$) :** The conducting power of all the ions produced by dissolving 1 mole of an electrolyte in solution.

Mathematically, $_M \quad k \quad V$

$$_M (\mathrm{S\,cm^2\,mol^{-1}}) \quad k \quad \frac{1000}{M} \qquad (M \quad \text{molarity})$$

Variation of conductivity and molar conductivity with concentration :
For weak and strong electrolytes

- Conductivity decreases with concentration.
- Molar conductivity increases with decrease in concentration.



Kohlrausch's Law

According to this law, at infinite dilution, when the dissociation is complete, each ion makes a definite contribution towards molar conductivity of the electrolyte irrespective of the nature of the other ions present.

The molar conductivity of an electrolyte at infinite dilution is the sum of the ionic conductivities of the cations and the anions each multiplied by the number of ions present in one formula unit of the electrolyte

The equivalent conductivity of an electrolyte at infinite dilution is the sum of the equivalent conductivities of the cations and anions.

$$e.g., A_xB_y. \qquad {}^0_M(A_xB_y) \quad x\ {}^0_M(A^y) \quad y\ {}^0_M(B^x)$$
$$ {}^0_{eq.}(A_xB_y) \quad {}^0_{eq.}(A^y) \quad {}^0_{eq.}(B^x)$$

Application

(a) Determination of equivalent/molar conductivities of weak electrolytes at infinite dilution

$$ {}^0_M(CH_3COOH) \quad {}^0_M(CH_3COONa) \quad {}^0_M(HCl) \quad {}^0_M(NaCl) \quad {}^0_M(NaCl)$$

(b) Determination of degree of dissociation ( ) and equilibrium constant (K) of an electrolyte at a given dilution

$$\frac{\text{Molar conductivity at concentration } C}{\text{Molar conductivity at infinite dilution}} \quad \frac{_M}{{}^0_M}$$

$$HA \rightleftharpoons H \quad A$$

Initial conc.	C	0	0
Conc. at equilibrium	$C \quad C$	C	C

For More Material Join: @JEEAdvanced_2024

$$K_a \quad \frac{C^{\ 2}}{1} \quad \frac{C \ \frac{M}{0}^{\ 2}}{1 \ \frac{M}{0}}$$
$$M$$

(c) Determination of the solubility of a sparingly soluble salt

Since the solution is saturated at infinite dilution $_0$ ^0_M and molarity solubility.

$$^0_M \quad \frac{k \quad 1000}{\text{molarity}}$$

$$\text{Solubility } (S) \quad \frac{k \quad 1000}{^0_M}$$


For sparingly soluble salt A_xB_y : Solubility product $K_{sp} \quad x^x y^y \quad S^{x \ y}$

Conductometric Titration

Titration in which end point can be determined by measuring conductivity at each addition of the titre solution (for burette) is called conductometric titration.


Titration of HCl (*aq*) with NaOH (*aq*) (Strong acid vs Strong base)

$$HCl(aq) \quad NaOH(aq) \qquad NaCl(aq) \quad H_2O(l)$$



Titration of CH_3COOH (*aq*) with NaOH (*aq*) (Weak acid vs Strong base)

$$CH_3COOH(aq) \quad NaOH(aq) \qquad CH_3COONa \quad H_2O(l)$$



For More Material Join: @JEEAdvanced_2024



Level 1

1. A cell reaction would be spontaneous if the cell potential and $_rG$ are respectively :
(a) positive and negative (b) negative, negative
(c) zero, zero (d) positive, zero

2. Which of the following statement is correct?
(a) Cathode is ve terminal in both, galvanic and electrolytic cells
(b) Anode is +ve terminal in both, galvanic and electrolytic cells
(c) Cathode and anode are –ve terminal in electrolytic and galvanic cell.
(d) Cathode and anode are +ve terminal in electrolytic and galvanic cell.

3. Electrolytes when dissolved in water dissociate into ions because :
(a) they are unstable.
(b) the water dissolves it.
(c) the force of repulsion increases.
(d) the force of electrostatic attraction is broken down by water.

4. The electric charge required for electrode deposition of one gram-equivalent of a substance is :
(a) one ampere per second (b) 96500 coulombs per second
(c) one ampere for one hour (d) charge on one mole of electrons

5. The amount of an ion liberated on an electrode during electrolysis does not depend upon :
(a) conductance of the solution (b) current strength
(c) time (d) electrochemical equivalent of the element

6. How many electrons are there in one coulomb electricity?
(a) 6.023 10^{23} (b) 1.64 10^{-24} (c) 6.24 10^{18} (d) 6.24 10^{-24}

7. How many coulombs are provided by a current of 0.010 mA in the calculator battery that can operate for 1000 hours?
(a) 1.0 (b) 10 (c) 0.010 (d) 36

8. How many minutes are required to deliver 3.21 10^{6} coulombs using a current of 500 A used in the commercial production of chlorine?
(a) 8.3 (b) 5.3 10^{4} (c) 6420 (d) 107

9. Passage of a current for 548 seconds through a silver coulometer results in the deposition of 0.746 g of silver. What is the current (in A)?
(a) 1.22 (b) 1.16 (c) 1.07 (d) 1.00

10. Electrolysis can be used to determine atomic masses. A current of 0.550 A deposits 0.55 g of a certain metal in 100 minutes. Calculate the atomic mass of the metal if eq. mass mol. mass/3
(a) 100 (b) 45.0 (c) 48.25 (d) 144.75

11. Beryllium occurs naturally in the form of beryl. The metal is produced from its ore by electrolysis after the ore has been converted to the oxide and then to the chloride. How many grams of Be(*s*) is deposited from a $BeCl_2$ solution by a current of 5.0 A that flows for 1.0 h?
(Atomic weight : Be 9)
(a) 0.840 (b) 1.68 (c) 1.42 (d) 1.08

For More Material Join: @JEEAdvanced_2024

12. How many minutes will it take to plate out 5.2 g of Cr from a $Cr_2(SO_4)_3$ solution using a current of 9.65 A? (Atomic mass : Cr = 52.0)
(a) 200 (b) 50 (c) 100 (d) 103

13. Calculate the current (in mA) required to deposit 0.195 g of platinum metal in 5.0 hours from a solution of $[PtCl_6^{2-}]$: (Atomic mass : Pt = 195)
(a) 310 (b) 31 (c) 21.44 (d) 5.36

14. How many Faradays are required to reduce 0.25 g of Nb (V) to the metal? (Atomic mass : Nb = 93)
(a) 2.7 10^{-3} (b) 1.3 10^{-2} (c) 2.7 10^{-2} (d) 7.8 10^{-3}

15. One gm metal M^3 was discharged by the passage of 1.81 10^{23} electrons. What is the atomic mass of metal?
(a) 33.35 (b) 133.4 (c) 66.7 (d) None of these

16. Total charge required for the oxidation of two moles Mn_3O_4 into MnO_4^{2-} in presence of alkaline medium is :
(a) 5 F (b) 10 F (c) 20 F (d) None of these

17. The electrolytic decomposition of dilute sulphuric acid with platinum electrode, cathodic reaction is :
(a) reduction of H (b) oxidation of SO_4^2 (c) reduction SO_3^2 (d) oxidation of H_2O

18. Which one of the following metals can not be obtained on electrolysis of aqueous solution of its salts ?
(a) Mg (b) Ag (c) Cu (d) Cr

19. A highly concentrated solution of potassium sulphate in water is electrolysed using inert electrodes. The products at the cathode and anode are respectively.
(a) H_2, O_2 (b) O_2, H_2 (c) O_2, Na (d) $H_2, S_2O_8^2$

20. The passage of current through a solution of certain electrolyte results in the evolution of $H_2(g)$ at cathode and $Cl_2(g)$ at anode. The electrolytic solution is :
(a) Water (b) *aq*. H_2SO_4 (c) *aq*. NaCl (d) *aq*. $CuCl_2$

21. When an aqueous solution of H_2SO_4 is electrolysed, the product at anode is :
(a) H (b) OH (c) SO_4^2 (d) O_2

22. An aqueous solution of Na_2SO_4 is electrolysed using Pt electrodes.The products at the cathode and anode are respectively :
(a) H_2, SO_2 (b) O_2, NaOH (c) H_2, O_2 (d) O_2, SO_2

23. The electrolysis of a solution resulted in the formation of $H_2(g)$ at the cathode and $O_2(g)$ at the anode. The solution is :
(a) AgCl(*aq*) (b) $H_2SO_4(aq)$
(c) highly concentrated NaCl(*aq*) solution (d) $CuCl_2(aq)$

24. If Pt is used as cathode in the electrolysis of aqueous NaCl solution, the ion produced at cathode is :
(a) H (b) Na (c) OH (d) Cl

25. A dilute aqueous solution of $CuSO_4$ is electrolysed using platinum electrodes.The products at the anode and cathode are :
(a) O_2, H_2 (b) H_2, O_2 (c) O_2, Cu (d) $S_2O_8^2$, H_2

For More Material Join: @JEEAdvanced_2024

26. What products are formed during the electrolysis of concentrated aqueous solution of sodium chloride?
(I) $Cl_2(g)$ at anode (II) NaOH as electrolyte (III) $H_2(g)$ at cathode
(a) I only (b) I and II only (c) I and III only (d) I, II and III

27. Which of the following aqueous solution produces metal after electrolysis ?
(a) $K_2Cr_2O_7$ (b) $KMnO_4$ (c) CH_3COONa (d) $CuCl_2$

28. How much time is required for complete decomposition of 4 moles of water using 4 ampere?
(a) 3.86 10^5 sec (b) 1.93 10^5 sec (c) 96500 sec (d) 48250 sec

29. An aqueous solution containing 1 *M* each of Au^{3+}, Cu^{2+}, Ag^+, Li^+ is being electrolysed by using inert electrodes. The value of standard reduction potentials are :
$E_{Ag^+/Ag}$ 0.80 V, $E_{Cu^{2+}/Cu}$ 0.34 V, and $E_{Au^{3+}/Au}$ 1.50 V, $E_{Li^+/Li}$ – 3.03 V,
With increasing voltage, the sequence of deposition of metals on the cathode will be :
(a) Li, Cu, Ag, Au (b) Cu, Ag, Au (c) Au, Ag, Cu (d) Au, Ag, Cu, Li

30. If 0.50 L of a 0.60 *M* $SnSO_4$ solution is electrolysed for a period of 30.0 min using a current of 4.60 A. If inert electrodes are used, what is the final concentration of Sn^{2+} remaining in the solution? [at. mass of Sn = 119]
(a) 0.342 *M* (b) 0.544 *M* (c) 0.389 *M* (d) 0.514 *M*

31. A 100.0 mL dilute solution of Ag is electrolysed for 15.0 minutes with a current of 1.25 mA and the silver is removed completely. What was the initial $[Ag^+]$?
(a) 2.32 10^{-1} (b) 2.32 10^{-4} (c) 2.32 10^{-3} (d) 1.16 10^{-4}

32. A 250.0 mL sample of a 0.20 *M* Cr^3 is electrolysed with a current of 96.5 A. If the remaining $[Cr^{3+}]$ is 0.1 *M*, the duration of process is :
(a) 25 sec (b) 225 sec (c) 150 sec (d) 75 sec

33. The element indium is to be obtained by electrolysis of a molten halide of the element. Passage of a current of 3.20 A for a period of 40.0 min results in formation of 3.05 g of In. What is the oxidation state of indium in the halide melt? (Atomic mass of In 114.8)
(a) 3 (b) 2 (c) 5 (d) 1

34. An electrolysis of a oxytungsten complex ion using 1.10 A for 40 min produces 0.838 g of tungsten. What is the charge on tungsten in the material? (Atomic mass of *W* 184)
(a) 6 (b) 2 (c) 4 (d) 1

35. In the electrolysis of aqueous NaCl, what volume of $Cl_2(g)$ is produced in the time that it takes to liberate 5.0 liter of $H_2(g)$? Assume that both gases are measured at STP.
(a) 5.0 (b) 2.50 (c) 7.50 (d) 10.0

36. How many grams of Cr are deposited in the electrolysis of solution of $Cr(NO_3)_3$ in the same time that it takes to deposit 0.54 g of Ag in a silver coulometer arranged in series with the $Cr(NO_3)_3$ cell? (Atomic mass : Cr = 52.0; Ag = 108)
(a) 0.0866 (b) 0.0288 (c) 0.173 (d) 0.220

37. In the electrolysis of a $CuSO_4$ solution, how many grams of Cu are plated out on the cathode in the time that it takes to liberate 5.6 litre of $O_2(g)$, measured at 1 atm and 273 K, at the anode?
(a) 31.75 (b) 14.2 (c) 4.32 (d) None of these

38. Ammonium perchlorate, NH_4ClO_4, used in the solid fuel in the booster rockets on the space shuttle, is prepared from sodium perchlorate, $NaClO_4$, which is produced commercially by the

For More Material Join: @JEEAdvanced_2024

electrolysis of a hot, stirred solution of sodium chloride. How many faradays are required to produce 1.0 kg of sodium perchlorate?

$$NaCl + 4H_2O \quad NaClO_4 + 4H_2$$

(a) 40.3 (b) 18.3 (c) 31.6 (d) 65.3

39. In the commercial preparation of aluminum, aluminum oxide (Al_2O_3) is electrolysed at 1000 C. How many coulombs of electricity are required to give 54 kg of aluminum? Assume following reaction takes place at cathode :

$$Al^{3+} + 3e^- \quad Al$$

(a) 17.3 10^8 (b) 3.21 10^7 (c) 1.82 10^4 (d) 57.6 10^7

40. When molten lithium chloride (LiCl) is electrolysed, lithium metal is formed at the cathode. If current efficiency is 75% then how many grams of lithium are liberated when 1930 C of charge pass through the cell ? (Atomic mass of Li = 7)

(a) 0.105 (b) 0.120 (c) 0.28 (d) 0.240

41. Sodium metal is produced commercially by the electrolysis of molten sodium chloride and chlorine is produced as a by product. How many litres of chlorine at 1.8 atm and 27 C will be produced if a current of 1.0 10^3 A is passed through NaCl (*l*) for 9.65 hr.?

(a) 2463 (b) 460 (c) 1800 (d) 1231.6

42. $H_2(g)$ and $O_2(g)$, can be produced by the electrolysis of water. What total volume (in L) of O_2 and H_2 are produced at 1 atm and 273 K when a current of 30 A is passed through a K_2SO_4 (*aq*) solution for 193 min.?

(a) 20.16 (b) 40.32 (c) 60.48 (d) 80.64

43. The cost of 2 Rs/kWh of operating an electric motor for 10 hours takes 10 amp at 110 V is :

(a) 79200 Rs (b) 22000 Rs (c) 220 Rs (d) 22 Rs

44. A 1 *M* solution of H_2SO_4 is electrolysed. Select correct statement in respect of products obtained at anode and cathode respectively :

Given : $2SO_4^{2-} \quad S_2O_8^{2-} + 2e^-; E \quad 2.01\,V$

$H_2O(l) \quad 2H^+(aq) \quad 1/2\,O_2(g) + 2e^-; E \quad 1.23\,V$

(a) concentration of H_2SO_4 remain constant; H_2, O_2
(b) concentration of H_2SO_4 increases; O_2, H_2
(c) concentration of H_2SO_4 decreases; O_2, H_2
(d) concentration of H_2SO_4 remains constant; $S_2O_8^{2-}, H_2$

45. Cadmium amalgam is prepared by electrolysis of a solution of $CdCl_2$ using a mercury cathode. How long should a current of 4 A be passed in order to prepare 10% by mass Cd in Cd—Hg amalgam on cathode of 4.5 g Hg? (atomic mass of Cd = 112)

(a) 400 sec (b) 215.40 sec (c) 861.6 sec (d) 430.8 sec

46. Use of electrolysis is in :

(a) electrorefining (b) electroplating
(c) both (a) and (b) (d) None of these

47. When a solution of $AgNO_3$ (1 *M*) is electrolysed using platinum anode and copper cathode, what are the products obtained at two electrodes?

Given : $E_{Cu^{2+}|Cu}$ 0.34 volt; $E_{O_2, H^+|H_2O}$ 1.23 volt; $E_{H^+|H_2}$ 0.0 volt;

For More Material Join: @JEEAdvanced_2024

$E_{Ag^+|Ag}$ 0.8 volt

(a) Cu Cu^{2+} at anode; Ag^+ Ag at cathode

(b) H_2O O_2 at anode; Cu^2 Cu at cathode

(c) H_2O O_2 at anode; Ag Ag at cathode

(d) NO_3^- NO_2 at anode; Ag Ag at cathode

48. Which of the following statements is correct about Galvanic cell ?

(a) It converts chemical energy into electrical energy.

(b) It converts electrical energy into chemical energy.

(c) It converts metal from its free state to the combined state.

(d) It converts electrolyte into individual ions.

49. E for $Cl_2(g)$ $2e^-$ $2Cl^-(aq)$ is 1.36 V; E for $Cl^-(aq.)$ $1/2Cl_2(g) + e^-$ is :

(a) 1.36 V (b) 1.36 V (c) 0.68 V (d) 0.68 V

50. When two half-cells of electrode potential of E_1 and E_2 are combined to form a half cell of electrode potential E_3, then

(when n_1, n_2 and n_3 are no. of electrons exchanged in first, second and combined half-cells) :

(a) E_3 E_2 E_1

(b) E_3 $\dfrac{E_1 n_1 \quad E_2 n_2}{n_3}$

(c) E_3 $\dfrac{E_1 n_1 \quad E_2 n_2}{n_3^2}$

(d) E_3 E_1 E_2

51. The function of a salt bridge is to :

(a) maintain electrical neutrality of both half cells

(b) increase the cell potential at the positive electrode

(c) decrease the cell potential at the negative electrode

(d) eliminate the impurities present in the electrolyte

52. Saturated solution of KNO_3 with agar-agar is used to make 'salt bridge' because :

(a) size of K is greater than that of NO_3^-

(b) velocity of NO_3^- is greater than that of K^+

(c) velocities of K and NO_3^- are nearly the same

(d) both velocity and sizes of K and NO_3^- ions are nearly same

53. A salt bridge may contain :

(a) a saturated solution of KCl and agar-agar

(b) a saturated solution of KNO_3 and agar-agar

(c) a saturated solution of NH_4NO_3 and agar-agar

(d) all of these

54. The nature of curve of E_{cell} *vs.* log K_c is :

(a) straight line (b) parabola (c) hyperbola (d) elliptical curve

55. Consider the following equations for a cell reaction

$A \quad B \rightleftharpoons C \quad D$; E^o x volt, K_{eq} K_1

$2A \quad 2B \rightleftharpoons 2C \quad 2D$; E^o y volt, K_{eq} K_2

For More Material Join: @JEEAdvanced_2024

then :

(a) $x \quad y, K_1 \quad K_2$ (b) $x \quad 2y, K_1 \quad 2K_2$


(c) $x \quad y, K_1^2 \quad K_2$ (d) $x^2 \quad y, K_1^2 \quad K_2$

56. Which graph correctly correlates E_{cell} as a function of concentrations for the cell


$$Zn(s) + 2Ag^+(aq) \qquad Zn^{2+}(aq) + 2Ag(s), \quad E_{cell} \quad 1.56 \text{ V}$$

Y-axis : E_{cell}, X-axis : $\log_{10} \dfrac{[Zn^{2+}]}{[Ag^+]^2}$


(a)




(b)



(c)



(d)



57. The Nernst equation $E \quad E \quad RT/nF \ln Q$ indicates that the Q will be equal to equilibrium constant K_c when :

(a) $E \quad E$ (b) $RT/nF = 1$ (c) E zero (d) $E \quad 1$

58. The cell reaction $2Ag \ (aq) \quad H_2(g) \qquad 2H \ (aq) \quad 2Ag(s)$, is best represented by :

(a) $Ag(s) \mid Ag \ (aq) \mid\mid H \ (aq) \mid H_2(g) \mid Pt(s)$ (b) $Pt(s) \mid H_2(g) \mid H \ (aq) \mid\mid Ag \ (aq) \mid Ag(s)$

(c) $Ag(s) \mid Ag \ (aq) \mid\mid H_2(g) \mid H \ (aq) \mid Pt(s)$ (d) $Ag \ (aq) \mid Ag(s) \mid\mid H_2(g) \mid H \ (aq)$

59. The cell reaction $Hg_2Cl_2(s) \quad Cu(s) \qquad Cu^2 \ (aq) \quad 2Cl \ (aq) \quad 2Hg(l)$, is best represented by :

(a) $Cu(s) \mid Cu^2 \ (aq) \mid\mid Hg_2Cl_2(s) \mid Hg(l)$

(b) $Cu(s) \mid Cu^2 \ (aq) \mid\mid Hg(l) \mid HgCl_2(s)$

(c) $Cu(s) \mid Cu^2 \ (aq) \mid\mid Cl \ (aq) \mid Hg_2Cl_2(s) \mid Hg(l) \mid Pt(s)$

(d) $Hg_2Cl_2(s) \mid Cl \ (aq) \mid\mid Cu^2 \ (aq) \mid Cu(s)$

60. The cell reaction $Cr_2O_7^2 \ (aq) \quad 14H \ (aq) \quad 6Fe^2 \ (aq) \qquad 6Fe^3 \ (aq) \quad 2Cr^3 \ (aq) \quad 7H_2O(l)$, is best represented by :

(a) $Pt(s) \mid Fe^2 \ (aq), Fe^3 \ (aq) \mid\mid Cr_2O_7^2 \ (aq), Cr^3 \ (aq) \mid Pt(s)$

(b) $Pt(s) \mid Cr_2O_7^2 \ (aq), Cr^3 \ (aq) \mid\mid Fe^3 \ (aq), Fe^2 \ (aq) \mid Pt(s)$

(c) $Fe^2 \ (aq) \mid Fe^3 \ (aq) \mid\mid Cr_2O_7^2 \ (aq) \mid Cr^3 \ (aq)$

(d) $Cr_2O_7^2 \ (aq) \mid Cr^3 \ (aq) \mid\mid Fe^3 \ (aq) \mid Fe^2 \ (aq)$

61. Select the correct cell reaction of the cell $Ag(s) \mid Ag \ (aq) \mid\mid Cu^2 \ (aq) \mid Cu(s)$:

(a) $2Ag(s) \quad Cu(s) \qquad Cu^2(aq) \quad 2Ag \ (aq)$

(b) $Cu(s) \quad 2Ag \ (aq) \qquad Cu^2 \ (aq) \quad 2Ag(s)$

For More Material Join: @JEEAdvanced_2024

(c) $2Ag(s)$ Cu^{2} (aq) $Cu(s)$ $2Ag$ (aq)

(d) Cu^{2} (aq) $2Ag$ (aq) $2Ag(s)$ $Cu(s)$

62. Select the correct cell reaction of the cell $Pt(s)|Cl_2(g)|Cl$ $(aq)||Ag$ $(aq)|Ag(s)$:

(a) $Cl_2(g)$ Ag (aq) $Ag(s)$ $2Cl$ (aq)

(b) $Cl_2(g)$ $Ag(s)$ $2Cl$ (aq) Ag (aq)

(c) $2Cl$ (aq) $2Ag$ (aq) $2Ag(s)$ $Cl_2(g)$

(d) $AgCl(s)$ Ag (aq) Cl (aq)

63. Standard electrode potential of SHE at 298 K is :

(a) 0.05 V (b) 0.10 V (c) 0.50 V (d) 0.00 V

64. The e.m.f. of the following galvanic cells :

(a) $Zn|Zn^{2+}(1\,M)||Cu^{2+}(1M)|Cu$ (b) $Zn|Zn^{2+}(0.1\,M)||Cu^{2+}(1\,M)|Cu$

(c) $Zn|Zn^{2+}(1\,M)||Cu^{2+}(0.1\,M)|Cu$ (d) $Zn|Zn^{2+}(0.1\,M)||Cu^{2+}(0.1\,M)|Cu$

are represented by $E_{1,}$ E_2, E_3 and E_4 respectively. Which of the following statement is true?

(a) E_1 E_2 E_3 E_4 (b) E_3 E_2 E_1 E_4

(c) E_3 E_1 E_4 E_2 (d) E_2 E_1 E_4 E_3

65. Based on the cell notation for a spontaneous reaction, at the anode :

$$Ag(s)|AgCl(s)|Cl^{-}(aq)||Br^{-}(aq)|Br_2(l)|C(s)$$

(a) AgCl gets reduced (b) Ag gets oxidized

(c) Br^{-} gets oxidized (d) Br_2 gets reduced

66. Given the listed standard electrode potentials, what is E for the cell :

$4BiO^{+}(aq) + 3N_2H_5^{+}(aq)$ $4Bi(s) + 3N_2(g) + 4H_2O(l) + 7H^{+}(aq)$

$N_2(g) + 5H^{+}(aq) + 4e^{-}$ $N_2H_5^{+}(aq), E = 0.23$ V

$BiO^{+}(aq) + 2H^{+}(aq) + 3e^{-}$ $Bi(s) + H_2O(l),$ $E = +0.32$ V

(a) 0.55 (b) 0.34 (c) 1.88 (d) 0.09

67. What is the standard electrode potential for the reduction of HClO?

$HClO(aq) + H^{+}(aq) + 2e^{-}$ $Cl^{-}(aq) + H_2O(l)$

Given : $Cr^{2+}(aq)$ $Cr^{3+}(aq) + e^{-},$ E 0.41 V

$HClO(aq) + H^{+}(aq) + 2Cr^{2+}(aq)$ $2Cr^{3+}(aq) + Cl^{-}(aq)$ $H_2O(l), E$ 1.80

(a) 1.39 (b) 1.54 (c) 1.22 (d) 0.90

68. The E for the following cell is 0.34 V. $In(s)|In(OH)_3(aq)||SbO_2^{-}(aq)|Sb(s)$.

Using E 1.0 V for the $In(OH)_3|$ In couple, calculate E for the $SbO_2^{-}|$ Sb half-reaction :

(a) 1.34 (b) 0.66

(c) +0.82 (d) 0.66

69. From the following half-cell reactions and their standard potentials, what is the smallest possible standard e.m.f. for spontaneous reactions?

$PO_4^{3-}(aq)$ $2H_2O(l)$ $2e^{-}$ HPO_3^{2} $3OH^{-}(aq);$ E 1.05 V

$PbO_2(s)$ $H_2O(l)$ $2e^{-}$ $PbO(s)$ $2OH^{-}(aq);$ E 0.28 V

For More Material Join: @JEEAdvanced_2024

$IO_3^-(aq)\quad 2H_2O(l)\quad 4e^-\qquad IO^-(aq)\quad 4OH^-(aq);\quad E^\circ = 0.56\ V$

(a) +0.00 (b) +0.74 (c) +0.56 (d) +0.28

70. Determine which substance is the best reducing agent in Q. no. 69 :

(a) HPO_3^{2-} (b) PO_4^{3-} (c) IO^- (d) IO_3^-

71. Which substance is the best oxidizing agent in Q. no. 69?

(a) IO_3^- (b) IO^- (c) PbO (d) PO_4^{3-}

72. Consider the following half-cell reactions and associated standard half-cell potentials and determine the maximum voltage that can be obtained by combination resulting in spontaneous processes :

$AuBr_4^-(aq)\quad 3e^-\qquad Au(s)\quad 4Br^-(aq);\quad E^\circ = 0.86\ V$

$Eu^{3+}(aq)\quad e^-\qquad Eu^{2+}(aq);\quad E^\circ = 0.43V$

$Sn^{2+}(aq)\quad 2e^-\qquad Sn(s);\quad E^\circ = 0.14\ V$

$IO^-(aq)\quad H_2O(l)\quad 2e^-\qquad I^-(aq)\quad 2OH^-(aq);\quad E^\circ = 0.49\ V$

(a) 0.72 (b) 1.54 (c) 1.00 (d) +1.35

73. The position of some metals in the electrochemical series in decreasing electropositive character is Mg > Al > Zn > Cu > Ag. What will happened if copper spoon is used to stirred a solution of aluminium nitrate ?

(a) The spoon gets coated with aluminium.

(b) An alloy of aluminium and copper is formed.

(c) No reaction occurs

(d) The solution starts turning blue

74. Zn can displace :

(a) Mg from its aqueous solution (b) Cu from its aqueous solution

(c) Na from its aqueous solution (d) Al from its aqueous solution

75. Based on the following informations arrange four metals A, B, C and D in order of decreasing ability to act as reducing agents :

(I) Only A, B and C react with $1\,M$ HCl to give $H_2(g)$

(II) When C is added to solutions of the other metal ions, metallic B and D are formed

(III) Metal C does not reduce A^{n+} .

(a) $C\ \ A\ \ B\ \ D$ (b) $C\ \ A\ \ D\ \ B$ (c) $A\ \ C\ \ D\ \ B$ (d) $A\ \ C\ \ B\ \ D$

76. When an aqueous solution of $CuSO_4$ is stirred with a silver spoon then :

(a) Cu will be formed (b) Ag will be formed

(c) Cu^{2+} will be deposited (d) None of these

77. Based on the following informations arrange four metals, A, B, C and D in order of increasing ability to act as reducing agents :

(I) Only C react with 1 M HCl to give $H_2(g)$

(II) When A is added to solution of the other metal ions, metallic D is formed but not B or C

(a) $D\ \ A\ \ C\ \ B$ (b) $A\ \ D\ \ C\ \ B$ (c) $B\ \ D\ \ A\ \ C$ (d) $D\ \ A\ \ B\ \ C$

78. In the reaction :

$4Fe(s)\quad 3O_2(g)\qquad 4Fe^{3+}(aq)\quad 6O^{2-}(aq)$

For More Material Join: @JEEAdvanced_2024

which of the following statement is incorrect ?

(a) It is a redox reaction (b) Fe is reducing agent

(c) O_2 is an oxidizing agent (d) Fe is reduced to Fe^3

79. Which of the following is displaced by Fe ?

(a) Ag (b) Zn (c) Na (d) All of these

80. The standard potential at 25°C for the following half reactions is given :

$Zn^2 \quad 2e \quad Zn ; E^o \quad 0.762$ V

$Mg^2 \quad 2e \quad Mg ; E^o \quad 2.37$ V

When Zinc dust is added to the solution of $MgCl_2$.

(a) $ZnCl_2$ is formed (b) Mg is precipitated

(c) Zn dissolved in the solution (d) No reaction takes place

81. The element which can displace three other halogens from their compound is :

(a) F (b) Cl (c) Br (d) I

82. Using the standard half-cell potential listed, calculate the equilibrium constant for the reaction :

$Co(s) \quad 2H \ (aq) \quad Co^2 \ (aq) \quad H_2(g)$ at 298 K

$Co^2 \ (aq) \quad 2e \quad Co(s) \quad E \quad 0.277$ V

(a) $2.3 \quad 10^9$ (b) $4.8 \quad 10^4$ (c) $4.8 \quad 10^7$ (d) $4.8 \quad 10^{11}$

83. The E at 25 C for the following reaction is 0.22 V. Calculate the equilibrium constant at 25 C :

$H_2(g) \quad 2AgCl(s) \quad 2Ag(s) \quad 2HCl(aq)$

(a) $2.8 \quad 10^7$ (b) $5.2 \quad 10^8$ (c) $5.2 \quad 10^6$ (d) $5.2 \quad 10^3$

84. Electrode potential of the half cell $Pt(s) | Hg(l) | Hg_2Cl_2(s) | Cl \ (aq)$ can be increased by :

(a) Increasing [Cl] (b) Decreasing [Cl]

(c) Increasing $Hg_2Cl_2(s)$ (d) Decreasing $Hg(l)$

85. The equilibrium constant for the following general reaction is 10^{30}. Calculate E for the cell at 298 K.

$2X_2(s) \quad 3Y^2 \ (aq) \quad 2X_2^3 \ (aq) \quad 3Y(s)$

(a) +0.105 V (b) +0.2955 V (c) 0.0985 V (d) 0.2955 V

86. A solution containing H and D ions is in equilibrium with a mixture of H_2 and D_2 gases at 25 C. If the partial pressures of both gases are 1.0 atm, find the ratio of [D]/[H] :

(Given : $E_{D^+/D_2} \quad 0.003$ V)

(a) 1.23 (b) 1.12 (c) 0.11 (d) 1.0

87. The E at 25 C for the following reaction is 0.55 V. Calculate the G in kJ/mol :

$4BiO \ (aq) \quad 3N_2H_5^+ \quad 4Bi(s) \quad 3N_2(g) \quad 4H_2O(l) \quad 7H \ (aq)$

(a) 637 (b) 424 (c) 106 (d) 318.5

88. Use the following standard electrode potentials, calculate G in kJ/mol for the indicated reaction :

$5Ce^4 \ (aq) \quad Mn^2 \ (aq) \quad 4H_2O(l) \quad 5Ce^3 \ (aq) \quad MnO_4(aq) \quad 8H \ (aq)$

$MnO_4^-(aq) \quad 8H \ (aq) \quad 5e^- \quad Mn^2 \ (aq) \quad 4H_2O(l); \quad E \quad 1.51$ V

For More Material Join: @JEEAdvanced_2024

$Ce^{4}\ (aq)\quad e\qquad Ce^{3}\ (aq)\qquad E\quad 1.61\,V$

(a) 9.65 (b) 24.3 (c) 48.25 (d) 35.2

89. Consider an electrochemical cell in which the following reaction occurs and predict which changes will decrease the cell voltage :

$Fe^{2}\ (aq)\quad Ag\ (aq)\qquad Ag(s)\quad Fe^{3}\ (aq)$

(I) decrease the $[Ag^{+}]$ (II) increase in $[Fe^{3+}]$ (III) increase the amount of Ag

(a) I (b) II and III (c) II (d) I and II

90. Consider the following equation for an electrochemical cell reaction. Which of the following changes in condition will increase the cell voltage?

$H_2(g)\quad PbCl_2(s)\qquad Pb(s)\quad 2H^{+}(aq)\quad 2Cl\ (aq)$

(I) addition of concentrated $HClO_4$ in the cell solution

(II) increase the pressure of $H_2(g)$ (III) increase the amount of $Pb(s)$

(a) III (b) I and II (c) II and III (d) II

91. The standared electrode potential for the following reaction is +1.33 V. What is the potential at pH 2.0?

$Cr_2O_7^{2-}(aq, 1M)\quad 14H\ (aq)\quad 6e\qquad 2Cr^{3}\ (aq, 1M)\quad 7H_2O(l)$

(a) +1.820 V (b) +1.990 V (c) +1.608 V (d) +1.0542 V

92. The standard electrode potential for the following reaction is 0.57 V. What is potential at pH = 12.0?

$TeO_3^{2-}(aq, 1M)\quad 3\,H_2O(l)\quad 4e\qquad Te(s)\quad 6\,OH\ (aq)$

(a) 0.17 V (b) 0.21 V (c) 0.39 V (d) +1.95 V

93. $Co|Co^{2+}(C_2)||Co^{2+}(C_1)|Co$; for this cell, G is negative if :

(a) C_2 C_1 (b) $C_1 > C_2$ (c) $C_1 = C_2$ (d) unpredictable

94. What will be the emf for the given cell?

$Pt|H_2(g, P_1)|H\ (aq)|H_2(g, P_2)|Pt$

(a) $\frac{RT}{F}\ln\frac{P_1}{P_2}$ (b) $\frac{RT}{2F}\ln\frac{P_1}{P_2}$ (c) $\frac{RT}{F}\ln\frac{P_2}{P_1}$ (d) None of these

95. For the electrochemical cell $Pt(s)|H_2(g)|_{1\ atm}|H\ (1M)||Cu^{2}\ (1M)|Cu(s)$, which one of the following statements is true ?

(a) H ions are formed at anode and Cu is deposited at cathode.

(b) H_2 liberated at cathode and Cu is deposited at anode.

(c) Oxidation occurs at cathode.

(d) Reduction occurs at anode.

96. In a concentration cell the same reagents are present in both the anode and the cathode compartments, but at different concentrations. Calculate the emf of a cell containing 0.040 M. Cr^{3+} in one compartment and 1.0 M Cr^{3+} in the other if Cr electrodes are used in both.

(a) 0.028 V (b) 0.249 V (c) 0.083 V (d) 0.125 V

97. A 1.0 M solution of Cd^{2+} is added to excess iron and the system is allowed to reach equilibrium. What is the concentration of Cd^{2+} ?

$Cd^{2+}(aq)\quad Fe(s)\qquad Cd(s)\quad Fe^{2}\ (aq);\quad E\quad 0.037$

For More Material Join: @JEEAdvanced_2024

(a) 0.195 (b) 0.097 (c) 0.053 (d) 0.145

98. The measured voltage for the reaction with the indicated concentrations is 1.50 V. Calculate E .

$$Cr(s) \quad 3Ag \ (aq, 0.10\ M) \qquad 3Ag(s) \quad Cr^{3} \ (aq, 0.30\ M)$$

(a) 1.35 (b) 1.40 (c) 1.65 (d) 1.55

99. Calculate the standard voltage that can be obtained from an ethane oxygen fuel cell at 25 C.

$$C_2H_6(g) \quad 7/2\,O_2(g) \qquad 2CO_2(g) \quad 3H_2O(l); \quad G \quad 1467\ kJ/mol$$

(a) +0.91 (b) +0.54 (c) +0.72 (d) +1.08

100. $I_2(s)|I$ $(0.1M)$ half cell is connected to a H $(aq)|H_2$ (1 bar)|Pt half cell and e.m.f. is found to be 0.7714 V. If $E_{I_2|I}$ 0.535 V, find the pH of $H^+|H_2$ half-cell.

(a) 1 (b) 3 (c) 5 (d) 7

101. What is the SRP for $S^{2-}|CuS|Cu$ half-cell?

Given : $K_{sp}(CuS)$ 10^{-35} and $E_{Cu^{2+}|Cu} = 0.34$ V

(a) 1.034 V (b) 1.0 V (c) 0.694 V (d) 0.694 V

102. Given the following standard electrode potentials, the K_{sp} for $PbBr_2$ is :

$$PbBr_2(s) \quad 2e \qquad Pb(s) \quad 2Br \ (aq); \quad E \quad 0.248\ V$$

$$Pb^{2} \ (aq) \quad 2e \qquad Pb(s)\ ; \qquad E \quad 0.126\ V$$

(a) 7.4 10^{-5} (b) 4.9 10^{-14} (c) 5.2 10^{-6} (d) 2.3 10^{-13}

103. The standard free energy change for the following reaction is 210 kJ/mol. What is the standard cell potential?

$$2H_2O_2(aq) \qquad 2H_2O(l) \quad O_2(g)$$

(a) +0.752 (b) +1.09 (c) +0.420 (d) +0.640

104. At equilibrium :

(a) E°_{cell} 0, G° 0 (b) E_{cell} 0, G 0

(c) both are correct (d) none is correct

105. The E at 25 C for the following reaction at the indicated concentrations is 1.50 V. Calculate the G in kJ/mol 25 C :

$$Cr(s) \quad 3Ag \ (aq, 0.1\,M) \qquad 3Ag(s) \quad Cr^{3} \ (aq, 0.1\,M)$$

(a) 140.94 (b) 295 (c) 212 (d) 422.83

106. If $E_{Au\ /Au}$ is 1.69 V and $E_{Au^{3+}/Au}$ is 1.40 V, then $E_{Au^{+}/Au^{3+}}$ will be :

(a) 0.19 V (b) 2.945 V (c) 1.255 V (d) None of these

107. Consider the following standard electrode potentials and calculate the equilibrium constant at 25 C for the indicated disproportionation reaction :

$$3\,Mn^{2+}(aq) \qquad Mn(s) + 2\,Mn^{3+}(aq)$$

$$Mn^{3+}(aq) \quad e \qquad Mn^{2} \ (aq); \quad E \quad 1.51\,V$$

$$Mn^{2} \ (aq) \quad 2e \qquad Mn(s); \qquad E \quad 1.185\,V$$

(a) 1.2 10^{-43} (b) 2.4 10^{-73} (c) 6.3 10^{-92} (d) 1.5 10^{-62}

For More Material Join: @JEEAdvanced_2024

108. A galvanic cell is composed of two hydrogen electrodes, one of which is a standard one. In which of the following solutions should the other electrode be immersed to get maximum e.m.f. :
(a) 0.1 M HCl (b) 0.1 M H_2SO_4 (c) 0.1 M NH_4OH (d) 0.01 M HCOOH

109. $Ag|AgCl|Cl^-(C_2)||Cl^-(C_1)|AgCl|Ag$ for this cell G is negative if :
(a) C_1 C_2 (b) $C_1 > C_2$ (c) C_2 C_1 (d) Both (a) and (c)

110. By how much is the oxidizing power of $Cr_2O_7^{2-}|Cr^{3+}$ couple decreased if the H concentration is decreased from 1 M to 10^{-3} M at 25 C?
(a) 0.001 V (b) 0.207 V (c) 0.441 V (d) 0.414 V

111. The temperature coefficient of a cell whose operation is based on the reaction

$$Pb(s) \quad HgCl_2(aq) \quad PbCl_2(aq) \quad Hg(l) \text{ is :}$$

$$\left(\frac{dE}{dT}\right)_p \quad 1.5 \quad 10^{-4}\,VK^{-1} \text{ at 298 K}$$

The change in entropy (in J/K mol) during the operation is :
(a) 8627 (b) 57.9 (c) 28.95 (d) 14.475

112. Thermodynamic efficiency of a cell is given by :
(a) $\frac{H}{G}$ (b) $\frac{nFE}{G}$ (c) $\frac{nEF}{H}$ (d) nFE°

113. Calculate the value of equilibrium constant (K_f) for the reaction :

$$Zn^{2+}(aq) \quad 4OH^-(aq) \rightleftharpoons Zn(OH)_4^{2-}(aq)$$

Given : $Zn^{2+}(aq) \quad 2e^- \quad Zn(s); \quad E \quad 0.76$ V

$Zn(OH)_4^{2-}(aq) \quad 2e \quad Zn(s) \quad 4OH^-(aq); \quad E \quad 1.36$ V

$$2.303\frac{RT}{F} \quad 0.06$$

(a) 10^{10} (b) $2 \quad 10^{10}$ (c) 10^{20} (d) none of these

114. Which of the following statement is false for fuel cells ?
(a) They are more efficient (b) They are free from pollution
(c) They run till reactants are active (d) Fuel burned with O_2

115. When a lead storage battery is charged it acts as :
(a) a fuel cell (b) an electrolytic cell
(c) a galvanic cell (d) a concentration cell

116. The metal that forms a self-protecting film of oxide to prevent corrosion is :
(a) Na (b) Al (c) Cu (d) Au

117. Rusting of iron is catalyzed by which of the following ?
(a) Fe (b) Zn (c) O_2 (d) H

118. Which of the following is a highly corrosive salt ?
(a) Hg_2Cl_2 (b) $HgCl_2$ (c) $FeCl_2$ (d) $PbCl_2$

119. The Zn acts as sacrificial or cathodic protection to prevent rusting of iron because :
(a) E°_{OP} of Zn E°_{OP} of Fe (b) E°_{OP} of Zn E°_{OP} of Fe
(c) E°_{OP} of Zn E°_{OP} of Fe (d) Zn is cheaper than iron

For More Material Join: @JEEAdvanced_2024

120. In electrochemical corrosion of metals, the metal undergoing corrosion :
(a) acts as anode (b) acts as cathode
(c) undergoes reduction (d) None

121. When an acid cell is charged, then :
(a) voltage of cell increases (b) resistance of cell increases
(c) electrolyte of cell dilutes (d) None of these

122. Electrolytic conduction is due to the movement of :
(a) electrons (b) ions
(c) atoms (d) electrons as well as ions

123. Molten sodium chloride conducts electricity due to the presence of :
(a) free electron (b) free ions
(c) free molecules (d) atoms of sodium and chlorine

124. Pure water does not conduct electricity because it :
(a) is neutral (b) is readily decomposed
(c) is almost totally unionized (d) has a low boiling point

125. The relation among conductance (G), specific conductance ( ) and cell constant (l/A) is :
(a) G $\frac{l}{A}$ (b) G $\frac{A}{l}$ (c) G $\frac{l}{A}$ (d) G Al

126. If x is specific resistance (in S^{-1}cm) of the electrolyte solution and y is the molarity of the solution, then $_m$ (in $S\,cm^2mol^{-1}$) is given by :
(a) $\frac{1000x}{y}$ (b) $1000\frac{y}{x}$ (c) $\frac{1000}{xy}$ (d) $\frac{xy}{1000}$

127. Equivalent conductivity can be expressed in terms of specific conductance ( ) and concentration (N) in gram equivalent dm^{-3} as :
(a) N (b) $\frac{1000}{N}$ (c) $\frac{N}{1000}$ (d) N 1000

128. Resistance of a decimolar solution between two electrodes 0.02 meter apart and 0.0004 m^2 in area was found to be 50 ohm. Specific conductance ( ) is :
(a) 0.1 Sm^{-1} (b) 1 $S\,m^{-1}$ (c) 10 $S\,m^{-1}$ (d) 4 10^{-4} $S\,m^{-1}$

129. Resistance of 0.1 M KCl solution in a conductance cell is 300 ohm and conductivity is 0.013 Scm^{-1}. The value of cell constant is :
(a) 3.9 cm^{-1} (b) 39 m^{-1} (c) 3.9 m^{-1} (d) None of these

130. Dissociation constant of a weak acid (HA) in terms of $_m$ and $_m$ is :
(a) K_a $\frac{C\ _m}{(\ _m\ \ )}$ (b) K_a $\frac{C\ ^2_m}{_m(\ _m\ \ _m)}$
(c) K_a $\frac{C(\ _m)^2}{_m(\ _m\ \ _m)}$ (d) None of these

131. When a concentrated solution of an electrolyte is diluted
(a) its specific conductance increases
(b) its equivalent conductivity decreases
(c) its specific conductivity decreases and equivalent conductivity increases
(d) both specific and equivalent conductivity increase

For More Material Join: @JEEAdvanced_2024

132. Molar conductivity of a solution of an electrolyte AB_3 is 150 Scm^2mol^{-1}. If it ionises as AB_3 A^3 $3B^-$, its equivalent conductivity will be :

(a) 150 (in Scm^2eq^{-1}) (b) 75 (in Scm^2eq^{-1})

(c) 50 (in Scm^2eq^{-1}) (d) 80 (in Scm^2eq^{-1})

133. Equivalent conductivity of $Fe_2(SO_4)_3$ is related to molar conductivity by the expression :

(a) $_{eq}$ $_m$ (b) $_{eq}$ $_m/3$ (c) $_{eq}$ 3 $_m$ (d) $_{eq}$ $_m/6$

134. The limiting equivalent conductivity of NaCl, KCl and KBr are 126.5, 150.0 and 151.5 $S\,cm^2$ eq^{-1}, respectively. The limiting equivalent ionic conductivity for Br is 78 $S\,cm^2eq^{-1}$. The limiting equivalent ionic conductivity for Na^+ ions would be :

(a) 128 (b) 125 (c) 49 (d) 50

135. The specific conductance of a saturated solution of silver bromide is Scm^{-1}. The limiting ionic conductivity of Ag^+ and Br ions are x and y, respectively. The solubility of silver bromide in gL^{-1} is : (molar mass of AgBr = 188)

(a) $\frac{1000}{x\ y}$ (b) $\frac{}{x\ y}$ 188 (c) $\frac{1000\ \ 188}{x\ y}$ (d) $\frac{x\ y}{}\ \frac{1000}{188}$

136. The resistance of 0.1 *N* solution of formic acid is 200 ohm and cell constant is 2.0 cm^{-1}. The equivalent conductivity (in Scm^2eq^{-1}) of 0.1 *N* formic acid is :

(a) 100 (b) 10 (c) 1 (d) none of these

137. A conductance cell was filled with a 0.02 *M* KCl solution which has a specific conductance of 2.768 10 ^{3} ohm ^{1} cm ^{1}. If its resistance is 82.4 ohm at 25°C, the cell constant is :

(a) 0.2182 cm ^{1} (b) 0.2281 cm ^{1} (c) 0.2821 cm ^{1} (d) 0.2381 cm ^{1}

138. The ionic conductivity of Ba^{2+} and Cl at infinite dilution are 127 and 76 $ohm^{-1}\,cm^2eq^{-1}$ respectively. The equivalent conductivity of $BaCl_2$ at infinity dilution (in $ohm^{-1}cm^2eq^{-1}$) would be :

(a) 203 (b) 279 (c) 101.5 (d) 139.5

139. Unit of ionic mobility is :

(a) $mV^{-1}s^{-1}$ (b) $m^2V^{-2}s^{-1}$ (c) $m^2V^{-1}s^{-1}$ (d) $m^{-2}Vs^{-1}$

140. $_{AgCl}$ can be obtained :

(a) by extraplotation of the graph and $\sqrt{C}$ to zero concentration

(b) by known values of of $AgNO_3$, HCl and HNO_3

(c) both (a) and (b)

(d) None of these

141. The conductance of a salt solution (*AB*) measured by two parallel electrodes of area 100 cm^2 separated by 10 cm was found to be 0.0001 $^{-1}$. If volume enclosed between two electrode contain 0.1 mole of salt, what is the molar conductivity (Scm^2mol^{-1}) of salt at same concentration :

(a) 10 (b) 0.1 (c) 1 (d) none of these

For More Material Join: @JEEAdvanced_2024


142. The conductivity of a strong electrolyte :
(a) increases on dilution (b) decreases on dilution
(c) does not change with dilution (d) depends upon density of electrolytes

143. The increase in equivalent conductivity of a weak electrolyte with dilution is due to :
(a) increase in degree of dissociation and decrease in ionic mobility
(b) decrease in degree of dissociation and decrease in ionic mobility
(c) increase in degree of dissociation and increase in ionic mobility
(d) decrease in degree of dissociation and increase in ionic mobility

144. Strong electrolytes are those which :
(a) conduct electricity (b) dissolve readily in water
(c) dissociate into ions at high dilution (d) completely dissociate into ions

145. The electric conduction of a salt solution in water depends on the :
(a) size of its molecules (b) shape of its molecules
(c) size of solvent molecules (d) extent of its ionization


146. A graph was plotted between molar conductivity of various electrolytes (NaCl, HCl and NH_4OH) and $\sqrt{C}$ (in $mol\,L^{-1}$). Correct set is :





(a) I(NaCl), II(HCl), III(NH_4OH)
(b) I(HCl), II(NaCl), III(NH_4OH)
(c) I(NH_4OH), II(NaCl), III(HCl)
(d) I(NH_4OH), II(HCl), III(NaCl)


147. Which of the following is arranged in increasing order of ionic mobility ?
(a) I Br Cl F (b) F Cl Br I
(c) F I Cl Br (d) F Cl I Br

148. $HNO_3(aq)$ is titrated with NaOH(aq) conductometrically, graphical representation of the titration is :

(a)



(b)



(c)



(d)



For More Material Join: @JEEAdvanced_2024

149. Which of the following plots will obtained for a conductometric titration of strong acid against a weak base?


(a) 


(b) 


(c) 


(d) 

150. Conductometric titration curve of a equimolar mixture of a HCl and HCN with NaOH(*aq*) is :

(a) 

(b) 

(c) 

(d) 

For More Material Join: @JEEAdvanced_2024



Level 2

1. In the Hall process, aluminium is produced by the electrolysis of molten Al_2O_3. How many second would it take to produce enough aluminum by the Hall process to make a case of 24 cans of aluminum soft-drink, if each can uses 5.0 g of Al, a current of 9650 amp is employed, and the current efficiency of the cell is 90.0% :

(a) 203.2 (b) 148.14 (c) 333 (d) 6.17

2. 108 g fairly concentrated solution of $AgNO_3$ is electrolysed by using 0.1 F charge. The mass of resulting solution is :

(a) 94 g (b) 11.6 g (c) 96.4 g (d) None of these

3. The electrolysis of acetate solution produces ethane according to reaction :

$$2CH_3COO^- \quad C_2H_6(g) \quad 2CO_2(g) \quad 2e$$

The current efficiency of the process is 80%. What volume of gases would be produced at 27 C and 740 torr, if the current of 0.5 amp is passed through the solution for 96.5 min?

(a) 6.0 L (b) 0.60 L (c) 1.365 L (d) 0.91 L

4. A layer of chromium metal 0.25 mm thick is to be plated on an auto bumper with a total area of 0.32 m^2 from a solution containing CrO_4^{2-}? What current flow is required for this electroplating if the bumper is to be plated in 60 s ? The density of chromium metal is 7.20 g/cm^3.

(a) 4.9 10^3 A (b) 1.78 10^3 A (c) 5.3 10^4 A (d) 10.69 10^6 A

5. 100 mL of 0.05 *M* $CuSO_4(aq)$ solution was electrolysed using inert electrodes by passing current till the pH of the resulting solution was 2. The solution after electrolysis was neutralized and then treated with excess KI and formed I_2 titrated with 0.04 *M* $Na_2S_2O_3$. Calculate the required volume (in mL) of $Na_2S_2O_3$:

(a) 112.5 mL (b) 100 mL (c) 125 mL (d) None of these

6. If the equilibrium constant for the reaction H (aq) $OH^-(aq) \rightleftharpoons H_2O(l)$ is 10^{13} at certain temperature then what is the E for the reaction, $2H_2O(l)$ $2e^- \rightleftharpoons H_2(g)$ $2OH^-(aq)$

Given : $\frac{2.303\,RT}{F}$ 0.066

(a) 1.230 V (b) 0.858 V (c) 0.80 V (d) 0.8274 V

7. A fuel cell develops an electrical potential from the combustion of butane at 1 bar and 298 K

$$C_4H_{10}(g) \quad 6.5O_2(g) \quad 4CO_2(g) \quad 5H_2O(l); \quad {}_rG \quad 2746\ kJ/mol$$

What is E of a cell?

(a) 4.74 V (b) 0.547 V (c) 4.37 V (d) 1.09 V

8. The cell $Pt|H_2(g, 0.1\,bar)|H^+$ (aq), pH $X||Cl^-(aq, 1\,M)|Hg_2Cl_2|Hg|Pt$,

has e.m.f. of 0.5755 V at 25 C. The SOP of calomel electrode is 0.28V, then pH of solution will be :

(a) 11 (b) 4.5 (c) 5.5 (d) None of these

For More Material Join: @JEEAdvanced_2024

9. For a cell reaction $2H_2(g)$ $O_2(g)$ $2H_2O(l)$ $_rS_{298}$ 0.32 kJ/K. What is the value of $_fH_{298}(H_2O, l)$?

Given : $O_2(g)$ $4H$ (aq) $4e$ $2H_2O(l)$; E 1.23 V

(a) 285.07 kJ/mol (b) 570.14 kJ/mol
(c) 285.07 kJ/mol (d) None of these

10. What is the potential of an electrode which originally contained 0.1 M NO_3^- and 0.4 M H^+ and which has been treated by 80% of the cadmium necessary to reduce all the NO_3^- to $NO(g)$ at 1 bar?

Given : NO_3^- $4H$ $3e$ $NO + 2H_2O$; E 0.96 V; $\log 2 = 0.3$

(a) 0.84 V (b) 1.08 V
(c) 1.23 V (d) 1.36 V

11. The standard reduction potential of normal calomel electrode and reduction potential of saturated calomel electrodes are 0.27 and 0.33 volt respectively. What is the concentration of Cl^- in saturated solution of KCl?

(a) 0.1 M (b) 0.01 M (c) 0.001 (d) None

12. Determine the potential of the following cell :

$Pt|H_2(g, 0.1\,bar)|H$ $(aq, 10^{-3}\ M)||MnO_4^-(aq, 0.1\,M)$,
Mn^2 $(aq, 0.01\,M)$, H $(aq, 0.01\,M)|Pt$

Given : $E_{MnO_4^-|Mn^{2+}}$ 1.51 V

(a) 1.54 V (b) 1.48 V (c) 1.84 V (d) none of these

13. Copper reduces NO_3^- into NO and NO_2 depending upon concentration of HNO_3 in solution. Assuming $[Cu^{2+}] = 0.1\,M$, and P_{NO} P_{NO_2} 10^{-3} bar, at which concentration of HNO_3, thermodynamic tendency for reduction of NO_3^- into NO and NO_2 by copper is same?

[Given : $E_{Cu^{2+}|Cu}$ 0.34 volt, $E_{NO_3^-|NO}$ 0.96 volt, $E_{NO_3^-|NO_2}$ 0.79 volt]

(a) $10^{1.23}\ M$ (b) $10^{0.56}\ M$ (c) $10^{0.66}\ M$ (d) $10^{0.12}\ M$

14. For the cell, $Pt|Cl_2(g, 0.4\,bar)|Cl$ $(aq, 0.1\,M)||Cl$ $(aq, 0.01\,M)|Cl_2(g, 0.2\,bar)|Pt$

The measured potential at 298 K is:

(a) 0.051 V (b) 0.051 V (c) 0.102 V (d) 0.0255 V

15. The chlorate ion can disproportionate in basic solution according to reaction,

$$2ClO_3^- \rightleftharpoons ClO_2^- + ClO_4^-$$


What is the equilibrium concentration of perchlorate ions from a solution initially at 0.1 M in chlorate ions at 298 K?

Given : $E_{ClO_4^-|ClO_3^-}$ 0.36 V and $E_{ClO_3^-|ClO_2^-}$ 0.33 V at 298 K

(a) 0.019 M (b) 0.024 M (c) 0.1 M (d) 0.19 M

For More Material Join: @JEEAdvanced_2024

16. A cell diagram shown below contains one litre of buffer solution of $HA(pK_a = 4)$ and NaA in both compartments. What is the cell e.m.f.?



(a) 0.03 V (b) 0.06 V (c) 0.06 V (d) None of these

17. Given the cell : $Cd(s)|Cd(OH)_2(s)|NaOH(aq, 0.01\,M)|H_2(g, 1\,bar)|Pt(s)$ with E_{cell} 0.0 V. If $E_{Cd^{2+}|Cd}$ 0.39 V, then K_{sp} of $Cd(OH_2)$ is :

(a) 0.1 (b) 10^{-13} (c) 10^{-15} (d) None of these

18. Calculate the e.m.f. (in V) of the cell :

$$Pt|H_2(g)|BOH(aq)||HA(aq)|H_2(g)|Pt;$$
$$0.1\ bar \quad 1\ M \qquad 0.1\ M \quad 1\ bar$$

Given : $K_a(HA) = 10^{-7}$, $K_b(BOH) = 10^{-6}$

(a) 0.39 V (b) 0.36 V (c) 0.93 V (d) None of these

19. Calculate the potential of a half cell having reaction : $Ag_2S(s)$ $2e^-$ $\rightleftharpoons$ $2Ag(s)$ $S^{2-}(aq)$ in a solution buffered at pH = 3 and which is also saturated with $0.1\,M\,H_2S(aq)$:
[Given: $K_{sp}(Ag_2S)$ 10^{-49}, K_{a_1} K_{a_2} 10^{-21}]

(a) 1.18 (b) 0.19 (c) 0.19 V (d) none of these

20. The conductivity of 0.1 *N* NaOH solution is 0.022 S cm^{-1}. When equal volume of 0.1 *N* HCl solution is added, the conductivity of resultant solution is decreased to 0.0055 S cm^{-1}. The equivalent conductivity in $Scm^2 equivalent^{-1}$ of NaCl solution is :

(a) 0.0055 (b) 0.11 (c) 110 (d) None of these

21. In above question after formation of NaCl, further 0.1 *N* HCl is added, the volume of which is double to that of the first portion added, the conductivity increases to 0.018 Scm^{-1}. The value of $_{eq}$(HCl) is [assume no change in equivalent conductivity of NaCl (*aq*)] :

(a) 330 S $cm^2 eq^{-1}$ (b) 305 S $cm^2 eq^{-1}$
(c) 415 S $cm^2 eq^{-1}$ (d) 360 S $cm^2 eq^{-1}$

22. Given the following molar conductivities at infinite dilution ($_m$), HCl, 426 $^{-1}$ $cm^2 mol^{-1}$; NaCl, 126 $^{-1}$ $cm^2 mol^{-1}$; NaC (sodium crotonate), 83 $^{-1} cm^2 mol^{-1}$. What is the dissociation constant of crotonic acid, if the conductivity of a 0.001 *M* crotonic acid solution is 3.83 10^{-5} $^{-1} cm^{-1}$?

(a) 10^{-5} (b) 1.11 10^{-5} (c) 1.11 10^{-4} (d) 0.01

For More Material Join: @JEEAdvanced_2024

23. Equivalent conductivity of $BaCl_2, H_2SO_4$ and HCl, are x_1, x_2 and x_3 $Scm^{-1}eq^{-1}$ at infinite dilution. If conductivity of saturated $BaSO_4$ solution is $x\,Scm^{-1}$, then K_{sp} of $BaSO_4$ is :

(a) $\dfrac{500x}{(x_1 \quad x_2 \quad 2x_3)}$ (b) $\dfrac{10^6 x^2}{(x_1 \quad x_2 \quad 2x_3)^3}$

(c) $\dfrac{2.5 \quad 10^5 x^2}{(x_1 \quad x_2 \quad x_3)^2}$ (d) $\dfrac{0.25x^2}{(x_1 \quad x_2 \quad x_3)^2}$

24. The conductivity of $0.001\,M$ Na_2SO_4 solution is $2.6 \quad 10^{-4}$ Scm ^{1} and increases to $7.0 \quad 10^{-4}$ Scm^{-1}, when the solution is saturated with $CaSO_4$. The molar conductivities of Na^+ and Ca^{2+} are 50 and $120\,Scm^2mol^{-1}$, respectively. Neglect conductivity of used water.

What is the solubility product of $CaSO_4$?

(a) $4 \quad 10^{\ 6}$ (b) $1.57 \quad 10^{-3}$

(c) $4 \quad 10^{\ 4}$ (d) $2.46 \quad 10^{-6}$

25. The dissociation constant of a weak acid is $1.6 \quad 10^{-5}$ and the molar conductivity at infinite dilution is $380 \quad 10^{-4}\,S\,m^2mol^{-1}$. If the cell constant is $0.01\,m^{-1}$ then conductance of 0.01 M acid solution is :

(a) $1.52 \quad 10^{-5}$ S (b) 1.52 S

(c) $1.52 \quad 10^{-3}$ S (d) $1.52 \quad 10^{-4}$ S

26. Three electrolytic cells X, Y, Z containing solution of $1M$ each NaCl, $AgNO_3$ and $CuSO_4$ respectively are connected in series combination. During electrolysis 21.6 gm of silver deposits at cathode in cell Y. Which is incorrect statement.

(a) 6.35 gm copper deposits at cathode in cell Z

(b) 2.24 litre Cl_2 is liberated (at 1 atm and 273 K) at anode in cell X

(c) 2.24 litre O_2 is liberated (at 1 atm and 273 K) at anode in cell Y

(d) 2.24 litre H_2 is liberated (at 1 atm and 273 K) at cathode in cell X

27. During electrolysis of $H_2SO_4(aq)$ with high charge density, $H_2S_2O_8$ formed as by product. In such electrolysis 22.4 L $H_2(g)$ and 8.4 L $O_2(g)$ liberated at 1 atm and 273 K at electrode. The moles of $H_2S_2O_8$ formed is :

(a) 0.25 (b) 0.50 (c) 0.75 (d) 1.00

28. $Zn(s)\,|\,Zn(CN)_4^{2}\ (0.5M), CN\ (0.01)\,||\,Cu(NH_3)_4^{2}\ (0.5M), NH_3(1M)\,|\,Cu(s)$

Given: K_f of $Zn(CN)_4^{\ 2} \quad 10^{16}$, K_f of $Cu(NH_3)_4^{2} \quad 10^{12}$,

$E^{\circ}_{Zn|Zn^{\ 2}} \quad 0.76\,V$; $E^{\circ}_{Cu^{\ 2}|Cu} \quad 0.34\,V$, $\dfrac{2.303RT}{F} \quad 0.06$


The emf of above cell is :

(a) 1.22 V (b) 1.10 V (c) 0.98 V (d) None of these

For More Material Join: @JEEAdvanced_2024

Level 3

Passage 1



A Galvanic cell consist of three compartment as shown in figure. The first compartment contain $ZnSO_4(1M)$ and III compartment contain $CuSO_4(1M)$. The mid compartment contain $NaNO_3(1M)$. Each. compartment contain 1 L solution :

$$E_{Zn^{2}/Zn} \quad 0.76; \quad E_{Cu^{2}/Cu} \quad 0.34$$

1. The concentration of Zn^{2+} in first compartment after passage of 0.1 F charge will be :
(a) 1 *M* (b) 1.05 *M* (c) 1.025 *M* (d) 0.5 *M*

2. The concentration of NO_3^- in mid compartment after passage of 0.1 F of charge will be :
(a) 0.95 *M* (b) 0.90 *M* (c) 0.975 *M* (d) 1.05 *M*

3. The concentration of SO_4^{2-} ion in III compartment after passage of 0.1 F of charge will be :
(a) 1.05 *M* (b) 1.025 *M* (c) 0.95 *M* (d) 0.975 *M*

Passage 2

The cell potential (E_{cell}) of a reaction is related as $G \quad nF\,E_{cell}$, where G represents max. useful electrical work
n no. of moles of electrons exchanged during the reaction
for reversible cell reaction $d(G) \quad (\ _rV)dp \quad (\ _rS).\,dT$
at constant pressure $d(G) \quad (\ _rS)\ dT$
∵ At constant pressure

$$G \quad H \quad T \quad S \qquad \ldots(1)$$

$$G \quad H \quad T\,\frac{d(G)}{dT}_p \qquad \ldots(2)$$

$\left(\frac{dE_{cell}}{dT}\right)_p$ is known as temperature coefficient of the e.m.f of the cell

For More Material Join: @JEEAdvanced_2024

1. The temperature coefficient of the e.m.f. of cell, $\left(\frac{dE}{dT}\right)_p$ is given by :

(a) $\frac{nF}{S}$ (b) $\frac{S}{nF}$ (c) $\frac{S}{nFT}$ (d) nFE

2. At 300 K, H for the reaction

$Zn(s)$ $AgCl(s)$ $ZnCl_2(aq)$ $2Ag(s)$ is

218 kJ/mol while the e.m.f. of the cell was 1.015 V. $\left(\frac{dE}{dT}\right)_p$ of the cell is :

(a) $4.2 \quad 10^{-4}\,VK^{-1}$ (b) $3.81 \quad 10^{-4}\,VK^{-1}$

(c) $0.11\,VK^{-1}$ (d) $7.62 \quad 10^{-4}\,VK^{-1}$

3. Calculate S for the given cell reaction in Q. no. 2 :

(a) 73.53 J/K mol (b) 83.53 J/K mol (c) 100 J/K mol (d) none of these



Passage 3

Molar conductivity ($_m$) is defined as conducting power of the ions produced by 1 mole of an electrolyte in a solution. $_m$ $\frac{}{C}$ where is conductivity (in S cm ^{1}) and C is molar concentration (in mol/cm^3)

The molar conductivity of 0.04 M solution of $MgCl_2$ is 200 S cm^2 mol^{-1} at 298 K. A cell with electrodes that are 2.0 cm^2 in surface area and 0.50 cm apart is filled with $MgCl_2$ solution.

1. Conductance of $MgCl_2$ solution is :

(a) $8 \quad 10^{-3}$ S (b) 32 S (c) 0.032 S (d) None of these

2. How much current will flow when the potential difference between the two electrodes is 5.0 V?

(a) 156.25 A (b) 0.16 A (c) 160 A (d) None of these



Passage 4

In a hydrogen oxygen fuel cell, electricity is produced. In this process $H_2(g)$ is oxidised at anode and $O_2(g)$ reduced at cathode.

Given : Cathode $O_2(g)$ $2H_2O(l)$ $4e$ $4OH^-(aq)$

Anode $H_2(g)$ $2OH^-(aq)$ $2H_2O(l)$ $2e$

4.48 litre H_2 at 1 atm and 273 K oxidised in 9650 sec.

1. The current produced is (in amp) :

(a) 1 A (b) 2 A (c) 4 A (d) 8 A

2. The mass of water produced is :

(a) 7.2 g (b) 3.6 g (c) 1.8 g (d) 0.9 g

3. If current produced in fuel cell, is used for the deposition of Cu^{2+} in 1 L, 2 M $CuSO_4(aq)$ solution for 241.25 sec using Pt electrode, the pH of solution after electrolysis is :

(a) 1 (b) 2 (c) 3 (d) 4

For More Material Join: @JEEAdvanced_2024

Passage 5

A saturated solution in Agx (K_{sp} 3 10 ^{12}) and Agy (K_{sp} 10 ^{12}) has conductivity 0.4 10 ^{6} ^{1}cm ^{1}.

Given : Limiting molar conductivity of Ag 60 ^{1}cm^{2}mol ^{1}

Limiting molar conductivity of x 90 ^{1}cm^{2}mol ^{1}

1. The conductivity of y is (in ^{1} cm ^{1}) :

(a) 1.45 10 ^{7} (b) 1.45 10 ^{5}

(c) 1.45 10 ^{9} (d) None of these

2. The limiting molar conductivity of y is (in ^{1} cm^{2} mol ^{1}) :

(a) 290 (b) 2900 (c) 2.90 (d) None of these

One or More Answers is/are Correct

1. If the e.m.f of a galvanic cell is negative, it implies that :

(a) the cell reaction is spontaneous

(b) the cell reaction is non-spontaneous

(c) the cell reaction is exothermic

(d) the cell is working in reverse direction

2. Select correct statement(s) about electrolysis :

(a) Electric current is used to drive a non-spontaneous reaction

(b) G is positive for chemical process during electrolysis

(c) Cations and anions are moved toward the anode and cathode respectively

(d) Over voltage is generally associated with evolution of O_2 gas

3. If the half-cell reaction A e^- A^- has a large negative reduction potentials, it follows that :

(a) A is readily reduced (b) A is readily oxidised

(c) A^- is readily reduced (d) A^- is readily oxidised

4. Which of the following statement is correct?

If $E_{Cu^2|Cu}$ 0.34 V and $E_{Sn^{2+}|Sn}$ 0.136 V, $E_{H^+|H_2}$ 0.0 V

(a) Cu^2 ions can be reduced by $H_2(g)$ (b) Cu can be oxidized by H

(c) Sn^{2+} ions can be reduced by H_2 (d) Sn can be oxidized by Cu^2

5. The oxidation potential of hydrogen half-cell will be negative if :

(a) p(H_2) 1 atm and [H] 1M (b) p(H_2) 1 atm and $[H^+] = 2\,M$

(c) p(H_2) 0.2 atm and [H] 1M (d) p(H_2) 0.2 atm and $[H^+] = 0.2\,M$

6. Which of the following arrangement will produce oxygen at anode during electrolysis?

(a) Dilute H_2SO_4 with Pt electrodes

(b) Fused NaOH with inert electrodes

For More Material Join: @JEEAdvanced_2024

(c) Dilute H_2SO_4 with Cu electrodes
(d) Concentrated aq. NaCl with Pt electrodes

7. When an aqueous concentrated solution of lithium chloride is electrolysed using inert electrodes :
(a) Cl_2 is liberated at the anode
(b) Li is deposited at the cathode
(c) as the current flows, pH of the solution around the cathode remains constant
(d) as the current flows, pH of the solution around the cathode increases

8. Oxygen and hydrogen gas are produced at the anode and cathode during the electrolysis of fairly concentrated aqueous solution of :
(a) K_2SO_4 (b) $AgNO_3$ (c) H_2SO_4 (d) NaOH

9. During the purification of copper by electrolysis :
(a) the anode used is made of copper ore
(b) pure copper is deposited on the cathode
(c) the impurities such as Ag, Au present in solution as ions
(d) concentration of $CuSO_4$ solution remains constant during dissolution of Cu

10. When a lead storage battery is discharged :
(a) SO_2 is evolved
(b) lead sulphate is produced at both electrodes
(c) sulphuric acid is consumed
(d) water is formed

11. Which of the following is characteristic of the cathode in a voltaic cell?
(a) It may gain weight during reaction
(b) Electrons flow to it through the external circuit
(c) It is where oxidation occurs
(d) It receives electrons from ions in solution

12. In an electrochemical process, a salt bridge is used :
(a) to maintain electrical neutrality in each solution
(b) to complete the external circuit so that current can flow for long time
(c) to mix the solution of anodic and cathodic compartment
(d) to supply voltage

13. For a reaction in a galvanic cell the value of G at certain temperature is not necessarily equal to :
(a) nFE (b) $RT \ln K$ (c) $T \quad S \quad H$ (d) zero

14. Standard electrode potential of two half-reactions are given below :

$$Fe^{2+} \rightleftharpoons Fe \quad ; \quad E \quad 0.44\ V$$

$$Fe^{3+} \rightleftharpoons Fe^{2+} \quad ; \quad E \quad 0.77\ V$$

If Fe^{2+}, Fe^{3+} and Fe are kept together :
(a) the concentration of Fe^{3+} increases (b) the concentration of Fe^{3+} decreases
(c) the mass of Fe increases (d) the concentration of Fe^{2} decreases

For More Material Join: @JEEAdvanced_2024

15. Which of the following statements are correct regarding to galvanic cell?
(a) A reaction is spontaneous from left to right if E_{cell} 0
(b) A reaction occurs from right to left if E_{cell} 0
(c) If the system is at equilibrium no net reaction occurs
(d) E_{cell} is temperature-independent

16. Which of the following are concentration cells?
(a) $\underset{P_1}{Pt}|H_2(g)|HCl|\underset{P_2}{H_2}(g)|Pt$
(b) $Cd,\underset{a_1}{(Hg)}|\underset{(c)}{Cd^{2}}|\underset{a_2}{(Hg)},Cd$
(c) $Zn(s)|\underset{c_1}{Zn^{2}}||\underset{c_2}{Cu^{2+}}|Cu$
(d) $Ag|AgCl|\underset{c_1}{Cl^{-}}(aq)||\underset{c_2}{Br}(aq)|AgBr|Ag$

17. In electrolyte concentration cell :
(a) the electrode material and the solution in both half-cells are composed of the same substances
(b) only the concentrations of solutions of the same substances is different
(c) E_{cell} 0
(d) the Nernst equation reduces to E_{cell} $\frac{0.0591}{n}$ log Q at 25 C

18. The standard electrode potential of a metal-metal ion ($Ag|Ag^{+}$) and metal-sparingly soluble salt anion ($Ag|AgCl|Cl^{-}$) are related as :
(a) $E_{Ag^{+}|Ag}$ $E_{Cl^{-}|AgCl|Ag}$ $\frac{RT}{F}\ln K_{sp}$
(b) $E_{Cl^{-}|AgCl|Ag}$ $E_{Ag^{+}|Ag}$ $\frac{RT}{F}\ln K_{sp}$
(c) $E_{Cl^{-}|AgCl|Ag}$ $E_{Ag^{+}|Ag}$ $\frac{RT}{F}\ln\frac{[Cl^{-}]}{K_{sp}}$
(d) $E_{Cl^{-}|AgCl|Ag}$ $E_{Ag^{+}|Ag}$ $\frac{RT}{F}\ln\frac{K_{sp}}{[Cl^{-}]}$

19. Which of the following units is correctly matched?
(a) SI units of conductivity is $S\,m^{-1}$
(b) SI units of molar conductivity is $S\,cm^{2}mol^{-1}$
(c) SI unit of conductance is S^{-1}
(d) All of these

20. Which of the following statements is/are correct?
(a) The conductance of one cm^{3} (or 1 $unit^{3}$) of a solution is called specific conductance
(b) Specific conductance increases while molar conductivity decreases on progressive dilution
(c) The limiting equivalent conductivity of weak electrolyte cannot be determined exactly by extraplotation of the plot of $_{eq}$ against $\sqrt{c}$
(d) The conductance of metals is due to the movement of free electrons

21. Which is/are correct statement ?
(a) No corrosion takes place in vacuum
(b) Corrosion is protected by electroplating
(c) During rusting Fe_2O_3 xH_2O formed
(d) In presence of electrolyte, corrosion takes place with greater rate

22. A dilute solution of KCl was placed between two Pt electrodes 10 cm apart across which a potential difference of 10 volt was applied. Which is/are correct statement (Given : molar conductivity of K at infinite dilution is 96.5 $Scm^{2}\,mol^{-1}$.)

For More Material Join: @JEEAdvanced_2024

(a) Ionic mobility of K is 10^{-3} cm^2 sec^{-1} volt^{-1}.

(b) The speed of K is 10^{-3} cm sec^{-1}.

(c) Distance travelled by K in 5 10^3 sec is 5 cm.

(d) The potential gradient is 1.0 volt cm^{-1}.

23. Given : $Pt(s) \mid \underset{1\text{ bar}}{H_2(g)} \mid 0.1M\ NH_4OH(aq) \,||\, 0.1M\ CH_3COOH(aq) \mid \underset{1\text{ bar}}{H_2(g)} \mid Pt(s)$

$pK_b(NH_4OH)$ 5; $pK_a(CH_3COOH)$ 5; $\frac{2.303RT}{F}$ 0.06

Volume of $0.1M\ NH_4OH$ in anode half cell 100 mL,

Volume of $0.1M\ CH_3COOH$ in cathode half cell 100 mL

Which is/are correct statement ?

(a) The emf of given cell is 0.48 V.

(b) The emf of given cell is 0.36 V when 50 mL, 0.1 *M* NaOH added to cathode compartment

(c) The emf of given cell is 0.36 V when 50 mL 0.1 *M* HCl added to anode compartment

(d) The emf of given cell is 0.192 V when 100 mL 0.1 *M* NaOH added to anode compartment

24. Given : $Pt(s) \mid \underset{P_1\text{ atm}}{Cl_2(g)} \mid Cl\ (C_1) \,||\, Cl\ (C_2) \mid \underset{P_2\text{ atm}}{Cl_2(g)} \mid Pt(s)$

Identify in which of following condition working of cell takes place :

(a) C_1 C_2 and P_1 P_2 (b) P_2 P_1 and C_1 C_2

(c) C_1 C_2 and P_1 P_2 (d) P_2 P_1 and C_1 C_2

25. 1000 mL $1M\ CuSO_4(aq)$ is electrolysed by 9.65 A current for 100 sec using Pt-electrode. Which is/are correct statement ?

(a) Blue colour intensity decreases during electrolysis.

(b) Blue colour intensity remains constant if Cu-electrode used.

(c) pH of solution is 8 after electrolysis.

(d) 28 mL of CH_4 at 1 atm and 273 K required for its combustion by O_2, liberated during electrolysis.

Match the Column

Column–I and **Column–II** contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II

1.

Column–I	Column–II
(A) Dilute solution of HCl	(P) O_2 evolved at anode
(B) Dilute solution of NaCl	(Q) H_2 evolved at cathode
(C) Concentrated solution of NaCl	(R) Cl_2 evolved at anode
(D) Fairly concentrated solution of $AgNO_3$	(S) Ag deposition at cathode

For More Material Join: @JEEAdvanced_2024

2.

Column–I	Column–II
(A) If SOP of substance lies between 1.23 to 0.81 V	(P) Oxidation of substance is not possible
(B) If SOP of substance lies between 0.81 V to –0.40 V	(Q) Oxidation possible only in acidic medium
(C) If SOP is less than –1.23 V	(R) Oxidation possible in any medium
(D) If SOP is greater than 0.40 V	(S) Oxidation easily takes place

3.

Column–I	Column–II (SRP)
(A) $F_2 \quad 2e \rightleftharpoons 2F$	(P) 0.54
(B) $Cl_2 \quad 2e \rightleftharpoons 2Cl$	(Q) 1.09
(C) $Br_2 \quad 2e \rightleftharpoons Br$	(R) 1.36
(D) $I_2 \quad 2e \rightleftharpoons 2BI$	(S) 2.87

4.

Column–I	Column–II
(A) $Pt\|Fe^{3+}, Fe^{2+}$	(P) Metal-metal ion half-cell
(B) $Pt\|H_2\|H^+$	(Q) Gas-gas ion half-cell
(C) $Pt\|Hg\|Hg_2^{2+}$	(R) Oxidation-reduction half-cell
(D) $Pb\|PbSO_4\|SO_4^{2-}$	(S) Metal-sparingly soluble salt half-cell

5.

Column–I (Property)	Column–II (Unit)
(A) Conductance	(P) Sm^{1}
(B) Conductivity	(Q) $S\ ^{1}m$
(C) Molar conductivity	(R) $Sm^{2}mol^{-1}$
(D) Resistivity	(S) S

6.

Column–I (Ion)	Column–II (Molar Conductivity)
(A) H^+	(P) 350
(B) Na^+	(Q) 50
(C) Li	(R) 39
(D) Cs	(S) 77

7.

Column-I	Column-II
(A) Galvanic cell	(P) Used in space craft
(B) Electrolytic cell	(Q) No transformation of electrical energy into chemical energy

For More Material Join: @JEEAdvanced_2024

(C) Dead battery	(R) Cell reaction is spontaneous
(D) Fuel cell	(S) Cell reaction is non-spontaneous

Assertion-Reason Type Questions



Each question contains STATEMENT-1 (Assertion) and STATEMENT-2 (Reason).
Examine the statements carefully and mark the correct answer according to the instructions given below :

(A) If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1

(B) If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1

(C) If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

(D) If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

1. STATEMENT-1 : E_{cell} is negative for electrolytic cell.
STATEMENT-2 : G is ve for electrolytic cell.

2. STATEMENT-1 : When 2 faraday of charge is passed through 0.1 M $H_2SO_4(aq)$, 11.2 litre O_2 evolved at STP.
STATEMENT-2 : Molecular mass of oxygen is 32.

3. STATEMENT-1 : Copper is dissolved at anode and deposited at cathode when Cu electrodes are used and electrolyte is 1 M $CuSO_4(aq)$ solution.
STATEMENT-2 : SOP of Cu is less than SOP of water and SRP of Cu is greater than SRP of water.

4. STATEMENT-1 : 1 coulomb charge deposits 1 g-equivalent of a substance.
STATEMENT-2 : 1 faraday is charge on 1 mole of electrons.

5. STATEMENT-1 : If SRP of substance is 0.3 V, its reduction is possible at cathode.
STATEMENT-2 : Reduction potential of water lies between 0 to 0.8274 V at 25 C.

6. STATEMENT-1 : If SRP of substance is 0.5 V then reduction of substance is possible only in basic medium.
STATEMENT-2 : SRP of water is 0.8274 V and at reduction potential is zero at pH = 7

7. STATEMENT-1 : The voltage of mercury cell remains constant for longer period of time.
STATEMENT-2 : It is because net cell reaction does not involve ions.

8. STATEMENT-1 : Lead storage battery is a galvanic cell without salt bridge.
STATEMENT-2 : A secondary cell is rechargeable cell.

9. STATEMENT-1 : The SRP of three metallic ions A, B, C are 0.3, – 0.5, 0.8 volt respectively, so oxidizing power of ions is C A B.
STATEMENT-2 : Higher the SRP, higher the oxidizing power.

10. STATEMENT-1 : If SOP of substance is less than 1.23 V and over voltage 0 V, then its oxidation in its aqueous solution is not possible at 298 K.
STATEMENT-2 : Standard reduction potential (SRP) of water is 1.23 V.

11. STATEMENT-1 : We cannot add the electrode potential in order to get electrode potential of third electrode if no. of moles of electrons exchanged are not same.

For More Material Join: @JEEAdvanced_2024

STATEMENT-2 : Electrode potential is an extensive property.

12. STATEMENT-1 : E_{cell} 0 for a chloride ion concentration cell.

STATEMENT-2 : For this concentration cell where E_{cell} $\frac{RT}{nF}\ln\frac{[Cl^-]_{LHS}}{[Cl^-]_{RHS}}$

13. STATEMENT-1 : If $\left(\frac{dE_{cell}}{dT}\right)_p$ 0 for a cell reaction then S is positive.

STATEMENT-2 : S $nFT\left(\frac{dE}{dT}\right)_p$

14. STATEMENT-1 : Molar conductivity increases with decrease in concentration for weak electrolytes.

STATEMENT-2 : No. of ions increases and no. of ions per unit volume decreases due to dilution.

15. STATEMENT-1 : Conductivity decreases with the decreases in concentration both the weak and strong electrolytes.

STATEMENT-2 : No. of ions per unit volume linearly decreases in both electrolytes.

Subjective Problems

1. How many faradays are required for reduction of 1 mol $C_6H_5NO_2$ into $C_6H_5NH_2$?

2. What is the equivalent mass of O_2 in the following reaction, H_2O $\frac{1}{2}O_2$ $2e$ $2OH$?

3. The amount of electricity which releases 2.0 g of gold from a gold salt is same as that which dissolves 0.967 g of copper anode during the electrolysis of copper sulphate solution. What is the oxidation number of gold in the gold ion ? (At. mass of Cu 63.5; Au 197)

4. When a molten salt was electrolysed for 5 min with 9.65 A current, 0.18 g of the metal was deposited. Calculate the Eq. mass of metal.

5. During the electrolysis of a concentrated brine solution, calculate the moles of chlorine gas produced by the passage of 4F electricity.

6. Calculate the cell potential (in V) if G 96.5 kJ/mol and n 1.

7. If K_c for the reaction

$Cu^{2}(aq)$ $Sn^{2}(aq)$ $Sn^{4}(aq)$ $Cu(s)$

at 25 C is represented as 2.6 10^{y} then find the value of y.

(Given : $E_{Cu^{2}|Cu}$ 0.34 V ; $E_{Sn^{4}|Sn^{2}}$ 0.15 V)

8. If G for the half cell $MnO_4|MnO_2$ in an acid solution is xF then find the value of x.

(Given : $E_{MnO_4|Mn^{2}}$ 1.5 V ; $E_{MnO_2|Mn^{2}}$ 1.25 V)

9. If the equilibrium constant for the reaction $Cd^{2}(aq)$ $4NH_3(aq) \rightleftharpoons Cd(NH_3)_4^{2}(aq)$ is 10^{x} then find the value of x.

(Given : $E_{Cd^{2}|Cd}$ 0.4 V ; $E_{Cd(NH_3)_4^{2}|Cd}$ 0.61 V)

For More Material Join: @JEEAdvanced_2024

10. At what pH oxidation potential of water is 0.81 V ?

11. The resistance of a conductivity cell containing 0.001 M KCl solution at 298 K is 1500 . What is the cell constant (in mm ^{1}) if the conductivity of 0.001 M KCl solution is 2 10^{3} S mm ^{1}.

12. Molar conductivities at infinite dilution of KCl, HCl and CH_3COOK are 0.013, 0.038 and 0.009 $Sm^2\,mol^{1}$ respectively at 291K. If conductivity of 0.001 M CH_3COOH is 2.72 10^{3} Sm^{1} then find % degree of dissociation of CH_3COOH.

13. A solution containing 1 M $XSO_4(aq)$ and 1 M $YSO_4(aq)$ is electrolysed. If conc. of X^2 is 10^{z} M when deposition of Y^2 and X^2 starts simultaneously, calculate the value of Z.

Given : $\frac{2.303RT}{F}$ 0.06

$E^{\circ}_{X^2|X}$ 0.12 V ; $E^{\circ}_{Y^2|Y}$ 0.24 V

14. Molar conductivity of aqueous solution of HA is 200 $Scm^2\,mol^{1}$, pH of this solution is 4. Calculate the value of pK_a(HA) at 25°C.

Given : $_M$(NaA) 100 Scm^2mol^{1}; $_M$(HCl) 425 Scm^2mol^{1};

$_M$(NaCl) 125 $Scm^2\,mol^{1}$

15. The standard reduction potential of a silver chloride electrode (metal-sparingly soluble salt electrode) is 0.209 V and for silver electrode is 0.80 V. If the moles of AgCl that can dissolve in 10 L of a 0.01 M NaCl solution is represented as 10^{Z} then find the value of Z.

For More Material Join: @JEEAdvanced_2024

ANSWERS

Level 1

1.	(a)	**2.**	(c)	**3.**	(d)	**4.**	(d)	**5.**	(a)	**6.**	(c)	**7.**	(d)	**8.**	(d)	**9.**	(a)	**10.**	(c)
11.	(a)	**12.**	(b)	**13.**	(c)	**14.**	(b)	**15.**	(d)	**16.**	(c)	**17.**	(a)	**18.**	(a)	**19.**	(d)	**20.**	(c)
21.	(d)	**22.**	(c)	**23.**	(b)	**24.**	(c)	**25.**	(c)	**26.**	(d)	**27.**	(d)	**28.**	(b)	**29.**	(c)	**30.**	(d)
31.	(d)	**32.**	(d)	**33.**	(a)	**34.**	(a)	**35.**	(a)	**36.**	(a)	**37.**	(a)	**38.**	(d)	**39.**	(d)	**40.**	(a)
41.	(a)	**42.**	(c)	**43.**	(d)	**44.**	(b)	**45.**	(b)	**46.**	(c)	**47.**	(c)	**48.**	(a)	**49.**	(b)	**50.**	(b)
51.	(a)	**52.**	(c)	**53.**	(d)	**54.**	(a)	**55.**	(c)	**56.**	(b)	**57.**	(c)	**58.**	(b)	**59.**	(c)	**60.**	(a)
61.	(c)	**62.**	(c)	**63.**	(d)	**64.**	(d)	**65.**	(b)	**66.**	(a)	**67.**	(a)	**68.**	(d)	**69.**	(d)	**70.**	(a)
71.	(a)	**72.**	(d)	**73.**	(c)	**74.**	(b)	**75.**	(d)	**76.**	(d)	**77.**	(d)	**78.**	(d)	**79.**	(a)	**80.**	(d)
81.	(a)	**82.**	(a)	**83.**	(a)	**84.**	(a)	**85.**	(b)	**86.**	(b)	**87.**	(a)	**88.**	(c)	**89.**	(d)	**90.**	(d)
91.	(d)	**92.**	(c)	**93.**	(b)	**94.**	(b)	**95.**	(a)	**96.**	(a)	**97.**	(c)	**98.**	(d)	**99.**	(d)	**100.**	(b)
101.	(c)	**102.**	(a)	**103.**	(b)	**104.**	(b)	**105.**	(d)	**106.**	(d)	**107.**	(c)	**108.**	(c)	**109.**	(c)	**110.**	(d)
111.	(c)	**112.**	(c)	**113.**	(c)	**114.**	(d)	**115.**	(b)	**116.**	(b)	**117.**	(d)	**118.**	(b)	**119.**	(b)	**120.**	(a)
121.	(a)	**122.**	(b)	**123.**	(b)	**124.**	(c)	**125.**	(b)	**126.**	(c)	**127.**	(b)	**128.**	(b)	**129.**	(a)	**130.**	(b)
131.	(c)	**132.**	(c)	**133.**	(d)	**134.**	(d)	**135.**	(c)	**136.**	(a)	**137.**	(b)	**138.**	(a)	**139.**	(c)	**140.**	(b)
141.	(b)	**142.**	(b)	**143.**	(c)	**144.**	(d)	**145.**	(d)	**146.**	(b)	**147.**	(b)	**148.**	(a)	**149.**	(c)	**150.**	(d)

Level 2

1.	(b)	**2.**	(c)	**3.**	(d)	**4.**	(d)	**5.**	(a)	**6.**	(b)	**7.**	(d)	**8.**	(c)	**9.**	(a)	**10.**	(a)
11.	(a)	**12.**	(b)	**13.**	(c)	**14.**	(a)	**15.**	(a)	**16.**	(b)	**17.**	(c)	**18.**	(a)	**19.**	(c)	**20.**	(c)
21.	(b)	**22.**	(b)	**23.**	(c)	**24.**	(a)	**25.**	(b)	**26.**	(c)	**27.**	(a)	**28.**	(c)				

For More Material Join: @JEEAdvanced_2024

Level 3

Passage–1:	**1.**	(c)	**2.**	(a)	**3.**	(d)		
Passage–2:	**1.**	(b)	**2.**	(b)	**3.**	(a)		
Passage–3:	**1.**	(c)	**2.**	(b)				
Passage–4:	**1.**	(c)	**2.**	(b)	**3.**	(b)		
Passage–5:	**1.**	(a)	**2.**	(a)				

One or More Answers is/are correct

1. (b,d) **2.** (a,b,d) **3.** (d) **4.** (a,d) **5.** (b,c) **6.** (a,b) **7.** (a,d) **8.** (a, c, d)
9. (a,b,d) **10.** (b,c,d) **11.** (a,b) **12.** (a) **13.** (b,d) **14.** (b) **15.** (a,b,c) **16.** (a,b,d)
17. (a,b,c,d) **18.** (b) **19.** (a) **20.** (a,c,d) **21.** (a,b,c,d) **22.** (a,b,c,d) **23.** (a,b,c) **24.** (a,b)
25. (a,b,d)

Match the Column

1.	A	P, Q;	B	P, Q;	C	Q, R;	D	P, S
2.	A	Q;	B	R;	C	P;	D	S, R
3.	A	S;	B	R;	C	Q;	D	P
4.	A	R;	B	Q;	C	P;	D	S
5.	A	S;	B	P;	C	R;	D	Q
6.	A	P;	B	Q;	C	R;	D	S
7.	A	Q, R;	B	S;	C	Q;	D	P, Q, R

Assertion-Reason Type Questions

1. (A) **2.** (B) **3.** (C) **4.** (D) **5.** (A) **6.** (C) **7.** (A) **8.** (B) **9.** (A) **10.** (C)
11. (C) **12.** (B) **13.** (C) **14.** (A) **15.** (C)

Subjective Problems

1. 6	**2.** 8	**3.** 3	**4.** 6	**5.** 2	**6.** 1	**7.** 6	**8.** 5	**9.** 7	**10.** 7
11. 3	**12.** 8	**13.** 4	**14.** 4	**15.** 7					

For More Material Join: @JEEAdvanced_2024

Hints and Solutions

Level 1

1. (a) For spontaneous

$$_rG \quad nFE \quad 0$$

$$E \quad 0$$

2. (c) Cathode and anode are ( ) ve terminals in electrolytic and galvanic cell.

3. (d) Electrostatic attraction is broken down by polar solvent (water).

4. (d) $A \quad e \quad A$
$1 \text{ mole} \qquad 1 \text{ mole}$

Charge of 1 mole electron is required for deposition of 1 equivalent of substance.

5. (a) According to stoichiometry liberated amount of ion depends upon $\because w \quad \frac{EIt}{F}$

(i) Electrolysis time (ii) Current strength

(ii) Electrochemical equivalent

6. (c) 1 F = 96500 C,

$$1 \text{ C} = \frac{6.023 \quad 10^{23}}{96500} \quad 6.24 \quad 10^{18}$$

7. (d) Charge $0.01 \quad 10^{3} \quad 1000 \quad 60 \quad 60 \quad 36$

8. (d) $3.21 \quad 10^{6} \quad 500 \quad t$

$$t \quad \frac{3.21 \quad 10^{6}}{500 \quad 60} \quad 107 \text{ min.}$$

9. (a) $W \quad ZIt$

$$0.746 \quad \frac{108}{1 \quad F} \quad 548 \quad I \qquad I \quad 1.22$$

10. (c) $\frac{0.55}{M} \quad 3 \quad \frac{0.55 \quad 100 \quad 60}{96500}$

$$M \quad 48.25 \text{ g/mol}$$

11. (a) $w \quad \frac{9}{2} \quad \frac{5 \quad 1 \quad 60 \quad 60}{96{,}500} \quad 0.840 \text{ g}$

12. (b) $5.2 \quad \frac{52}{3} \quad \frac{9.65}{96{,}500} \quad t$ (sec.),

t(sec) 3000

t(min) 50

13. (c) $\frac{W}{E} \quad \frac{I \quad t}{96500}; \frac{0.195}{195} \quad 4 \quad \frac{5 \quad 60 \quad 60 \quad I}{96500},$

$I \quad 21.44$

14. (b) $0.25 \quad \frac{93}{5} \quad \frac{Q}{F}$

$$Q \quad 1.3 \quad 10^{2}$$

15. (d) When 1 mole of e^{-} is passed, mass of substance deposited

$$\frac{1 \quad 6.023 \quad 10^{23}}{1.81 \quad 10^{23}} \quad 3.34 \text{ g} \quad \text{eq. wt}$$

atomic mass of metal = $3.34 \quad 3$

10.03 g

16. (c) 1 mole Mn_3O_4 lose $6 \quad \frac{8}{3} \quad 3 \quad 10$ mole e^{-}; so total charge required $2 \quad 10 \quad 20$ F

17. (a) $2H \quad 2e \quad H_2$

18. (a) $\because \quad E^{\circ}_{Mg^{2} \,|Mg} \quad E^{\circ}_{Cr^{3} \,|Cr} \quad E^{\circ}_{Cu^{2} \,|Cu} \quad E^{\circ}_{Ag \,|Ag}$

19. (d) $\because \quad E^{\circ}_{K \,|K} \quad E^{\circ}_{H \,|H_2}$

$E^{\circ}_{H_2O|O_2} \quad E^{\circ}_{SO_4^{2} \,|S_2O_8^{2}}$

(High concentrate solution)

20. (c) $\because \quad E^{\circ}_{Na \,|Na} \quad E^{\circ}_{H \,/H_2}$

$E^{\circ}_{Cu^{2} \,|Cu} \quad E^{\circ}_{H \,/H_2}$

21. (d) $E^{\circ}_{SO_4^{2} \,|S_2O_8^{2}} \quad E^{\circ}_{H_2O|O_2}$

22. (c) $E^{\circ}_{Na \,|Na} \quad E^{\circ}_{H \,|H_2}; E^{\circ}_{SO_4^{2} \,|S_2O_8^{2}} \quad E^{\circ}_{H_2O|O_2}$

23. (b) In electrolysis of concentrated Cl , $Cl_2(g)$ is evolved in place of $O_2(g)$.

25. (c) $E^{\circ}_{Cu^{2} \,|Cu} \quad E^{\circ}_{H_2O|H_2}; E^{\circ}_{SO_4^{2} \,|S_2O_8^{2}} \quad E^{\circ}_{H_2O|O_2}$

26. (d) Anode : $2Cl\ (aq) \qquad Cl_2(g) \quad 2e$

Cathode: $2H_2O \quad 2e \qquad H_2(g) \quad 2OH\ (aq)$

Net reaction

$2Na\ (aq) \quad 2Cl\ (aq) \quad 2H_2O\ (l)$

$Cl_2(g) \quad H_2(g) \quad 2OH\ (aq) \quad 2Na\ (aq)$

27. (d) $\because \quad E^{\circ}_{K \,|K} \quad E^{\circ}_{H_2O|H_2}; E^{\circ}_{Na \,|Na} \quad E^{\circ}_{H_2O|H_2};$

$E^{\circ}_{Cu^{2} \,|Cu} \quad E^{\circ}_{H_2O|H_2}$

28. (b) Let t sec be used; No. of faradays $\frac{4 \quad t}{96500}$

No. of moles of $H_2O = 4$, no. of equivalents of $H_2O = 4 \quad 2$

$$\frac{4 \quad t}{96500} \quad 4 \quad 2$$

$$t \quad 1.93 \quad 10^{5}$$

For More Material Join: @JEEAdvanced_2024

29. (c) only Au^{3+}, Ag^{+} and Cu^{2+} will deposit at cathode.
Li will not deposit at cathode because SRP of water is 0.8274 V so after Cu^{2+}; H_2 will evolve at cathode.

30. (d) $\frac{W}{M}$ 2 $\frac{4.6 \quad 30 \quad 60}{96500}$
moles of Sn^{2+} reduced 0.043
Initial moles of $Sn^{2+} = 0.5 \quad 0.6 \quad 0.30$;
remaining moles of $Sn^{2+} = 0.30 - 0.043$
$[Sn^{2+}] = \frac{0.257}{0.5} = 0.514\ M$

31. (d) No. of moles of
Ag $\frac{15 \quad 60 \quad 1.25 \quad 10^{3}}{96,500} \quad \frac{1}{1}$
$0.0116 \quad 10^{3}$
[Ag] $\frac{1.16 \quad 10^{5}}{\frac{100}{1000}}$ $1.16 \quad 10^{-4}$

32. (d) Initial moles of $Cr^{3+} = 0.25 \quad 0.2 \quad 0.05$;
final moles of $Cr^{3+} = 0.25 \quad 0.1 \quad 0.025$
moles of Cr^{3+} reduced,
$0.05 \quad 0.025 \quad 0.025$;
eq. of Cr^{3+} reduced, $0.025 \quad 3 = \frac{t \quad 96.5}{96500}$;
t 75 sec

33. (a) 3.05 $\frac{114.8}{n}$ $\frac{3.2 \quad 40 \quad 60}{96,500}$
n 3

34. (a) $\frac{W}{M}$ n $\frac{I \quad t}{96500}$; $\frac{0.838}{184}$ n
$\frac{40 \quad 60 \quad 1.1}{96500}$
n 6

35. (a) No. of equivalent of H_2 No. of equivalent of Cl_2
$\frac{5 \quad 1}{22.4}$ 2 $\frac{V_{Cl_2}}{22.4}$ 2
$V_{Cl_2(g)}$ 5L

36. (a) No. of equivalent of Ag $\frac{0.54}{108}$ 1 No. of equivalent of Cr $\frac{W_{Cr}}{52}$ 3
W_{Cr} $\frac{0.54}{108} \quad \frac{52}{3}$ 0.0866

37. (a) No. of equivalent of Cu $\frac{5.6 \quad 1}{22.4}$ 4
$\frac{W_{Cu}}{63.5}$ 2, W_{Cu} 31.75

38. (d) no. of eq no. of F $\frac{1000}{122.5}$ 8 65.3

39. (d) 54 10^{3} $\frac{27}{3} \quad \frac{x}{96,500}$
x $57.9 \quad 10^{7}$

40. (a) $\frac{W}{7}$ $\frac{1930 \quad 0.75}{96500}$; W 0.105 gm

41. (a) equivalents of Cl_2 produced
$\frac{1000 \quad 9.65 \quad 3600}{96500}$ 360
moles of Cl_2 180
so V $\frac{nRT}{P}$ $\frac{180 \quad 0.0821 \quad 300}{1.8}$
2463 L

42. (c) No. of equivalent of H_2 produced Eq. of O_2
$\frac{I \quad t}{96500}$ $\frac{30 \quad 193 \quad 60}{96500}$
= 3.6
vol. of O_2 and H_2 produced are
3.6 5.6 + 3.6 11.2
= 60.48 litre

43. (d) 1 watt = 1 J/sec; 1 kWh = $\frac{V \quad I \quad t}{1000}$ (in hr)
Total kWh 10^{-3} 110 10 10 11
Total cost 11 2 22 Rs.

44. (b) $E^{\circ}_{SO_4^{2-}|S_2O_8^{2-}}$ $E^{\circ}_{H_2O|O_2}$

45. (b) 10 g Cd should present with 4.5 g Hg.
mass of Cd required with 4.5 g of Hg at cathode
$\frac{10}{90}$ 4.5 0.5; ($Cd^{2} + 2e^{-}$ Cd)
$\frac{0.5}{112}$ 2 $\frac{4 \quad t}{96500}$
t 215.40 sec

46. (c) Use of electrolysis is in (i) Electrorefining (ii) Electroplating

47. (c) $E^{\circ}_{Ag|Ag}$ $E^{\circ}_{H|H_2}$, $E_{Cu^{2}/Cu}$ $E_{O_2,H/H_2O}$

48. (a) Chemical energy into electrical energy.

49. (b) G_1° G_2°
2 $FE^{\circ}_{Cl|Cl_2}$ 2 $FE^{\circ}_{Cl_2|Cl}$
1.36 V

For More Material Join: @JEEAdvanced_2024

50. (b) $G_3^\circ \quad G_1^\circ \quad G_2^\circ$

$n_3FE_3^\circ \quad n_1FE_1^\circ \quad n_2FE_2^\circ$

$E_3^\circ \quad \dfrac{n_1E_1^\circ \quad n_2E_2^\circ}{n_3}$

51. (a) Maintain electrical neutrality of both half cells.

52. (c) Velocities of ions are nearly the same.

53. (d) A salt bridge contains strong electrolyte with agar-agar.

54. (a) $\log K_C \quad \dfrac{nFE}{2.303\,RT}$

$y \quad mx$

55. (c) $G_1^\circ \quad 1 \quad F(x) \quad RT \ln K_1$

$G_2^\circ \quad 1 \quad F(y) \quad RT \ln K_2$

$G_2^\circ \quad 2 \quad G_1^\circ \quad 2RT \ln K_1$

$K_2 \quad K_1^2, \; x \quad y$

56. (b) $E \quad E \quad \dfrac{0.0591}{2} \log \dfrac{[Zn^{2}]}{[Ag\ ]^2}$

$y \quad C \quad mx$

57. (c) When $Q \quad K_C$, $E \quad 0$

58. (b) $Pt(s)\,|H_2(g)\,|H\ (aq)\,||Ag\ (aq)\,|Ag(s)$,

Anode : $H_2|H$, Cathode : $Ag\ |Ag$

59. (c) $Cu(s)\,|Cu^{2}\ (aq)\,||Cl\ (aq)\,|Hg_2Cl_2(s)\,|Hg(l)\,|Pt(s)$,

Anode : $Cu|Cu^{2}$

Cathode : $Cl\ |Hg_2Cl_2|Hg$

60. (a) $Pt(s)\,|Fe^{2}\ (aq), Fe^{3}\ (aq)\,||Cr_2O_7^{2}\ (aq), Cr^{3}\ (aq)\,|Pt(s)$

Anode : $Fe^{2}\ |Fe^{3}$

Cathode : $Cr_2O_7^{2}\ |Cr^{3}$

61. (c) Anode : $2Ag(s) \rightleftharpoons 2Ag\ (aq) \quad 2e$

Cathode : $Cu^{2}\ (aq) \quad 2e \rightleftharpoons Cu(s)$

Net reaction : $2Ag(s) \quad Cu^{2}\ (aq) \rightleftharpoons 2Ag\ (aq) \quad Cu(s)$

62. (c) Anode : $Cl_2(g) \rightleftharpoons 2Cl\ (aq) \quad 2e$

Cathode : $2Ag\ (aq) \quad 2e \rightleftharpoons 2Ag(s)$

Net reaction : $Cl_2(g) \quad 2Ag\ (aq) \rightleftharpoons 2Cl\ (aq) \quad 2Ag(s)$

63. (d) $E_{H_2|H}^\circ \quad E_{H_2|H}^\circ \quad 0V$

64. (d) $E \quad E \quad \dfrac{0.0591}{2} \log \dfrac{[Zn^{2}]}{[Cu^{2}]}$

As Q , E

$Q_1 \quad 1, \; Q_2 \quad 0.1, \; Q_3 \quad 10, \; Q_4 \quad 1$

$E_2 \quad E_1 \quad E_4 \quad E_3$

65. (b) Ag becomes oxidized and Br becomes reduced.

66. (a) $N_2(g) \quad 5H\ (aq) \quad 4e \quad N_2H_5(aq)$,

$E \quad 0.23\,V \quad \ldots(1)$

$BiO\ (aq) \quad 2H\ (aq) \quad 3e$

$Bi(S) \quad H_2O\,(l)$

$E \quad 0.32\,V \quad \ldots(2)$

[Reaction (1)] 3 [Reaction (2)] 4, we get

$4BiO\ (aq) \quad 3N_2H_5(aq)$

$4Bi\,(s) \quad 3N_2(g) \quad 4H_2O(l) \quad 7H\ (aq)$

$G \quad 3(\ 4FE\ ) \quad 4(\ 3FE\ )$

$12FE \quad 12F(\ 0.23) \quad 12F(0.32)$

$E \quad 0.23 \quad 0.32 \quad 0.55$

67. (a) $Cr^{2}\ (aq) \quad Cr^{3}\ (aq) \quad e$, $E \quad 0.41\,V$

$\ldots(1)$

$HClO(aq) \quad H\ (aq) \quad 2Cr^{2}\ (aq)$

$2Cr^{3}\ (aq) \quad Cl\ (aq) \quad H_2O(l), E \quad 1.8 \quad \ldots(2)$

[Reaction (1)] 2 Reaction (2)] we get.

$HClO(aq) \quad H\ (aq) \quad 2e \quad Cl\ (aq) \quad H_2O(l)$

$G \quad 2FE \quad 2(\ 1 \quad FE\ )$

$2FE \quad 2F(1.8) \quad 2F(0.41)$

$E \quad 1.8 \quad 0.41 \quad 1.39$

68. (d) $0.34 \quad 1 \quad E_{SbO_2|sb}^\circ$

$E_{SbO_2|Sb}^\circ \quad 0.34 \quad 1 \quad 0.66$

69. (d) For spontaneous $E \quad 0$,

minimum possible spontaneous $E \quad 0.28$

70. (a) For best reducing agent, E_{Red}° is minimum.

71. (a) For best oxidizing agent, E_{Red}° is maximum.

72. (d) Maximum voltage $E \quad 0.49 \quad 0.86 \quad 1.35$

73. (c) $E_{Al^3\ |Al}^\circ \quad E_{Cu^2\ |Cu}^\circ ; E_{Cu|Cu^2}^\circ \quad E_{Al^3\ |Al}^\circ$ ve (Non-spontaneous)

74. (b) $E_{Zn^2\ |Cu}^\circ \quad E_{Cu^2\ |Cu}^\circ$;

$E_{Cu^2\ |Cu}^\circ \quad E_{Zn|Zn^2}^\circ$ ve

75. (d) D does not react with HCl, so not easily oxidized at compared to A, B and C

C reduces B as well as D, but unable to reduce A

so A is oxidized easily as compared to other metals

76. (d) $E_{Cu^2\ |Cu}^\circ \quad E_{Ag\ |Ag}^\circ ; E_{Cu^2\ |Cu}^\circ \quad E_{Ag\ |Ag}^\circ$ ve (Non-spontaneous)

77. (d) From (I) information, C is strong reducing agent.

For More Material Join: @JEEAdvanced_2024

From (II) information, A is strong reducing agent than D, B and C reducing agent than A.

$D \quad A \quad B \quad C$

78. (d) $4Fe(s) \quad 3O_2(g) \quad 4Fe^3 \quad 6O^2\ (aq)$

Fe is reducing agent, O_2 is oxidizing agent.

79. (a) $E^\circ_{Na\ |Na} \quad E^\circ_{Zn^2\ |Zn} \quad E^\circ_{Fe^2\ |Fe} \quad E^\circ_{Ag\ |Ag}$

$E^\circ_{Fe|Fe^2} \quad E^\circ_{Ag\ |Ag}$ ve (Spontaneous)

80. (d) $E^\circ_{Zn|Zn^2} \quad E^\circ_{Mg^2\ |Mg} \quad 1.608$

(Non-spontaneous)

81. (a) $E^\circ_{F_2|F} \quad E^\circ_{Cl_2|Cl} \quad E^\circ_{Br_2|Br} \quad E^\circ_{I_2|I}$

82. (a) $K \quad 10^{\frac{2(0.277)}{0.0591}} \quad 2.3 \quad 10^9$

83. (a) $E \quad \frac{0.0591}{2}\log K_{eq};\ \log K_{eq} \quad \frac{2 \quad 0.22}{0.0591}$

$7.44 \quad \text{or} \quad K_{eq} \sim 2.8 \quad 10^7$

84. (a) $E \quad E \quad \frac{0.0591}{2}\log\frac{1}{[Cl\ ]^2};$ as $[Cl\ ], E$

85. (b) $K \quad 10^{\frac{6 \quad FE}{2.303RT}}$

$\log 10^{30} \quad \frac{6 \quad E}{0.0591} \quad E \quad 0.2955\ V$

86. (b) $E_{cell} \quad \frac{0.0591}{2}\log K_{eq}$

$$0.003 = \frac{0.0591}{2}\log\frac{[D^+]^2}{[H^+]^2}$$

$$= \frac{[D^+]}{[H^+]} = 1.123$$

87. (a) $G \quad 12 \quad F \quad E \quad 636.9\ kJ \quad 637\ kJ$

88. (c) $E_{cell} \quad E_{RP(RHS)} \quad E_{RP(LHS)}$

$1.61 \quad 1.51 \quad 0.10\ V$

$G \quad nFE \quad 5 \quad 96500 \quad 0.10\ J$

$G \quad 48.25\ kJ$

89. (d) $E \quad E \quad \frac{0.0591}{1}\log\frac{[Fe^3\ ]}{[Ag\ ][Fe^2\ ]}$

as $Q \quad \{[Ag\ ], [Fe^3\ ]\}, E$

90. (d) $E \quad E \quad \frac{0.0591}{2}\log\frac{[HCl]^2}{P_{H_2}}$

as $Q \quad \{(P_{H_2}, [H\ ]\}E$

91. (d) $E_{Cr_2O_7^{2-}|Cr^{3+}} \quad E_{Cr_2O_7^{2-}|Cr^{3+}} \quad \frac{0.0591}{6}$

$\log\frac{[Cr^{3+}]^2}{[Cr_2O_7^{2-}]} \quad \frac{1}{[H^+]^{14}}$

$E_{Cr_2O_7^{2-}|Cr^{3+}} \quad 1.33 \quad \frac{0.0591}{6}\log\frac{1}{(0.01)^{14}}$

$1.0542\ V$

92. (c) $E \quad 0.57 \quad \frac{0.0591}{4}\log\frac{[OH\ ]^6}{[TeO_3^2\ ]}$

$0.57 \quad \frac{0.06}{4}\log(10^{\ 2})^6 \quad 0.39$

93. (b) $Co^2\ (aq) \rightleftharpoons Co^2\ (aq), \quad Q \quad \frac{C_2}{C_1}$

$C_1 \qquad C_2$

If $\frac{C_2}{C_1} \quad 1, \quad G \quad 0$

94. (b) $\underset{P_1}{H_2(g)} \rightleftharpoons \underset{P_2}{H_2(g)}$

$E \quad \frac{RT}{F}\ln\frac{P_1}{P_2}$

95. (a) $A: \quad H_2(g) \rightleftharpoons 2H\ (aq) \quad 2e$

$C: \quad Cu^2\ (aq) \quad 2e \rightleftharpoons Cu(s)$

96. (a) $E \quad \frac{0.0591}{3}\log\frac{1}{0.04} \quad 0.028$

97. (c) $Cd^{2+}(aq) \quad Fe(s) \rightleftharpoons Cd(s) \quad Fe^2\ (aq)$

At eqm. $1 \quad x \qquad\qquad x$

At equilibrium $E \quad \frac{0.0591}{2}\log\frac{Fe^{2+}}{Cd^{2+}}$

$$0.037 = \frac{0.0591}{2}\log\frac{x}{1 \quad x}$$

$x \quad [Fe^{2+}] \quad 0.947\ M$

$[Cd^{2+}] = 0.053\ M$

98. (d) $1.50 \quad E \quad \frac{0.0591}{3}\log\frac{0.3}{(0.1)^3}$

$E \quad 1.55$

99. (d) $G \quad nFE$

$E \quad \frac{1467 \quad 10^3}{(\ 14 \quad 96{,}500)} \quad 1.08$

100. (b) The cell reaction is

$H_2(g) \quad I_2(s) \rightleftharpoons 2H^+(aq) \quad 2I^-(aq)$

$$0.7714 = 0.535 \quad \frac{0.0591}{2}\log\frac{[H^+]^2[I^-]^2}{P_{H_2}}$$

$pH = 3$

101. (c) One design a cell in which the half cell reaction is

For More Material Join: @JEEAdvanced_2024

$CuS(s) \quad 2e^- \rightleftharpoons Cu(s) \quad S^{2-}(aq)$ occurs

so $Cu(s) \rightleftharpoons Cu^{2} \quad 2e^- \qquad G_1$

$$CuS + 2e^- \rightleftharpoons Cu(s) \quad S^{2-}(aq) \qquad G_2$$

$$CuS(s) \rightleftharpoons Cu^{2}\,(aq) \quad S^{2-}(aq) \qquad G_3$$

from $G_3 \quad G_1 \quad G_2$

$$E_{S^{2-}|CuS|Cu} \quad E_{Cu^{2}\ Cu} \quad \frac{0.0591}{2} \log K_{sp}$$

$$0.34 + \frac{0.0591}{2} \log (10^{-35})$$

$$0.694 \text{ V}$$

102. (a) $Pb \rightleftharpoons Pb^{2} \quad 2e \ , \quad G_1^{\circ}$

$$Pb^{2} \quad 2Br^{-} \rightleftharpoons PbBr_2(s), \quad G_2^{\circ}$$

$$Pb(s) \quad 2Br\ (aq) \rightleftharpoons PbBr_2(s) \quad 2e$$

$$G_3^{\circ} \quad 2F(0.126) \quad RT \ln \frac{1}{K_{sp}}$$

$$2F(0.248) \quad 2F(0.126) \quad RT \ln \frac{1}{K_{sp}}$$

$$\ln \frac{1}{K_{sp}} \quad \frac{2F}{RT}[0.248 \quad 0.126]$$

$$\log \frac{1}{K_{sp}} \quad \frac{2\ (0.122)}{(0.0591)} \quad 4.128$$

$$K_{sp} \quad 10^{-4.128} \text{ or } 7.4 \quad 10^{-5}$$

103. (b) $n \quad 2$ for given reaction so $G \quad nFE$

$$210 \quad 1000 \quad 2 \quad 96500 \quad E$$

$$E \quad 1.09 \text{ V}$$

104. (b) At equilibrium, $G \quad 0$, $E_{cell} \quad 0$

105. (d) $G \quad G \quad RT \ln Q$

$$nFE \quad 2.303\ RT \log \frac{[Cr^{3+}]}{[Ag^{+}]^3}$$

$$G \quad 3 \quad 96500 \quad 1.50 + 8.314 \quad 298 \quad 2.303 \log \frac{(0.1)}{(0.1)^3}$$

$$422838.3 \text{ J or } 422.83 \text{ kJ}$$

106. (d) $Au \quad e^- \quad Au(s) \quad E \quad 1.69 \text{ V} \quad \ldots(1); \quad G_1$

$Au^{3} \quad 3e^- \quad Au(s) \quad E \quad 1.40 \text{ V} \quad \ldots(2); \quad G_2$

From (2)-(1) $Au^{3} \quad 2e^- \quad Au^{+}; \quad G_3 \quad G_2 \quad G_1$

$$2 \quad F \quad E \quad 3 \quad F \quad 1.40 \quad 1 \quad 1.69 \quad F;$$

$$E \quad 1.255 \text{ V}$$

$E_{Au^{+}|Au^{3+}} \quad 1.255 \text{V}$

107. (c) $2Mn^{2+} \quad 2Mn^{3+} + 2e^-, \quad G_1$

$$Mn^{2+} + 2e^- \quad Mn, \qquad G_2$$

$$3Mn^{2+}(aq) \quad Mn(s) \quad 2Mn^{3+}(aq)$$

$$2 \quad F \quad E_3 \quad 2 \quad F \quad [\ 1.51]$$

$$2 \quad F \quad (\ 1.185)$$

$E_3 \quad 2.695$

$$E_3 \quad \frac{0.0591}{2} \log K_{eq}; \quad K_{eq} \sim 6.3 \quad 10^{-92}$$

108. (c) For given concentration cell;

$$E_{cell} \quad \frac{RT}{F} \log \frac{[H^{+}]_{RHS}}{[H^{+}]_{LHS}}$$

for max. emf $[H^{+}]_{LHS}$ should be min. and that is for $NH_4OH(aq)$

109. (c) Given cell is electrolyte concentration cell

$$E_{cell} \quad \frac{RT}{F} \ln \frac{[Ag^{+}]_{RHS}}{[Ag^{+}]_{LHS}}$$

where $[Ag^{+}] = \frac{K_{sp}}{[Cl^{-}]}$. For spontaneous chemical change conc. of Ag^{+} of cathodic compartment must be greater than anodic compartment.

110. (d) half cell reaction is

$Cr_2O_7^{2-} \quad 14\ H \quad 6e \quad 2Cr^{3+} + 7H_2O$

$E_{Cr_2O_7^{2-}|Cr^{3+}} \quad E_{Cr_2O_7^{2-}|Cr^{3+}}$

$$\frac{0.0591}{n} \log \frac{[Cr^{3+}]^2}{[Cr_2O_7^{2-}][H^{+}]^{14}}$$

$$\frac{0.0591}{6} \log [10^{-3}]^{14} \quad 0.414 \text{ V}$$

Oxidising power reduced = 0.414 V

111. (c) $S \quad nF \left(\frac{dE}{dT}\right)_p \quad 2 \quad 96500 \quad 1.5 \quad 10^{-4}$

$$28.95 \text{ J/K - mol}$$

112. (c) Thermodynamic efficiency of a cell is given by the ratio of Gibbs function change to enthalpy change in the overall cell reaction.

113. (c) $E_{cell} \quad E_{RP(RHS)} \quad E_{RP(LHS)}$

$$0.76 \quad (\ 1.36) \quad 0.6$$

$_rG \quad RT \ln K_{eq};$

For More Material Join: @JEEAdvanced_2024

or $\log K_{eq} \quad \frac{nFE}{RT \quad 2.303}$

$\frac{2 \quad 0.6}{0.06} \quad 20 ; K \quad 10^{20}$

114. (d) Combustion takes place.

115. (b) $\because \quad G \quad 0$ (Electrolytic cell)

116. (b) When aluminium surfaces are exposed to the atmosphere a thin invisible oxidise skin forms immediately, which protects the metal from further oxidation.

[$\because \ E^{\circ}_{OP}$ of Al is much longer]

117. (d) Acidic medium.

118. (b) $E^{\circ}_{Hg^2 \ |Hg} \quad E^{\circ}_{Hg_2^2 \ |Hg} \quad E^{\circ}_{Fe^2 \ |Fe} \quad E^{\circ}_{Pb^2 \ |Pb}$

119. (b) E°_{OP} of Zn E°_{OP} of Fe

120. (a) Metal acts as anode

121. (a) Voltage of cell increases

122. (b) Movement of ions

123. (b) Free ions

124. (c) Water is weak electrolyte.

125. (b) $G \quad K \quad \frac{A}{l}$

126. (c) $_m \quad \frac{1000}{M} \quad \frac{1}{} \quad \frac{1}{M} \quad 1000;$

$_m \quad \frac{1000}{xy}$

127. (b) $_{eq} \quad \frac{K \quad 1000}{N}$

128. (b) $G^* \quad \frac{l}{A} \quad \frac{0.02}{0.0004} \quad 50 \ m^{-1}$

$G \quad \frac{1}{R} \quad \frac{1}{50} \quad 0.02 \ S$

$G.G^* \quad 50 \quad 0.02 = 1 \ S \ m^{-1}$

129. (a) Conductance $(G) \quad \frac{1}{R} \quad \frac{1}{300};$

$\because \quad G \quad \frac{l}{A}$

Cell constant $\frac{l}{A} \quad \frac{0.013}{1} \quad 300 \quad 3.9 \ cm^{-1}$

130. (b) $HA(aq) \rightleftharpoons H \ (aq) \quad A^-(aq)$

At eqm. $c(1 - \quad) \qquad c \qquad c$

$K_a \quad \frac{c \quad ^2}{1}$; where $\frac{}{} \quad _m$

$K_a \quad \frac{c \quad \frac{m}{m}^2}{1 \quad \frac{m}{m}}$

$\frac{c \quad ^2_m}{_m(\quad _m \quad _m)}$

131. (c) On dilution number of ions decrease in unit volume hence specific conductance decreases. But separation between ions also increase hence equivalent conductance increases.

132. (c) $_{eq} \quad \frac{150}{3} \quad 50 \ S \ cm^2 \ eq^{-1}$

133. (d) $_m \quad n. \ _{eq}$; when n total cationic/Anionic charge

$\because \quad n \quad 6; \qquad _{eq} \quad _m/6$

134. (d) $_m(NaBr) \quad _m(NaCl) \quad _mKBr \quad _m(KCl)$

$_m(Na^+) \quad _m(Br^-) \quad 126.5 \quad 151.5 \quad 150$

$_m(Na^+) = 50$

135. (c) $_m \quad \frac{}{M} \quad 1000; (x \quad y) \quad \frac{}{M} \quad 1000$

$M \quad \frac{1000}{(x \quad y)}$

Solubility (in gL^{-1}) $\frac{1000 \quad 188}{x \quad y}$

136. (a) $G.G^* \quad \frac{1}{200} \ 2 \quad 0.01;$

$_{eq} \quad \frac{}{N} \quad 1000$

$\frac{0.01}{0.1} \quad 1000 \quad 100$

137. (b) $R \quad \frac{l}{A}$

$82.4 \quad \frac{1}{2.768} \quad 10^3 \quad \frac{l}{A}$

$\frac{l}{A} \quad 0.2281$

138. (a) Equivalent conductivity of $BaCl_2$

$127 \quad 76 \quad 203$

139. (c) Ionic mobility $(m^2 v^{-1} s^{-1})$

$\frac{\text{Velocity}}{\text{Voltage / Distance}}$

140. (b) $AgCl \quad AgNO_3 \quad HCl \quad HNO_3$

For More Material Join: @JEEAdvanced_2024

141. (b) $G^* \quad \frac{l}{a} \quad \frac{10}{100} \quad 0.1; G \quad 0.0001S$

$V \quad 100 \quad 10 \quad 1000\ cm^3 \quad 1$ litre

$GG^* \quad 0.1 \quad 0.0001 = 10^{-5}$

$m \quad \frac{1000}{M}$

$\frac{(0.1 \quad 0.0001) \quad 1000}{0.1} \quad 0.1\ Scm^2 mol^{-1}$

142. (b) On dilution, no. of ions decrease in per unit volume.

143. (c) On dilution degree of dissociation of a weak electrolyte increase, hence increase in ionic mobility.

144. (d) Completely dissociate into ions.

145. (d) Electric conduction of a salt solution in water depends on the extent of its ionization.

146. (b) Ionic molar conductivity of H is very high and NH_4OH is a weak electrolyte.

147. (b) Hydrated size order F Cl Br I

Ionic mobility $\frac{1}{\text{Hydrated size}}$

148. (a) Molar conductivity of H and OH are very high as compare to other ions.

Initially conductance of solution sharply decreases due to consumption of free H . After complete neutralization further slightly increases due to presence of OH .

149. (c) Before neutralization conductance of solution sharply decreases due to consumption of free H . After neutralization further conductance almost remains same due to formation of buffer solution.

150. (d) Molar conductivity of H and OH are very high as compared to other ions.

Initially conductance of solution sharply decreases due to consumption of free H then increases due to formation of salt (NaCN) and after complete neutralization further sharply increases due to presence of OH

Level 2

1. (b) No. of equivalent of aluminum, $\frac{W}{E} \quad \frac{I \quad t}{96500}$

$\frac{24 \quad 5}{27} \quad 3 \quad \frac{9650 \quad 0.9 \quad t}{96500}$

$t \quad 148.14$ sec

2. (c) At Anode:

$2H_2O(l) \quad 4H\ (aq) \quad O_2(g) \quad 4e^-$

At Cathode : $Ag\ (aq) \quad e \quad Ag(s)$

Eq. of O_2 evolved Eq. of Ag formed 0.1

Total loss in mass $\frac{0.1 \quad 32}{4} \quad 0.1 \quad 108$

$0.8 + 10.8 \quad 11.6$

mass of final solution $108 \quad 11.6 \quad 96.4$ g

3. (d) Equivalents of CO_2 produced $\frac{(I \quad) \quad t}{96500}$

$\frac{0.5 \quad 0.8 \quad 96.5 \quad 60}{96500} \quad 0.024$

moles of CO_2 ($n \quad 1$) produced 0.024

moles of C_2H_6 ($n \quad 2$) produced

$\frac{0.024}{2} \quad 0.012$

Total moles of gases produced 0.036

$V_{gases} \quad \frac{nRT}{P} \quad \frac{0.036 \quad 0.0821 \quad 300}{\frac{740}{760}}$

0.91 litre

4. (d) Total volume of metal layer

$0.25 \quad 0.32 \quad 10^{-3}$

$0.08 \quad 10^{-3}\ m^3$

Total mass of chromium layer

$0.08 \quad 7.20 \quad 10^6 \quad 10^{-3} \quad 576$ g

$\frac{576}{52} \quad 6 \quad \frac{I \quad 60}{96500}$

$I \quad 10.69 \quad 10^4$ A

5. (a) Initial m moles of $Cu^{2+} = 5$;

m-eq. or m-moles of H produced

$100 \quad 10^{-2} \quad 1$

m-moles of Cu^2 converted into

Cu $\frac{1}{2} = 0.5$

m-moles of Cu^2 remaining in solution

$5 \quad 0.5 = 4.5$

$2Cu^{2+} + 4I^- \quad Cu_2I_2 + I_2$

and $I_2 + 2NaS_2O_3 \quad 2NaI + Na_2S_4O_6$

m-moles of Cu^2 remaining m-moles of $Na_2S_2O_3$

$4.5 = 0.04 \quad V \quad V \quad 112.5$ mL

For More Material Join: @JEEAdvanced_2024

6. (b) $2H_2O(l) \rightleftharpoons 2H^+(aq) \quad 2OH\ (aq); \qquad G_1$

$2H\ (aq) \quad 2e \rightleftharpoons H_2(g); \qquad G_2$

$2H_2O(l) \quad 2e \rightleftharpoons H_2(g) \quad 2OH\ (aq); \quad {}_rG_3$

$${}_rG_3 \quad G_1 \quad G_2$$

$$2 \quad F \quad E \quad RT \ln (10^{\ 26}) \quad 2 \quad F \quad 0$$

$$E \quad \frac{0.066}{2} \log (10^{\ 26}) \quad 0.858\ V$$

7. (d) To find n we break the cell reaction into two half cell reactions

Anode : $13H_2O(l) \quad 6.5\,O_2(g) + 26H^+(aq.) \quad 26e$

Cathode : $4CO_2(g) \quad 26H^+(aq.) \quad 26e \quad C_4H_{10}(g) \quad 8H_2O(l)$

$$E \quad \frac{G}{nF} \quad \frac{(\ 2746) \quad 1000}{26 \quad 96500} \quad 1.09\ V$$

8. (c) For normal calomel electrode $E_{RP} \quad E_{RP}$;

$$E_{cell} \quad (0.28 - 0) \quad \frac{0.0591}{2} \log \frac{[H^+]^2}{P_{H_2}}$$

$$0.5755 = 0.28 + 0.0591\ pH + \frac{0.0591}{2} \log (0.1)$$

$$pH = 5.5$$

9. (a) We know

$$2H_2(g) \quad 4H \quad 4e\ ; \quad E \quad 0.0\ V$$

$$O_2(g) \quad 4H\ (aq) \quad 4e \quad 2H_2O(l)\ ; E \quad 1.23\ V$$

$$2H_2(g) \quad O_2(g) \quad 2H_2O(l)\ ;\ E_{cell} \quad 1.23\ V$$

$$G_{298} \quad nFE \quad 4 \quad 96500 \quad 1.23 \quad 474.78\ kJ$$

$${}_rH_{298} \quad {}_rG_{298} \quad T.\ {}_rS_{298}$$

$$= (-474.78) \quad 298 \quad (-0.32)$$

$$570.14\ kJ/mol$$

$${}_fH_{298} \quad 285.07\ kJ/mol$$

10. (a) After addition of Cd and its oxidation into Cd^{2+}.

$$NO_3^-(aq) \quad 4H^+(aq) \quad 3e \quad NO(g) \quad 2H_2O(l)$$

$$0.1 \quad x \quad 0.4 \quad 4x;$$

where $x \quad 0.08$

$[NO_3^-]$ remaining $\quad 0.02\ M$; $[H^+]$ remaining $\quad 0.08\ M$

$$E_{NO_3^-|NO} \quad E_{NO_3^-|NO} \quad \frac{0.0591}{3} \log \frac{1}{[NO_3^-][H^+]^4}$$

$$0.96 - \frac{0.0591}{3} \log \frac{1}{(0.02)(0.08)^4} = 0.84\ V$$

For More Material Join: @JEEAdvanced_2024

11. (a) E for normal calomel electrode E for saturated calomel electrode

$$0.33 = 0.27 - \frac{0.06}{2}\log [Cl^-]^2; \quad [Cl^-] \quad 0.1\,M$$

12. (b) Anode : $H_2(g)$ $2H^+(aq)$ $2e$; E 0.0 V

Cathode : $MnO_4(aq)$ $8H^+(aq)$ $5e$ $Mn^{2+}(aq)$ $4H_2O(l)$; E 1.51 V

E_{cell} $E_{RP(RHS)}$ $E_{RP(LHS)}$

$$1.51 \quad \frac{0.06}{5}\log\frac{[Mn^{2+}]}{[MnO_4^-][H^+]^8} \quad 0 \quad \frac{0.06}{2}\log\frac{P_{H_2}}{[H^+]^2}$$

$$1.51 \quad \frac{0.06}{5}\log\frac{0.01}{0.1 \quad (10^{-2})^8} \quad \frac{0.06}{2}\log \quad \frac{0.1}{(10^{-3})^2}$$

$= 1.48$ V

13. (c) $3Cu(s)$ $8H^+$ $2NO_3^-$ $3\,Cu^{2+}$ $2NO$ $4H_2O$

$Cu(s)$ $4H^+$ $2NO_3^-$ Cu^{2+} $2NO_2$ $2H_2O$

Let concentration of HNO_3 is X so $[H^+] = X$

and $[NO_3^-]$ X

$$E_{NO_3^-|NO} \quad E_{Cu^{2+}|Cu} \quad E_{NO_3^-|NO_2} \quad E_{Cu^{2+}|Cu}$$

or $$E_{NO_3^-|NO} \quad E_{NO_3^-|NO_2}$$

$$0.96 - \frac{0.0591}{3}\log\frac{10^{-3}}{x^5} \quad 0.79 \quad \frac{0.0591}{1}\log \quad \frac{10^{-3}}{x^3}$$

$$0.62 = \frac{0.0591}{6}\log\frac{10^{-9}}{x^{10}} \quad 0.45 \quad \frac{0.0591}{2}\log\frac{10^{-9}}{x^6}$$

$$\log x \quad 0.657 \sim 0.66 \text{ or } x \quad 10^{+0.66}$$

14. (a) Anode : $2Cl^-(aq)$ $Cl_2(g)$ $2e^-$

Cathode : $Cl_2(g)$ $2e^-$ $2Cl^-(aq)$

Cell reaction : $2\,Cl^-(aq)$ $Cl_2(g)$ $2\,Cl^-(aq)$ $Cl_2(g)$

$$E_{cell} \quad \frac{0.06}{2}\log\frac{[Cl^-]^2_{RHS}\cdot(P_{Cl_2})_{LHS}}{[Cl^-]^2_{LHS}\cdot(P_{Cl_2})_{RHS}} \qquad (\because E_{cell} \quad 0)$$

$$\frac{0.06}{2}\log\frac{(0.01)^2}{(0.1)^2} \quad \frac{0.4}{0.2} \quad 0.051 \text{ V}$$

15. (a) $H_2O(l) + ClO_3^-(aq)$ $ClO_4^-(aq) + 2H^+(aq) + 2e$

$2H^+(aq)$ $ClO_3(aq) + 2e$ $2ClO_2^-(aq) + H_2O(l)$

$2ClO_3^-(aq)$ $ClO_2^-(aq) + ClO_4^-(aq)$

E_{cell} 0.33 0.36 = 0.03; E_{cell} $\frac{RT}{2F}\ln K$

For More Material Join: @JEEAdvanced_2024

$$0.03 \quad \frac{0.06}{2} \log K \quad \text{or} \quad K \quad 0.1$$

$$\underset{0.1}{2ClO_3^-} \rightleftharpoons \underset{x}{ClO_4^-} + \underset{x}{ClO_2^-}$$

$$\frac{x^2}{(0.1-2x)^2} \quad \frac{1}{10}; \; 3.16x \quad 0.1 \quad 2x$$

$$5.16x \quad 0.1$$

$$x \quad 0.1/5.16 = 0.0193 \quad 1.9 \quad 10^{-2}$$

16. (b)

$$E_{cell} \quad \frac{0.06}{1} \log \frac{[H^+]_{RHS}}{[H^+]_{LHS}}$$

or

$$E_{cell} \quad 0.06\,[(pH)_{LHS} \quad (pH)_{RHS}]$$

$$(pH)_{LHS} \quad pK_a \quad \log \frac{[A^-]}{[HA]}$$

$$4 \quad \log \frac{1}{0.1} \quad 5; \quad (pH)_{RHS} = 4$$

$$E_{cell} \quad 0.06\,(5 \quad 4) \quad 0.06 \text{ V}$$

17. (c)

$$E_{cell} \quad E_{H^+|H_2} \quad E_{OH^-|Cd(OH)_2|Cd}$$

$$E_{H^+|H_2} \quad E_{Cd^{2+}|Cd}$$

or

$$E_{cell} \quad E \quad \frac{0.06}{2} \log \frac{[Cd^{2+}]}{[H^+]^2}$$

$\because$

$$E_{cell} \quad 0,$$

$$E_{cell} \quad 0.03 \log \frac{[Cd^{2+}][OH^-]^2}{K_w^2}$$

$$\log \frac{K_{sp}}{K_w^2} \quad \frac{0.39}{0.03} \quad 13$$

$$K_{sp} \quad 10^{13} \quad (10^{-14})^2 \quad 10^{-15}$$

18. (a)

$$E_{cell} \quad \frac{0.06}{2} \log \frac{[H^+]^2_{RHS}[P_{H_2}]_{LHS}}{[H^+]^2_{LHS}[P_{H_2}]_{RHS}}; \quad \because \quad 1$$

$$[OH^-] = \sqrt{K_b \quad C}; \; [OH^-] = 10^{\quad 3}$$

or

$$[H^+]_{LHS} \quad 10^{\quad 11}$$

$$[H^+]_{RHS} \quad \sqrt{K_a \quad C} \quad 10^{-4}$$

$$E_{cell} \quad \frac{0.06}{2} \log \frac{(10^{\quad 4})^2 \quad 0.1}{(10^{\quad 11})^2} \quad 1 \quad 0.39 \text{ V}$$

For More Material Join: @JEEAdvanced_2024

19. (c) $H_2S \rightleftharpoons 2H^+ + S^{2-}$

$K_{a_1}.K_{a_2} \quad \frac{[H^+]^2[S^{2-}]}{[H_2S]}$

$10^{-21} \quad 0.1 \quad (10^{-3})^2[S^{2-}]$

$[S^{2-}] \quad 10^{-16}$

$E_{S^{-2}|Ag_2S|Ag} \quad E_{Ag^+|Ag} \quad \frac{0.06}{2}\log\frac{[S^{2-}]}{K_{sp}}$

$= 0.80 - \frac{0.06}{2}\log\frac{10^{-16}}{10^{-49}}$

$0.80 \quad \frac{0.06}{2} \quad 33 \quad 0.19$ V

20. (c) Normality of resultant solution

$\frac{0.1 \quad V}{V \quad V} \quad 0.05\ N$

$_{eq} \quad 1000 \quad \frac{0.0055}{0.05} \quad 110$ S cm^{2}eq^{-1}

21. (b) Resultant conc. of NaCl $\frac{0.1 \quad V}{4\ V} \quad 0.025\ N$

and HCl $\frac{0.1 \quad 2\ V}{4\ V} \quad 0.05\ N$

$_{solution}^{total} \quad \frac{N(NaCl).\ _{eq}^{NaCl}(NaCl)}{1000} \quad \frac{N(HCl).\ _{eq}(HCl)}{1000}$

$0.018 = \frac{0.025 \quad 110}{1000} \quad \frac{0.05.\ _{eq}(HCl)}{1000}$

$_{eq}(HCl) \quad \frac{18 \quad 2.75}{0.05} \quad 305$ S cm^{2}eq.

22. (b) The molar conductivity of the dissociated form of crotonic acid is

$_m(HC) \quad _m(HCl) \quad _m(NaC) \quad _m(NaCl)$

$(426 \quad 83 \quad 126)$ $^{-1}$ cm^{2}mol^{-1}

$383 \quad ^{-1}$ cm^{2}mol^{-1}

The molar conductivity of HC $\frac{1000}{m}$

$_m(HC) \quad \frac{3.83 \quad 10^{-5} \quad ^{-1}\text{cm}^{-1}}{0.001} \quad 1000$

$= 38.3 \quad ^{-1}$ cm^{2}mol^{-1}

The degree of dissociation,

$\frac{_m(HC)}{_m(HC)} \quad \frac{(38.3 \quad ^{-1}\text{ cm}^2\text{mol}^{-1})}{(383 \quad ^{-1}\text{ cm}^2\text{mol}^{-1})} \quad 0.1$

$K_a \quad \frac{C^{\ 2}}{1} \quad \frac{(10^{-3})(0.1)^2}{1 \quad 0.1} \quad 1.11 \quad 10^{-5}$

23. (c) $_m(BaSO_4) \quad 2 \quad _{eq}(BaSO_4)$

$_{eq}(BaSO_4) \quad _{eq}(Ba^{2\ }) \quad _{eq}(SO_4^{2\ })$

$_{eq}(BaCl_2) \quad _{eq}(H_2SO_4) \quad _{eq}(HCl)$

$_{eq}(BaSO_4) \quad x_1 \quad x_2 \quad x_3$

$_m \quad 2(x_1 \quad x_2 \quad x_3)$

for sparingly soluble salt

$_m \quad \frac{}{M} \quad 1000$

or $M \quad \frac{x}{2(x_1 \quad x_2 \quad x_3)} \quad 1000$

$\frac{500x}{(x_1 \quad x_2 \quad x_3)}$

$K_{sp} \quad M^2 \quad \frac{2.5 \quad 10^5 x^2}{(x_1 \quad x_2 \quad x_3)^2}$

24. (a) Conductivity of Na_2SO_4 $2.6 \quad 10^{-4}$

$_m(Na_2SO_4) \quad \frac{1000 \quad 2.6 \quad 10^{-4}}{0.001}$

260 S cm^{2}mol^{-1}

$_m(SO_4^{2-}) \quad _m(Na_2SO_4) \quad 2 \quad _m(Na\ )$

$260 \quad 2 \quad 50$

160 S cm^{2}mol^{-1}

Conductivity of $CaSO_4$ solution

$7 \quad 10^{-4} \quad 2.6 \quad 10^{-4}$

$= 4.4 \quad 10^{-4}$ S cm^{-1}

$_m(CaSO_4) \quad _m(Ca^{2\ }) \quad _m(SO_4^{2-})$

$120 \quad 160$

280 S cm^{2}mol^{-1}

Solubility $C \quad \frac{1000}{_m} \quad \frac{1000 \quad 4.4 \quad 10^{-4}}{280}$

$1.57 \quad 10^{-3}\ M$

$K_{sp} \quad [Ca^{2+}][SO_4^{2-}]_{total}$

$(0.00157)(0.00157 + 0.001)$

$= 4.0 \quad 10^{-6}\ M^2$

25. (b) $K_a \quad \frac{c^{\ 2}}{1} \quad 1.6 \quad 10^{\ 5} \quad \frac{0.01^{\ 2}}{1}$

$\sqrt{\frac{1.6 \quad 10^{-5}}{0.01}} \quad 0.04$

$\frac{_m}{_m}; \quad _m \quad 0.04 \quad 380 \quad 10^{\ 4}$

$15.2 \quad 10^{\ 4}$ S m^{2}mol^{-1}

$_m \quad C \quad 15.2 \quad 10^{\ 4} \quad 10^3 \quad 10^{\ 2}$

$(\because 1\text{ m}^3 \quad 1000\text{ litre})$

$k \quad G.G^*$ and $G^* \quad 0.01\text{ m}^{-1}$,

For More Material Join: @JEEAdvanced_2024

where $G^* = \frac{l}{A}$

$G = \frac{1.52 \times 10^{-2}}{0.01} = 1.52$ S

26. (c) Cell X : At anode Cl_2 gas evolved; At cathode H_2 gas evolved.

Cell Y : At anode O_2 gas evolved; At cathode Ag deposited.

Cell Z : At anode O_2 gas evolved; At cathode Cu deposited

Eqn. of Ag deposited $= \frac{W}{E} = \frac{21.6}{108} = 0.20$

Eqn. of O_2 gas evolved; Vol. of O_2 gas evolved $= 0.2 \times 5.6 = 1.12$ L

Vol. of H_2 evolved $=$ Vol. of Cl_2 evolved $= 0.2 \times 11.2 = 2.24$ L

27. (a) Eq. of H_2 = Eq. of O_2 = Eq. of $H_2S_2O_8$

$2 \times \frac{22.4}{22.4} = \frac{4 \times 8.4}{22.4} = 2 \times n$

$2 = 1.5 = 2n \quad n = 0.25$

Level 3

Passage–1

1. (c), **2.** (a), **3.** (d)

Equivalent of Zn^{2+} produced $= 0.1$ or moles of $Zn^{2+} = \frac{0.1}{2} = 0.05$

+ve charge increases in first compartment so due to interaction and maintain electrical neutrality Zn^{2+} move toward II compartment and NO_3^- move towards first compartment. Solution is always electrically neutral so charge of 1 Zn^{2+} is neutralized by 2 NO_3^-.

$[Zn^{2+}]$ in first compartment

$= 1 + \frac{0.05}{2} = 1.025\ M$

Concentration of NO_3^- in second compartment

$= 1 - 0.05 = 0.95\ M$

In third compartment moles of Cu^{2+} reduced

$= \frac{0.05}{2} = 0.025$

Relatively ve charge increased so SO_4^{2-} and Na move toward opposite direction to maintain electrical neutrality

$[SO_4^{2-}]_{remaining} = 1 - \frac{0.025}{2} = 0.975\ M$

Passage–2

1. (b) From equation 1 and 2, $S = \frac{d(\ G)}{dT}$

$S = \frac{d(nFE)}{dT} = nF\frac{dE}{dT}$ or $\frac{dE}{dT}_p = \frac{S}{nF}$

2. (b) From equation 2;

$H = nF\frac{dE}{dT}_p T + nFE_{cell}$

$218 \times 1000 = 2 \times 96500 \times 300\frac{dE}{dT}_p + 2 \times 96500 \times 1.015$

$\frac{dE}{dT}_p = 3.81 \times 10^{-4}\ VK^{-1}$

3. (a) $S = nF\frac{dE}{dT}_p = 2 \times 96500 \times 3.81 \times 10^{-4}$

$= 75.53$ J/mol-K

Passage–3

2. (b) $_m.C = \frac{_m.M}{1000}$

$\frac{200 \times 0.04}{1000} = 8 \times 10^3\ Scm^{-1}$

$G = \frac{l}{A} = 8 \times 10^{-3} \quad G = \frac{0.50}{2}$

or $G = 0.032$ S; $V = IR = \frac{I}{G}$

$I = 5 \times 0.032 = 0.16$ A

Passage-4

1. (c) $2 \times \frac{4.48}{22.4} \times \frac{1}{2} = \frac{I \times 9650}{96,500}$

$I = 4$ amp.

2. (b) Mass of $H_2O = \frac{4.48}{22.4} \times 18 = 3.6$ g

3. (b) $2H_2O(l) \rightarrow 4H^+(aq) + O_2(g) + 4e^-$

$2Cu^{2+} + 4e^- \rightarrow 2Cu(s)$

$2Cu^{2+} + 2H_2O(l) \rightarrow 4H^+(aq) + O_2(g) + 2Cu(s)$

$Moles_{Cu} = 2 \times \frac{4 \times 241.25}{96,500}$

$Mole_H = 2 \times$ mole of Cu $= 0.005 \times 2 = 0.01$

pH $= 2$

Passage-5

$Ag_xS(s) \rightleftharpoons Ag_{(x-y)}^{+}(aq) + S_x^{2-}(aq)$

$Ag_yS(s) \rightleftharpoons Ag_{(x-y)}^{+}(aq) + S_y^{2-}(aq)$

For More Material Join: @JEEAdvanced_2024

$K_{sp}(AgX) \quad x(x \quad y) \quad 3 \quad 10^{\quad 12}$ ...(1)

$K_{sp}(AgY) \quad y(x \quad y) \quad 10^{\quad 12}$ ...(2)

$K_{(AgX \quad AgY)} \quad K_{Ag} \quad K_x \quad K_y \quad 0.4 \quad 10^{\quad 6}$ ...(3)

$_{Ag} \quad \dfrac{1000 \quad K_{Ag}}{(x \quad y)} \quad 60$ ...(4)

$_{x} \quad \dfrac{1000 \quad K_{x}}{x} \quad 90$ ...(5)

$K_y \quad 1.45 \quad 10^{\quad 7}$

$_{y} \quad \dfrac{1000 \quad 1.45 \quad 10^{\quad 7}}{5 \quad 10^{\quad 7}} \quad 290$


$x \quad y \quad 2 \quad 10^{\quad 6}$ ...(6)

$\dfrac{x}{y} \quad 3$ ...(7)

$x \quad 1.5 \quad 10^{\quad 6}, \; y \quad 5 \quad 10^{\quad 7}, \; K_{Ag} \quad 1.2 \quad 10^{\quad 7}$

$K_x \quad 1.35 \quad 10^{\quad 7} \quad \dfrac{4}{3}x \quad 2 \quad 10^{\quad 6}$

Subjective Problems

1. $6e^- +$ 

Required charge for reduction of 1 mole $C_6H_5NO_2$ 6F

2. Equivalent mass $O_2 \quad \dfrac{32}{4} \quad 8$

3. $\dfrac{0.967}{63.5} \quad 2 \quad \dfrac{2}{197} \quad n_f$

$n_f \quad 3$

4. $\dfrac{0.18}{\text{Eq. mass}} \quad \dfrac{5 \quad 60 \quad 9.65}{96500}$, Eq. mass 6

5. 2 mole $Cl_2 \quad \dfrac{4F}{F}$

mole $Cl_2 \quad 2$

6. $G \quad 96.5 \quad 10^3 \quad 1 \quad 96500 \quad E$

$E \quad 1$

7. $K_C \quad 10^{\frac{2(0.34 \quad 0.15)}{0.0591}} \quad 2.6 \quad 10^6$

8. $4H \quad MnO_4 \quad 3e \rightleftharpoons MnO_2 \quad 2H_2O$

$G \quad 3 \quad F \quad \dfrac{1.5 \quad 5 \quad 2 \quad 1.25}{3}$

$5F \; ; \quad x \quad 5$

9. $Cd^{2} \quad 2e \rightleftharpoons Cd \; , \quad G_1^\circ$

$Cd \quad 4NH_3 \rightleftharpoons [Cd(NH_3)_4]^2 \quad 2e \; , \quad G_2^\circ$

$Cd^{2} \quad 4NH_3 \rightleftharpoons [Cd(NH_3)_4]^2 \; , \quad G_3^\circ$

$G_1^\circ \quad G_2^\circ \quad G_3^\circ$

$2 \quad F \quad (\ 0.40) \quad 2 \quad F(\ 0.61) \quad RT \ln K_f$

$\ln K_f \quad \dfrac{2F}{RT}(0.61 \quad 0.4)$

$2 \quad F(\ 0.4) \quad 2F(0.61)$

$\log K_f \quad \dfrac{2 \quad 0.21}{0.0591} \quad 7 \qquad K_f \quad 10^7$

10. $2H_2O(l) \qquad 4H \; (aq). \; O_2(g) \quad 4e$

$0.81 \quad 1.23 \quad \dfrac{0.0591}{4}\log [H \;]^4$

pH 7

11. $1500 \quad \dfrac{1}{2 \quad 10^{\quad 3}} \quad \dfrac{l}{A}$

Cell constant $1500 \quad 2 \quad 10^{\quad 3} \quad 3$

12. $CH_3COOH \qquad HCl \qquad CH_3COOK \qquad KCl$

$0.038 \quad 0.009 \quad 0.013$

0.034

% ionisation of $CH_3COOH \quad — \quad 100$

$\dfrac{2.72 \quad 10^{\quad 3}}{0.034} \quad 100 \quad 8$

13. $0.12 \quad \dfrac{0.0591}{2}\log \quad \dfrac{1}{x} \qquad 0.24$

$\log \dfrac{1}{x} \quad \dfrac{0.12 \quad 2}{0.06} \quad 4$

$x \quad 10^{\quad 4}$

14. $_M(HA) \qquad _M(HCl) \qquad _M(NaA) \qquad _M(NaCl)$

$425 \quad 100 \quad 125 \quad 400 \; S\,cm^{\,2}\,mol^{\quad 1}$

pH 4, $[H \;] \quad 10^{\quad 4} \qquad C$

$\dfrac{\quad m}{m} \quad \dfrac{200}{400} \quad 0.5;$

$K_a \quad \dfrac{(C \quad)}{(1 \quad)} \quad \dfrac{10^{\quad 4}(0.5)}{(1 \quad)} \quad 10^{\quad 4}; \; pK_a \quad 4$

15. $E_{Cl^{\quad}/AgCl/Ag} \quad E_{Ag/Ag} \quad \dfrac{0.0591}{1}\log K_{sp}$

$0.209 \quad 0.80 \quad \dfrac{0.0591}{1}\log K_{sp}$

$K_{sp} \quad 10^{\quad 10}$; Let solubility of AgCl in 0.01 M NaCl solution is x

$10^{\quad 10} \quad x(x \quad 0.01)$

$x \quad 10^{\quad 8}$

Moles of AgCl dissolved in

$10\,L \quad 10^{\quad 8} \quad 10 \quad 10^{\quad 7}$

For More Material Join: @JEEAdvanced_2024



9

DILUTE SOLUTIONS

Vapour Pressure : The pressure exerted by the vapours of liquid which are in equilibrium with it at a given temperature.

- Vapour pressure is not a surface phenomena.
- Vapour pressure increases with increase in temperature.

$$X(l) \rightleftharpoons X(g)$$

$$K_p \quad P_{x(g)}$$

$$\ln\frac{K_{P_2}}{K_{P_1}} \quad \ln\frac{P_2}{P_1} \quad \frac{H_{\text{vap.}}}{R} \quad \frac{1}{T_1} \quad \frac{1}{T_2} \ ;$$

where $H_{\text{vap.}}$ is molar enthalpy of vaporisation (j/mol or cal/mol)

Raoult's Law

(i) Liquid solution containing non volatile solute :

Vapour pressure of liquid solution Mole fraction of solvent in liquid solution.

$$P_{\text{Solution}} \quad X_{\text{Solvent}}$$

$$P_{\text{Solution}} \quad P_{\text{Solvent}} \cdot X_{\text{Solvent}}$$

Where P_{Solvent} Vapour pressure of pure liquid solvent, X_{Solvent} Mole fraction of solvent.

For binary liquid solution ; $X_{\text{Solute}} \quad X_{\text{Solvent}} \quad 1$

Relative lowering of vapour pressure $\frac{P \quad P}{P} \quad X_{\text{Solute}}$

(ii) Vapour Pressure of Solution containing volatile solute (B) and volatile solvent (A) :

The equilibrium vapour pressure of a volatile component is linearly proportional to the mole fraction of that component in liquid phase.

For More Material Join: @JEEAdvanced_2024

A solution contains volatile solute (B) and volatile solvent (A) :

$P_A \quad P_A X_A$ and $P_B \quad P_B X_B$

$P \quad P_A X_A \quad P_B X_B$

Where :

P_A and P_B are partial vapour pressure of A and B at given temperature.

P_A vapour pressure of pure A. ; P_B vapour pressure of pure B.

X_A mole fraction of A in liquid phase. ; X_B mole fraction of B in liquid phase.

y_A mole fraction of A in vapour phase ; y_B mole fraction of B in vapour phase.

Partial vapour pressure of 'A' mole fraction of 'A' in vapour phase total vapour pressure.

$P_A \quad y_A P$ and $P_B \quad y_B P$

$P_A X_A \quad y_A P$ and $P_B X_B \quad y_B P$

$X_A \quad X_B \quad 1$

$\frac{y_A P}{P_A} \quad \frac{y_B P}{P_B} \quad 1$; $\frac{y_A}{P_A} \quad \frac{y_B}{P_B} \quad \frac{1}{P}$

$y_A \quad \frac{P_A X_A}{P_A X_A \quad P_B X_B}$; $y_B \quad \frac{P_B X_B}{P_A X_A \quad P_B X_B}$

Ideal Solutions

The solutions which obey Raoult's Law are called ideal solutions. For ideality :

$F_{\text{Solute .....Solvent}} \quad F_{\text{Solute..... Solute}} \quad (F$ Intermolecular attractive interactions)

$F_{\text{Solvent .....Solvent}}$

$H_{\text{mix.}} \quad 0$; $V_{\text{mix.}} \quad 0$

$S_{\text{mix.}} \quad 0$; $G_{\text{mix.}} \quad 0$

Examples : Benzene & Toluene, Hexane & Heptane, C_2H_5Br & C_2H_5I

Non-ideal solutions : The mixture which do not follow Raoult's law is known as non-ideal solutions.

Types of Real or Non-ideal Solutions

- Non-ideal solution with positive deviation.
- Non-ideal solution with negative deviation.

Solution shows positive deviation

$P_{T,\text{exp}} \quad (P_A X_A \quad P_B X_B)$

A - - - -B A - - - -A

B - - - -B

$H_{\text{mix.}}$ ve

$V_{\text{mix.}}$ ve

$S_{\text{mix.}}$ ve

$G_{\text{mix.}}$ ve

Ex. $H_2O \quad CH_3OH$

$H_2O \quad C_2H_5OH$

$CHCl_3 \quad CCl_4$

Solution shows negative deviation

$P_{T,\text{exp}} \quad (P_A X_A \quad P_B X_B)$

A - - - -B A - - - -A

B - - - -B

$H_{\text{mix.}}$ ve

$V_{\text{mix.}}$ ve

$S_{\text{mix.}}$ ve

$G_{\text{mix.}}$ ve

Ex. $H_2O \quad CH_3COOH$

$CHCl_3 \quad CH_3COCH_3$

$H_2O \quad HNO_3$

For More Material Join: @JEEAdvanced_2024

Azeotropic Mixture. During distillation, the mole fraction of more volatile component in vapour state is higher than that in liquid state. This makes distillation possible. However, there exist some solutions for particular compositions of which the mole fraction of components in liquid and vapour state is same. Thus, no distillation is possible, a mixture and it is termed as **azeotropic mixture.**

Azeotropes are binary mixtures having the same composition in liquid and vapour phase and boil at a constant temperature.

The solutions which show a large positive deviation from Raoult's law form minimum boiling azeotrope at a specific composition.

The solutions which show a large negative deviation from Raoult's law form maximum boiling azeotrope at a specific composition.

Completely Immiscible Liquids : When they are distilled, they distil in the ratio of their vapour pressure at that temperature.

$$P \quad P_A \quad P_B$$

For vapour of pure liquid we can apply : $P_A V \quad n_A RT$ and $P_B V \quad n_B RT$

So, $$\frac{P_A}{P_B} \quad \frac{n_A}{n_B}$$

When A and B are distilled mass ratio is given as $\frac{w_B}{w_A} \quad \frac{P_B \quad M_B}{P_A \quad M_A}$

Colligative Properties : Colligative Properties depend on the number of solute particles irrespective to their nature.

- ❖ Relative lowering of vapour pressure
- ❖ Elevation of boiling point
- ❖ Depression of freezing point
- ❖ Osmotic pressure

Normal Colligative Properties : When neither association nor dissociation of solute particles take place.

(i) Relative Lowering of Vapour Pressure : $\frac{P \quad P}{P} \quad X_{\text{Solute}}$

(ii) Elevation of boiling point : $T_b \quad K_b m$

(iii) Depression of freezing point : $T_f \quad K_f m$

(iv) Osmotic pressure : CRT

(i) Relative lowering of Vapour Pressure :

$$\frac{P \quad P}{P} \quad X_{\text{Solute}} \quad \frac{n}{n \quad N}$$

Where, n moles of solute, N moles of solvent


$$\frac{P \quad P}{P} \quad \frac{n}{N}$$

Molality (m) $$\frac{P \quad P}{P} \quad \frac{1000}{M(\text{in g/mol})}$$

Where, M molar mass of solvent.

For More Material Join: @JEEAdvanced_2024

Ostwald-Walker Method : Experimental measure of relative lowering in V.P.



Loss in mass of solution containers P

Loss in mass of solvent containers $(P \quad P)$

gain in mass of dehydrating agent P

$\frac{P \quad P}{P}$ $\frac{\text{Loss in mass of solvent}}{\text{Gain in mass of dehydrating agent}}$

(ii) Elevation in Boiling Point :

$T_b \quad K_b m$

where $T_b \quad T_b \quad T_b$

K_b Boiling point elevation constant or ebullioscopic constant

$K_b \quad \frac{RT_b^{\,2}}{1000\,L_v}$

L_v Latent heat of vapourisation per gram

$K_b \quad \frac{MRT_b^{\,2}}{1000 \quad H_{\text{Vap}}}$

H_{vap} enthalpy of vapourisation per mole

M molar mass of solvent (in g/mol)

T_b Boiling point of solvent

(iii) Depression in Freezing point :

$T_f \quad K_f m$

where $T_f \quad T_f \quad T_f$

K_f Freezing point depression constant or cryoscopic constant

$K_f \quad \frac{RT_f^{\,2}}{1000\,L_f}$

L_f Latent heat of fusion per gram

$K_f \quad \frac{MRT_f^{\,2}}{1000 \quad H_{\text{Fus}}}$

H_{Fus} enthalpy of fusion per mole

M molar mass of solvent (in g/mol)

T_f freezing point of solvent

(iv) Osmotic Pressure ( ) : The hydrostatic pressure built up on the solution which just stops osmosis. Alternatively, it may be defined as the pressure which must be applied to the concentrated solution in order to stop osmosis.

For dilute solutions $CRT \quad h \; g$

where C is the total molar concentration of all the free species present in the solution, h is the height developed by the column of the concentrated solution and is the density of the solution in the column.

On the basis of osmotic pressure, the solutions can be classified in three classes.

- **Isotonic solutions:** Two solutions having same osmotic pressures are called isotonic solutions.

(This implies $C_1 \quad C_2$ at same temperature).

- **Hypertonic and hypotonic solution :** When two solutions are being compared, then the solution with higher osmotic pressure is termed as hypertonic. The solution with lower osmotic pressure is termed as **hypotonic.**

For More Material Join: @JEEAdvanced_2024

❖ **Abnormal Colligative Properties :** When solute particle associated or dissociated in solvent.

i Vant Hoff factor $\dfrac{\text{Actual moles of solute}}{\text{Moles of solute without dissociation or association}}$

$\dfrac{\text{Observed or experimental colligative properties}}{\text{Theoretical or calculated colligative properties}}$

(i) Relative lowering of vapour pressure : $\dfrac{P \quad P}{P} \quad \dfrac{i.n_{\text{solute}}}{i.n_{\text{solute}} \quad n_{\text{solvent}}}$

(ii) Elevation of boiling point : $T_b \quad i.K_b m$

(iii) Depression of freezing point : $T_f \quad i.K_f m$

(iv) Osmotic pressure : $i.CRT$

For dissociation : $i \quad 1 \quad (n \quad 1)$ For association : $i \quad 1 \quad \dfrac{1}{n} \quad 1$

Degree of dissociation $\dfrac{i \quad 1}{n \quad 1}$ Degree of association $\dfrac{1 \quad i}{1 \quad 1/n}$

Where n Total number of particles of solute after dissociation or association.

Henry Law

This law deals with dissolution of gas in liquid. It states that mass of any gas dissolved in any solvent per unit volume is proportional to pressure of gas in equilibrium with liquid.

(i) $m \quad k \quad P$

Where, m mass of gas dissolved per unit volume.

k proportionality constant.

P pressure of gas.

(ii) $P \quad K_H . X_{\text{gas}}$

Where, P Pressure of gas.

K_H Henry's constant

X_{gas} Mole fraction of gas

For More Material Join: @JEEAdvanced_2024



Level 1

1. The vapour pressure of a given liquid will decrease if :
(a) surface area of liquid is decreased
(b) the volume of liquid in the container is decreased
(c) the volume of the vapour phase is increased
(d) the temperature is decreased

2. The normal boiling point of water is 373 K. Vapour pressure of water at temperature T is 19 mm Hg. If enthalpy of vaporisation is 40.67 kJ/mol, then temperature T would be
(Use : log 2 0.3, R : 8.3 JK ^{1} mol ^{1}):
(a) 250 K (b) 291.4 K (c) 230 K (d) 290 K

3. A sample of liquid H_2O of mass 18.0 g is injected into an evacuated 7.6 L flask maintained at 27.0°C. If vapour pressure of H_2O at 27°C is 24.63 mm Hg, what weight percentage of the water will be vapourised when the system comes to equilibrium? Assume water vapours behaves as an ideal gas. The volume occupied by the liquid water is negligible compared to the volume of the container:
(a) 1% (b) 10% (c) 18% (d) 20%

4. Raoult's law is obeyed by each constituent of a binary liquid solution when:
(a) the forces of attractions between like molecules are greater than those between unlike molecules
(b) the forces of attractions between like molecules are smaller than those between unlike molecules
(c) the forces of attractions between like molecules are identical with those between unlike molecules
(d) the volume occupied by unlike molecules are different

5. For a binary ideal liquid solution, the total vapour pressure of the solution is given as:
(a) P_{total} P_A° $(P_A^\circ$ $P_B^\circ)\,X_B$ (b) P_{total} P_B° $(P_A^\circ$ $P_B^\circ)\,X_A$
(c) P_{total} P_B° $(P_B^\circ$ $P_A^\circ)\,X_A$ (d) P_{total} P_B° $(P_B^\circ$ $P_A^\circ)\,X_B$

6. For an ideal binary liquid solution with P_A P_B, which relation between X_A (mole fraction of A in liquid phase) and Y_A (mole fraction of A in vapour phase) is correct ?
(a) Y_A Y_B (b) X_A X_B (c) $\frac{Y_A}{Y_B}$ $\frac{X_A}{X_B}$ (d) $\frac{Y_A}{Y_B}$ $\frac{X_A}{X_B}$

7. An ideal solution is formed by mixing two volatile liquids A and B. X_A and X_B are the mole fractions of A and B respectively in the solution and Y_A and Y_B are the mole fractions of A and B respectively in the vapour phase. A plot of $\frac{1}{Y_A}$ along y-axis against $\frac{1}{X_A}$ along x-axis gives a straight line. What is the slope of the straight line ?
(a) $\frac{P_B}{P_A}$ (b) $\frac{P_A}{P_B}$ (c) P_B P_A (d) P_A P_B

For More Material Join: @JEEAdvanced_2024

8. For a dilute solution, Raoult's law states that :
(a) the lowering of vapour pressure is equal to the mole fraction of solute
(b) the relative lowering of vapour pressure is equal to the mole fraction of solute
(c) the relative lowering of vapour pressure is proportional to the amount of solute in solution
(d) the vapour pressure of the solution is equal to the mole fraction of solvent

9. The solubility of a specific non-volatile salt is 4 g in 100 g of water at 25°C. If 2.0 g, 4.0 g and 6.0 g of the salt added of 100 g of water at 25°C, in system X, Y and Z. The vapour pressure would be in the order:
(a) $X \quad Y \quad Z$ (b) $X \quad Y \quad Z$ (c) $Z \quad X \quad Y$ (d) $X \quad Y \quad Z$

10. The boiling point of C_6H_6, CH_3OH, $C_6H_5NH_2$ and $C_6H_5NO_2$ are 80 C, 65°C, 184°C and 212°C respectively. Which will show highest vapour pressure at room temperature :
(a) C_6H_6 (b) CH_3OH (c) $C_6H_5NH_2$ (d) $C_6H_5NO_2$

11. 6.0 g of urea (molecular mass = 60) was dissolved in 9.9 moles of water. If the vapour pressure of pure water is P , the vapour pressure of solution is:
(a) 0.10 P (b) 1.10 P (c) 0.90 P (d) 0.99 P

12. An ideal solution was found to have a vapour pressure of 80 torr when the mole fraction of a non-volatile solute was 0.2. What would be the vapour pressure of the pure solvent at the same temperature?
(a) 64 torr (b) 80 torr (c) 100 torr (d) 400 torr

13. The vapour pressure of an aqueous solution of sucrose at 373 K is found to be 750 mm Hg. The molality of the solution at the same temperature will be :
(a) 0.26 (b) 0.73 (c) 0.74 (d) 0.039

14. Estimate the lowering of vapour pressure due to the solute (glucose) in a 1.0 M aqueous solution at 100°C :
(a) 10 torr (b) 18 torr (c) 13.45 torr (d) 24 torr

15. Calculate the mass of non-volatile solute having molecular mass 40, which should be dissolved in 57 gm octane to reduce its vapour pressure to 80%:
(a) 47.2 g (b) 5 g (c) 106.2 g (d) None of these

16. Equal mass of a solute are dissolved in equal mass of two solvents A and B and formed very dilute solution. The relative lowering of vapour pressure for the solution B has twice the relative lowering of vapour pressure for the solution A. If M_A and M_B are the molecular mass of solvents A and B respectively, then:
(a) $M_A \quad M_B$ (b) $M_B \quad 2 \quad M_A$ (c) $M_A \quad 4M_B$ (d) $M_A \quad 2M_B$

17. An ideal solution has two components A and B. A is more volatile than B, *i.e.*, $P_A \quad P_B$ and also $P_A \quad P_{\text{total}}$. If X_A and Y_A are mole fractions of components A in liquid and vapour phases, then:
(a) $X_A \quad Y_A$ (b) $X_A \quad Y_A$ (c) $X_A \quad Y_A$ (d) Data insufficient

18. At 25°C, the vapour pressure of pure liquid A (mol. mass 40) is 100 torr, while that of pure liquid B is 40 torr, (mol. mass 80). The vapour pressure at 25°C of a solution containing 20 g of each A and B is:
(a) 80 torr (b) 59.8 torr (c) 68 torr (d) 48 torr


For More Material Join: @JEEAdvanced_2024

19. Two liquids *A* and *B* form ideal solutions. At 300 K, the vapour pressure of solution containing 1 mole of *A* and 3 mole of *B* is 550 mm Hg. At the same temperature, if one more mole of *B* is added to this solution, the vapour pressure of the solution increases by 10 mm Hg. Determine the vapour pressure of *A* and *B* in their pure states (in mm Hg):
(a) 400, 600 (b) 500, 500 (c) 600, 400 (d) None of these

20. Two liquids *A* and *B* have vapour pressure in the ratio $P_A : P_B$ 1:3 at a certain temperature. Assume *A* and *B* form an ideal solution and the ratio of mole fractions of *A* to *B* in the vapour phase is 4 : 3, then the mole fraction of *B* in the solution at the same temperature is :
(a) $\frac{1}{5}$ (b) $\frac{2}{3}$ (c) $\frac{4}{5}$ (d) $\frac{1}{4}$


21. Two liquids *A* and *B* have P_A and P_B in the ratio of 1 : 3 and the ratio of number of moles of *A* and *B* in liquid phase are 1 : 3. Then mole fraction of '*A*' in vapour phase in equilibrium with the solution is equal to:
(a) 0.1 (b) 0.2 (c) 0.5 (d) 1.0

22. Based on the given diagram, which of the following statements regarding the homogenous solutions of two volatile liquids are correct?



(1) Plots *AD* and *BC* show that Raoult's law is obeyed for the solution in which *B* is a solvent and *A* is the solute and as well as for that in which *A* is solvent and *B* is solute.
(2) Plot *CD* shows that Dalton's law of partial pressures is obeyed by the binary solution of components *A* and *B*.
(3) *EF* *EG* *EH*; and *AC* and *BD* correspond to the vapour pressures of the pure solvents *A* and *B* respectively.
(a) Only 1 (b) 2 and 3 (c) 1 and 3 (d) All

23. Which represents correct difference when non-volatile solute is present in an ideal solution?




(a) I, II, III (b) I, III (c) II, III (d) I, II

For More Material Join: @JEEAdvanced_2024

24. Select correct statement :
(a) Solution has more molecular randomness than a pure solvent. The entropy change between solution and solid is larger than the entropy change between pure solvent and solid
(b) Heat of fusion of solution and solvent are similar since similar forces of intermolecular forces are involved
(c) Solution containing sugar freezes at a lower temperature than pure water
(d) All are correct statements


25. Select correct statement?
(a) Heats of vaporisation for a pure solvent and for a solution are similar because similar intermolecular forces between solvent molecules must be overcome in both cases
(b) Entropy change between solution and vapour is smaller than the entropy change between pure solvent and vapour
(c) Boiling point of the solution is larger than that of the pure solvent
(d) All are correct statements

26. The vapour pressure curves of the same solute in the same solvent are shown below. The curves are parallel to each other and do not intersect. The concentrations of solutions are in order of:



(a) I < II < III (b) I = II = III (c) I > II > III (d) I > III > II

27. Boiling point composition diagram of the liquid-vapour equilibrium for *A* and *B* is shown in the figure. If a binary liquid mixture of *A* and *B* is distilled fractionally, which of the following would be correct observation?



(a) Composition of the still (residue) will approach pure liquid *B* only
(b) Composition of the distillate will approach pure *A* only
(c) Composition of distillate and residue will approach pure *A* and *B* respectively
(d) Neither of the component can be obtained in pure state

28. The boiling point of an azeotropic mixture of water–ethanol is less than that of both water and ethanol. Then:
(a) the mixture will show negative deviation from Raoult's law
(b) the mixture will show positive deviation from Raoult's law
(c) the mixture will show no deviation from Raoult's law
(d) this mixture can be considered as true solution


For More Material Join: @JEEAdvanced_2024

29. Formation of a solution from two components can be considered as :
(i) Pure solvent separated solvent molecules, H_1
(ii) Pure solute separated solute molecules, H_2
(iii) separated solvent and solute molecules solution, H_3
Solution so formed will be ideal if :
(a) H_{Soln} H_1 H_2 H_3 (b) H_{Soln} H_1 H_2 H_3
(c) H_{Soln} H_1 H_2 H_3 (d) H_{Soln} H_3 H_1 H_2

30. Total vapour pressure of mixture of 1 mol X (P_X 150 torr) and 2 mol Y (P_Y 300 torr) is 240 torr. In this case:
(a) there is a negative deviation from Raoult's law
(b) there is a positive deviation from Raoult's law
(c) there is no deviation from Raoult's law
(d) can not be decided

31. In a mixture of A and B, components show positive deviation when:
(a) A B interaction is stronger than A A and B B interaction
(b) A B interaction is weaker than A A and B B interaction
(c) V mix 0, S mix 0
(d) V mix 0, S mix 0

32. A liquid mixture having composition corresponding to point Z in the figure shown is subjected to distillation at constant pressure. Which of the following statement is correct about the process?



(a) The composition of distillate differs from the mixture
(b) The boiling point goes on changing
(c) The mixture has highest vapour pressure than for any other composition
(d) Composition of an azeotrope alters on changing the external pressure

33. Which will form maximum boiling azeotrope ?
(a) C_6H_6 $C_6H_5CH_3$ solution (b) HNO_3 H_2O solution
(c) C_2H_5OH H_2O solution (d) *n*-hexane and *n*-heptane

34. Total vapour pressure of mixture of 1 mole of volatile component A (P_A 100 mm Hg) and 3 mole of volatile component B (P_B 80 mm Hg) is 90 mm Hg. For such case:
(a) there is positive deviation from Raoult's law
(b) boiling point has been lowered
(c) force of attraction between A and B is weaker than that between A and A or between B and B
(d) all the above statements are correct

For More Material Join: @JEEAdvanced_2024

35. The azeotropic mixture of water (B.P. = 100 C) and HCl (B.P. = 86°C) boils at about 120°C. During fractional distillation of this mixture, it is possible to obtain :
(a) pure HCl (b) pure H_2O
(c) pure H_2O as well as pure HCl (d) neither H_2O nor HCl

36. Azeotropic mixture of water and C_2H_5OH boils at 351 K. By distilling the mixture it is possible to obtain :
(a) Pure C_2H_5OH only (b) Pure water only
(c) Neither C_2H_5OH nor water (d) Both water and C_2H_5OH in pure state

37. An azeotropic mixture of two liquids has a boiling point higher than either of them when it:
(a) shows positive deviation from Raoult's law
(b) shows negative deviation from Raoult's law
(c) shows ideal behaviour
(d) is saturated

38. If two liquids *A* (P_A 100 torr) and *B* (P_B 200 torr) which are completely immiscible with each other (each one will behave independently of the other) are present in a closed vessel, the total vapour pressure of the system will be:
(a) less than 100 torr (b) greater than 200 torr
(c) between 100 to 200 torr (d) 300 torr

39. When a liquid that is immiscible with water was steam distilled at 95.2°C at a total pressure of 748 torr, the distillate contained 1.25 g of the liquid per gram of water. The vapour pressure of water is 648 torr at 95.2°C, what is the molar mass of liquid?
(a) 7.975 g/mol (b) 166 g/mol (c) 145.8 g/mol (d) None of these

40. Water and chlorobenzene are immiscible liquids. Their mixture boils at 89°C under a reduced pressure of 7.7 10^4 Pa. The vapour pressure of pure water at 89°C is 7 10^4 Pa. Mass per cent of chlorobenzene in the distillate is :
(a) 50 (b) 60 (c) 78.3 (d) 38.46

41. Which of the following is not a colligative property?
(a) Vapour pressure (b) Depression in f.pt.
(c) Elevation in b.pt. (d) Osmotic pressure

42. The degree of dissociation of an electrolyte is and its van't Hoff factor is *i*. The number of ions obtained by complete dissociation of 1 molecule of the electrolyte is :
(a) $\frac{i \quad 1}{\quad}$ (b) $i \quad 1$ (c) $\frac{i \quad 1}{\quad}$ (d) $\frac{i \quad 1}{1}$

43. One mole of a solute *A* is dissolved in a given volume of a solvent. The association of the solute take place as follows:

$$nA \rightleftharpoons A_n$$

If is the degree of association of *A*, the van't Hoff factor *i* is expressed as:
(a) $i \quad 1$ (b) $i \quad 1 \quad \frac{}{n}$ (c) $i \quad \frac{1 \quad \frac{}{n}}{1}$ (d) $i \quad 1$

44. The van't Hoff factor *i* for an electrolyte which undergoes dissociation and association in solvent are respectively:
(a) greater than one and less than one (b) less than one and greater than one
(c) less than one and less than one (d) greater than one and greater than one

For More Material Join: @JEEAdvanced_2024

45. Which solution has the highest vapour pressure?
(a) 0.02 *M* NaCl at 50°C (b) 0.03 *M* sucrose at 15°C
(c) 0.005 *M* $CaCl_2$ at 50°C (d) 0.005 *M* $CaCl_2$ at 25°C

46. An aqueous solution is 1.00 molal in KI. Which change will cause the vapour pressure of the solution to increase ?
(a) addition of water (b) addition of NaCl
(c) addition of Na_2SO_4 (d) Addition of 1.0 molal KI

47. Four solutions of K_2SO_4 with the concentrations 0.1 *m*, 0.01 *m*, 0.001 *m* and 0.0001 *m* are available. The maximum value of colligative property corresponds to :
(a) 0.0001 *m* solution (b) 0.001 *m* solution (c) 0.01 *m* solution (d) 0.1 *m* solution

48. Moles of Na_2SO_4 to be dissolved in 12 mole water to lower its vapour pressure by 10 mm Hg at a temperature at which vapour pressure of pure water is 50 mm is:
(a) 1.5 mole (b) 2 mole (c) 1 mole (d) 3 mole

49. A very diluted saturated solution of a sparingly soluble salt X_3Y_4 has a vapour pressure of 20 mm Hg at temperature *T*, while pure water exerts a pressure of 20.0126 mm Hg at the same temperature. Calculate molality (*m*) at temperature *T*:
(a) 6.3 10 ^{4} (b) 3.5 10 ^{2} (c) 5 10 ^{3} (d) None of these

50. When 1 mole of a solute is dissolved in 1 kg of H_2O, boiling point of solution was found to be 100.5°C. K_b for H_2O is:
(a) 0.5 (b) 100 (c) 100.5 (d) 95.5

51. Chloroform, $CHCl_3$, boils at 61.7°C. If the K_b for chloroform is 3.63°C/molal, what is the boiling point of a solution of 15.0 kg of $CHCl_3$ and 0.616 kg of acenaphthalene, $C_{12}H_{10}$?
(a) 61.9 (b) 62.0 (c) 52.2 (d) 62.67

52. A compound has the empirical formula $C_{10}H_8Fe$. A solution of 0.26 g of the compound in 11.2 g of benzene (C_6H_6) boils at 80.26°C. The boiling point of benzene is 80.10°C; the K_b is 2.53°C/molal. What is the molecular formula of the compound?
(a) $C_{30}H_{24}Fe_3$ (b) $C_{10}H_8Fe$ (c) C_5H_4Fe (d) $C_{20}H_{16}Fe_2$

53. A solution of 0.640 g of azulene in 100.0 g of benzene boils at 80.23°C. The boiling point of benzene is 80.10°C; and K_b is 2.53°C/molal. What is the molecular mass of azulene?
(a) 108 (b) 99 (c) 125 (d) 134

54. One molal solution of a carboxylic acid in benzene shows the elevation of boiling point of 1.518 K. The degree of association for dimerization of the acid in benzene is (K_b for benzene 2.53 K kg mol ^{1}) :
(a) 60% (b) 70% (c) 75% (d) 80%

55. The boiling point elevation constant for toluene is 3.32 K kg mol ^{1}. The normal boiling point of toluene is 110.7°C. The enthalpy of vaporisation of toluene would be nearly:
(a) 17.0 kJ mol ^{1} (b) 34.0 kJ mol ^{1} (c) 51.0 kJ mol ^{1} (d) 68.0 kJ mol ^{1}

56. Which one of the following aqueous solutions will exhibit highest boiling point ?
(a) 0.015 *M* urea (b) 0.01 *M* KNO_3 (c) 0.10 *M* Na_2SO_4 (d) 0.015 *M* glucose

57. Calculate the percentage degree of dissociation of an electrolyte XY_2 (Normal molar mass 164) in water if the observed molar mass by measuring elevation in boiling point is 65.6:
(a) 75% (b) 25% (c) 65% (d) None of these

For More Material Join: @JEEAdvanced_2024

58. If the elevation in boiling point of a solution of non-volatile, non-electrolytic and non-associating solute in solvent (K_b x K.kg.mol^{-1}) is y K, then the depression in freezing point of solution of same concentration would be
(K_f of the solvent z K.kg.mol^{-1})
(a) $\frac{2xz}{y}$ (b) $\frac{yz}{x}$ (c) $\frac{xz}{y}$ (d) $\frac{yz}{2x}$

59. When a solution containing non-volatile solute freezes, which equilibrium would exist?
(a) solid solvent $\rightleftharpoons$ liquid solvent (b) solid solute $\rightleftharpoons$ liquid solution
(c) solid solute $\rightleftharpoons$ liquid solvent (d) solid solvent $\rightleftharpoons$ liquid solution

60. Bromoform has a normal freezing point of 7.734°C and K_f 14.4°C/m. A solution of 2.60 g of an unknown substance in 100 g of bromoform freezes at 5.43°C. What is the molecular mass of the unknown substance?
(a) 16.25 (b) 162.5 (c) 100 (d) none of these

61. C_6H_6 freezes at 5.5°C. At what temperature will a solution of 10.44 g of C_4H_{10} in 200 g of C_6H_6 freeze? K_f (C_6H_6) 5.12°C/m
(a) 4.608°C (b) 0.892°C (c) 5.5°C (d) none of these

62. How much ethyl alcohol must be added to 1.0 L of water so that solution will not freeze at –4°F? (K_f 1.86°C/m)
(a) < 20 g (b) < 10.75 g (c) < 494.5 g (d) > 494.5 g

63. The freezing point of a solution of 2.40 g of biphenyl ($C_{12}H_{10}$) in 75.0 g of benzene (C_6H_6) is 4.40°C. The normal freezing point of benzene is 5.50°C. What is the molal freezing point constant (°C/m) for benzene?
(a) 5.3 (b) 5.1 (c) 4.6 (d) 4.8

64. A solution containing 1.8 g of a compound (empirical formula CH_2O) in 40 g of water is observed to freeze at –0.465°C. The molecular formula of the compound is (K_f of water 1.86 kg K mol^{-1}) :
(a) $C_2H_4O_2$ (b) C_3H_6 (c) $C_4H_8O_4$ (d) $C_6H_{12}O_6$

65. Freezing point of the following equilibrium, liquid solvent $\rightleftharpoons$ solid solvent is :
(a) $\frac{H \quad G}{T \quad S}$ (b) $\frac{H}{S}$ (c) $\frac{G}{S}$ (d) $\frac{S}{H}$

66. Freezing point of a solution is smaller than freezing point of a solvent. It is due to :
(a) H of solution and solvent is almost identical since intermolecular forces between solvent molecules are involved
(b) S of solution (between solution and solid) is larger than that of the S of solvent (between solvent and solid)
(c) S of the solution is smaller than that of the solvent
(d) H of the solution is much higher than of solvent but S of solution is smaller than that of the solvent

67. When 36.0 g of a solute having the empirical formula CH_2O is dissolved in 1.20 kg of water, the solution freezes at –0.93°C. What is the molecular formula of the solute? (K_f 1.86°C kg mol^{-1})
(a) C_2H_4O (b) $C_2H_2O_2$ (c) $C_2H_4O_3$ (d) $C_2H_4O_2$

For More Material Join: @JEEAdvanced_2024

68. Calculate the molecular mass of a substance whose 7.0% by mass solution in water freezes at –0.93°C. The cryoscopic constant of water is 1.86°C kg mol ^{1}:
(a) 140 g mol ^{1} (b) 150.5 g mol ^{1}
(c) 160 g mol ^{1} (d) 155 g mol ^{1}

69. Camphor is often used in molecular mass determination because
(a) it is readily available (b) it has a very high cryoscopic constant
(c) it is volatile (d) it is solvent for organic substances

70. For 1 molal solution of each compound minimum freezing point will be assuming complete ionisation in each case:
(a) $[Fe(H_2O)_6]Cl_3$ (b) $[Fe(H_2O)_5Cl]Cl_2$ H_2O
(c) $[Fe(H_2O)_4Cl_2]Cl$ $2H_2O$ (d) $[Fe(H_2O)_3Cl_3]$ $3H_2O$

71. For 1 molal solution of each compound maximum freezing point will be assuming complete ionisation in each case:
(a) $[Fe(H_2O)_6]Cl_3$ (b) $[Fe(H_2O)_5Cl]Cl_2$ H_2O
(c) $[Fe(H_2O)_4Cl_2]Cl$ $2H_2O$ (d) $[Fe(H_2O)_3Cl_3]$ $3H_2O$

72. $PtCl_4$ $6H_2O$ can exist as a hydrated complex. Its 1 molal aq. solution has depression in freezing point of 3.72. Assume 100% ionisation and $K_f(H_2O)$ 1.86°C mol ^{1} kg, then complex is :
(a) $[Pt(H_2O)_6]Cl_4$ (b) $[Pt(H_2O)_4Cl_2]Cl_2$ $2H_2O$
(c) $[Pt(H_2O)_3Cl_3]Cl$ $3H_2O$ (d) $[Pt(H_2O)_2Cl_4]$ $4H_2O$

73. A complex is represented as $CoCl_3xNH_3$. Its 0.1 molal solution in water shows T_f 0.558 K. K_f for H_2O is 1.86 K molality ^{1}. Assuming 100% ionisation of complex and co-ordination number of Co is six, calculate formula of complex:
(a) $[Co(NH_3)_6]Cl_3$ (b) $[Co(NH_3)_5Cl]Cl_2$
(c) $[Co(NH_3)_4Cl_2]Cl$ (d) none of these

74. The freezing point of equimolal aqueous solutions will be highest for:
(a) $C_6H_5NH_3Cl$ (b) $Ca(NO_3)_2$ (c) $La(NO_3)_2$ (d) $C_6H_{12}O_6$

75. The freezing point of a 4% aqueous solution of '*A*' is equal to the freezing point of 10% aqueous solution of '*B*'. If the molecular mass of '*A*' is 60, then the molecular mass of '*B*' will be:
(a) 160 (b) 90 (c) 45 (d) 180

76. Depression in freezing point of 0.01 molal aqueous HCOOH solution is 0.02046. 1 molal aqueous urea solution freezes at –1.86°C. Assuming molality equal to molarity, pH of HCOOH solution is:
(a) 2 (b) 3 (c) 4 (d) 5

77. When mercuric iodide is added to the aqueous solution of KI, then the:
(a) freezing point is raised (b) freezing point is lowered
(c) freezing point does not change (d) boiling point does not change

78. Dimer of acetic acid in liquid benzene is in equilibrium with acetic acid monomer at certain temperature and pressure. If 25% of the dimer molecules are separated out then
(a) freezing point of the solution reduces (b) average molar mass of solute increases
(c) boiling point of solution increases (d) molar mass of solute decreases

For More Material Join: @JEEAdvanced_2024

79. The temperature of a city was found to be –9.3°C. A car was used, whose radiator was filled with 5 L of water. What minimum quantity of antifreezing agent ethylene glycol were added to water of radiator in order to use the car for travelling? (K_f of water 1.86 K kg mol ^{1})

(a) 3200 g (b) 1670 g (c) 1550 g (d) 2100 g

80. The cryoscopic constant of water is 1.86 K kg mol ^{1}. A 0.01 molal acetic acid solution produces a depression of 0.0194°C in the freezing point. The degree of dissociation of acetic acid is:

(a) zero (b) 0.043 (c) 0.43 (d) 1

81. If 1m solution of benzoic acid in benzene has a freezing point depression of 3.84°C. (K_f 5.12°C mol ^{1} kg) and boiling point elevation of 2.53°C (K_b 2.53°C mol ^{1} kg), then select the correct statement/s :

Statement I : there is dimer formation when undergoing freezing

Statement II : there is no change when undergoing boiling

Statement III : reverse of I and II

Statement IV : dimer formation in freezing and boiling state

(a) I, II (b) II, III (c) III, I (d) only I

82. In a 0.5 molal solution KCl, KCl is 50% dissociated. The freezing point of solution will be (K_f 1.86 K kg mol ^{1}):

(a) 274.674 K (b) 271.60 K (c) 273 K (d) none of these

83. A 1.0 g sample of $Co(NH_2CH_2CH_2NH_2)_3Cl_3$ is dissolved in 25.0 g of water and the freezing point of the solution is –0.87°C. How many ions are produced per mole of compound? The K_f of water is 1.86°C/molal

(a) 2 (b) 3 (c) 4 (d) 5

84. An aqueous solution contains 3% and 1.8% by mass. urea and glucose respectively. What is the freezing point of solution? (K_f 1.86 C/m)

(a) –1.172°C (b) –2.27°C (c) –1.5°C (d) none of these

85. Phenol associates in benzene to a certain extent in dimerisation reaction. A solution containing 0.02 kg of phenol in 1.0 kg of benzene has its freezing point depressed 0.69 K. Hence, degree of association of phenol dimerized will be :

[$K_f(C_6H_6)$ 5.12 K kg mol ^{1}]

(a) 0.63 (b) 0.73 (c) 0.83 (d) 0.93

86. Assuming complete ionisation, the solution having maximum freezing point will be:

(a) 1 *M* CaF_2 (b) 1.5 *M* $Al_2(SO_4)_3$ (c) 2 *M* NaCl (d) 1 *M* $AgNO_3$

87. In a 0.2 molal aqueous solution of a weak acid H*X* the degree of dissociation is 0.25. The freezing point of the solution will be nearest to: (K_f 1.86 K kg mol ^{1})

(a) –0.26°C (b) 0.465°C (c) –0.48°C (d) –0.465°C

88. An aqueous solution of 0.01 *M* KCl causes the same elevation in boiling point as an aqueous solution of urea. The concentration of urea solution is:

(a) 0.01 *M* (b) 0.005 *M* (c) 0.02 *M* (d) 0.04 *M*

89. When some NaCl was dissolved in water, the freezing point depression was numerically equal to twice the molal f.p. depression constant. The relative lowering of vapour pressure of the solution is nearly :

(a) 0.036 (b) 0.018 (c) 0.0585 (d) 0.072

For More Material Join: @JEEAdvanced_2024

90. Which one of the following statements is false ?
(a) The correct order of osmotic pressure for 0.01 *M* aqueous solution of each compound is $BaCl_2 > KCl > CH_3COOH >$ sucrose
(b) Isotonic solutions are those solutions which have the same osmotic pressure
(c) Raoult's law states that the vapour pressure of a component over a solution is proportional to its mole fraction in liquid state
(d) Two sucrose solutions of same molality prepared in different solvents will have the same freezing point depression

91. 0.1 molal aqueous solution of an electrolyte AB_3 is 90% ionised. The boiling point of the solution at 1 atm is : ($K_{b(H_2O)}$ 0.52 K kg mol ^{1})
(a) 273.19 K (b) 374.92 K (c) 376.4 K (d) 373.19 K

92. Which of the following aqueous solutions has osmotic pressure nearest to pure solvent?
(a) Na_2SO_4 (b) $BaCl_2$ (c) $Al_2(SO_4)_3$ (d) $C_{12}H_{22}O_{11}$

93. 0.1*M* NaCl and 0.05*M* $BaCl_2$ solutions are separated by a semi-permeable membrane in a container. For this system, choose the correct answer :
(a) There is no movement of any solution across the membrane
(b) Water flows from $BaCl_2$ solution towards NaCl solution
(c) Water flows from NaCl solution towards $BaCl_2$ solution
(d) Osmotic pressure of 0.1*M* NaCl is lower than the osmotic pressure of $BaCl_2$ (assume complete dissociation)

94. Two aqueous solutions, *A* and *B*, are separated by a semi-permeable membrane. The osmotic pressure of solution *A* immediately begins to decrease. Which of the following statement is true ?
(a) The solvent molecules are moving from the solution of higher osmotic pressure to that of lower osmotic pressure.
(b) The initial osmotic pressure of solution *B* is greater than that of solution *A*.
(c) Solvent molecules are moving from solution *B* into solution *A*.
(d) Both (a) and (b) are true statements.

95. Which one of the following pairs of solution can we expect to be isotonic at the same temperature?
(a) 0.1 *M* urea and 0.1 *M* NaCl (b) 0.1 *M* urea and 0.2 *M* $MgCl_2$
(c) 0.1 *M* NaCl and 0.1 *M* Na_2SO_4 (d) 0.1 *M* $Ca(NO_3)_2$ and 0.1 *M* Na_2SO_4

96. The empirical formula of a non-electrolyte is CH_2O. A solution containing 3 g L ^{1} of the compound exerts the same osmotic pressure as that of 0.05*M* glucose solution. The molecular formula of the compound is :
(a) CH_2O (b) $C_2H_4O_2$ (c) $C_4H_8O_4$ (d) $C_3H_6O_3$

97. A semipermeable membrane used in the measurement of osmotic pressure of a solution allows the passage of:
(a) solute molecules through it (b) solvent molecules through it
(c) both solvent and solute molecules (d) either solvent or solute

98. In the case of osmosis, solvent molecules move from :
(a) higher vapour pressure to lower vapour pressure
(b) higher concentration to lower concentration

For More Material Join: @JEEAdvanced_2024

(c) lower vapour pressure to higher vapour pressure
(d) higher osmotic pressure to lower osmotic pressure

99. The osmotic pressures of equimolar solutions of urea, $BaCl_2$ and $AlCl_3$ will be in the order :
(a) $AlCl_3 > BaCl_2 >$ urea (b) $BaCl_2 > AlCl_3 >$ urea
(c) urea $> BaCl_2 > AlCl_3$ (d) $BaCl_2 >$ urea $> AlCl_3$

100. Which of the following solutions will have maximum osmotic pressure? Assume 90% dissociation of each salt:
(a) decimolar aluminium sulphate
(b) decimolar barium chloride solution
(c) decimolar sodium sulphate solution
(d) solution of equal volumes of decimolar barium chloride and decimolar sodium sulphate solutions

101. Consider 0.1 *M* solutions of two solutes *X* and *Y*. The solute *X* behaves as a univalent electrolyte while the solute *Y* dimerises in solution. Which of the following statements are correct regarding these solutions?
(1) The boiling point of the solution of *X* will be higher than that of *Y*
(2) The osmotic pressure of the solution of *Y* will be lower than that of *X*
(3) The freezing point of the solution of *X* will be lower than that of *Y*
(4) The relative lowering of vapour pressure of both the solutions will be the same
Select the correct answer from the option given below :
(a) 1, 2 and 3 (b) 2, 3 and 4
(c) 1, 2 and 4 (d) 1, 3 and 4

102. If M_{normal} is the normal molecular mass and is the degree of ionization of $K_3[Fe(CN)_6]$, then the abnormal molecular mass of the complex in the solution will be:
(a) M_{normal} (1 2) ^{1} (b) M_{normal} (1 3) ^{1}
(c) M_{normal} (1) ^{1} (d) equal to M_{normal}

103. Equal volumes of 0.1 *M* urea and 0.1 *M* glucose are mixed. The mixture will have:
(a) lower osmotic pressure (b) same osmotic pressure
(c) higher osmotic pressure (d) none of these

104. A 5% (w/V) solution of cane sugar (molecular mass 342) is isotonic with 1% (w/V) solution of a substance *X*. The molecular mass of *X* is:
(a) 34.2 (b) 171.2 (c) 68.4 (d) 136.8

105. Insulin is dissolved in a suitable solvent and the osmotic pressure ( ) of solutions of various concentrations (g/cm^3) C is measured at 20°C. The slope of a plot of against C is found to be 4.65 10 ^{3}. The molecular mass of the insulin (g/mol) is:
(a) 3 10^5 (b) 9 10^5
(c) 4.5 10^5 (d) 5.16 10^6

106. An aqueous solution of sucrose ($C_{12}H_{22}O_{11}$) having a concentration of 34.2 gram/litre has an osmotic pressure of 2.38 atmospheres at 17°C. For an aqueous solution of glucose ($C_6H_{12}O_6$) to be isotonic with this solution, its concentration should be :
(a) 34.2 gram per litre (b) 17.1 gram per litre
(c) 18.0 gram per litre (d) 36.0 gram per litre

For More Material Join: @JEEAdvanced_2024

107. Which of the following experimental methods is adopted to determine osmotic pressure?
(a) Berkley-Hartely's method (b) Beckmann's method
(c) Landsberger's method (d) Differential method

108. Based upon the technique of reverse osmosis the approximate pressure required to desalinate sea water containing 2.5% (mass/volume) KNO_3 at 27°C will be
(a) 10.5 atm (b) 21 atm (c) 12.2 atm (d) 6.09 atm


109. A 1% (mass/vol) KCl solution is ionised to the extent of 80%. The osmotic pressure at 27°C of the solution will be:
(a) 6.95 atm (b) 5.94 atm
(c) 2.71 atm (d) 3.30 atm


110. Osmotic pressure of blood is 7.40 atm, at 27°C. Number of moles of glucose to be used per litre for an intravenous injection that is to have same osmotic pressure of blood is:
(a) 0.3 (b) 0.2 (c) 0.1 (d) 0.4


111. The relationship between osmotic pressures ($_1$, $_2$ and $_3$) at a definite temperature when 1 g glucose, 1 g urea and 1 g sucrose are dissolved in 1 litre of water is (assume i 1 for all):
(a) $_1$ $_2$ $_3$ (b) $_3$ $_1$ $_2$
(c) $_2$ $_1$ $_3$ (d) $_2$ $_3$ $_1$


112. van't Hoff proved that osmotic pressure ( ) is a colligative property. For an ideal solution, osmotic pressure ( ) is helpful to determine that molecular mass of solute using M_B $\frac{W_B RT}{V}$

Relation can be expressed by the curve (C concentration):

(a) 

(b) 

(c) 

(d) 

113. A solution containing 4.0 g of PVC in 2 litre of dioxane (industrial solvent) was found to have an osmotic pressure 3.0 10^{-4} atm at 27°C. The molar mass of the polymer (g/mol) will be :
(a) 1.6 10^4 (b) 1.6 10^5 (c) 1.6 10^3 (d) 1.6 10^2

114. The osmotic pressures of 0.010 M solutions of KI and of sucrose ($C_{12}H_{22}O_{11}$) are 0.432 atm and 0.24 atm respectively. The van't Hoff factor for KI is :
(a) 1.80 (b) 0.80 (c) 1.2 (d) 1.0

115. What is the correct sequence of osmotic pressure of 0.01 M *aq.* solution of :
(1) $Al_2(SO_4)_3$ (2) Na_3PO_4
(3) $BaCl_2$ (4) Glucose
(a) 4 2 3 1 (b) 3 4 2 1
(c) 3 4 1 2 (d) 1 2 3 4


For More Material Join: @JEEAdvanced_2024

116. 1.0 molar solution of the complex salt, $CrCl_3$ $6H_2O$, displays an osmotic pressure of $3RT$. 0.5 L of the same solution on treatment with excess of $AgNO_3$ solution will yield (assume 1) :
(a) 0.5 mole of AgCl (b) 1.0 mole of AgCl (c) 1.5 mole of AgCl (d) 3.0 mole of AgCl

117. A 0.010 g sample of $Cr(NH_3)_4(SO_4)Cl$ is dissolved in 25.0 mL of water and the osmotic pressure of the solution is 59.1 torr at 25°C. How many moles of ions are produced per mole of compound?
(a) 1 (b) 4 (c) 2 (d) 3


118. Which of the following aqueous solutions should have the highest osmotic pressure?
(a) 0.011 *M* $AlCl_3$ at 50°C (b) 0.03 *M* NaCl at 25°C
(c) 0.012 *M* $(NH_4)_2SO_4$ at 25°C (d) 0.03 *M* NaCl at 50°C


119. $X_3Y_2(i$ 5) when reacted with $A_2B_3(i$ 5) in aqueous solution gives brown colour. These are separated by a semipermeable membrane *AB* as shown. Due to osmosis there is:





(a) brown colour formation in side *X* (b) brown colour formation in side *Y*
(c) formation in both of the sides *X* and *Y* (d) no brown colour formation

120. Which of the following curves represents the Henry's law?

(a)


(b)


(c)


(d)


121. According to Henry's law, the solubility of a gas in a given volume of liquid increases with increase in:
(a) temperature (b) pressure
(c) Both (a) and (b) (d) None of these

122. At 300 K, 40 mL of $O_3(g)$ dissolves in 100 g of water at 1.0 atm. What mass of ozone dissolved in 400 g of water at a pressure of 4.0 atm at 300 K?
(a) 0.1 g (b) 1.24 g
(c) 0.48 g (d) 4.8 g

For More Material Join: @JEEAdvanced_2024

123. 1 kg of water under a nitrogen pressure of 1 atmosphere dissolves 0.02 gm of nitrogen at 293 K. Calculate Henry's law constant :

(a) 7.7 10^4 atm (b) 7.7 10^3 atm

(c) 2 10^5 atm (d) 2 10^2 atm

124. According to Henry's law, the partial pressure of gas (P_g) is directly proportional to mole fraction of gas in liquid solution, P_{gas} $K_H . X_{gas}$, where K_H is Henry's constant. Which is incorrect ?

(a) K_H is characteristic constant for a given gas-solvent system

(b) Higher is the value of K_H, lower is solubility of gas for a given partial pressure of gas

(c) K_H has temperature dependence

(d) K_H decreases with increase of temperature

125. At 760 torr pressure and 20°C temperature, 1 L of water dissolves 0.04 gm of pure oxygen or 0.02 gm of pure nitrogen. Assuming that dry air is composed of 20% oxygen and 80% nitrogen (by volume), the masses (in g/L) of oxygen and nitrogen dissolved by 1 L of water at 20°C exposed to air at a total pressure of 760 torr are respectively :

(a) 0.008, 0.016 (b) 0.016, 0.008

(c) 0.16, 0.08 (d) 0.04, 0.02

For More Material Join: @JEEAdvanced_2024

Level 2

1. Two beaker *A* and *B* present in a closed vessel. Beaker *A* contains 152.4 g aqueous solution of urea, containing 12 g of urea. Beaker *B* contains 196.2 g glucose solution, containing 18 g of glucose. Both solutions allowed to attain the equilibrium. Determine mass % of glucose in its solution at equilibrium :

(a) 6.71 (b) 14.49 (c) 16.94 (d) 20

2. Two components *A* and *B* form an ideal solution. The mole fractions of *A* and *B* in ideal solution are X_A and X_B, while that of in vapour phase, these components have their mole fractions as Y_A and Y_B. Then, the slope and intercept of plot of $\frac{1}{Y_A}$ *vs.* $\frac{1}{X_A}$ will be :

(a) $\frac{P_A}{P_B}, \frac{P_B \quad P_A}{P_B}$ (b) $\frac{P_B}{P_A}, \frac{P_A \quad P_B}{P_A}$ (c) $\frac{P_B}{P_A}, \frac{P_B}{P_B \quad P_A}$ (d) $P_A \quad P_B, \frac{P_A}{P_B}$

3. At 48°C, the vapour pressure of pure CS_2 is 850 torr. A solution of 2.0 g of sulphur in 100 g of CS_2 has a vapour pressure 844.9 torr. Determine the atomicity of sulphur molecule:

(a) 1 (b) 2 (c) 4 (d) 8


4. An ideal solution contains two volatile liquids *A* (P 100 torr) and *B* (P 200 torr). If mixture contain 1 mole of *A* and 4 mole of *B* then total vapour pressure of the distillate is:


(a) 150 (b) 180 (c) 188.88 (d) 198.88


5. The vapour pressure of two pure liquids *A* and *B*, that form an ideal solution are 100 and 900 torr respectively at temperature *T*. This liquid solution of *A* and *B* is composed of 1 mole of *A* and 1 mole of *B*. What will be the pressure, when 1 mole of mixture has been vaporized?


(a) 800 torr (b) 500 torr (c) 300 torr (d) None of these

6. Which of the following represents correctly the changes in thermodynamic properties during the formation of 1 mole of an ideal binary solution :

(a)


(b)


(c)


(d)


For More Material Join: @JEEAdvanced_2024

7. A saturated solution of XCl_3 has a vapour pressure 17.20 mm Hg at 20°C, while pure water vapour pressure is 17.25 mm Hg. Solubility product (K_{sp}) of XCl_3 at 20°C is:
(a) 9.8 10 ^{2} (b) 10 ^{5} (c) 2.56 10 ^{6} (d) 7 10 ^{5}

8. A certain non-volatile electrolyte contains 40% carbon, 6.7% hydrogen and 53.3% oxygen. An aqueous solution containing 5% by mass of the solute boils at 100.15°C. Determine the molecular formula of the compound (K_b 0.51°C/m):
(a) HCHO (b) CH_3OH (c) C_2H_5OH (d) $C_6H_{12}O_6$

9. A 0.10 *M* solution of a monoprotic acid (d 1.01 g/cm ^{3}) is 5% dissociated What is the freezing point of the solution? The mol. mass of the acid is 300 and $K_f(H_2O)$ 1.86°C/m:
(a) –0.189°C (b) –0.194°C (c) –0.199°C (d) None of these

10. An aqueous solution boils at 101°C. What is the freezing point of the same solution? (Given : K_f 1.86°C/m and K_b 0.51°C/m)
(a) 3.647°C (b) –3.647°C (c) –0.364°C (d) None of these

11. An industrial waste water is found to contain 8.2% Na_3PO_4 and 12% $MgSO_4$ by mass in solution. If % ionisation of Na_3PO_4 and $MgSO_4$ are 50 and 60 respectively then its normal boiling point is [$K_b(H_2O)$ 0.50 K kg mol ^{1}] :
(a) 102.3°C (b) 103.35°C (c) 101.785°C (d) None of these

12. Ratio of $\frac{T_b}{K_b}$ of 10 g AB_2 and 14 g A_2B per 100 g of solvent in their respective, solution (AB_2 and A_2B both are non-electrolytes) is 1 mol/kg in both cases. Hence, atomic wt. of *A* and *B* are respectively :
(a) 100, 40 (b) 60, 20 (c) 20, 60 (d) None of these

13. The freezing point of a solution containing 0.2 g of acetic acid in 20.0 g benzene is lowered by 0.45°C. Calculate the degree of dimerization of acetic acid in benzene. K_f for benzene is 5.12 K mol ^{1} kg :
(a) 0.527 (b) 0.80 (c) 0.945 (d) None of these

14. If boiling point of an aqueous solution is 100.1°C, what is its freezing point? Given, enthalpy of fusion and vaporisation of water are 80 cal g ^{1} and 540 cal g ^{1} respectively:
(a) 0.361°C (b) –0.361°C (c) –3.61°C (d) None of these

15. 100 g of $C_6H_{12}O_6$ (*aq*.) solution has vapour pressure is equal to 40 torr at certain temperature. Vapour pressure of $H_2O(l)$ is 40.18 torr at same temperature. If this solution is cooled to –0.93°C, what mass of ice will be separated out? (K_f 1.86 kg mol ^{1})
(a) 95.5 g (b) 4.5 g (c) 45.5 g (d) 47.8 g

16. 1.0 g of a monobasic acid H*A* in 100 g water lowers the freezing point by 0.155 K. If 0.75 g, of same acid requires 25 mL of *N*/5 NaOH solution for complete neutralisation then %, degree of ionization of acid is (K_f of H_2O 1.86 K kg mol ^{1}) :
(a) 20% (b) 25% (c) 40% (d) 50%

17. 0.1 *M* KI and 0.2 *M* $AgNO_3$ are mixed in 3 : 1 volume ratio. The depression of freezing point of the resulting solution will be [$K_f(H_2O)$ 1.86 K kg mol ^{1}] :
(a) 3.72 K (b) 1.86 K (c) 0.93 K (d) 0.279 K

For More Material Join: @JEEAdvanced_2024


18. If 0.1 M H_2SO_4 ($aq.$) solution shows freezing point –0.3906°C then what is the K_{a_2} for H_2SO_4?
(Assume m M and $K_{f(H_2O)}$ 1.86 K kg mol ^{1})
(a) 0.122 (b) 0.0122 (c) 1.11 10 ^{3} (d) None of these

19. A living cell contains a solution which is isotonic with 0.2 M glucose solution. What osmotic pressure develops when the cell is placed in 0.05 M $BaCl_2$ solution at 300 K?
(a) 1.23 atm (b) 3.69 atm (c) 6.15 atm (d) None of these

20. What is the osmotic pressure of 0.2 M HX ($aq.$) solution at 300 K?
(Given : $K_{a(HX)}$ 8 10 ^{5})
(a) 4.926 atm (b) 0.5024 atm (c) 5.024 atm (d) None of these

21. A solution contain 8 g of a carbohydrate in 100 g of water has a density 1.025 g/mL and an osmotic pressure of 5 atm at 27°C. What is the molar mass of the carbohydrate?
(a) 387 (b) 374 (c) 3740 (d) None of these

22. Study the following figure and choose the correct options. Assuming complete dissociation of electrolyte:



(a) There will be no net moment of any substance across the membrane
(b) $MgCl_2$ will flow towards the $Al_2(SO_4)_3$ solution
(c) $Al_2(SO_4)_3$ will flow towards the $MgCl_2$ solution
(d) The (osmotic pressure) of 0.1 M $MgCl_2$ is higher than the of 0.05 M $Al_2(SO_4)_3$

23. The total vapour pressure of a 4 mole % solution of NH_3 in water at 293 K is 50.0 torr. The vapour pressure of pure water is 17.0 torr at this temperature. Applying Henry's and Raoult's laws, calculate the total vapour pressure for a 5 mole % solution:
(a) 58.25 torr (b) 33 torr (c) 42.1 torr (d) 52.25 torr

24.-25. The vapour pressure of two pure liquids A and B which form an ideal solution are 500 and 800 torr respectively at 300 K. A liquid solution of A and B for which the mole fraction of A is 0.60 is contained in a cylinder closed by a piston on which the pressure can be varied. The solution is slowly vaporized at 300 K by decreasing the applied pressure.

24. The composition of vapour when first bubble formed is:
(a) y_A 0.6; y_B 0.4 (b) y_A 0.48; y_B 0.52
(c) y_A 0.52; y_B 0.48 (d) y_A 0.5; y_B 0.5

25. What is the composition of last droplet of liquid remaining in equilibrium with vapour?
(a) x_A 0.6; x_B 0.4 (b) x_A 0.5; x_B 0.5
(c) x_A 0.7; x_B 0.3 (d) x_A 0.3; x_B 0.7


For More Material Join: @JEEAdvanced_2024

Level 3

Passage 1

Lowering in vapour pressure is determined by Ostwald and Walker dynamic method. It is based on the principle, that when air is allowed to pass through a solvent or solution, it takes up solvent vapour with it to get itself saturated at that temperature

I and II are weighted separately before and after passing dry air. Loss in mass of each set, gives the lowering of vapour pressure. The temperature of air, the solution and the solvent is kept constant.



1. Loss in mass of solvent (w_{II}) will be proportional to:

(a) P P (b) P P (c) $\frac{P}{P}$ (d) P P

2. Gain in mass of anhydrous $CaCl_2$ is proportional to:

(a) P (b) P (c) P P (d) P P

3. $\frac{P \quad P}{P}$ is equal to:

(a) $\frac{w_I}{w_I \quad w_{II}}$ (b) $\frac{w_{II}}{w_I \quad w_{II}}$ (c) $\frac{w_I}{w_I \quad w_{II}}$ (d) $\frac{w_{II}}{w_I}$

4. Dry air was passed through 9.24 g of solute in 108 g of water and then through pure water. The loss in mass of solution was 3.2 g and that of pure water was 0.08 g. The molecular mass (g/mol) of solute is nearly :

(a) 50 (b) 62 (c) 70 (d) 80

Passage 2

A dilute solution contains 'x' moles of solute A in 1 kg of solvent with molal elevation constant K_b. The solute dimerises in the solution according to the following equation. The degree of association is ( ) :

$$2A \rightleftharpoons A_2$$

1. The van't Hoff factor will be:

(a) i 1 2 (b) i 1 $\frac{}{2}$ (c) i 1 $\frac{}{2}$ (d) i 1

2. The molecular mass observed will be:

(a) greater than actual molecular mass (b) lesser than actual molecular mass
(c) equal to the actual molecular mass (d) cannot be predicted by the data given

For More Material Join: @JEEAdvanced_2024

3. The degree of association is equal to:

(a) $\frac{(K_b x \quad T_b)}{T_b .2}$ (b) $\frac{2(K_b x \quad T_b)}{K_b x}$ (c) $2 \quad \frac{2 \quad T_b}{K_b x}$ (d) $\frac{T_b}{2K_b x}$

One or More Answers is/are Correct



1. Which of the following statement(s) is/are correct, if intermolecular forces in liquids *A, B* and *C* are in the order of $A \quad B \quad C$?

(a) *B* evaporates more readily than *A*
(b) *B* evaporates more readily than *C*
(c) *A* evaporates more readily than *C*
(d) all evaporate at same rate at same temperature.

2. When non-volatile solute is added to a pure solvent, the:

(a) vapour pressure of the solution becomes lower than the vapour pressure of the pure solvent
(b) rate of evaporation of solvent is reduced
(c) solute does not affect the rate of condensation
(d) none of these

3. The total vapour pressure of a binary solution is given by

$$P \quad (100 X_A \quad 260 X_B) \text{ mm Hg}$$

where, X_A and X_B are the mole fractions of components *A* and *B*. This indicates that the:

(a) vapour pressure of solution is less than the pure *B* component
(b) vapour pressure of solution is more than that of pure *A* component
(c) vapour pressure of pure *A* is 100 mm Hg and that of pure *B* is 260 mm Hg
(d) the vapour pressure of pure *A* and *B* are 260 mm Hg and 100 mm Hg respectively

4. Which of the following is correct for an ideal solution?

(a) $H_{mix} \quad 0$ and $V_{mix} \quad 0$ (b) $V_{mix} \quad 0$ and $S_{mix} \quad 0$
(c) $H_{mix} \quad 0$ and $S_{mix} \quad 0$ (d) $G_{mix} \quad 0$ and $S_{mix} \quad 0$

5. For a non-ideal solution with a negative deviation:

(a) H_{mix} ve (b) V_{mix} ve (c) S_{mix} ve (d) G_{mix} ve

6. A binary solution of liquids *A* and *B* will show positive deviation from Raoult's law if it fulfils the following condition:

(a) $P_A \quad X_A P_A$ and $P_B \quad X_B P_B$
(b) The intermolecular forces of $A \quad B \quad A \quad A, B \quad B$
(c) *H* mixing is positive
(d) *V* mixing is negative

7. Which of the following statements is/are correct about acetone and trichloromethane mixture?

(a) Mixtures of acetone and trichloromethane shows positive deviation from Raoult's law
(b) The forces of attraction acting between molecules of acetone and trichloromethane in a mixture are stronger than those acting between the molecules in pure acetone
(c) Pure acetone can be obtained by the careful fractional distillation of any mixture of acetone and trichloromethane
(d) When acetone and trichloromethane are mixed, the enthalpy change is negative

For More Material Join: @JEEAdvanced_2024

8. The azeotropic solutions of two miscible liquids:
(a) can be separated by simple distillation
(b) may show positive or negative deviation from Raoult's law
(c) are supersaturated solution
(d) behave like a single component and boil at a constant temperature

9. For exact determination of molecular mass through colligative properties measurement:
(a) solute must be volatile
(b) solution must be very dilute
(c) solution must be formed by similar nature of substances
(d) solute must not be dissociated or associated

10. For the depression in freezing point experiment, the correct statement(s) is/are:
(a) vapour pressure of pure solvent is more than that of solution
(b) vapour pressure of pure solvent is less than that of solution
(c) only solute molecules solidify at the freezing point
(d) only solvent molecules solidify at the freezing point

11. The cryoscopic constant value depends upon:
(a) the molar mass of the solute in the solution
(b) the molar mass of the solvent in the solution
(c) the enthalpy of fusion of the solvent
(d) the freezing point of the solvent

12. Consider 0.1 *M* solutions of two solutes *X* and *Y*. The solute *X* behaves as univalent electrolyte, while the solute *Y* dimerises in solution. Select correct statement(s) regarding these solutions:
(a) The boiling point of solution of '*X*' will be higher than that of '*Y*'
(b) The osmotic pressure of solution of '*Y*' will be lower than that of '*X*'
(c) The freezing point of solution of '*X*' will be lower than that of '*Y*'
(d) The relative lowering of vapour pressure of both the solution will be the same

13. Consider following solutions:
(I) I *M* glucose (aq) (II) 1 *M* sodium chloride (aq)
(III) 1 *M* acetic acid in benzene (IV) 1 *M* ammonium phosphate (aq)
(a) all are isotonic solutions (b) III is hypotonic of I, II, IV
(c) I, II, IV are hypertonic of III (d) IV is hypertonic of I, II, III

14. Which of the following statements is (are) incorrect?
(a) 0.1 *M* KCl solution will have the same osmotic pressure as 0.1 *M* glucose solution
(b) 0.1 *M* KCl solution will have the same boiling point as 0.1 *M* urea solution
(c) 0.1 *m* glucose and 0.1 *m* urea are isotonic
(d) 0.1 *m* $MgCl_2$ solution will have less relative lowering of vapour pressure than 0.1 *m* NaCl

15. Consider following solutions:
0.1 *m* $C_6H_5NH_3^+Cl$; 0.1 *m* KCl; 0.1 m Glucose; 0.1 *m* $Na_2C_2O_4\ 10H_2O$
(a) the solution with highest boiling point is 0.1 *m* $Na_2C_2O_4\ 10H_2O$
(b) the solution with highest freezing point is 0.1 *m* glucose
(c) 0.1 *m* $C_6H_5NH_3Cl$ and 0.1 *m* NaCl will have the same osmotic pressure
(d) 0.1 *m* glucose solution will have the lowest osmotic pressure

For More Material Join: @JEEAdvanced_2024

Match the Column



Column-I and **Column-II** contains four entries each. Entries of column-I are to be matched with some entries of column-II. One or more than one entries of column-I may have the matching with the same entries of column-II.

1.

Column-I	Column-II
(A) *n*-hexane + *n*-heptane	(P) Can be separated by fractional distillation
(B) Acetone + chloroform	(Q) Maximum boiling azeotrope
(C) Chlorobenzene and bromobenzene	(R) Cannot be separated by fractional distillation completely
(D) Ethanol + water	(S) Minimum boiling azeotrope

2.

Column-I	Column-II
(A) $_1$: 0.1 *M* glucose; $_2$: 0.1 *M* urea	(P) $_1$ and $_2$ are isotonic
(B) $_1$: 0.1 *M* NaCl; $_2$: 0.1 *M* Na_2SO_4	(Q) No net migration of solvent across the membrane
(C) $_1$: 0.1 *M* NaCl; $_2$: 0.1 *M* KCl	(R) $_1$ is hypertonic to $_2$
(D) $_1$: 0.1 *M* $CuSO_4$; $_2$: 0.1 *M* sucrose	(S) $_1$ is hypotonic to $_2$

3.

Column-I	Column-II
(A) 0.1 *M* $Al_2(SO_4)_3$	(P) Solution with highest boiling point
(B) 0.1 *M* $AlPO_4$	(Q) van't Hoff factor is greater than 1
(C) 0.1 *M* urea	(R) Solution with lowest osmotic pressure
(D) 0.1 *M* $MgCl_2$	(S) Solution with lowest freezing point

4.

Column-I (Solute)	Column-II (Van't Hoff factor, *i*)
(A) $AlCl_3$ if 0.8	(P) *i* 3.4
(B) $BaCl_2$ if 0.9	(Q) *i* 2.8
(C) Na_3PO_4 if 0.9	(R) *i* 3.8
(D) $K_4[Fe(CN)_6]$ if 0.7	(S) *i* 3.7

5.

Column-I	Column-II
(A) Elevation of B.P.	(P) Colligative property
(B) Osmotic pressure	(Q) Ebullioscopic constant
(C) Relative lowering in V.P.	(R) Berkeley-Heartley method
(D) Depression of F.P.	(S) Ostwald and Walker method

For More Material Join: @JEEAdvanced_2024

Assertion-Reason Type Questions



Each question contains STATEMENT-1 (Assertion) and STATEMENT-2 (Reason).

Examine the statement carefully and mark the correct answer according to the instructions given below:

(A) If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1

(B) If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1

(C) If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

(D) If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

1. STATEMENT-1 : An increase in surface area increases the rate of evaporation.
STATEMENT-2 : Stronger the intermolecular attraction forces, faster is the rate of evaporation at a given temperature.

2. STATEMENT-1 : An ideal solution obeys Raoult's law.
STATEMENT-2 : In an ideal solution, solute-solute as well as solvent-solvent, interactions are similar to solute-solvent interactions.

3. STATEMENT-1 : If a liquid solute more volatile than the solvent is added to the solvent, the vapour pressure of the solution is greater than vapour pressure of pure solvent.
STATEMENT-2 : Vapour pressure of solution is equal to vapour pressure of solvent.

4. STATEMENT-1 : V_{mix} and S_{mix} for an ideal solution is zero.
STATEMENT-2 : *A*...*B* interaction in an ideal solution are same as between *A*... *A* and *B*...*B*.

5. STATEMENT-1 : Elevation in boiling point will be high if the molal elevation constant of the liquid is high.
STATEMENT-2 : Elevation in boiling point is a colligative property.

6. STATEMENT-1 : The boiling point of 0.1 *M* urea solution is less than that of 0.1 *M* KCl solution.
STATEMENT-2 : Elevation of boiling point is directly proportional to the number of moles of non-volatile solute particles present in the solution.

7. STATEMENT-1 : The observed molar mass of acetic acid in benzene is more than the normal molar mass of acetic acid.
STATEMENT-2 : Molecules of acetic acid dimerise in benzene due to hydrogen bonding.

8. STATEMENT-1 : Addition of ethylene glycol to water lowers the freezing point of water, therefore, used as antifreeze substance.
STATEMENT-2 : Ethylene glycol is soluble in water.

9. STATEMENT-1 : Osmotic pressure is a colligative property.
STATEMENT-2 : Osmotic pressure is developed in a column due to osmosis.

10. STATEMENT-1 : Osmosis involves movement of solvent molecules from lower concentration to higher concentration.
STATEMENT-2 : Solutions having the same osmotic pressure are called isotonic solutions.

For More Material Join: @JEEAdvanced_2024

11. STATEMENT-1 : Isotonic solutions must have the same molar concentration.
STATEMENT-2 : Solutions which have the same osmotic pressure are known as isotonic solution.

12. STATEMENT-1 : Isotonic solutions do not show phenomenon of osmosis.
STATEMENT-2 : Isotonic solutions have same molar concentration at same temperature.

13. STATEMENT-1 : When dried fruits and vegetables are placed in water, they slowly get swollen.
STATEMENT-2 : It happens due to the phenomenon of osmosis.

14. STATEMENT-1 : Reverse osmosis is used to purify sea water.
STATEMENT-2 : Solvent molecules pass from concentrate solution to pure solvent through semipermeable membrane if high pressure ( ) is applied on solution side.

15. STATEMENT-1 : All solutes becomes more soluble in water at higher temperature.
STATEMENT-2 : Solubility of solute depends upon temperature.

16. STATEMENT-1 : Henry's law is always applicable for gases.
STATEMENT-2 : Raoult's law is a special case of Henry's law.

17. STATEMENT-1 : Increasing pressure on pure water decrease its freezing point.
STATEMENT-2 : Density of water is maximum at 273 K.

18. STATEMENT-1 : The molecular mass of acetic acid determined by depression in freezing point method in benzene and water was found to be different.
STATEMENT-2 : Water is polar and benzene is non-polar.

19. STATEMENT-1 : If red blood cells were removed from the body and placed in pure water, pressure inside the cell increases.
STATEMENT-2 : The concentration of the salt content in the cells increases.

20. STATEMENT-1 : Azeotrope is a binary mixture formed by ideal solutions.
STATEMENT-2 : Azeotrope boils with unchanged composition.

Subjective Problems



1. The vapour pressure of two pure liquids A and B are 5 and 10 torr respectively. Calculate the total pressure of the solution (in torr) obtained by mixing 2 mole of A and 3 mole of B.

2. The vapour pressure of two pure liquids A and B are 50 and 40 torr respectively. If 8 moles of A is mixed with x moles of B , then vapour pressure of solution obtained is 48 torr. What is the value of x.

3. The vapour pressure of a liquid solution containing A and B is 99 torr. Calculate mole % of B in vapour phase.
(Given : P_A 100 torr ; P_B 80 torr)

4. If 30 g of a solute of molecular mass 154 is dissolved in 250 g of benzene. What will be the elevation in boiling point of the resulting solution ?
(Given : $K_{b(C_6H_6)}$ 2.6 K kg mol ^{1})

For More Material Join: @JEEAdvanced_2024

5. Calculate elevation in boiling point for 2 molal aqueous solution of glucose.
(Given : $K_{b(H_2O)}$ 0.5 K kg mol ^{1})

6. Calculate depression of freezing point for 0.56 molal aq. solution of KCl.
(Given : $K_{f(H_2O)}$ 1.8 K kg mol ^{1}).

7. What is the maximum value of van't Hoff factor for $AlCl_3$?

8. A solution containing 500 g of a protein per litre is isotonic with a solution containing 3.42 g sucrose per litre. The molecular mass of protein in 5 10^{x}, hence x is.

9. An aqueous solution of urea has a freezing point of 0.515 C. Predict the osmotic pressure (in atm) of the same solution at 37 C.
(Given : $K_{f(H_2O)}$ 1.86 K kg mol ^{1})

10. 0.2 M aq. solution of KCl is isotonic with 0.2 M K_2SO_4 at same temperature.
What is the van't Hoff factor of K_2SO_4 ?

For More Material Join: @JEEAdvanced_2024

Answers

Level 1

1.	(d)	**2.**	(b)	**3.**	(a)	**4.**	(c)	**5.**	(b)	**6.**	(c)	**7.**	(a)	**8.**	(b)	**9.**	(d)	**10.**	(b)
11.	(d)	**12.**	(c)	**13.**	(c)	**14.**	(c)	**15.**	(b)	**16.**	(b)	**17.**	(c)	**18.**	(a)	**19.**	(a)	**20.**	(a)
21.	(a)	**22.**	(d)	**23.**	(a)	**24.**	(d)	**25.**	(d)	**26.**	(a)	**27.**	(c)	**28.**	(b)	**29.**	(a)	**30.**	(a)
31.	(b)	**32.**	(d)	**33.**	(b)	**34.**	(d)	**35.**	(d)	**36.**	(c)	**37.**	(b)	**38.**	(d)	**39.**	(c)	**40.**	(d)
41.	(a)	**42.**	(a)	**43.**	(c)	**44.**	(a)	**45.**	(c)	**46.**	(a)	**47.**	(d)	**48.**	(c)	**49.**	(c)	**50.**	(a)
51.	(d)	**52.**	(d)	**53.**	(c)	**54.**	(d)	**55.**	(b)	**56.**	(c)	**57.**	(a)	**58.**	(b)	**59.**	(d)	**60.**	(b)
61.	(b)	**62.**	(d)	**63.**	(a)	**64.**	(d)	**65.**	(b)	**66.**	(b)	**67.**	(d)	**68.**	(b)	**69.**	(b)	**70.**	(a)
71.	(d)	**72.**	(c)	**73.**	(b)	**74.**	(d)	**75.**	(a)	**76.**	(b)	**77.**	(a)	**78.**	(b)	**79.**	(c)	**80.**	(b)
81.	(a)	**82.**	(b)	**83.**	(c)	**84.**	(a)	**85.**	(b)	**86.**	(d)	**87.**	(d)	**88.**	(c)	**89.**	(a)	**90.**	(d)
91.	(d)	**92.**	(d)	**93.**	(b)	**94.**	(c)	**95.**	(d)	**96.**	(b)	**97.**	(b)	**98.**	(a)	**99.**	(a)	**100.**	(a)
101.	(a)	**102.**	(b)	**103.**	(b)	**104.**	(c)	**105.**	(d)	**106.**	(c)	**107.**	(a)	**108.**	(c)	**109.**	(b)	**110.**	(a)
111.	(c)	**112.**	(a)	**113.**	(b)	**114.**	(a)	**115.**	(d)	**116.**	(b)	**117.**	(c)	**118.**	(d)	**119.**	(d)	**120.**	(a)
121.	(b)	**122.**	(b)	**123.**	(a)	**124.**	(d)	**125.**	(a)										

Level 2

1.	(b)	**2.**	(b)	**3.**	(d)	**4.**	(c)	**5.**	(c)	**6.**	(c)	**7.**	(d)	**8.**	(d)	**9.**	(c)	**10.**	(b)
11.	(c)	**12.**	(b)	**13.**	(c)	**14.**	(b)	**15.**	(d)	**16.**	(b)	**17.**	(d)	**18.**	(b)	**19.**	(a)	**20.**	(c)
21.	(b)	**22.**	(d)	**23.**	(a)	**24.**	(b)	**25.**	(c)										

For More Material Join: @JEEAdvanced_2024



Level 3

Passage–1:	**1.** (a)	**2.** (b)	**3.** (b)	**4.** (b)			
Passage–2:	**1.** (b)	**2.** (a)	**3.** (b)				

One or More Answers is/are Correct

1.	(b,c)	**2.**	(a,b,c)	**3.**	(a,b,c)	**4.**	(a,b,d)	**5.**	(a,b,d)	**6.**	(a,b,c)	**7.**	(b,d)	**8.**	(b,d)
9.	(b,d)	**10.**	(a,d)	**11.**	(b,c,d)	**12.**	(a,b,c)	**13.**	(b,c,d)	**14.**	(a,b,d)	**15.**	(a,b,c,d)		

Match the Column

1.	A	P;	B	Q, R;	C	P;	D	R, S
2.	A	P, Q;	B	S;	C	P, Q;	D	R
3.	A	P, Q, S;	B	Q;	C	R;	D	Q
4.	A	P;	B	Q;	C	S;	D	R
5.	A	P, Q;	B	P, R;	C	P, S;	D	P

Assertion-Reason Type Questions

1.	(C)	**2.**	(A)	**3.**	(C)	**4.**	(D)	**5.**	(B)	**6.**	(A)	**7.**	(A)	**8.**	(B)	**9.**	(B)	**10.**	(B)
11.	(D)	**12.**	(A)	**13.**	(A)	**14.**	(B)	**15.**	(D)	**16.**	(D)	**17.**	(C)	**18.**	(A)	**19.**	(C)	**20.**	(D)

Subjective Problems

1.	(8	**2.**	2	**3.**	4	**4.**	2	**5.**	1	**6.**	2	**7.**	4	**8.**	4	**9.**	7	**10.**	2

For More Material Join: @JEEAdvanced_2024

Hints and Solutions

Level 1

1. (d) $x(l) \overset{Eq^m}{\rightleftharpoons} x(g)$, K_p Vp of liq, $_rH$ ve

T , Vp

2. (b) Given P_1 19 mm Hg, P_2 760 mm Hg;

$H_{vap.}$ 40670 J/mol

Applying Clausius-Clapeyron's equation

$$\log \frac{P_2}{P_1} \quad \frac{H_{vap}}{2.303 \quad R} \quad \frac{T_2 \quad T_1}{T_1 T_2}$$

$$\text{or} \quad \log \frac{760}{19} \quad \frac{40670}{2.303 \quad 8.3} \quad \frac{373 \quad T_1}{T_1 \quad 373}$$

on solving, we get T_1 291.4 K

3. (a) Initial mass of $H_2O(l)$ 18 g;

$$n_{H_2O}(g) \quad \frac{\frac{24.63}{760} \quad 7.6}{0.0821 \quad 300} \quad 0.01$$

% wt. of H_2O vaporised $\frac{0.01 \quad 18}{18}$ 100

1%

4. (c) Raoult's law is applicable for ideal solution.

5. (b) $P \quad P_A^\circ x_A \quad P_B^\circ x_B \quad P_B^\circ \quad x_A (P_A^\circ \quad P_B^\circ)$

$[\because x_B \quad 1 \quad x_A]$

6. (c) $y_A \quad \frac{P_A^\circ \quad x_A}{P_A^\circ x_A \quad P_B^\circ x_B} \quad \frac{x_A}{x_A \quad x_B \quad \frac{P_B^\circ}{P_A^\circ}}$

$[\because P_A^\circ \quad P_B^\circ]$

$$\frac{y_A}{x_A} \quad \frac{1}{x_A \quad x_B \quad \frac{P_B^\circ}{P_A^\circ}} \quad 1 \qquad y_A \quad x_A$$

…(1)

Similarly, $y_B \quad \frac{P_B^\circ \quad x_B}{P_A^\circ x_A \quad P_B^\circ x_B} \quad \frac{x_B}{x_B \quad x_A \quad \frac{P_A^\circ}{P_B^\circ}}$

…(2)

$$\frac{y_B}{y_A} \quad \frac{x_B}{x_A}$$

$$\frac{y_B}{y_B} \quad \frac{x_B}{x_B}$$

7. (a) $y_A \quad \frac{P_A^\circ x_A}{P_A^\circ x_A \quad P_B^\circ x_B} \quad \frac{P_A^\circ x_A}{P_B^\circ \quad (P_A^\circ \quad P_B^\circ) x_A}$

$$\frac{1}{y_A} \quad \frac{P_B^\circ x_B \quad P_A^\circ x_A}{P_A^\circ x_A} \quad 1 \quad \frac{P_B^\circ}{P_A^\circ} \quad \frac{1 \quad x_A}{x_A}$$

$$1 \quad \frac{P_B^\circ}{P_A^\circ} \quad \frac{1}{x_A} \quad \frac{P_B^\circ}{P_A^\circ}$$

Slope $\frac{P_B^\circ}{P_A^\circ}$

8. (b) $P_{Solution} \quad P_{Solution}^\circ X_{Solvent}$

$$\frac{P \quad P}{P} \quad X_{Solute}$$

9. (d) Solution X is unsaturated so v.p. will be more, solution Y and Z are saturated so v.p. of Y v.p. of Z and 2 gm of solute would be present in form of solid in system Z.

10. (b) Highest vapour pressure for lowest boiling point.

11. (d) $\frac{P \quad P}{P} \quad x_{solute}$ or $\frac{P \quad P}{P} \quad \frac{n}{n \quad N}$

$$\frac{0.1}{0.1 \quad 9.9} \qquad 0.01; \quad P \quad 0.99\,P$$

12. (c) 80 P (0.8) P 100 torr.

13. (c) Given P_A 750 mm Hg

$\because$ 373 K is boiling point of water.

Thus, P_A 760 mm Hg

$$m \quad \frac{P \quad P}{P} \quad \frac{1000}{M_{solvent}}$$

$$\frac{10}{750} \quad \frac{1000}{18} \qquad 0.74$$

14. (c) Normal boiling point of water is 100°C, hence

P (H_2O) 760 torr; $M(H_2O)$ 18 g/mol

$$m \quad \frac{P \quad P}{P} \quad \frac{1000}{M_{H_2O}}$$

P 746.5; P 13.45 torr

15. (b) $P_s \quad \frac{80}{100} \quad P$, w ?,

M_{solute} 40, w 114 g, $M_{solvent}$ 114

For More Material Join: @JEEAdvanced_2024

$\because \quad \frac{P - P_s}{P_s} = \frac{w}{M_{solute}} \frac{M_{solvent}}{W}$

or $\frac{P - (80/100)P}{(80/100)P} = \frac{w}{40} \frac{114}{57}$;

$w = 5$ g

16. (b) $\frac{P - P}{P} = \frac{n}{n + N} \frac{n}{N}$

(for very dilute solution)

$2 = \frac{n}{N_A} \frac{n}{N_B}$

$M_B = 2M_A$

17. (c) We know that

$Y_A = \frac{P_A X_A}{P_{total}}$ or $\frac{Y_A}{X_A} = \frac{P_A}{P_{total}}$

$\because \quad P_A < P_{total}$ so $\frac{Y_A}{X_A} < 1$ or $Y_A < X_A$

18. (a) Moles of $A = \frac{20}{40} = 0.5$;

moles of $B = \frac{20}{80} = 0.25$

$x_A = \frac{0.5}{0.5 + 0.25} = 0.67$; $x_B = 0.33$

$P_{total} = P_A X_A + P_B X_B$

$100 = 0.67 \times 40 + 0.33 \times P_B = 80$ torr

19. (a) Since, $P = X_A P_A + X_B P_B$, we have

$\frac{1}{1+3} P_A + \frac{3}{1+3} P_B = 550$ mm Hg;

$\frac{1}{1+4} P_A + \frac{4}{1+4} P_B = 560$ mm Hg

That is, $0.25 P_A + 0.75 P_B = 550$ mm Hg;

$0.20 P_A + 0.8 P_B = 560$ mm Hg

Solving for P_A and P_B, we get;

$P_A = 400$ mm Hg and $P_B = 600$ mm Hg

20. (a) $y_A = \frac{P_A}{P} = \frac{P_A x_A}{P}$ and $y_B = \frac{P_B x_B}{P}$;

$\frac{y_A}{y_B} = \frac{P_A}{P_B} \frac{x_A}{x_B} = \frac{4}{3} \frac{1}{3} \frac{x_A}{(1 - x_A)}$

$x_A = \frac{4}{5}$ or $x_B = \frac{1}{5}$


21. (a) $x_A = \frac{1}{1+3} = \frac{1}{4}$; $x_B = \frac{3}{4}$;

$\frac{y_A}{y_B} = \frac{P_A}{P_B} \frac{x_A}{x_B}$

$\frac{y_A}{(1 - y_A)} = \frac{1}{3} \frac{1}{3}$

$\frac{1}{9}$ or $y_A = \frac{1}{10}$

22. (d)



Plots AD show vapour pressure of B.

Plots BC show vapour pressure of A.

Plot CD show vapour pressure of liquid solution containing A and B volatile components.

$P = P_A + P_B$

$EH = EF + EG$

23. (a) $\because \quad (T_b)_{Solvent} < (T_b)_{Solution}$;

$S_{Solv.} < S_{Solution}$

$(\Delta H_{Solution}) > (\Delta H)_{Solvent}$

[Only solvent participate in vaporisation]

[Due to elevation in BP of solution]

24. (d) $(\Delta_f S)_{Solution} > (\Delta_f S)_{Solvent}$

[Since solution has more randomness than solvent]

$(\Delta_f H)_{Solution} \approx (\Delta_f H)_{Solvent}$

[Similar forces of intermolecular forces are involved in solution and solvent]

Due to presence of non-volatile solute (Sugar), freezing point of solution decrease.

25. (d) $(\Delta H_{Vap})_{Solvent} \approx (\Delta H_{Vap})_{Solution}$

[Because similar intermolecular forces are involved in both cases].

$\because \quad (T_b)_{Solvent} < (T_b)_{Solution}$

$(\Delta S_{Vap})_{Solution} > (\Delta S_{Vap})_{Solvent}$

26. (a) Elevation in boiling point $\propto$ concentration of a solution. Thus, the order of concentration of solutions is I < II < III.

For More Material Join: @JEEAdvanced_2024

27. (c) T_B° T_A° [From diagram] B Residue
A Distillate

28. (b) The mixture will show positive deviation from Raoult's law hence boiling point of an azeotropic mixture of water-ethanol is less than that of both water and ethanol.

29. (a) H_{Solution} H_1 H_2 H_3

30. (a) $P_{\text{Ideal solution}}$ $\frac{1}{3}$ 150 $\frac{2}{3}$ 300 250

$\because$ P_{Solution} $P_{\text{Ideal solution}}$; ( ) ve deviation.

31. (b) When A B interaction is weaker than A A and B B interaction, then mixture of A and B show positive deviation.

32. (d) On changing external pressure, composition of azeotrope alters.

33. (b) After mixing HNO_3 and H_2O, intermolecular force of attraction increase.

34. (d) $\because$

$P_{\text{Ideal solution}}$ $\frac{1}{4}$ 100 80 $\frac{3}{4}$ 85 mmHg

P_{Solution} (90 mm Hg)

Solution show positive deviation from Raoult's law.

35. (d) From an azeotropic mixture the components cannot be separated at boiling point because it is a constant temperature boiling mixture.

36. (c) See solution of Q. 35.

37. (b) Show negative deviation from Raoult's law.

38. (d) P P_A° P_B° 300 torr.

39. (c) For two immiscible liquid;

P_A P_{total} P_{H_2O} 748 648 100

$\frac{W_A}{W_B}$ $\frac{P_A M_A}{P_B M_B}$; M_A $\frac{1.25}{1}$ $\frac{648}{100}$ $\frac{18}{}$

145.8

40. (d) $\frac{W_A}{W_B}$ $\frac{P_A}{P_B}$ $\frac{M_A}{M_B}$ $\frac{0.7}{7}$ $\frac{112.5}{18}$

0.625

$\%$ $\frac{W_A}{W_A \quad W_B}$ 100 $\frac{0.625}{1.625}$ 100

38.46

41. (a) Lowering in vapour pressure is a colligative property.

42. (a) i 1 $(n$ 1) so n $\frac{i \quad 1}{}$

43. (c) i $\frac{C(1 \quad) \quad \frac{C}{n}}{C}$ i 1 $\frac{}{n}$

44. (a) For dissociation i 1 $(n$ 1) 1

For dissociation i 1 $\frac{1}{n}$ 1 1

45. (c) For high v.p. concentration of solute should be low and temperature should be high.

46. (a) On addition of strong electrolyte, vapour pressure decrease. On addition of water, vapour pressure increases due to increase in X_{Solvent}.

47. (d) Colligative property moles of non-volatile solute.

48. (c) $\frac{P \quad P}{P}$ $\frac{10}{50}$ $\frac{n}{n \quad 12}$ n 3

(Number of ions of solute), moles of K_2SO_4 1

49. (c) mi $\frac{P \quad P}{P}$ $\frac{1000}{M_{\text{solvent}}}$;

7 m $\frac{(20.0126 \quad 20) \quad 1000}{20 \quad 18}$;

m 5 $10^{\ 3}$

50. (a) T_b 100.5 100 K_b 1 K_b 0.5

51. (d) T_b $K_b.m$ 3.63 $\frac{0.616/154}{15}$ 1000;

T_b 61.7 0.968

62.67°C

52. (d) T_b 80.26, T_b 0.16;

0.16 2.53 $\frac{0.26 / M}{11.20}$ 1000; M 367

that is almost molar mass of $C_{20}H_{16}Fe_2$.

53. (c) 80.23 80.10 2.53 $\frac{0.64/x}{100}$ 10^3

x 125

54. (d) T_b (normal) $K_b m$ 2.53 1 2.53 K;

i $\frac{T_{b(\text{obs})}}{T_{b(\text{nor})}}$ $\frac{1.518}{2.53}$ 0.6

i 1 1 $\frac{1}{n}$; 0.8

55. (b) $H_{\text{vap.}}$ $\frac{M_{\text{solvent}} \quad R \quad T_{b(\text{solvent})}^{\circ 2}}{1000 \quad K_b}$;

$H_{\text{vap.}}$ $\frac{92 \quad 8.314 \quad (383.7)^2}{1000 \quad 3.32}$

33.91 kJ/mol 34 kJ/mol

For More Material Join: @JEEAdvanced_2024

56. (c) B.P. moles of non-volatile solute.

57. (a)

	XY_2	$\rightleftharpoons$	X^2	$2Y$
Initially	1		0	0
at equilibrium	1			2

Total no. of moles 1 2 1 2

i $\frac{\text{Normal molar mass}}{\text{Observed molar mass}}$; $\frac{1\ 2}{1}$ $\frac{164}{65.6}$

0.75; % 75%

58. (b) T_b K_b m

m $\frac{y}{x}$ ...(1)

T_f $K_f m$ ...(2)

T_f $\frac{y}{x}$ z

59. (d) Liquid solution $\rightleftharpoons$ solid solvent.

60. (b) T_f 2.304; m $\frac{2.304}{14.4}$ 0.16;

m $\frac{n_{\text{solute}}}{W_{\text{solvent}}}$ 1000

0.16 $\frac{2.6}{M}$ $\frac{1000}{100}$; M 162.5

61. (b) $n_{C_4H_{10}}$ $\frac{10.44}{58}$ 0.18;

m $\frac{0.18}{200}$ 1000 0.9

T_f 5.12 0.9 4.608;

T_f 5.5 4.608 0.892°C

62. (d) 4 F 20 C; T_f 20 C

T_f $K_f.m$ m $\frac{20}{1.86}$ 10.75

mass of C_2H_5OH required

10.75 46 494.5 g

63. (a) 4.4 5.5 K_f $\frac{2.4/154}{75}$ 10^3 K_f 5.3

64. (d) 0.465 1.86 $\frac{1.8/x}{40}$ 10^3

x 180 g/mol $C_6H_{12}O_6$

65. (b) T_f $\frac{H}{S}$

66. (b) $\because$ $\frac{H_f}{S_f}_{\text{Solution}}$ $\frac{H_f}{S_f}_{\text{Solvent}}$

67. (d) T_f $K_f.m$ 0.93 1.86 $\frac{36}{M}$ $\frac{1}{1.2}$,

M 60 and 30 n 60; n 2

so compound is $C_2H_4O_2$.

68. (b) 0.93 1.86 $\frac{7/x}{93/1000}$

x 150.5 g mol ^{1}.

69. (b) It has a very high cryoscopic constant.

70. (a) Maximum ions minimum T_f.

71. (d) Maximum T_f for minimum ions.

72. (c) 3.72 1.86 $[i\,m]$

i 2 $[Pt(H_2O)_3Cl_3]Cl$ $3H_2O$

73. (b) T_f K_f molality i;

0.558 1.86 0.1 i or i 3

Thus complex is $[CoCl.xNH_3].Cl_2$. Since co-ordination number of CO is six, thus complex is $[CO(NH_3)_5Cl]Cl_2$.

74. (d) Depression in freezing point, T_f iK_fm. The value of van't Hoff factor (i) is minimum for the glucose, which is a non-electrolyte. Hence, aqueous solution of glucose has highest freezing point.

75. (a) Molality of solution should be same m_1 m_2;

$\frac{4}{60}$ $\frac{1000}{96}$ $\frac{10}{M_B}$ $\frac{1000}{90}$ or M_B 160

76. (b) T_f $K_f.m$ 1.86 K_f 1

$K_f(H_2O)$ 1.86

for HCOOH; 0.02046 1.86 0.01(1);

0.1; so [H] 10^{3}; pH 3

77. (a) The reaction when KI is added in HgI_2

$HgI_2 + 2KI$ $K_2HgI_4 \rightleftharpoons 2K$ HgI_4^{2}

Since, number of ions decreased, so freezing point increases.

78. (b) T_f mK_fi

T_b mK_bi

M_{obs} $\frac{M}{i}$

i , So M_{obs} increases

79. (c) To use car; T_f T_f T_f 9.3 K

T_f $K_f m$ 9.3 1.86 m m 5

number of moles of ethylene glycol added to 5 L of water of radiator 5 5 25

or 25 62 1550 g

80. (b) Given K_1 1.86 K kg mol ^{1}, m 0.01,

T_f 0.0194 C

T_f $i.K_fm$; $\frac{0.0194}{1.86\ \ 0.01}$ i

or i 1.043 or 0.043

81. (a) 3.84 i 5.12 1

i 0.75 1

For More Material Join: @JEEAdvanced_2024

$2.53 \quad i \quad 2.53 \quad 1$

$i \quad 1$

82. (b) $T_f \quad (1 \quad 0.5) \quad 1.86 \quad 0.5 \quad 1.395$

$T_f \quad 273 \quad 1.395 \quad 271.60$

83. (c) $T_f \quad K_f.m.i;$

$0.87 \quad 1.86 \quad i \quad \frac{1}{345.5} \quad \frac{1000}{25}$; $i \quad 4$

84. (a) $T_f \quad K_f \quad \frac{n_1 \quad n_2}{W_{\text{solvent}}} \quad 1000$;

$$T_f \quad 1.86 \quad \frac{\frac{3}{60} \quad \frac{1.8}{180}}{95.2} \quad 1000 \qquad 1.172$$

$T_f \quad T_f \quad 1.172$ C; $T_f \quad 1.172$ C

85. (b) $T_f \quad K_f.m.i \qquad i \quad 0.633;$

$i \quad 1 \qquad \frac{}{2} \qquad 0.73$

86. (d) Maximum freezing point for minimum ion.

87. (d) $i \quad 1 \qquad 1.25$

$T_f \quad K_f.m.i \qquad 1.86 \quad 0.2 \quad 1.25$

or $T_f \quad 0.465$ or $T_f \quad 0.465$ C

88. (c) $0.01 \quad K_b \quad 2 \quad K_b \quad x \quad 1$

$x \quad 0.02 \, M$

89. (a) $\frac{P^o \quad P}{P^o} \quad \frac{in}{N} \quad \frac{2 \quad m \quad M_{\text{Solution}}}{10^3}$

$\frac{2 \quad 1 \quad 18}{1000} \quad \frac{9}{250} \quad 0.036$

$T_f \quad k_f(m), \frac{T_f}{K_f} \quad 2 \quad im, m \quad 1, i \quad 2$

90. (d) Depression in freezing point

$K_f m$ (as K_f , T_f)

K_f depends upon the solvent.

91. (d) $T \quad 100 \quad 0.1(1 \quad 3 \quad 0.9) \quad 0.52$

$T \quad 373.19$ K

92. (d) No. of ions for 1 mole solute.

93. (b) Water (solvent) flows from lower concentration to higher concentration.

94. (c) Solvent molecules are moving from solution *B* into solution *A* hence, osmotic pressure of solution *A* immediately begins to decrease.

95. (d) Solutions having same no. of ions are isotonic

Urea ($i \quad 1$), $NaCl(i \quad 2)$, $MgCl_2(i \quad 3)$, $Na_2SO_4(i \quad 3)$, $Ca(NO_3)_2(i \quad 3)$.

96. (b) Glucose Non-electrolyte

$0.05 \quad RT \quad \frac{3}{x} \, RT$

$x \quad 60$, $n \quad \frac{MFM}{EFM} \quad \frac{60}{30} \quad 2$

$M.F. \quad (E.F.)_2$, $C_2H_4O_2$

97. (b) A semipermeable membrane allow the passage of only solvent molecules.

98. (a) In osmosis, solvent molecules move from lower concentration (higher vapour pressure) to higher concentration (lower vapour pressure)

99. (a) i [Equal concentration]

$AlCl_3(i \quad 4)$, Urea $(i \quad 1)$, $BaCl_2(i \quad 3)$

100. (a) i [Equal concentration]

Aluminium sulphate $Al_2(SO_4)_3$ $(i \quad 5)$, Barium chloride $BaCl_2(i \quad 3)$,

Sodium sulphate $Na_2SO_4(i \quad 3)$.

101. (a) For $x(i \quad 2)$ and $Y(i \quad 1)$

$(T_b)_x \quad (T_b)_y$; $_x \quad _y$; $(T_f)_x \quad (T_f)_y$

102. (b) $K_3[Fe(CN)_6] \rightleftharpoons 3K^+ + [Fe(CN)_6]^{3-}$

At $t \quad 0 \quad 1 \qquad 0 \qquad 0$

After ionization $(1 \quad) \qquad 3 \qquad ; i \quad 1 \quad 3$

Since, $i \quad \frac{M_{\text{normal}}}{M_{\text{abnormal}}}$;

$\frac{1 \quad 3}{1} \quad \frac{M_{\text{normal}}}{M_{\text{abnormal}}}$


103. (b) $\because$ [Urea] [Glucose],

Isotonic solutions

104. (c) $_1 \quad _2$; $\frac{5}{342} \quad \frac{1000}{100} \quad \frac{1}{X} \quad \frac{1000}{100}$

$X \quad 68.4$

105. (d)



$\frac{W}{M} \quad \frac{RT}{V}$ or $\frac{}{C} \quad \frac{RT}{M}$

slope $\frac{RT}{M}$

or $M \quad \frac{0.082 \quad 293}{4.65 \quad 10^{-3} \quad 10^{-3}}$

$5.16 \quad 10^6$ g

For More Material Join: @JEEAdvanced_2024

106. (c) $\frac{34.2}{342}\ RT \quad x\ RT$

Concentration of glucose $0.1 \quad 180$ $18\ g/L$

107. (a) Berkley-Hartely's method is adopted to determine osmotic pressure.

108. (c) $P \quad \frac{2.5}{39+14+48} \quad \frac{1000}{100} \quad 2 \quad R \quad 300$

12.2 atm.

109. (b) $i.C.RT$

$1.8 \quad \frac{1}{74.5} \quad \frac{1}{0.1} \quad 24.63$

5.94

110. (a) $7.4 \quad n \quad 0.0821 \quad 300$

$n \quad 0.3$

111. (c) $iCRT$; Where, $i \quad 1$ for each of the solution

$\because \quad \frac{1}{\text{molecular mass of solute}}$;

$\overset{2}{\text{urea}}\ (60) \quad \overset{1}{\text{glucose}}\ (180) \quad \overset{3}{\text{sucrose}}\ (342)$

112. (a) CRT

$T\ [C \quad \text{const.}]$

$C\ [T \quad \text{const.}]$

113. (b) $M_B \quad \frac{W_B \quad R \quad T}{V} \quad \frac{4 \quad 0.0821 \quad 300}{3 \quad 10^{-4} \quad 2}$

$1.6 \quad 10^5$

114. (a) $i \quad \frac{\text{Number of particles in KI solution}}{\text{Number of particles in sucrose solution}}$

$\frac{(\text{KI solution})}{(\text{sucrose solution})} \quad \frac{0.432\ \text{atm}}{0.24\ \text{atm}} \quad 1.80$

115. (d) $iCRT$; $i \quad 5$ for $Al_2(SO_4)_3$,

$i \quad 4$ for Na_3PO_4,

$i \quad 3$ for $BaCl_2$, $i \quad 1$ for glucose

So osmotic pressure of :

Glucose $< BaCl_2 < Na_3PO_4 < Al_2(SO_4)_3$

116. (b) $iCRT, i \quad 3 \quad 1 \quad (n \quad 1)$;

For the salt 1, hence $n \quad 3$

$[Cr(H_2O)_5Cl]Cl_2\ H_2O + 2AgNO_3$

$[Cr(H_2O)_5Cl]^{2+} \quad 2AgCl(s)$

0.5 L of 1m salt $0.5 \quad 2 \quad 1.0$ mole of AgCl

117. (c) $\frac{59.1}{760} \quad \frac{0.01}{(52 \quad 4 \quad 17 \quad 96 \quad 35.5)}$

$\frac{1000}{25} \quad 0.0821 \quad 298\,(n)$

$n \quad 2$

118. (d) iCT; iCT is maximum for 0.03 M NaCl at 50 C.

119. (d) Only solvent molecules can passed through SPM so only dilution is possible.

120. (a) Henry's law is $m \quad K \quad P$; where, m mass of gas absorbed by given volume of the solvent.

P pressure of gas;

$\log m \quad \log K \quad \log P$

121. (b) $W_{Gas} \quad P_{Gas}$ at constant T.

122. (b) $\because\ n_{O_3}$ dissolved in 100 g water at 1.0 atm

$\frac{40 \quad 1 \quad 10^{-3}}{0.0821 \quad 300}$

n_{O_3} dissolved in 400 g water at 4 atm

$\frac{4 \quad 4 \quad 40 \quad 10^{-3}}{24.63}$

$W_{O_3} \quad \frac{640 \quad 10^{-3}}{24.63} \quad 48 \quad 1.2\ g$

123. (a) $P_{N_2} \quad (K_H)_{N_2} X_{Gas}$

$1 \quad (K_H)_{N_2} \quad \frac{0.02/28}{\frac{0.02}{28} \quad \frac{1000}{18}}$

$(K_H)_{N_2} \quad 7.7 \quad 10^4$ atm

124. (d) $P_{Gas} \quad \frac{1}{K} \quad n_{Solvent} \quad MM_{Gas} \quad X_{Gas} \quad K_H X_{Gas}$

$\because K_H \quad \frac{n_{Solvent} \quad MM_{Gas}}{K}$

As T , K_H .

125. (a) Mass of oxygen $\frac{20}{100} \quad 0.04 \quad 0.008$ g/L

Mass of nitrogen $\frac{80}{100} \quad 0.02 \quad 0.016$ g/L

Level 2

1. (b) Mole fraction of urea in its solution

$\frac{\frac{12}{60}}{\frac{12}{60} \quad \frac{140.4}{18}} \quad 0.025$

Mole fraction of glucose

For More Material Join: @JEEAdvanced_2024

$$\frac{\frac{18}{180}}{\frac{18}{180}+\frac{178.2}{18}} \quad 0.01$$

$\because$ Mole fraction of glucose is less so vapour pressure above the glucose solution will be higher than the pressure above urea solution, so some H_2O molecules will transfer from glucose to urea side in order to make the solutions of equal mole fraction to attain equilibrium, let x moles H_2O transferred

$$\frac{0.2}{0.2 \quad 7.8 \quad x} \quad \frac{0.1}{0.1 \quad 9.9 \quad x} \qquad x \quad 4$$

now mass of glucose solution

$196.2 \quad 4 \quad 18 \qquad 124.2$

wt. % of glucose $\frac{18}{124.2} \quad 100 \qquad 14.49$

2. (b) $\because \quad P_A \quad P_A X_A,$

$P_B \quad P_B X_B$

(P_A and P_B v.p. of pure A and B)

$$Y_A \quad \frac{P_A}{P_A \quad P_B}$$

$$\frac{P_A X_A}{P_A X_A \quad P_B (1 \quad X_A)}$$

$$Y_A \quad \frac{P_A X_A}{X_A (P_A \quad P_B) \quad P_B}$$

$$\frac{1}{Y_A} \quad \frac{P_A \quad P_B}{P_A} \quad \frac{P_B}{P_A} \quad \frac{1}{X_A}$$

So, slope is $\frac{P_B}{P_A}$ and intercept $\frac{P_A \quad P_B}{P_A}$

3. (d) $\frac{P \quad P}{P} \quad \frac{n}{N} \quad \frac{W}{M \quad N}$

$$\frac{850 \quad 844.9}{844.9} \quad \frac{2 \quad 76}{M \quad 100}$$

$M \quad 252; \quad n \quad \frac{252}{32} \quad 8$

atomicity of sulphur is 8

4. (c) $P \quad P_A x_A \quad P_B x_B \qquad 100 \quad \frac{1}{5} \quad 200 \quad \frac{4}{5}$

180 torr

y_A(composition of A vapour phase) $\frac{P_A x_A}{p}$

$\frac{20}{180} \qquad \frac{1}{9}, \quad Y_B \quad \frac{8}{9}$

For condensation $y_A \quad x_A^1; \quad y_B \quad x_B^1$

$P_{\text{total}} \quad P_A x_A^1 \quad P_B x_B^1;$

$P_{\text{total}} \quad 100 \quad \frac{1}{9} \quad \frac{8}{9} \quad 200 \qquad 188.88$ torr

5. (c) Let n_B mole of B present in 1 mole of mixture that has been vaporized. Thus, $y_B \quad \frac{n_B}{1}$

Mole fraction of B in the remaining liquid phase will be $x_B \quad \frac{1 \quad n_B}{1}$

$$x_B \quad \frac{P \quad P_T}{P_B \quad P_T} \qquad \ldots(1)$$

$[\because P \quad P_T \quad (P_B \quad P_T) x_B]$

and $y_B \quad \frac{P_B}{P} \qquad \frac{P_B x_B}{P} \qquad \ldots(2)$

After substitution of values of x_B and y_B in (1) and (2)

we get $1 \quad n_B \quad \frac{P \quad P_T}{P_B \quad P_T} \qquad \ldots(3)$

and $n_B \quad \frac{(1 \quad n_B) P_B}{P} \qquad \ldots(4)$

or $n_B \quad \frac{P_B}{P \quad P_B}$

so $1 \quad \frac{P_B}{P \quad P_B} \quad \frac{P \quad P_T}{P_B \quad P_T}$

$P \quad \sqrt{P_B \quad P_T} \quad \sqrt{100 \quad 900}$

300 torr

6. (c) For an ideal solution $H_{\text{mix}} \quad 0$ and S_{mix} is always positive so G_{mix} is negative.

7. (d) $m i \quad \frac{P \quad P}{P} \quad \frac{1000}{M_{\text{solvent}}};$

$$4.m \quad \frac{(17.25 \quad 17.20) \quad 1000}{17.2 \quad 18}; \quad m \quad 0.04$$

for dilute solution molality molarity or S;

$K_{sp} \qquad 27 S^4 \quad \sim \quad 7 \quad 10^{-5}$

8. (d) Mole ratio of C : H : O is 1 : 2 : 1 so empirical formula is CH_2O

$m \quad \frac{T_b}{K_b} \qquad \frac{0.15}{0.51} \qquad 0.294;$

$0.294 \quad \frac{50}{M} \quad \frac{1000}{950}; \quad M \quad 180$

$(CH_2O)_n \quad 180$ or $30 \quad n \quad 180$ or $n \quad 6;$

molecular formula is $C_6H_{12}O_6$.

For More Material Join: @JEEAdvanced_2024

9. (c) Mass of 1 litre of solution 1010 g;
mass of solvent 1010 300 0.1 980 g
$m \quad \frac{0.1}{0.98} \quad 0.102;$

$T_f \quad K_f.m.i \quad (1 \quad) K_f m$
$T_f \quad 1.05 \quad 1.86 \quad 0.102 \quad 0.199 \quad C;$
$T_f \quad 0 \quad 0.199 \quad 0.199 \quad C$

10. (b) $m \quad \frac{T_b}{K_b} \quad \frac{1}{0.51} \quad \frac{T_f}{K_f}$

$\frac{1}{0.51} \quad \frac{T_f}{1.86}$

$T_f \quad 3.647$ or $T_f \quad 3.647 \quad C$

11. (c) For Na_3PO_4, $i \quad 1 \quad 3 \quad 1 \quad 3 \quad 0.5$
$= 2.5$; for $MgSO_4$, $i \quad 1.6$
100 g solution contains 8.2 g Na_3PO_4 and 12 g $MgSO_4$

$T_b \quad K_b.m.i =$

$$K_b \frac{\text{effective no. of moles of } (Na_3PO_4 + MgSO_4)}{\text{mass of solvent (in g)}} 1000$$

$$T_b \quad 0.50 \frac{\frac{8.2}{164} \quad 2.5 \quad \frac{12}{120} \quad 1.6}{79.8} \quad 1000$$

$1.785°C$

$T_b \quad 100 \quad 1.785 \quad 101.785°C$

12. (b) $T_b \quad K_b \frac{w_{solute}}{W_{solvent}} \frac{1}{M_{solute}} \quad 1000$

$1 \quad \frac{1000 \quad 10}{100 \quad M_{AB_2}} \quad M_{AB_2} \quad 100$

similarly $M_{A_2B} \quad 140$
$100 \quad M_A \quad 2 \cdot M_B$
and $140 \quad 2 . M_A \quad M_B$;
$M_A \quad 60$ and $M_B \quad 20$

13. (c) Given $w \quad 0.2$ g, $W \quad 20$ g, $T \quad 0.45$;

$$T_f \quad \frac{1000 \quad K_f \quad w}{M \quad W}$$

or $0.45 \quad \frac{1000 \quad 5.12 \quad 0.2}{20 \quad M}$

M (observed) 113.78

Now for $2CH_3COOH \rightleftharpoons (CH_3COOH)_2$
Before association 1 0
After association 1 /2;
where is the degree of association.

$\frac{M_{normal}}{M_{observed}} \quad 1 \quad \frac{}{2};$

or $\frac{60}{113.78} \quad 1 \quad \frac{}{2}; \quad 0.945$

14. (b) Given $H_{fus.} \quad 80$ cal g ^{1},
$H_{vap.} \quad 540$ cal g ^{1}
We know, $T_b \quad K_b \quad m$
and $T_f \quad K_f \quad m;$
Also $K \quad \frac{RT^2}{1000 \quad H}$

$\frac{T_b}{T_f} \quad \frac{K_b}{K_f}$

$\frac{RT_b^2}{1000 \quad H_{vap.}} \quad \frac{1000 \quad H_{fus.}}{RT_f^2}$

$\frac{T_b}{T_f} \quad \frac{T_b^2}{T_f^2} \quad \frac{H_{fus.}}{H_{vap.}}$

$\frac{0.1}{T_f} \quad \frac{373}{273} \quad \frac{373}{273} \quad \frac{80}{540}$

$\because \quad T_f \quad 0.361$ so,
$T_f \quad 0.361 \quad C$

15. (d) Molality of solution $\frac{P \quad P}{P} \quad \frac{1000}{M}$

$\frac{40.18 \quad 40}{40} \quad \frac{1000}{18} \quad 0.25$

1000 g water present with 45 g glucose or 100 g solution has 4.31 g glucose and 95.69 g H_2O. Final molality is 0.5, 1000 g solvent contain 90 g glucose or 4.31 g glucose present with $\frac{1000}{90} \quad 4.31 \quad 47.88$ g H_2O

Mass of ice formed = 95.69 – 47.88 = 47.8 g

16. (b) $T_f \quad K_f \quad \frac{w_{acid}}{M_{acid} \quad W_{H_2O}} \quad 1000;$

M_{acid} (observed) 120

Normal molecular mass of acid can be calculated as

milli-equivalents of acid milli-equivalents of base

(for HA and NaOH, $N \quad M$)

$\frac{0.75}{M_{acid}} \quad 1000 \quad 25 \quad \frac{1}{5}$

$M_{acid\ (Normal)} \quad 150$

$i \quad \frac{150}{120} \quad 1.25$

$i \quad 1 \quad ; \quad 0.25$

$\% \quad 25$

For More Material Join: @JEEAdvanced_2024

17. (d) $KI(aq)$ $AgNO_3(aq)$ $AgI(s)$ $KNO_3(aq)$

	KI +	$AgNO_3$	KNO_3 +	AgI
Initial :	0.3 mole	0.2 mole	—	—
Final :	0.1 mole	—	0.2	0.2

Total moles of solute ions $(0.1 + 0.2)$ 2 ($\because$ i 2)

[Solute] $\frac{0.6}{4}$ M

T_f K_f $\frac{0.6}{4}$

1.86 $\frac{0.6}{4} = 0.279$ K

18. (b) T_f $K_f.m.i$ 0.3906 i 1.86 0.1, i 2.1

first step of H_2SO_4 is strong so

H_2SO_4 H^+ HSO_4;

0 C $C(1$ $)$

$HSO_4 \rightleftharpoons H$ SO_4^{2}

$C(1$ $)$ C C

i $\frac{C(1 \quad) \; C \quad C \quad C}{C}$ 2.1;

0.1

Now, $HSO_4 \rightleftharpoons H$ SO_4^{2}

$C(1$ $)$ $(C$ $C)$ C

K_{a_2} $\frac{C(1 \quad) \; C}{C(1 \quad)}$

$\frac{1.1 \quad 0.1 \quad 0.1}{0.9}$ 0.0122

19. (a) $(iC_1$ $i_2C_2)\,RT$

(1 0.2 3 0.05) 0.0821 300

1.23 atm

20. (c) $\sqrt{\frac{8 \quad 10^{-5}}{0.2}}$ 0.02;

i (1 1.02

$i.C.R.T$

1.02 0.2 0.0821 300

5.024 atm

21. (b) 8 g of carbohydrate is present with 100 g of water in solution

or 108 g solution contains 8 g of carbohydrate;

$V_{solution}$ $\frac{108}{1.025}$ 10^{-3} litre

$\frac{n}{V}$ R T

$\frac{8}{M}$ $\frac{1.025}{108}$ 1000 0.0821 300 5;

M 374 g/mol

22. (d) Only solvent molecule can pass through SPM.

Osmotic pressure iC (at constant T).

23. (a) The given data are

P_{water} 17.0 torr;

P_{total} (4 mole % solution)

P_{NH_3} P_{water} 50.0 torr

x_{NH_3} 0.04 and x_{water} 0.96

Now according to Raoult's law;

P_{water} x_{water} P_{water}

0.96 17.0 torr 16.32 torr

Now Henry's law constant for ammonia is

$K_H(NH_3)$ $\frac{P_{NH_3}}{x_{NH_3}}$ $\frac{33.68 \text{ torr}}{0.04}$ 842 torr

Hence, for 5 mole % solution, we have

P_{NH_3} $K_H(NH_3)\,x_{NH_3}$

(842 torr) (0.05) 42.1 torr

P_{water} $P_{water}\,x_{water}$

(17 torr) (0.95) 16.15 torr

Thus, P_{total} (5 mole % solution)

P_{NH_3} P_{water} 42.1 16.15 58.25 torr

24. (b) P $P_A x_A$ $P_B x_B$


500 0.6 800 0.4

620 torr

y_A $\frac{P_A}{P}$ $\frac{300}{620}$

0.48; y_B 0.52

25. (c) P_A 500; P_B 800 torr



When most of the liquid has vaporized x_A 0.6 (given) would be y_A 0.6

For More Material Join: @JEEAdvanced_2024

$$y_A \quad \frac{P_A x_A}{P_A x_A \quad P_B (1 \quad x_A)}$$

$$0.6 \quad \frac{500. x_A}{500 x_A \quad 800 (1 \quad x_A)}$$

$$x_A \quad 0.70; \quad x_B \quad 0.30$$

Level 3

Passage-1

1. (a) Loss of mass of solvent lowering in vp $(p \quad P)$.

2. (b) Gain in mass of anhydrous $CaCl_2$ P .

3. (b) $\frac{P \quad P}{P} \quad \frac{w_{II}}{w_I \quad w_{II}} \quad \because p \quad w_I \quad w_{II} \quad P \quad P \quad w_{II}$

4. (b) $\frac{w_{II}}{w_I \quad w_{II}} \quad \frac{9.24 / x}{\frac{9.24}{x} \quad \frac{108}{8}} \quad \frac{0.08}{3.28}$

$x \quad 62$

Passage-2

1. (b) $2A \rightleftharpoons A_2 \quad ; \quad n \quad 2$

$i \quad 1 \quad \frac{1}{n} \quad 1 \quad 1 \quad \frac{}{2}$

2. (a) $i \quad \frac{M_T}{M} \quad 1$

3. (b) $T_b \quad 1 \quad \frac{1}{2} \quad 1 \quad K_b x$

$$\frac{2 (K_b x \quad T_b)}{K_b x}$$

Subjective Problems

1. $P \quad \frac{2}{5} \quad 5 \quad \frac{3}{5} \quad 10 \quad 8$ torr

2. $48 \quad \frac{8}{8 \quad x} \quad 50 \quad \frac{x}{x \quad 8} \quad 40$

$x \quad 2$

3. $P \quad P_A^\circ \quad x_B (P_B^\circ \quad P_A^\circ)$

$99 \quad 100 \quad x_b (\quad 20) \quad x_B \quad \frac{1}{20}$

$y_B \quad \frac{80 \quad \frac{1}{20}}{99} \quad \frac{4}{99}$

% mole_B in vapour phase $\frac{4}{99} \quad 100 \quad 4$

4. $T_b \quad \frac{2.6 \quad 30 \quad 1000}{154 \quad 250} \quad 2$

5. $T_b \quad 0.5 \quad 2 \quad 1$

6. $T_f \quad 2 \quad 0.56 \quad 1.8$

7. For 100% ionization $i \quad 4$

8. For isotonic solution

$\frac{\overset{\text{Protein}}{500}}{x} \quad \frac{\overset{\text{Sucrose}}{3.42}}{342} \quad x \quad 5 \quad 10^4$

9. $T_f \quad 0.515 \quad 1.86 \quad m$


$m \quad 0.276$

$0.276 \quad 0.0821 \quad 310 \quad 7$ atm

10. $\overset{KCl}{0.2} \quad 2 \quad \overset{K_2SO_4}{0.2} \quad x$

$x \quad 2$

For More Material Join: @JEEAdvanced_2024



10

SOLID STATE

Types of Solids

S. No.	Crystalline solids	Amorphous solids
1.	They have a definite crystal shape.	They do not have a definite geometrical shape
2.	Long-range order	Short-range order
3.	They have sharp melting points	They do not have sharp melting points.
4.	Enthalpy of fusion is definite	Enthalpy of fusion is not definite
5.	They are hard and rigid solids.	They are comparatively soft and not very rigid.
6.	Possess definite crystal system (cubic, tetragonal, hexagonal, octahedral, etc.)	Crystal system absent ; no regular repeating unit
7.	They are true solid	They are pseudo solid
8.	They split into two pieces when cut with a sharp-edged tool ; newly generated surfaces are plane and smooth.	They cut into two pieces with irregular surfaces when cut with a sharp-edged tool.
9.	They are anisotropic (refractive index and electrical conductivity are different in different directions) (*e. g.*, NaCl, CsBr, CaF_2 and ZnS	They are isotropic. (physical properties are same in all directions) (*e. g.*, Rubber, plastic and glass)

For More Material Join: @JEEAdvanced_2024

Types of Crystalline Solids

Type of solid	Intermolecular forces	Properties	Examples
Ionic	Ion-ion forces	Brittle, hard high melting	NaCl, KCl, $MgCl_2$
Molecular	Dispersion forces/Dipole-Dipole H-bond	Soft, low melting non-conducting	H_2O, Br_2, CO_2, CH_4
Covalent network	Covalent bonds	Hard: High melting	C-Diamond, SiO_2
Metallic	Metallic bonds	Variable hardness and melting point conducting	Na, Zn, Cu, Fe

Crystal System

Crystal Systems		Bravais Lattice	Unit Cell Parameters	
			Intercepts	Crystal Angles
1.	Cubic	Primitive, face centered, body centered	$a \quad b \quad c$	90
2.	Orthorhombic	Primitive, face centered, body centered, end centered	$a \quad b \quad c$	90
3.	Rhombohedral	Primitive	$a \quad b \quad c$	90
4.	Monoclinic	Primitive, end centered	$a \quad b \quad c$	90 , 90
5.	Triclinic	Primitive	$a \quad b \quad c$	90
6.	Tetragonal	Primitive, body centered	$a \quad b \quad c$	90
7.	Hexagonal	Primitive	$a \quad b \quad c$	90 , 120

Some Important Formula

(i) $d \quad \frac{ZM}{N_0 a^3}$; (ii) Packing fraction $\quad \frac{Z \quad V_{\text{Atom}}}{V_{\text{Unit cell}}}$; (iii) Void fraction 1 Packing fraction ;

(iv) C.No. No. of first neighbours

where d density ; M Atomic mass of element ;

N_A Avogadro's number ; a Edge length of cube.

Cubic System

S.No.	Property	Simple Cubic Lattice	BCC Lattice	FCC Lattice
1.	Atomic radius (r), a edge length of cube	$r \quad \frac{a}{2}$	$r \quad \frac{a\sqrt{3}}{4}$	$r \quad \frac{a\sqrt{2}}{4}$
2.	No of atoms per unit cell (Z)	$Z \quad 1$	$Z \quad 2$	$Z \quad 4$
3.	Co-ordination No. (C.No.)	6	8	12
4.	Packing fraction (P.F.)	0.52	0.68	0.74

For More Material Join: @JEEAdvanced_2024

Types of Packing

S.No.	Property	Hexagonal close packing (HCP)	Cubic close packing (CCP)
1.	Co-ordination No. (C.No.)	12	12
2.	No. of atoms per unit cell (z)	z 6	z 4
3.	Packing fraction (P.F.)	P.F. 0.74	P.F. 0.74
4.	Type of packing	*ABAB*	*ABCABC* .....

Types of Voids

S. No.	Name of Void	r_{Void}/r_{Sphere}	Co-ordination Number (C.No.)
1.	Triangular void	0.155	3
2.	Tetrahedral void	0.225	4
3.	Octahedral void	0.414	6
4.	Cubic void	0.732	8

Radius Ratio

S.No.	Limiting radius ratio $\frac{r_{cation}}{r_{anion}}$	C.No.	Shape
1.	0.155	2	Linear
2.	0.155 — 0.225	3	Planar triangle
3.	0.225 — 0.414	4	Tetrahedral
4.	0.414 — 0.732	6	Octahedral
5.	0.732 — 0.999	8	Like body centred cubic

Types of Ionic Structure

S.No.	Name of Structure	Location of Particle
1.	Rock salt (*AB*) NaCl	*B* : ccp lattice ; *A* : Octahedral void
2.	Zinc blende (ZnS)	S^{2} : ccp lattice ; Zn^{2} : Alternate tetrahedral void
3.	CsCl	Cs : Cube centre ; Cl : Corner of cube
4.	Fluorite structure (CaF_2)	Ca^{2} : fcc lattice ; F : Tetrahedral void
5.	Antifluorite structure Na_2O	O^{2} : fcc lattice ; Na : Tetrahedral void

For More Material Join: @JEEAdvanced_2024


Bragg's Equation

$2d\sin \quad n$

n 1 First order reflection ; n 2 Second order reflection ; angle of reflection.

d distance between planes ; wavelength of x-ray

Defects in Crystals



(1) Stoichiometric defects

- **Vacancy:** When constituent particle missing, then vacancy defects occurs.
- **Frenkel defect :** The defect in which an ion is displaced from its regular position to an interstitial position creating a vacancy. Such defect is known as Frenkel defects or dislocation defect.
- **Schottky defect :** Defect in which a pair of one cation and one anion of equal valence is missing from an ionic crystal leading to a pair of vacant sites. Such defect is known as Schottky defect.

(2) Non-Stoichiometric defects :

- **Metal excess defect :** The defect occurs due to anionic vacancies or due to the presence of extra cations at the interstitial site.
- **Metal deficiency defect :** The defect occurs due to absence of positive ions from lattice site or extra interstitial negative ions.

Electrical Properties

The solids can be **conductors, insulators** or **semiconductors.**

- Conductors conduct electricity through movement of electrons (metals) or ions (electrolytes).
- The atomic orbitals of metals form molecular orbitals which are so close in energy that they form bands. If the gap between the filled valence band the next higher unoccupied conduction band overlaps, the electrons can jump and are called conductors. If this gap is more, the electrons can not jump and the substance behaves as an insulator.
- In semiconductors, this gap is small. For example, silicon and germanium. Their conductivity can be increased by adding impurity (doping) which can be electron rich or electron deficient.
- When increase in conductivity is due to the electron-rich impurity, it is called n-type semiconductor.
- When the increase in conductivity is due to electron-deficient impurity, the conductors are called *p*-type semiconductors.

For More Material Join: @JEEAdvanced_2024

Magnetic Properties

- **Paramagnetic :** Substance weakly attracted in magnetic field and magnetized in same direction is known as paramagnetic substance.
- **Diamagnetic :** Substance weakly repelled in magnetic field is known as diamagnetic substance
- **Ferromagnetic :** Substance attracted very strongly by magnetic field is known as ferromagnetic substance.
- **Antiferromagnetic :** Substance in which oppositely located domains cancel out each other's magnetic moment is known as Antiferromagnetic substance.
- **Ferrimagnetic :** Substance in which magnetic moment of domains are arranged in parallel and antiparallel directions in unequal numbers is known as ferrimagnetic substance.

For More Material Join: @JEEAdvanced_2024



Level 1

1. Which of the following statement is true for ionic solids?
(a) Ionic solids are soluble in non-polar solvent
(b) Under the electric field cations and anions acquire translatory motion in opposite directions
(c) Structural units have strong electrostatic force of attraction
(d) Structural units have dipole-dipole interactions

2. Which one is called pseudo solid ?
(a) CaF_2 (b) Glass (c) NaCl (d) All

3. Solids which do not show the same physical properties in different directions are called:
(a) pseudo solids (b) isotropic solids
(c) polymorphic solids (d) anisotropic solids

4. Graphite is an example of :
(a) ionic solid (b) covalent solid
(c) metallic crystal (d) none of these

5. Amorphous solids are:
(a) isotropic and supercooled liquids (b) anisotropic and supercooled liquids
(c) isoenthalpic and superheated liquids (d) isotropic and superheated solids

6. Crystals which are good conductor of electricity and heat are:
(a) ionic crystals (b) covalent crystals
(c) metallic crystals (d) molecular crystals

7. The bond length and bond angles in molecules in the solid state are calculated by:
(a) X-ray diffraction technique (b) neutrons bombardment
(c) protons bombardment (d) none of these

8. The unit cell with parameters 90 and $a \quad b \quad c$ is :
(a) cubic (b) triclinic (c) hexagonal (d) tetragonal

9. Triclinic crystal has the following unit cell parameters :
(a) $a \quad b \quad c$; 90 (b) $a \quad b \quad c$; 90
(c) $a \quad b \quad c$; 90 (d) $a \quad b \quad c$; 90 , 120

10. If all three interfacial angles defining the unit cell, are equal in magnitude, the crystal cannot be:
(a) rhombohedral (b) cubic (c) hexagonal (d) tetragonal

11. In a hexagonal crystal :
(a) 90 ; $a \quad b \quad c$ (b) 90 ; $a \quad b \quad c$
(c) 90 ; $a \quad b \quad c$ (d) 90 , 120 ; $a \quad b \quad c$

12. Orthorhombic crystal has the following unit cell parameters :
(a) $a \quad b \quad c$; 90 (b) $a \quad b \quad c$; 90
(c) $a \quad b \quad c$; 90 (d) $a \quad b \quad c$; 90 , 120

13. Which of the crystal systems contains the maximum number of Bravais lattices?
(a) Cubic (b) Hexagonal (c) Triclinic (d) Orthorhombic

For More Material Join: @JEEAdvanced_2024

14. The most unsymmetrical and symmetrical systems are, respectively :
(a) Tetragonal, Cubic (b) Triclinic , Cubic
(c) Rhombohedral, Hexagonal (d) Orthorhombic, Cubic

15. The crystal system of a compound with unit cell parameters,
a 0.328 nm, b 0.328 nm, c 0.527 nm and 90 is :
(a) cubic (b) tetragonal (c) monoclinic (d) rhombohedral

16. In the primitive cubic unit cell, the atoms are present at the:
(a) corners of the unit cell (b) centre of the unit cell
(c) centre of each face of the unit cell (d) one set of faces of the unit cell

17. In the body centered unit cell, the lattice points are present at the:
(a) corners of unit cell only
(b) corners and centre of unit cell
(c) corners and centre of each face of the unit cell
(d) corners and at one set of faces of unit cell

18. In the face centered unit cell, the lattice points are present at the:
(a) corners of unit cell only (b) corners and centre of the unit cell
(c) corners and face centres of the unit cell (d) face centres of the unit cell

19. The effective number of atoms per unit cell in a simple cube, face centred cube and body centred cube are respectively:
(a) 1, 4, 2 (b) 1, 2, 4 (c) 8, 14, 9 (d) 8, 4, 2

20. What would be the effective number of atoms per unit cell in end centered cubic unit cell, if this type of unit cell exist in nature ?
(a) 1 (b) 2 (c) 3 (d) 4

21. In the body centered cubic unit cell and simple cubic unit cell, the radius of atom in terms of edge length (a) of the unit cell is respectively:
(a) $\frac{a}{2}, \frac{a}{2\sqrt{2}}$ (b) $\frac{a}{\sqrt{2}}, \frac{a}{2}$ (c) $\frac{a}{2\sqrt{2}}, \frac{a}{2}$ (d) $\frac{\sqrt{3}\,a}{4}, \frac{a}{2}$

22. In the face centered cubic unit cell, the radius of atoms in terms of edge length (a) of unit cell is:
(a) $a/2$ (b) $a/\sqrt{2}$ (c) $a/2\sqrt{2}$ (d) $\sqrt{3}a/4$

23. The fraction of volume occupied by atoms in a primitive cubic unit cell is nearly:
(a) 0.48 (b) 0.52 (c) 0.55 (d) 0.68

24. The fraction of volume occupied by atoms in a body centered cubic unit cell is:
(a) 0.32 (b) 0.48 (c) 0.68 (d) 0.74

25. The fraction of volume occupied by atoms in a face centered cubic unit cell is:
(a) 0.32 (b) 0.48 (c) 0.68 (d) 0.74

26. Which of the following crystal lattice has the minimum empty space?
(a) Simple cubic (b) Body centred cubic
(c) Face centred cubic (d) Simple tetragonal

27. Which of the following has the smallest packing efficiency for atoms of a single type?
(a) Body centered cubic (b) Face centered cubic
(c) Simple cubic (d) None of these

28. Polonium crystallizes in a simple cubic structure. The edge of the unit cell is 0.236 nm. What is the radius of the polonium atom:
(a) 0.144 nm (b) 0.156 nm (c) 0.118 nm (d) 0.102 nm

For More Material Join: @JEEAdvanced_2024

29. Lithium crystallizes as body centered cubic crystals. If the length of the side of unit cell is 350 pm, the atomic radius of lithium is:
(a) 303.1 pm (b) 606.2 pm (c) 151.5 pm (d) 123.7 pm

30. Gold has a face centered cubic lattice with an edge length of the unit cube of 407 pm. The diameter of the gold atom is:
(a) 576.6 pm (b) 287.8 pm (c) 352.5 pm (d) 704.9 pm

31. The neon atoms has a radius of 160 pm. What is the edge of the unit cell of a face centered structure of neon?
(a) 490 pm (b) 320 pm (c) 453 pm (d) 481 pm

32. What are the number of atoms per unit cell and the number of nearest neighbours in a simple cubic structure?
(a) 1, 6 (b) 4, 12 (c) 2, 8 (d) 2, 6

33. What are the number of atoms per unit cell and the number of nearest neighbours in a face centered cubic structure?
(a) 4, 8 (b) 2, 8 (c) 2, 6 (d) 4, 12

34. What are the number of atoms per unit cell and the number of nearest neighbours in a body centered cubic structure?
(a) 4, 12 (b) 1, 6 (c) 2, 8 (d) 2, 5

35. Each edge of a cubic unit cell is 400 pm long. If atomic mass of the element is 120 and its density is 6.25 g/cm^3, the crystal lattice is : (use N_A 6 10^{23})
(a) primitive (b) body centered (c) face centered (d) end centered

36. Tungsten has an atomic radius of 0.136 nm. The density of tungsten is 19.4 g/cm^3. What is the crystal structure of tungsten? (Atomic mass W 184)
(a) Simple cubic (b) Body centered cubic
(c) Face centered cubic (d) None of these

37. The density of argon (face centered cubic cell) is 1.83 g/cm^3 at 20°C. What is the length of an edge a unit cell? (Atomic mass : Ar 40)
(a) 0.599 nm (b) 0.569 nm (c) 0.525 nm (d) 0.551 nm

38. The density of nickel (face centered cubic cell) is 8.94 g/cm^3 at 20°C. What is the radius of the atom? (Atomic mass : Ni 59)
(a) 0.124 nm (b) 0.136 nm (c) 0.149 nm (d) 0.110 pm

39. The density of krypton (face centered cubic cell) is 3.19 g/cm^3. What is the radius of the atom? (Atomic mass : Kr 84)
(a) 0.198 nm (b) 0.221 nm (c) 0.206 nm (d) 0.225 nm

40. The face centered cubic cell of platinum has an edge length of 0.392 nm. Calculate the density of platinum (g/cm^3) : (Atomic mass : Pt 195)
(a) 20.9 (b) 20.4 (c) 19.6 (d) 21.5

41. The body centered cubic cell of chromium has an edge length of 0.288 nm. Calculate the density of chromium (g/cm^3) : (Atomic mass : Cr 52.0)
(a) 6.80 (b) 7.60 (c) 6.60 (d) 7.23

42. An element crystallizes in a face centered cubic lattice and the edge of the unit cell is 0.559 nm. The density is 3.19 g/cm^3. What is the atomic mass?
(a) 87.6 (b) 79.9 (c) 85.5 (d) 83.9

For More Material Join: @JEEAdvanced_2024

43. The element crystallizes in a body centered cubic lattice and the edge of the unit cell is 0.351 nm. The density is 0.533 g/cm^3. What is the atomic mass?
(a) 12.0 (b) 6.94 (c) 9.01 (d) 10.8

44. An element X (At. mass 80 g/mol) has fcc structure. Calculate no. of unit cells in 8 gm of X:
(a) 0.4 N_A (b) 0.1 N_A (c) 4 N_A (d) none of these

45. Molybdenum (At. mass 96 g mol ^{1}) crystallizes as bcc crystal. If density of crystal is 10.3 g/cm^3, then radius of Mo atom is (use N_A 6 10^{23}):
(a) 111 pm (b) 314 pm (c) 135.96 pm (d) none of these

46. What is the coordination number of an atom for an element crystallizing with a cubic lattice? Calculate the corresponding coordination number for the simple, fcc and bcc lattices:
(a) 12 sc; 12 fcc; 8 bcc (b) 6 sc; 14 fcc; 8 bcc
(c) 8 sc; 12 fcc; 6 bcc (d) 6 sc; 12 fcc; 8 bcc

47. Which of the following layering pattern will have a void fraction of 0.260?
(a) ABCCBAABC (b) ABBAABBA (c) ABCABCABC (d) ABCAABCA

48. The most malleable metals (Cu, Ag, Au) have close-packing of the type :
(a) Hexagonal close-packing
(b) Cubic close-packing
(c) Body-centred cubic packing
(d) Malleability is not related to type of packing

49. The co-ordination number of a metal crystallising in a hexagonal close-packed structure is:
(a) 12 (b) 4 (c) 8 (d) 6

50. If the ratio of coordination no. of A to that of B is $x : y$, then the ratio of no. of atoms of A to that no. of atoms of B in unit cell is:
(a) $x : y$ (b) $y : x$ (c) $x^2 : y$ (d) $y : x^2$

51. The atomic radius of strontium (Sr) is 215 pm and it crystallizes with a cubic closest packing. Edge length of the cube is:
(a) 430 pm (b) 608.2 pm (c) 496.53 pm (d) none of these

52. By X-ray diffraction it is found that nickel (at mass 59 g mol ^{1}), crystallizes with ccp. The edge length of the unit cell is 3.5 Å. If density of Ni crystal is 9.0 g/cm^3, then value of Avogadro's number from the data is :
(a) 6.05 10^{23} (b) 6.11 10^{23} (c) 6.02 10^{23} (d) 6.023 10^{23}

53. Stacking of square close packed layers give rise to :
(a) bcc structure (b) fcc structure
(c) Simple cubic structure (d) hcp structure

54. In a hexagonal close packed (hcp) structure of spheres, the fraction of the volume occupied by the sphere is A. In a cubic close packed structure the fraction is B. The relation for A and B is :
(a) A B
(b) A B
(c) A B
(d) A is equal to the fraction in a simple cubic lattice.

For More Material Join: @JEEAdvanced_2024

55. The unit cell present in ABCABC, closest packing of atoms is:
(a) hexagonal (b) tetragonal
(c) face centered cubic (d) primitive cubic




56. The number of atoms present in a hexagonal close-packed unit cell is :
(a) 4 (b) 6 (c) 8 (d) 12

57. The unit cell present in ABAB, closest packing of atoms is:
(a) hexagonal (b) tetragonal
(c) face centered cubic (d) primitive cubic

58. The numbers of tetrahedral and octahedral holes in a hexagonal primitive unit cell are respectively:
(a) 8, 4 (b) 6, 12 (c) 2, 1 (d) 12, 6

59. Which one of the following schemes of ordering closed packed sheets of equal sized spheres do not generate closest packed lattice?
(a) ABCABC (b) ABACABAC (c) ABBAABBA (d) ABCBCABCBC

60. In the closest packing of atoms, there are:
(a) one tetrahedral void and two octahedral voids per atom
(b) two tetrahedral voids and one octahedral void per atom
(c) two of each tetrahedral and octahedral voids per atom
(d) one of each tetrahedral and octahedral void per atom

61. Which of the following figures represents the cross-section of an octahedral site?
(a)  (b)  (c)  (d) 

62. In which of the following pairs of structures, tetrahedral as well as octahedral holes are found?
(a) bcc and fcc (b) hcp and simple cubic
(c) hcp and ccp (d) bcc and hcp

63. The number of octahedral void in bcc structure is :
(a) 0 (b) 1 (c) 2 (d) 4

64. An ionic compound is expected to have octahedral structure if r_c / r_a (r_c r_a) lies in the range of:
(a) 0.414 to 0.732 (b) 0.732 to 0.82
(c) 0.225 to 0.414 (d) 0.155 to 0.225

65. An ionic compound is expected to have tetrahedral structure if r_c / r_a:
(a) lies in the range of 0.414 to 0.732 (b) lies in the range of 0.225 to 0.414
(c) lies in the range of 0.155 to 0.225 (d) is more than 0.732

66. An ionic compound is expected to have body centered type cubic unit cell if r_c / r_a:
(a) is greater than 0.732 (b) lies in the range of 0.414 to 0.732
(c) lies in the range of 0.255 to 0.414 (d) lies in the range of 0.155 to 0.225

67. In the closest packing of atoms A (radius : r_a), the radius of atom B that can be fitted into tetrahedral void is:
(a) 0.155 r_a (b) 0.225 r_a (c) 0.414 r_a (d) 0.732 r_a

For More Material Join: @JEEAdvanced_2024

68. In the closest packing of atoms A (radius : r_a), the radius of atom B that can be fitted octahedral void is:
(a) $1.155\, r_a$ (b) $0.225\, r_a$ (c) $0.414\, r_a$ (d) $0.732\, r_a$

69. How many nearest neighbours are there in an atom or ion for an octahedral hole of a closed packed structure?
(a) 4 (b) 6 (c) 8 (d) 12

70. How many "nearest" and "next nearest" neighbours, respectively, does potassium have in bcc lattice ?
(a) 8, 8 (b) 8, 6 (c) 6, 8 (d) 6, 6

71. In the closest packing of atoms:
(a) the size of tetrahedral void is greater than that of octahedral void
(b) the size of tetrahedral void is smaller than that of octahedral void
(c) the size of tetrahedral void is equal to that of octahedral void
(d) the size of tetrahedral void may be greater or smaller or equal to that of octahedral void depending upon the size of atoms

72. In the ionic compound AB the ratio $r_A : r_B$ is 0.414. Indicate the correct statement among the following:
(a) Cations form close packing and anions exactly fit into the octahedral voids
(b) Anions form close packing and cations occupy precisely half of the tetrahedral voids
(c) Anions form close packing and cations occupy precisely all the octahedral voids
(d) Anions form close packing and cations fit into the octahedral voids loosely

73. In the unit cell of KCl (NaCl type), Cl ions constitute ccp and K ions fall into the octahedral holes. These holes are:
(a) one at the centre and 6 at the centres of the faces
(b) one at the centre and 12 at the centres of the edges
(c) 8 at the centres of 8 small cubes forming the unit cell
(d) none of these

74. Which is incorrect statement ?
(a) In NaCl structure, tetrahedral voids are unoccupied
(b) In ZnS structure, octahedral voids are unoccupied
(c) In CaF_2 structure, all tetrahedral voids are occupied
(d) In Na_2O structure, all tetrahedral voids are unoccupied

75. If the radius of the anion in an ionic solid is 200 pm, what would be the radius of the cation that fits exactly into a cubic hole?
(a) 146.4 pm (b) 82.8 pm (c) 45 pm (d) None of these

76. The CsCl type structure is exhibited by alkali halides only when the radius of the cation is large enough to keep touching its eight nearest neighbour anions. Below what minimum ratio of cation to anion radii (r /r) this contact is prevented?
(a) 0.225 (b) 0.414
(c) 0.632 (d) 0.732

77. MgO crystallizes in a cubic type crystal system. The ionic radii for Mg^{2} and O^{2} are 0.066 and 0.140 nm respectively. One can conclude that the Mg^{2} ions occupy:
(a) a cubic hole in a simple cubic structure
(b) every tetrahedral hole in a close packed structure

For More Material Join: @JEEAdvanced_2024

(c) an octahedral hole in a cubic close packed structure
(d) every other tetrahedral hole in a close packed structure

78. The unit cell of diamond is made up of:
(a) 6 carbon atoms, 4 atoms constitute ccp and two atoms occupy half of octahedral voids
(b) 8 carbon atom, 4 atoms constitute ccp and 4 atoms occupy all the octahedral voids
(c) 8 carbon atoms, 4 atoms form fcc lattice and 4 atoms occupy half of the tetrahedral voids alternately
(d) 12 carbon atoms, 4 atoms form fcc lattice and 8 atoms occupy all the tetrahedral holes

79. In diamond, the coordination number of carbon is:
(a) four and its unit cell has eight carbon atoms
(b) four and its unit cell has six carbon atoms
(c) six and its unit cell has four carbon atoms
(d) four and its unit cell has four carbon atoms

80. Predict coordination number of the cation in crystals of the following compounds:
1. MgO : r_c 0.65 Å; r_a 1.40 Å, 2. MgS : r_c 0.65 Å; r_a 1.84 Å
(a) 6, 4 (b) 4, 6 (c) 3, 4 (d) 6, 8

81. In a cubic unit cell, seven of the eight corners are occupied by atoms *A* and centres of faces are occupied by atoms *B*. The general formula of the compound is :
(a) A_7B_6 (b) A_7B_{12} (c) A_7B_{24} (d) $A_{24}B_7$

82. CaS exists in a cubic close packed arrangement of S^{2} ions in which Ca^{2} ions occupy 1/2 of the available tetrahedral holes. How many Ca^{2} and S^{2} ions are contained in the unit cell?
(a) 1, 1 (b) 2, 4 (c) 4, 4 (d) 4, 2

83. In the spinel structure, oxides ions are cubical-closest packed whereas 1/8th of tetrahedral voids are occupied by A^{2} cation and 1/2 of octahedral voids are occupied by B^{3} cations. The general formula of the compound having spinel structure is:
(a) $A_2B_2O_4$ (b) AB_2O_4 (c) $A_2B_4O_2$ (d) $A_4B_2O_2$

84. If the anions (*A*) form hexagonal closest packing and cations (*C*) occupy only 2/3 octahedral voids in it, then the general formula of the compound is:
(a) CA (b) CA_2 (c) C_2A_3 (d) C_3A_2

85. In a solid, oxide ions are arranged in ccp, cations *A* occupy $(1/8)^{th}$ of the tetrahedral voids and cations *B* occupy $(1/4)^{th}$ of the octahedral voids. The formula of the compound is:
(a) ABO_4 (b) AB_2O_3 (c) A_2BO_4 (d) AB_4O_4

86. In a face centered cubic arrangement of *A* and *B* atoms whose *A* atoms are at the corner of the unit cell and *B* atoms at the face centers. One of the *B* atoms missing from one of the face in unit cell. The simplest formula of compound is:
(a) AB_3 (b) A_8B_5 (c) A_2B_5 (d) $AB_{2/5}$

87. An alloy of copper, silver and gold is found to have copper constituting the ccp lattice. If silver atoms occupy the edge centres and gold is present at body centre, the alloy will have the formula:
(a) Cu_4Ag_2Au (b) Cu_4Ag_4Au (c) Cu_4Ag_3Au (d) CuAgAu

For More Material Join: @JEEAdvanced_2024

88. Which of the following statements is correct in the rock-salt structure of an ionic compound?
(a) Co-ordination number of cation is four and anion is six
(b) Co-ordination number of cation is six and anion is four
(c) Co-ordination number of each cation and anion is four
(d) Co-ordination number of each cation and anion is six

89. Which of the following statements is correct in the body centered type cubic structure of an ionic compound?
(a) Co-ordination number of each cation and anion is two
(b) Co-ordination number of each cation and anion is four
(c) Co-ordination number of each cation and anion is six
(d) Co-ordination number of each cation and anion is eight

90. Which of the following statements is correct in the zinc blende type structure of an ionic compound?
(a) Co-ordination number of each cation and anion is two
(b) Co-ordination number of each cation and anion is four
(c) Co-ordination number of each cation and anion is six
(d) Co-ordination number of each cation and anion is eight

91. Which of the following expression is correct in case of a sodium chloride unit cell (edge length, a)?
(a) $r_c \quad r_a \quad a$ (b) $r_c \quad r_a \quad a/2$
(c) $r_c \quad r_a \quad 2a$ (d) $r_c \quad r_a \quad \sqrt{2}a$

92. In an ideal closest rock salt structure (edge length, a), which of the following expressions is correct?
(a) $r_a \quad \sqrt{2}a$ (b) $r_a \quad a/\sqrt{2}$
(c) $r_a \quad a/2\sqrt{2}$ (d) $r_a \quad a/4$

93. Which of the following expressions is correct in case of a CsCl unit cell (edge length, a)?
(a) $r_c \quad r_a \quad a$ (b) $r_c \quad r_a \quad a/\sqrt{2}$
(c) $r_c \quad r_a \quad \sqrt{3}a/2$ (d) $r_c \quad r_a \quad a/2$

94. In NaCl the centres of two nearest like-charged ions are present at a distance of:
(a) $\frac{1}{2}a\sqrt{2}$ (b) $\frac{1}{2}a$
(c) $\frac{\sqrt{3}}{2}a$ (d) $\frac{1}{\sqrt{2}}2a$

95. In sodium chloride crystal, the number of next nearest neighbours of each Na ions is:
(a) 8 Cl ions (b) 12 Na ions
(c) 12 Cl ions (d) 24 Cl ions

96. In an ionic compound $A \quad X$, the radii of A and X ions are 1.0 pm and 2.0 pm, respectively. The volume of the unit cell of the crystal AX will be :
(a) $27\,\text{pm}^3$ (b) $64\,\text{pm}^3$
(c) $125\,\text{pm}^3$ (d) $216\,\text{pm}^3$

For More Material Join: @JEEAdvanced_2024


97. The coordination number of cation and anion in fluorite CaF_2 and anti-fluorite Na_2O are respectively:
(a) 8 : 4 and 6 : 3 (b) 6 : 3 and 4 : 4
(c) 8 : 4 and 4 : 8 (d) 4 : 8 and 8 : 4

98. Select the incorrect statement in a CsCl crystal:
(a) Cs forms a simple cubic lattice, Cl forms a simple cubic lattice
(b) Cl occupies body centre of Cs
(c) Cs occupies body centre of Cl
(d) It is impossible for Cl to occupy body centre of Cs because the body centre void of Cs is smaller than Cl ion size

99. The radius of a divalent cation A^2 is 94 pm and of divalent anion B^2 is 146 pm. The compound AB has :
(a) Rock salt structure (b) Zinc blende structure
(c) Antifluorite structure (d) Caesium chloride like structure

100. A binary solid (AB) has a rock salt structure. If the edge length is 400 pm, and radius of cation is 80 pm the radius of anion is:
(a) 100 pm (b) 120 pm
(c) 250 pm (d) 325 pm

101. For a solid with the structure shown in Fig., the coordination numbers of the points A and B, respectively, are



(a) 6, 8 (b) 8, 8 (c) 6, 6 (d) 4, 6

102. An ionic compound AB_2 has fluorite type structures. If the radius B is 200 pm, then the ideal radius of A would be:
(a) 82.8 pm (b) 146.4 pm (c) 40 pm (d) 45 pm

103. In which of the following structures, the anion has maximum coordination number ?
(a) NaCl (b) ZnS (c) CaF_2 (d) Na_2O

104. CsCl has bcc structure with Cs at the centre and Cl ion at each corner. If r_{Cs} is 1.69 Å and r_{Cl} is 1.81 Å, what is the edge length of the cube?
(a) 3.50 Å (b) 3.80 Å
(c) 4.04 Å (d) 4.50 Å

105. CsBr has bcc type structure with edge length 4.3 pm. The shortest inter ionic distance in between Cs and Br is:
(a) 3.72 pm (b) 1.86 pm (c) 7.44 pm (d) 4.3 pm

For More Material Join: @JEEAdvanced_2024

106. If the radius of Cl ion is 181 pm, and the radius of Na ion is 101 pm then the edge length of unit cell is:
(a) 282 pm (b) 285.71 pm
(c) 512 pm (d) 564 pm

107. Ammonium chloride crystallizes in a body centered cubic lattice with edge length of unit cell equal to 387 pm. If the size of Cl ion is 181 pm, the size of NH_4 ion would be:
(a) 116 pm (b) 154 pm
(c) 174 pm (d) 206 pm

108. Salt *AB* has a zinc blende structure. The radius of A^2 and B^2 ions are 0.7 Å and 1.8 Å respectively. The edge length of *AB* unit cell is:
(a) 2.5 Å (b) 5.09 Å
(c) 5 Å (d) 5.77 Å

109. Transition metals, when they form interstitial compounds, the non-metals (H, B, C, N) are accomodated in :
(a) voids or holes in cubic-packed structure (b) tetrahedral voids
(c) octahedral voids (d) all of these

110. In diamond, each carbon atom is bonded to four other carbon atoms tetrahedrally. Alternate tetrahedral voids are occupied by carbon atoms. The number of carbon atoms per unit cell is :
(a) 4 (b) 6
(c) 8 (d) 12

111. Which of the following statements for crystals having Schottky defect is not correct?
(a) Schottky defect arises due to the absence of a cation and anion from the position which it is expected to occupy
(b) Schottky defect are more common in ionic compounds with high co-ordination numbers
(c) The density of the crystals having Schottky defect is larger than that of the perfect crystal
(d) The crystal having Schottky defect is electrically neutral as a whole

112. Which is correct statement ?
(a) When temperature increases then number of defects decreases.
(b) Schottky defect occurs when radius of cation is smaller
(c) Frenkel defect occurs when radius of cation is smaller
(d) None of these

113. Which of the following statements for crystals having Frenkel defect is not correct?
(a) Frenkel defects are observed where the difference in sizes of cations and anions is large
(b) The density of crystals having Frenkel defect is lesser than that of a pure perfect crystal
(c) In an ionic crystal having Frenkel defect may also contain Schottky defect
(d) Usually alkali halides do not have Frenkel defect

114. When anion leaves the normal lattice site and electron occupies interstitial sites in its crystal lattice, it is called:
(a) Schottky defect (b) Frenkel defect
(c) Metal excess defect (d) Stoichiometric defect

115. Which of the following defects does KBr show?
(a) Frenkel (b) Schottky
(c) Metal excess (d) Metal deficiency

For More Material Join: @JEEAdvanced_2024

116. Dopping of AgCl crystals with $CdCl_2$ results in:
(a) Schottky defect (b) Frenkel defect
(c) Substitutional cation vacancy (d) Formation of F-centres

117. NaCl shows Schottky defects and AgCl Frenkel defects. Their electrical conductivity is due to the:
(a) motion of ions and not the motion of electrons
(b) motion of electrons and not the motion of ions
(c) lower coordination number of NaCl
(d) higher coordination number of AgCl

118. Which one of the following crystals does not exhibit Frenkel defect?
(a) AgBr (b) AgCl
(c) CsCl (d) ZnS

119. Select the incorrect statement:
(a) Stoichiometry of crystal remains uneffected due to Schottky defect
(b) Frenkel defect is usually shown by ionic compounds having low coordination number
(c) F-centres generation is responsible factor for imparting the colour to the crystal
(d) Density of crystal always increases due to substitutional impurity defect

120. In diamond, carbon atom occupy fcc lattice points as well as alternate tetrahedral voids. If edge length of the unit cell is 356 pm, then diameter of carbon atom is:
(a) 77.07 pm (b) 154.14 pm
(c) 251.7 pm (d) 89 pm

121. When NaCl is dopped with 10^{-5} mole % of $SrCl_2$, what is the no. of cationic vacancies?
(a) $10^{-5} N_A$ (b) $10^{-7} N_A$
(c) $2 \times 10^{-7} N_A$ (d) None of these

122. The composition of a sample of Wustite is $Fe_{0.93}O$. What is the percentage of iron present as Fe^{3+} in total iron?
(a) 15.05% (b) 25%
(c) 35% (d) 45%

123. A certain sample of cuprous sulphide is found to have composition $Cu_{1.8}S$, because of incorporation of Cu^{2+} ions in the lattice. What is the mole % of Cu^{2+} in total copper content in this crystal?
(a) 99.8% (b) 11.11%
(c) 88.88% (d) None of these

124. Ferrimagnetism is in :
(a) (b)
(c) (d) none of these

125. Fe_3O_4 is ferrimagnetic at room temperature but at 850 K it becomes :
(a) diamagnetic (b) ferromagnetic
(c) non-magnetic (d) paramagnetic

For More Material Join: @JEEAdvanced_2024

Level 2

1. When heated above 916°C, iron changes its bcc crystalline form to fcc without the change in the radius of atom. The ratio of density of the crystal before heating and after heating is:
(a) 1.069 (b) 0.918 (c) 0.725 (d) 1.231

2. $TlAl(SO_4)_2 \; xH_2O$ is bcc with 'a' 1.22 nm. If the density of the solid is 2.32 g/cc, then the value of x is (Given : N_A 6 10^{23}; at. mass : Tl 204, Al 27, S 32).
(a) 2 (b) 4 (c) 47 (d) 70

3. In an atomic bcc lattice what fraction of edge is not covered by atoms?
(a) 0.32 (b) 0.16 (c) 0.134 (d) 0.268

4. The packing efficiency of a simple cubic crystal with an interstitial atom exactly fitting at the body center is:
(a) 0.48 (b) 0.52 (c) 0.73 (d) 0.91

5. An atomic solid crystallizes in a body centre cubic lattice and the inner surface of the atoms at the adjacent corner are separated by 60.3 pm. If the atomic mass of A is 48, then density of the solid, is nearly:
(a) 2.7 g/cc (b) 5.07 g/cc (c) 3.5 g/cc (d) 1.75 g/cc

6. Sodium (Na 23) crystallizes in bcc arrangement with the interfacial separation between the atoms at the edge 53.6 pm. The density of sodium crystal is:
(a) 2.07 g/cc (b) 2.46 g/cc (c) 1.19 g/cc (d) none of these


7. The density of solid argon (Ar 40 g/mol) is 1.68 g/mL at 40 K. If the argon atom is assumed to be a sphere of radius 1.50 10^{-8} cm, what % of solid Ar is apparently empty space? (use N_A 6 10^{23})
(a) 35.64 (b) 64.36 (c) 74 (d) none of these

8. A body centered cubic lattice is made up of hollow spheres of B. Spheres of solid A are present in hollow spheres of B. Radius of A is half of radius of B. What is the ratio of total volume of spheres of B unoccupied by A in a unit cell and volume of unit cell?
(a) $\frac{7\sqrt{3}}{64}$ (b) $\frac{7\sqrt{3}}{128}$ (c) $\frac{7.}{24}$ (d) none of these

9. First three nearest neighbour distances for primitive cubic lattice are respectively (edge length of unit cell a):
(a) $a, \sqrt{2}a, \sqrt{3}a$ (b) $\sqrt{3}a, \sqrt{2}a, a$ (c) $a, \sqrt{2}a, 2a$ (d) $a, \sqrt{3}a, 2a$

10. First three nearest neighbour distances for body centered cubic lattice are respectively:
(a) $\sqrt{2}a, a, \sqrt{3}a$ (b) $\frac{a}{\sqrt{2}}, a, \sqrt{3}a$ (c) $\frac{\sqrt{3}a}{2}, a, \sqrt{2}a$ (d) $\frac{\sqrt{3}a}{2}, a, \sqrt{3}a$

11. Given : The unit cell structure of compound is shown below.



For More Material Join: @JEEAdvanced_2024

The formula of compound is :

(a) $A_8B_{12}C_5$ (b) AB_2C_3 (c) $A_2B_2C_5$ (d) ABC_5

12. The density of a pure substance '*A*' whose atoms are in cubic close pack arrangement is 1 g/cc. If all the tetrahedral voids are occupied by '*B*' atom, what is the density of resulting solid in g/cc. [Atomic mass (*A*) 30 g/mol and atomic mass (*B*) 50 g/mol]

(a) 3.33 (b) 4.33 (c) 2.33 (d) 5.33

13. In a planar tetra-atomic molecule, XY_3, *X* is at the centroid of the equilateral triangle formed by the atoms, *Y*. If the *X*—*Y* bond distance is 1 Å, what is the distance between the centres of any two *Y* atoms?

(a) $1/\sqrt{3}$ Å (b) $\sqrt{2}$ Å

(c) $\sqrt{3}$ Å (d) $1/\sqrt{2}$ Å

14. How many unit cells are present in 5.0 gm of crystal *AB* (formula mass of *AB* 40) having rock salt type structure? (N_A Avogadro's no.)

(a) N_A (b) $\frac{N_A}{10}$

(c) $4\ N_A$ (d) none of these

15. The density of CaF_2 (fluorite structure) is 3.18 g/cm^3. The length of the side of the unit cell is:

(a) 253 pm (b) 344 pm

(c) 546 pm (d) 273 pm

16. A crystal of lead (II) sulphide has NaCl structure. In this crystal the shortest distance between a Pb^2 ion and S^2 ion is 297 pm. What is the volume of unit cell in lead sulphide?

(a) 209.6 10 ^{24} cm^3 (b) 207.8 10 ^{23} cm^3

(c) 22.3 10 ^{23} cm^3 (d) 209.8 10 ^{23} cm^3

17. CdO has NaCl structures with density 8.27 g/cc. If the ionic radius of O^2 is 1.24 Å, determine ionic radius of Cd^2 :

(a) 1.5 Å (b) 1.1 Å

(c) 1.9 Å (d) 1.5 Å

18. KCl crystallizes in the same type of lattice as does NaCl. Given that r_{Na}/r_{Cl} 0.50 and r_{Na}/r_K 0.70, calculate the ratio of the side of the unit cell for KCl to that for NaCl:

(a) 1.143 (b) 1.224

(c) 1.414 (d) 0.875

19. Ferrous oxide has a cubic structure and edge length of the unit cell is 5.0 Å. Assuming the density of ferrous oxide to be 3.84 g/cm^3, the no. of Fe^2 and O^2 ions present in each unit cell be : (use N_A 6 10^{23}) :

(a) 4 Fe^2 and 4 O^2 (b) 2 Fe^2 and 2 O^2

(c) 1 Fe^2 and 1 O^2 (d) 3 Fe^2 and 4 O^2

20. If an element (at. mass 50) crystallises in fcc lattice, with a 0.50 nm. What is the density of unit cell if it contains 0.25% Schottky defects (use N_A 6 10^{23})?

(a) 2.0 g/cc (b) 2.66 g/cc

(c) 3.06 g/cc (d) none of these

For More Material Join: @JEEAdvanced_2024

21. An element X (atomic mass 24 gm/mol) forms a face centered cubic lattice. If the edge length of the lattice is 4 10^{-8} cm and the observed density is 2.40 10^{3} kg/m^{3}, then the percentage occupancy of lattice points by element X is : (Use N_A 6 10^{23}) :

(a) 96 (b) 98

(c) 99.9 (d) none of these

22. In fcc lattice, A, B, C, D atoms are arranged at corner, face center, octahedral void and tetrahedral void respectively, then the body diagonal contains:

(a) $2A, C, 2D$ (b) $2A, 2B, 2C$


(c) $2A, 2B, D$ (d) $2A, 2D$

23. The distance between an octahedral and tetrahedral void in fcc lattice would be:

(a) $\sqrt{3}a$ (b) $\frac{\sqrt{3}a}{2}$

(c) $\frac{\sqrt{3}a}{3}$ (d) $\frac{\sqrt{3}a}{4}$

24. A_2B molecules (molar mass 259.8 g/mol) crystallises in a hexagonal lattice as shown in figure. The lattice constants were a 5 Å and b 8 Å. If density of crystal is 5 g/cm^{3} then how many molecules are contained in given unit cell? (use N_A 6 10^{23})



(a) 6 (b) 4 (c) 3 (d) 2

25. Graphite has h.c.p. arrangements of carbon atoms and the parallel planes are 3.35 Å apart. Determine density of graphite:

(a) 2.46 g/cc (b) 0.41 g/cc (c) 1 g/cc (d) 1.41 g/cc

26. How many effective Na and Cl ions are present respectively in a unit cell of NaCl solid (Rock salt structure) if all ions along line connecting opposite face centres are absent?

(a) 3, 3 (b) $\frac{7}{2}, 4$ (c) $\frac{7}{2}, \frac{7}{2}$ (d) $4, \frac{7}{2}$

27. A crystal is made of particles X and Y. X forms fcc packing and Y occupies all the octahedral voids. If all the particles along one body diagonal are removed then the formula of the crystal would be:

(a) X_4Y_3 (b) X_5Y_4 (c) X_4Y_5 (d) none of these

For More Material Join: @JEEAdvanced_2024

28. Select right expression for determining packing fraction (P. F.) of NaCl unit cell (assume ideal), if ions along an edge diagonal are absent:


(a) P.F. $\dfrac{\frac{4}{3}\ (r^3\ \ r^3)}{16\sqrt{2}\ r^3}$ (b) P.F. $\dfrac{\frac{4}{3}\ \ \frac{5}{2}r^3\ \ 4r^3}{16\sqrt{2}\ r^3}$

(c) P.F. $\dfrac{\frac{4}{3}\ \ \frac{5}{2}r^3\ \ r^3}{16\sqrt{2}\ r^3}$ (d) P.F. $\dfrac{\frac{4}{3}\ \ \frac{7}{2}r^3\ \ r^3}{16\sqrt{2}\ r^3}$

29. A crystal is made of particles X, Y and Z. X forms fcc packing. Y occupies all the octahedral voids of X and Z occupies all the tetrahedral voids of X. If all the particles along one body diagonal are removed then the formula of the crystal would be:

(a) XYZ_2 (b) X_2YZ_2 (c) $X_8Y_4Z_5$ (d) $X_5Y_4Z_8$

30. A crystal is made of particles A and B. A forms fcc packing and B occupies all the octahedral voids. If all the particles along the plane as shown in figure are removed, then, the formula of the crystal would be:



(a) AB (b) A_5B_7 (c) A_7B_5 (d) none of these

31. In the rock salt AB, if C introduced in tetrahedral voids such that no distortion occurs, then formula of resultant compound is :

(a) ABC (b) ABC_2 (c) A_4B_4C (d) ABC_8

32. Given length of side of hexagonal unit cell is $\dfrac{100}{\sqrt{2}}$ pm. The volume of hexagonal unit cell is (in pm^3) :

(a) 8 10^6 (b) 1.5 10^6 (c) 64 10^6 (d) 36 10^6

For More Material Join: @JEEAdvanced_2024

Level 3

Passage 1

Packing fraction of a unit cell is defined as the fraction of the total volume of the unit cell occupied by the atom(s).

$$\text{P.F.} \quad \frac{\text{Volume of the atom(s) present in a unit cell}}{\text{Volume of unit cell}} \quad \frac{Z \quad \frac{4}{3} \quad r^3}{a^3}$$

and % of empty space 100 P.F. 100

where Z effective number of atoms in a cube

r radius of an atom

a edge length of the cube

1. % of empty space in body centered cubic unit cell is nearly:

(a) 52.36 (b) 68 (c) 32 (d) 26

2. Packing fraction in face centered cubic unit cell is:

(a) 0.7406 (b) 0.6802 (c) 0.5236 (d) None of these

Passage 2

Density of a unit cell is represented as

$$\frac{\text{Effective no. of atom(s)} \quad \text{Atomic mass / formula mass}}{\text{Volume of a unit cell}} \quad \frac{Z.M.}{N_A.a^3}$$

where, mass of unit cell mass of effective no. of atom(s) or ion(s).

M At. mass/formula mass

N_A Avogadro's no. 6.023 10^{23}

a edge length of unit cell

1. Silver crystallizes in a fcc lattice and has a density of 10.6 g/cm^3. What is the length of an edge of the unit cell?

(a) 40.7 nm (b) 0.2035 nm (c) 0.101 nm (d) 4.07 nm

2. An element crystallizes in a structure having fcc unit cell of an edge 200 pm. Calculate the density, if 100 g of this element contains 12 10^{23} atoms:

(a) 41.66 g/cm^3 (b) 4.166 g/cm^3 (c) 10.25 g/cm^3 (d) 1.025 g/cm^3

3. The density of KBr is 2.75 g/cm ^{3}. The length of the edge of the unit cell is 654 pm. To which type of cubic crystal, KBr belongs?

(a) simple cubic (b) bcc (c) fcc (d) none of these

For More Material Join: @JEEAdvanced_2024

Passage 3

A spinel is an important class of oxides consisting of two types of metal ions with the oxide ions arranged in ccp layers. The normal spinel has one-eighth of the tetrahedral holes occupied by one type of metal ion and one-half of the octahedral holes occupied by another type of metal ion. Such a spinel is formed by Mg^{2}, Al^{3} and O^{2}. The neutrality of the crystal is being maintained.

1. The formula of the spinel is:
(a) Mg_2AlO_4 (b) $MgAl_2O_4$ (c) $Mg_3Al_2O_6$ (d) none of these

2. Type of hole occupied by Al^{3} ions is:
(a) tetrahedral (b) octahedral (c) both (a) and (b) (d) none of these

3. Type of hole occupied by Mg^{2} ions is:
(a) tetrahedral (b) octahedral (c) both (a) and (b) (d) none of these


4. If oxide ion is replaced by $X^{8/3}$, the number of anionic vacancy per unit cell is:
(a) 1 (b) 2 (c) 3 (d) 3/4

Passage 4

Ionic lattice has two major points defects, (1) Schottky (2) Frenkel defects. Schottky defects occurs due to the cation-anion pair's missing from the lattice sites. Frenkel defects occurs when an ion leaves its lattice site and fits into an interstitial space. The neutrality of the crystal is being maintained and we considered all losses from interstitial positions.


1. Which defect decreases density of the crystal?
(a) Frenkel defect (b) Schottky defect (c) Both (a) and (b) (d) None of these

2. Structure shown here represents:



(a) Schottky defect (b) Frenkel defect
(c) Metal excess defect (d) None of these

3. Structure shown here represents:



(a) Schottky defect (b) Frenkel defect (c) Both defects (d) None of these

For More Material Join: @JEEAdvanced_2024



Passage 5

Doping means introduction of small amount of impurities like phosphorus, arsenic or boron into the pure crystal. In pure silicon, there are four valencies used in bonding with other four adjacent silicon atoms. When a silicon crystal is doped with a group-15 element (with five valence electrons) such as P, As, or Bi, the structure of the crystal lattice remains unchanged. Out of the five valence electrons of group-15 doped element, four electrons are used in normal covalent bonding with silicon while fifth electron is delocalised and thus conducts electricity.

Doping a silicon crystal with a group-13 element (with three valence electrons) such as B, Al, Ga or In products a semiconductor with three electrons in dopant. The place where fourth electron is missing is called an electron vacancy or hole. Such hole can move through the crystal like a positive charge giving rise conduction of electricity.

1. No. of valence electrons in silicon are:
(a) 3 (b) 4 (c) 5 (d) 6

2. Silicon that has been doped with group-15 elements is called:
(a) *p*-type semiconductor (b) *n*-type semiconductor
(c) electron vacancy or hole (d) none of these

3. Silicon that has been doped with group-13 elements is called:
(a) *p*-type semiconductor (b) *n*-type semiconductor
(c) electron vacancy or hole (d) none of these



Passage 6

Metallic Gold crystallise in fcc lattice and the length of cubic unit cell is 407 pm.
(Given : Atomic mass of Gold 197, N_A 6 10^{23})

1. The density if it have 0.2% Schottky defect is (in gm/cm^3) :
(a) 4.86 (b) 9.72 (c) 19.48 (d) 19.44

2. The distance between next neighbours is :
(a) 407 pm (b) $407\sqrt{2}$ (c) $\frac{407}{\sqrt{2}}$ (d) $\frac{407\sqrt{3}}{2}$



Passage 7

In diamond structure, carbon atoms form fcc lattice and 50% tetrahedral voids occupied by carbon atoms. Every carbon atoms is surrounded tetrahedrally by four carbon atoms with bond length 154 pm. Germanium, silicon and grey tin also crystallise in same way as diamond.
(N_A 6 10^{23})

1. The mass of diamond unit cell is :
(a) 96 amu (b) 96 gm (c) 144 amu (d) 144 gm

2. The side length of diamond unit cell is (in pm) :
(a) 154 (b) 1422.63 (c) 711.32 (d) 355.66

For More Material Join: @JEEAdvanced_2024

3. The density of diamond unit cell is (in gm/cm^3) :
(a) 28.48 (b) 0.0556 (c) 0.445 (d) 3.56

One or More Answers is/are Correct



1. Select the correct statement(s):
(a) Co-ordination no. of an atom at a lattice point in simple cubic arrangement is 6
(b) Co-ordination no. of an atom at octahedral site is 8
(c) Co-ordination no. of an atom at a lattice point in hcp arrangement is 6
(d) Co-ordination no. of an atom at octahedral site is 6

2. Packing fraction of an identical solid sphere is 74% in:
(a) simple cubic structure (b) fcc structure
(c) hcp structure (d) bcc structure

3. In fcc structure octahedral voids are present at:
(a) edge centers (b) face centers (c) body centers (d) corners

4. If the radius of Na is 95 pm and that of Cl ions is 181 pm then:
(a) co-ordination no. of Na is 6 (b) co-ordination no. of Cl is 8
(c) length of the unit cell is 552 pm (d) length of the unit cell is 380 pm

5. Select the correct statement(s):
(a) The co-ordination number of each type of ion in a CsCl crystal is twelve
(b) A metal that crystallizes in a bcc structure has a co-ordination number of twelve
(c) A unit cell of an ionic crystal shares some of its ions with other unit cells
(d) The length of the unit cell in NaCl in 552 pm (given that r_{Na} 85 pm and r_{Cl} 181 pm)

6. The hcp and ccp structure for a given element would be expected to have:
(a) the same co-ordination number (b) the same density
(c) the same packing fraction (d) all of the above

7. Select the correct statement(s) for the rock-salt structure (NaCl) :
(a) The tetrahedral voids are smaller than the octahedral sites
(b) The octahedral voids are occupied by cations and the tetrahedral sites are empty
(c) The radius ratio (r_c/r_a) is 0.225
(d) The radius ratio (r_c/r_a) is 0.732

8. Select the correct statement(s):
(a) The ionic crystal of AgBr may have Schottky defect
(b) The unit cell having crystal parameters, a b c, 90 , 120 is hexagonal
(c) In ionic compounds having Frenkel defect the ratio r $/r$ is high
(d) The co-ordination number of Na ion in NaCl is 6

9. Which of the following represents the closest packed arrangement of uniform solid spheres:
(a) simple cubic unit cell (b) body centered cubic unit cell
(c) face centered cubic unit cell (d) hcp unit cell

For More Material Join: @JEEAdvanced_2024

10. Select the correct statement(s):
(a) A cubic system possesses a total of 23 elements of symmetry
(b) A cubic system contains centre of symmetry, planes of symmetry as well as axes of symmetry
(c) For triclinic system a b c and 90
(d) The total no. of Bravais space lattice belonging to all the seven crystals are 14

11. Select the correct statement(s):
(a) Co-ordination no. of Cs and Cl are 8, 8 in CsCl crystal
(b) If radius ratio (r_c / r_a) 0.225 then shape of compound must be linear
(c) If radius ratio (r_c / r_a) lies between 0.414 to 0.732 then shape of ionic compound may be square planer (Ex. $PtCl_4^{2}$)
(d) If radius ratio is less than 0.155 then shape of compound is linear

12. Select the correct statement(s):
(a) CsCl changes to NaCl structure on heating
(b) NaCl changes to CsCl structure on applying pressure
(c) Co-ordination number decreases on applying pressure
(d) Co-ordination number increases on heating

13. Select the correct statement(s):
(a) A NaCl type AB crystal lattice can be interpreted to be made up of two individual fcc type unit lattice of A and B fused together in such a manner that the corner of one unit lattice becomes the edge centre of the other
(b) In a fcc unit cell, the body center is an octahedral void
(c) In an scc lattice, there can be no octahedral void
(d) In an scc lattice, the body center is the octahedral void

14. In a AB unit cell (Rock salt type) assuming A forming fcc:
(a) The nearest neighbour of A is $6B$ ion (b) The nearest neighbour of B is $6A$ ion
(c) The second neighbour of A is $12A$ (d) The packing fraction of AB crystal is $\frac{\sqrt{3}}{8}$

15. Amorphous solids:
(a) do not have sharp melting points
(b) are isotropic
(c) have same physical properties in all directions
(d) are supercooled liquids

16. Which is/are correct statement about zinc blende structure ?
(a) The number of first neighbours of S^{2} is 4
(b) The maximum distance between Zn^{2} is $\frac{a\sqrt{3}}{2}$, where 'a' edge length of unit cell
(c) If all tetrahedral voids are occupied by Zn^{2} then C.N. of S^{2} is 8.
(d) If all tetrahedral voids are occupied by Zn^{2} then C.N. change from 4 : 4 to 8 : 8.

17. Compound X_2Y have antifluorite structure. Which is/are correct statement ?
(a) The minimum distance between X is $\frac{a}{2}$, where 'a' edge length of unit cell

For More Material Join: @JEEAdvanced_2024

(b) The co-ordination number ratio of x and y is 8 : 4
(c) If X removed from alternate tetrahedral void then CN is 4 : 4.
(d) If X removed from alternate tetrahedral void then CN is 4 : 8

18. Which is/are correct statement ?
(a) Packing fraction in 2D-hcp is 0.785
(b) Packing fraction in *AAA* ..... is 0.52
(c) Packing fraction in *ABAB* ..... is 0.74
(d) Void fraction in *ABCABC* ..... is 0.26

19. Which is/are correct statement ?
(a) In simple cubic close packed arrangement no octahedral void is present at edge centre.
(b) In fcc unit cell octahedral void and tetrahedral void are vacant.
(c) Packing fraction : simple cubic unit cell < bcc unit cell < fcc unit cell
(d) Size of void: cubic void > octahedral void > tetrahedral void

20. Given : Radius of A^2 100 pm ; Radius of C 240 pm ; Radius of B^2 300 pm ; Radius of D 480 pm Which is/are correct statement ?
(a) Coordination number of A^2 in comp *AB* is 4
(b) Coordination number of A^2 in comp *AB* is 6.
(c) Coordination number of C in comp *CD* is 6
(d) Coordination number of C in comp *CD* is 8.

Match the Column

Column-I and Column-II contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

1.

Column-I	Column-II
(A) Tetragonal and Hexagonal	(P) are two crystal systems
(B) Cubic and Rhombohedral	(Q)
(C) Monoclinic and Triclinic	(R) a b c
(D) Cubic and Orthorhombic	(S) a b c

2.

Column-I	Column-II
(A) If radius ratio x $\frac{r_c}{r_a}$ 0.155	(P) Co-ordination no. is 8
(B) If 0.225 x 0.414	(Q) Co-ordination no. is 4
(C) If 0.414 x 0.732	(R) Co-ordination no. is 6
(D) If 0.732 x 1	(S) Co-ordination no. is 2

For More Material Join: @JEEAdvanced_2024

3.

Column-I (Shape of Compound)	Column-II (Co-ordination No.)
(A) Linear	(P) 6
(B) Triangular planar	(Q) 4
(C) Square planar	(R) 2
(D) Octahedral	(S) 3

4.

Column-I	Column-II
(A) Rock salt structure	(P) general formula is AB
(B) Zinc blende structure	(Q) general formula is AB_3
(C) Fluorite structure	(R) general formula is A_2B
(D) Anti fluorite structure	(S) general formula is AB_2

5.




Column-I	Column-II
(A) Co-ordination no. of Ca^2 and F^- in fluorite structure	(P) 8, 4
(B) C.No. of Zn^2 and S^{2-} in zinc blende structure	(Q) 8, 8
(C) C.No. of Cs and Cl^- in CsCl (bcc type) structure	(R) 4, 8
(D) C.No. of Li and O^{2-} in antifluorite structure	(S) 4, 4

6.

Column-I **[Bravais Lattice(s)]**	Column-II **(Crystal System)**
(A) Primitive, face centered, body centered, end centered	(P) Cubic
(B) Primitive, face centered, body centered	(Q) Orthorhombic
(C) Primitive, body centered	(R) Hexagonal
(D) Primitive only	(S) Tetragonal

For More Material Join: @JEEAdvanced_2024

7.

Column-I	Column-II
(A) Magnetic moment in a paramagnetic substance	(P) 
(B) Magnetic moment in a ferromagnetic	(Q) 
(C) Magnetic moment in a antiferromagnetic	(R) 
(D) Magnetic moment in a ferrimagnetic	(S) 

8.

Column-I (Structure)	Column-II (Edge length of unit cell)
(A) Rock salt (NaCl)	(P) $a \quad (r_{Cation} \quad r_{Anion})$
(B) CsCl	(Q) $a \quad \frac{4}{\sqrt{3}}(r_{Cation} \quad r_{Anion})$
(C) Zinc blende (ZnS)	(R) $a \quad \frac{2}{\sqrt{3}}(r_{Cation} \quad r_{Anion})$
(D) Anti fluorite (Na_2O)	(S) $a \quad 2(r_{Cation} \quad r_{Anion})$

9.





Column-I (Ideal structure)	Column-II (Packing fraction)
(A) Rock salt	(P) 0.729
(B) Zinc blende	(Q) 0.756
(C) Fluorite	(R) 0.793
(D) CsCl	(S) 0.748

10.

Column-I (Structure)	Column-II (Voids occupied)
(A) Rock salt	(P) 100% tetrahedral voids occupied by cation
(B) Zinc blende	(Q) 100% tetrahedral voids occupied by anion
(C) Fluorite	(R) 100% octahedral voids occupied by cation
(D) Anti fluorite (Na_2O)	(S) 50% tetrahedral voids occupied by cation

For More Material Join: @JEEAdvanced_2024

11.

Column-I (Void representation)	Column-II (Name of void)
(A) 	(P) Tetrahedral void
(B) 	(Q) Octahedral void
(C) 	(R) Cubic void
(D) 	(S) Triangular void

Assertion-Reason Type Questions

Each question contains STATEMENT-1 (Assertion) and STATEMENT-2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below:

(A) If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1

(B) If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1

(C) If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

(D) If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

1. **STATEMENT-1 :** In any ionic solid [*MX*] with Schottky defects, the number of positive and negative ions are the same.

STATEMENT-2 : Equal number of cation and anion vacancies are present.

2. **STATEMENT 1 :** Amorphous solids are isotropic.

STATEMENT 2 : Amorphous solids lack a regular three-dimensional arrangement of atoms.

For More Material Join: @JEEAdvanced_2024

3. **STATEMENT-1 :** Diamond is a covalent solid.
   STATEMENT-2 : The co-ordination number of each carbon atom in diamond is 4.
4. **STATEMENT-1 :** In NaCl structure, Na ions occupy octahedral holes and Cl ions occupy ccp.
   STATEMENT-2 : The distance of the nearest neighbours in NaCl structure is $a/2$ where a is the edge length of the cube.
5. **STATEMENT-1 :** For fluorite structure, the F ions occupy tetrahedral void and Ca^{2} ions in ccp
   STATEMENT-2 : The radius ratio of fluorite structure is 0.414.
6. **STATEMENT-1 :** In the body centered cubic type structure of CsCl the arrangement of Cs is present in central octahedral void.
   STATEMENT-2 : In CsCl, the Cs ions are at body centered position and Cl are at corners.
7. **STATEMENT-1 :** Electrical conductivity of semi-conductors increases with increasing temperature.
   STATEMENT-2 : With increase in temperature, number of electrons which jumps from the valence bond to the conduction band in semi-conductors increase.
8. **STATEMENT-1 :** FeO is non-stoichiometric with formula $Fe_{0.95}O$.
   STATEMENT-2 : Some Fe^{2} ions are replaced by Fe^{3} as $3Fe^{2}$ $2Fe^{3}$ to maintain electrical neutrality.
9. **STATEMENT-1 :** The number of tetrahedral voids is double the number of octahedral voids.
   STATEMENT-2 : The size of the tetrahedral void is half of that of the octahedral void.
10. **STATEMENT-1 :** Due to Frenkel defect there is no effect on density of a solid.
    STATEMENT-2 : Ions shift from lattice sites to interstitial sites in Frenkel defect.
11. **STATEMENT-1 :** Antiferromagnetic substances become paramagnetic on heating to high temperature.
    STATEMENT-2 : Heating results in spins of electrons becoming random.

Subjective Problems



1. In seven possible crystal system how many crystal system have ?
2. In seven possible crystal system how many crystal system have more than one Bravais lattice ?
3. How many effective no. of atoms present in bcc unit cell ?
4. In solid X atoms goes to corner of the cube and two alternate face centre. Calculate effective number of atom of X in unit cell ?
5. Find the distance (in pm) between the body centered atom and one corner atom in an element (a 2.32 pm)

For More Material Join: @JEEAdvanced_2024

6. The structure of MgO is similar to NaCl. What is the co-ordination number of Mg ?

7. How many number of formula units in a unit cell of CaF_2 ?

8. What is the co-ordination number of sodium in Na_2O ?

9. Density of Li crystal is $0.53\ g/cm^3$. The edge length of Li unit-call is 3.5 Å. Find out the number of Li atoms in a unit cell. (N_A 6.0 10^{23} mol ^{1}, M 6.94 g mol ^{1})

10. What is the co-ordination number of Cl in CsCl structure ?

11. In cubic system how many atoms arrangement exist in nature ?

12. The ionic radii of A and B are 1.7 Å and 1.8 Å respectively. Find the co-ordination number of A .

13. If edge fraction unoccupied in ideal anti-fluorite structure is x. Calculate the value of Z. Where

Z $\frac{x}{0.097}$

14. Ionic solid Na A crystallise in rock salt type structure. 2.592 gm of ionic solid salt NaA dissolved in water to make 2 litre solution. The pH of this solution is 8. If distance between cation and anion is 300 pm, calculate density of ionic solid (in gm/cm^3).

(Given : pK_w 13, pK_a(HA) 5, N_A 6 10^{23})

15. Calculate the value of $\frac{Z}{10}$. Where

Z Co-ordination number of 2D-square close packing

Co-ordination number of 2D-hcp

Co-ordination number of 3D-square close packing

Co-ordination number of 3D, *ABCABC* ..... Packing

Co-ordination number of 3D, *ABAB* ..... packing

For More Material Join: @JEEAdvanced_2024

Answers

Level 1

1.	(c)	**2.**	(b)	**3.**	(d)	**4.**	(b)	**5.**	(a)	**6.**	(c)	**7.**	(a)	**8.**	(d)	**9.**	(c)	**10.**	(c)
11.	(d)	**12.**	(c)	**13.**	(d)	**14.**	(b)	**15.**	(b)	**16.**	(a)	**17.**	(b)	**18.**	(c)	**19.**	(a)	**20.**	(b)
21.	(d)	**22.**	(c)	**23.**	(b)	**24.**	(c)	**25.**	(d)	**26.**	(c)	**27.**	(c)	**28.**	(c)	**29.**	(c)	**30.**	(b)
31.	(c)	**32.**	(a)	**33.**	(d)	**34.**	(c)	**35.**	(b)	**36.**	(b)	**37.**	(c)	**38.**	(a)	**39.**	(a)	**40.**	(d)
41.	(d)	**42.**	(d)	**43.**	(b)	**44.**	(d)	**45.**	(c)	**46.**	(d)	**47.**	(c)	**48.**	(b)	**49.**	(a)	**50.**	(b)
51.	(b)	**52.**	(b)	**53.**	(c)	**54.**	(a)	**55.**	(c)	**56.**	(b)	**57.**	(a)	**58.**	(d)	**59.**	(c)	**60.**	(b)
61.	(d)	**62.**	(c)	**63.**	(a)	**64.**	(a)	**65.**	(b)	**66.**	(a)	**67.**	(b)	**68.**	(c)	**69.**	(b)	**70.**	(b)
71.	(b)	**72.**	(c)	**73.**	(b)	**74.**	(d)	**75.**	(a)	**76.**	(d)	**77.**	(c)	**78.**	(c)	**79.**	(a)	**80.**	(a)
81.	(c)	**82.**	(c)	**83.**	(b)	**84.**	(c)	**85.**	(a)	**86.**	(c)	**87.**	(c)	**88.**	(d)	**89.**	(d)	**90.**	(b)
91.	(b)	**92.**	(c)	**93.**	(c)	**94.**	(a)	**95.**	(b)	**96.**	(d)	**97.**	(c)	**98.**	(d)	**99.**	(a)	**100.**	(b)
101.	(c)	**102.**	(d)	**103.**	(d)	**104.**	(c)	**105.**	(a)	**106.**	(d)	**107.**	(b)	**108.**	(d)	**109.**	(d)	**110.**	(c)
111.	(c)	**112.**	(c)	**113.**	(b)	**114.**	(c)	**115.**	(b)	**116.**	(c)	**117.**	(a)	**118.**	(c)	**119.**	(d)	**120.**	(b)
121.	(b)	**122.**	(a)	**123.**	(b)	**124.**	(c)	**125.**	(d)										

Level 2

1.	(b)	**2.**	(c)	**3.**	(c)	**4.**	(c)	**5.**	(d)	**6.**	(c)	**7.**	(b)	**8.**	(d)	**9.**	(a)	**10.**	(c)
11.	(b)	**12.**	(b)	**13.**	(c)	**14.**	(d)	**15.**	(c)	**16.**	(a)	**17.**	(b)	**18.**	(a)	**19.**	(a)	**20.**	(b)
21.	(a)	**22.**	(a)	**23.**	(d)	**24.**	(d)	**25.**	(b)	**26.**	(a)	**27.**	(b)	**28.**	(b)	**29.**	(d)	**30.**	(a)
31.	(b)	**32.**	(b)																

For More Material Join: @JEEAdvanced_2024

Level 3

Passage–1:	**1.**	(c)	**2.**	(a)				
Passage–2:	**1.**	(a)	**2.**	(a)	**3.**	(c)		
Passage–3:	**1.**	(b)	**2.**	(b)	**3.**	(a)	**4.**	(a)
Passage–4:	**1.**	(b)	**2.**	(b)	**3.**	(d)		
Passage–5:	**1.**	(b)	**2.**	(b)	**3.**	(a)		
Passage–6:	**1.**	(d)	**2.**	(a)				
Passage–7:	**1.**	(a)	**2.**	(d)	**3.**	(d)		

One or More Answers is/are correct

1.	(a,d)	**2.**	(b,c)	**3.**	(a,c)	**4.**	(a,c)	**5.**	(c)	**6.**	(a,c)	**7.**	(a,b)	**8.**	(a,b,d)
9.	(c,d)	**10.**	(a,b,c,d)	**11.**	(a,c,d)	**12.**	(a,b)	**13.**	(a,b,c)	**14.**	(a,b,c)	**15.**	(a,b,c,d)	**16.**	(a,b,c)
17.	(a,c)	**18.**	(a,b,c,d)	**19.**	(a,b,c,d)	**20.**	(a,c)								

Match the Column

1.	A	P;	B	P, S;	C	P, R;	D	P, Q
2.	A	S;	B	Q;	C	Q, R;	D	P
3.	A	R;	B	S;	C	Q;	D	P
4.	A	P;	B	P;	C	S;	D	R
5.	A	P;	B	S;	C	Q;	D	R
6.	A	Q;	B	P;	C	S;	D	R
7.	A	P;	B	R;	C	Q;	D	S
8.	A	S;	B	R;	C	Q;	D	Q
9.	A	R;	B	S;	C	Q;	D	P
10.	A	R;	B	S;	C	Q;	D	P
11.	A	Q;	B	Q;	C	P;	D	R

Assertion-Reason Type Questions

1. (A)	**2.** (A)	**3.** (B)	**4.** (B)	**5.** (C)	**6.** (D)	**7.** (A)	**8.** (A)	**9.** (C)	**10.** (A)
11. (A)									

Subjective Problems

1.	5	**2.**	4	**3.**	2	**4.**	2	**5.**	2	**6.**	6	**7.**	4	**8.**	4	**9.**	2	**10.**	8
11.	3	**12.**	8	**13.**	3	**14.**	4	**15.**	4										

For More Material Join: @JEEAdvanced_2024

Hints and Solutions

Level 1

13. (d) Orthorhombic contains
Simple, Body centre, Face centre & End centre bravais lattices.

14. (b) Cubic-symmetrical
$90^\circ, a = b = c$
Triclinic-unsymmetrical
$90^\circ, a \neq b \neq c$

15. (b) Tetragonal $a = b \neq c$, 90°

19. (a) Simple cubic cell "Z" $8 \times \frac{1}{8} = 1$
Face centred cubic unit cell
$Z = 8 \times \frac{1}{8} + 6 \times \frac{1}{2} = 4$

20. (b) Effective no. of atoms in End centred unit cell
$8 \times \frac{1}{8} + 2 \times \frac{1}{2} = 2$
atoms are present at corners as well as at end face's centre of unit cell.

21. (d) Simple cubic $a = 2R$ Body centred cubic $\sqrt{3}a = 4R$

22. (c) Face centred cubic $\sqrt{2}a = 4R$

23. (b) Packing fraction (simple cubic) $\frac{1 \times \frac{4}{3} R^3}{a^3}$
$\frac{}{6} = 0.52$
"$a = 2R$"

24. (c) Body centred cubic $\frac{2 \times \frac{4}{3} R^3}{a^3} = \frac{\sqrt{3}}{8}$
$\sqrt{3}a = 4R$

25. (d) Face centred cubic $\frac{4 \times \frac{4}{3} R^3}{a^3} = 0.74$
$\sqrt{2}a = 4R$

26, 27. (c, c)
Packing efficiency
In simple cubic $= 0.52$
Body centred cubic $= 0.68$
Face centred cubic $= 0.74$

28. (c) $a = 2R$
$R = \frac{0.236}{2} = 0.118$ nm

29. (c) $\sqrt{3}a = 4R$

30. (b) $\sqrt{2}a = 4R$

31. (c) $\sqrt{2}a = 4R$
$a = 2\sqrt{2}R$

32. (a) $Z = 8 \times \frac{1}{8} = 1$
Nearest neighbour $= 6$

33. (d) $Z = 8 \times \frac{1}{8} + 6 \times \frac{1}{2} = 4$
Nearest neighbour $= 12$

34. (c) $Z = 8 \times \frac{1}{8} = 1 = 2$
Nearest neighbour $= 8$

35. (b) $D = \frac{ZM}{N_A a^3} = 6.25 = \frac{Z \times 12}{6.02 \times 10^{23} \times (4 \times 10^{-8})^3}$
$Z = 2$ body centred

36. (b) $D = \frac{ZM}{N_A a^3}$ $Z = 2$

37. (c) $D = \frac{ZM}{N_A a^3} = \frac{4 \times 40}{6.02 \times 10^{23} \times a^3} = 1.83$
$a = 0.525$ nm

38. (a) $D = \frac{ZM}{N_A a^3}$ $Z = 4$
$\sqrt{2}a = 4R$

39. (a) Refer Q.no. 38

40. (d) Refer Q.no. 38

41. (d) $D = \frac{ZM}{N_A a^3}$ $Z = 2$

42. (d) Refer Q.no. 38

43. (b) Refer Q.no. 38

44. (d) Mole of 'x' $= \frac{8}{80} = 0.1$
No. of atoms $= 0.1 N_A$
No. of unit cell $= \frac{0.1 N_A}{4} = 0.025 N_A$

For More Material Join: @JEEAdvanced_2024

45. (c) $D \quad \frac{ZM_w}{N_A a^3}$

$Z \quad 2, \; M_w \quad 96, \; D \quad 10.3 \text{ g/cm}^3$

$\sqrt{3}a \quad 4R \quad \frac{\sqrt{3}\,a}{4}$

46. (d) Coordination no. of

simple cubic 6

Face centred cubic 12

Body centred cubic 8

47. (c) F.C.C. unit cell *ABCABCABC* .....

packing efficiency 74 %

50. (b) C.N. of $A \quad x(B)$

C.N. of $B \quad y(A)$

Formula A_yB_x

51. (b) $D \quad \frac{ZM}{N_A a^3}$

$Z \quad 4, \; a \quad 2\sqrt{2}R$

52. (b) $D \quad \frac{ZM}{N_A a^3}$

$Z \quad 4, \; a \quad 3.5 \quad 10^{-8}$ cm, $Mw \quad 59$

54. (a) Packing efficiency of HCP & CCP is same.

56. (b) $Z \quad 12 \quad \frac{1}{6} \quad 2 \quad \frac{1}{2} \quad 3 \quad 6$

58. (d) No. of octahedral hole in a unit cell

effective no. of atoms per unit cell

Z

No. of tetrahedral holes 2 octahedral holes.

59. (c) If sheet is repeated than packing is poor.

60. (b) Refer Q. no. 58

62. (c) Tetrahedral & octahedral holes are present in hcp & ccp.

63. (a) Refer Q. no. 62

67. (b) $\frac{r_c}{r_a} \quad 0.225$ tetrahedral holes

68. (c) $\frac{r_c}{r_a} \quad 0.414$ octahedral holes

69. (b) an octahedral holes is formed by 6 sphere.

70. (b) No. of nearest neighbour 8 (All body centre atom w.r.t. corner atom)

No. of next nearest neighbour 6 (No. of corner atoms along edge w.r.t. to any corner).

71. (b) Refer Q. no. 67, 68

72. (c) Refer Q. no. 68

73. (b) Octahedral sites in fcc are present at each edge centre as well as at body centre.

74. (d) In Na_2O all tetrahedral void are occupied.

75. (a) $\frac{r_c}{r_a} \quad 0.732$

$r_c \quad 0.732 \quad 200 \quad 146.4$ pm

76. (d) for cubic void, $\frac{r_c}{r_a} \quad 0.732$

77. (c) $\frac{r_c}{r_a} \quad \frac{0.066}{0.140} \quad 0.47$

So, cation (Mg^{2}) fitted in octahedral holes of O^{2} are present at fcc lattice point.

78. (c) In diamond C-atoms are present in fcc lattice points as well as in alternate tetrahedral void.

79. (a) Refer Q. no. 78

80. (a) For MgO, $\frac{r_c}{r_a} \quad \frac{0.65}{1.4} \quad 0.47$

Mg^{2} is in octahedral void so C.N. 6

For MgS, $\frac{r_c}{r_a} \quad \frac{0.65}{1.84}$

Mg^{2} is in tetrahedral void so C.N. 4

81. (c) No. of $A \quad 7 \quad \frac{1}{8} \quad \frac{7}{8}$

No. of $B \quad 6 \quad \frac{1}{2} \quad 3$

$A_{7/8}B_3 \quad A_7B_{24}$

82. (c) No. of $S^{2} \quad 8 \quad \frac{1}{8} \quad 6 \quad \frac{1}{2} \quad 4$

No. of $Ca^{2} \quad \frac{1}{2} \quad 8 \quad 4$

83. (b) No. of $A^{2} \quad \frac{1}{8} \quad 8 \quad 1$

No. of $B^{3} \quad \frac{1}{2} \quad 4 \quad 2$

No. of $O^{2} \quad 8 \quad \frac{1}{8} \quad 6 \quad \frac{1}{2} \quad 4$

(AB_2O_4)

84. (c) No. of $C \quad 6 \quad \frac{2}{3}$

No. of $A \quad 6$

(C_2A_3)

85. (a) No. of $O^{2} \quad 8 \quad \frac{1}{8} \quad 6 \quad \frac{1}{2} \quad 4$

For More Material Join: @JEEAdvanced_2024

No. of A $\frac{1}{8}$ 8 1

No. of B $\frac{1}{4}$ 4 1

86. (c) No. of A 8 $\frac{1}{8}$ 1

No. of B 5 $\frac{1}{2}$ $\frac{5}{2}$

A_2B_5

87. (c) No. of Cu 8 $\frac{1}{8}$ 6 $\frac{1}{2}$ 4

No. of Ag 12 $\frac{1}{4}$ 3

No. of Au 1

$Cu_4Ag_3Au_1$

88. (d) In NaCl structure

Cl ions are present at fcc lattice point


Na ions are present in all octahedral void.

89. (d) In B.C.C. ionic structure

Cation is present in cubic void

Anions are present at each corner of cubic

90. (b) In ZnS, S^2 ions are present in F.C.C. Lattice point Zn^2 ions are present in alternate tetrahedral void.

91. (b) "edge" 

r r $\frac{a}{2}$

92. (c) In ideal arrangement of anions

$\sqrt{2}a$ $4r_a$

r_a $\frac{a}{2\sqrt{2}}$

93. (c) r_c r_a $\frac{\sqrt{3}a}{2}$

Anion are at corners and cations are at body centre.

94. (a) Distance b/w two nearest Cl $\frac{a}{\sqrt{2}}$

Distance b/w two nearest Na $\frac{a}{\sqrt{2}}$

95. (b) w.r.t. body centre Na ion

all 12 Na ions of edge centre.

96. (d) $\frac{r}{r}$ $\frac{1}{2}$ 0.5 F.C.C. lattice point

Cations are at octahedral void

(r r) $\frac{a}{2}$ 3

a 6

volume $(6)^3$ 216 pm^3

97. (c) In CaF_2 cations are present at F.C.C. lattice points and anions are at all tetrahedral void, antiflourites structure is reverse arrangement of CaF_2.

98. (d) Refer Q. no. 89.

99. (a) $\frac{r_c}{r_a}$ $\frac{94}{146}$ 0.64

100. (b) r_c r_a $\frac{a}{2}$

102. (d) $\frac{r_A}{r_B}$ 0.225

r_A 45

103. (d) In Na_2O O^2 are at F.C.C. lattice point

Na are at all tetrahedral void

C.N. of O^2 8

104 (c) r_c r_A $\frac{\sqrt{3}a}{2}$

a $\frac{2}{\sqrt{3}}$ [1.69 1.81]

105. (a) r_c r_a $\frac{\sqrt{3}a}{2}$

106. (d) r_c r_a $\frac{a}{2}$

107. (b) r_c r_a $\frac{\sqrt{3}a}{2}$

108. (d) r_c r_a $\frac{\sqrt{3}a}{4}$

110. (c) Refer Q. no. 78

111. (c) Due to missing of ions. Density decreases in Schottky defect.

112. (c) In Frenkel defect ions are shifted in its interstitial site but density remains unchanged.

113. (b) Refer Q.no. 112

115. (b) Alkali metal halide shows schottky defect.

116. (c) Two Ag ions will replaced by one Cd^2, so there is one vacancy for each Cd^2

117. (a) In Schottky defect ions are missing from their position and in Frenkel defect ions are shifted in their interstitial site from their lattice point.

For More Material Join: @JEEAdvanced_2024

118. (c) Refer Q. no. 115.

119. (d) Density of crystal always increases due to substitutional impurity defect.

120. (b) $\frac{\sqrt{3}a}{4}$ $(2R)$

121. (b) No. of vacancy No. of Sr^{2}

No of Sr^{2} in 1 mole NaCl 10^{-7} N_A

122. (a) Fe^{3} Fe^{2}

$x\%$ $(100 - x)\%$

$\frac{200}{93}$ $\frac{3x}{100}$ $\frac{2(100 - x)}{100}$

$\%Fe^{3}$ 15.05%

123. (b) Cu^{2} Cu

$x\%$ $(100 - x)\%$

$\frac{20}{18}$ $\frac{2x}{100}$ $\frac{(100 - x)}{100}$

x 11.11

Level 2

1. (b) $_1$ $\frac{2 \quad 56}{\left(\frac{4r}{\sqrt{3}}\right)^{3}}$; $_2$ $\frac{4 \quad 56}{(2\sqrt{2}r)^{3}}$

$\frac{1}{2}$ 0.918

2. (c) 2.32 $\frac{2 \quad M}{6 \quad 10^{23} (1.22)^{3} \quad 10^{-21}}$

M 1264 x 47

3. (c) Edge not covered by atom a $2r$

also in bcc, $4r$ $\sqrt{3}a$

Therefore, edge not covered

a $\frac{\sqrt{3}}{2}a$ a $\frac{2 - \sqrt{3}}{2}$

Fraction of edge not covered by atoms

$\frac{2 - \sqrt{3}}{2}$ 0.134

4. (c) In a simple cubic crystal, a $2r$

Let the radius of the interstitial atom in simple cubic structure be x, then

$2(r + x)$ $\sqrt{3}a$ $\sqrt{3}$ $2r$

So, x $\sqrt{3}r$ r $0.732\,r$

Packing fraction $\frac{\frac{4}{3} r^{3} \quad \frac{4}{3} (0.732\,r)^{3}}{(2r)^{3}}$

$\frac{\frac{4}{3} r^{3} [1 \quad (0.732)^{3}]}{8r^{3}}$

$\frac{}{6}[1 \quad (0.732)^{3}]$ 0.73

5. (d) Given a $2r$ 60.3 and for bcc, $4r$ $\sqrt{3}a$

a $\frac{\sqrt{3}}{2}a$ 60.3 a 450 pm

Density ( ) $\frac{2 \quad 48}{6.023 \quad 10^{23} \quad (4.5)^{3} \quad 10^{-24}}$

1.75 g/cc

6. (c) a $2r$ 53.6 pm …(i)

also $4r$ $\sqrt{3}a$

a $\frac{\sqrt{3}}{2}a$ 53.6

a $\frac{53.6 \quad 2}{2 \quad \sqrt{3}}$ 400 pm

Density ( ) $\frac{2 \quad 23}{6.023 \quad 10^{23} \quad 4^{3} \quad 10^{-24}}$

1.19 g/cc

7. (b) Vol. of all atoms in 1.68 g of Ar

$\frac{1.68}{40}$ 6 10^{23} $\frac{4}{3}$ $\frac{22}{7}$ $(1.50^{3} \quad 10^{-8})$

0.3564

Vol. of solid Ar 1 cm^{3}

% of empty space (1 0.3564) 100

64.36

8. (d) Effective no. of atoms of B present in a unit cell 2

Total volume of B unoccupied by A in a unit cell

$2 \quad \frac{4}{3}(R^{3} \quad r^{3})$

$\frac{7 \quad R^{3}}{3}$ $\because r \quad \frac{R}{2}$


Volume of unit cell a^{3}

$\left(\frac{4R}{\sqrt{3}}\right)^{3}$ $\frac{64}{3\sqrt{3}} R^{3}$ $(\because \sqrt{3}a \quad 4R)$

Desired ratio $\frac{\frac{7 \quad R^{3}}{3}}{\frac{64}{3\sqrt{3}} R^{3}}$ $\frac{7 \quad \sqrt{3}}{64}$

For More Material Join: @JEEAdvanced_2024

9. (a)




AB a (nearest)

BD $\sqrt{2}a$ (next-nearest)

CE $\sqrt{3}a$ (next-next-nearest)

10. (c)



AB $\frac{\sqrt{3}a}{2}$ (nearest)

AC a (next-nearest)

CD $\sqrt{2}a$ (next-next-nearest)

11. (b) For atoms of A $8 \quad \frac{1}{8} \quad 1$

For atoms of B $8 \quad \frac{1}{4} \quad 2$

For atoms of C $4 \quad \frac{1}{2} \quad 1 \quad 3$

12. (b) Let volume of fcc unit cell V

$$_A \quad \frac{4 \quad M_A}{N_A . V}$$


$$_B \quad \frac{8 \quad M_B}{N_A . V}$$

$$\frac{_A}{_B} \quad \frac{M_A}{2M_B} \quad \frac{30}{2 \quad 50} \quad 0.3$$

$$_B \quad \frac{_A}{0.3} \quad 3.33 \text{ g/cc}$$

Total density $_A \quad _B \quad 4.33$ g/cc

13. (c)



$$\frac{BE}{BC} \quad \cos 30 \quad \frac{\sqrt{3}}{2}$$

Distance between the centres of two B atoms

$2 \quad BE \quad \sqrt{3}$ Å

14. (d) 4 formula units of AB are present in a unit cell

no. of unit cell $\frac{4}{40} \quad \frac{N_A}{4}$

$0.025 \; N_A$

15. (c) Use $\frac{Z.M}{N_A . a^3}$

$Z \quad 4; \quad M \quad 40 \quad 2 \quad 19$

16. (a) $a \quad 2(r \quad r)$

$2 \quad 297$ pm

Volume of unit cell a^3

$209.6 \quad 10^{-24}$ cm^3

17. (b) $8.27 \quad \frac{4 \quad 128}{6.023 \quad 10^{23} \; a^3}$

$a \quad 46.8 \quad 10^{-9}$ cm $\quad 4.68$ Å

$2 \, [r_{O^{2-}} \quad r_{Cd^{2+}}]$

$r_{Cd^{2+}} \quad 1.1$ Å

18. (a) $\frac{r_{Na^+} \quad r_{Cl^-}}{r_{Cl^-}} \quad 1.5$

and $\frac{r_{K^+}}{r_{Cl^-}} \quad \frac{5}{7}$ or $\frac{r_{K^+} \quad r_{Cl^-}}{r_{Cl^-}} \quad \frac{12}{7}$

$$\frac{a_1}{a_2} \quad \frac{2(r_{K^+} \quad r_{Cl^-})}{2(r_{Na^+} \quad _{Cl^-})}$$

$$\frac{12}{7} \quad \frac{1}{1.5} \quad 1.143$$

19. (a) Mass of unit cell $d \quad V$

$(5 \quad 10^{-8})^3 \quad 3.84 \quad 6 \quad 10^{23}$

288 a.m.u.

Mass of formula unit of FeO $56 \quad 16 \quad 72$ a.m.u.

no. of formula unit $\frac{288}{72} \quad 4$

20. (b) Density of crystals decrease due to Schottky defects effective no. of atoms per unit cell in given lattice

$$4 \quad 1 \quad \frac{0.25}{100} \quad 3.99$$

$$\frac{50 \quad 3.99}{6 \quad 10^{23} \, (0.50 \quad 10^{-7})^3}$$

$$\frac{50 \quad 3.99}{6 \quad 125 \quad 0.1}$$

2.66 g/cm^3

For More Material Join: @JEEAdvanced_2024

21. (a) Theoretical density $\frac{Z.M}{N_A.a^3}$

$$\frac{4 \quad 24 \quad 10^{-3}}{6 \quad 10^{23} \quad (4 \quad 10^{-10})^3}$$

$$2.5 \quad 10^3 \text{ kg/m}^3$$

$$\% \text{ occupancy} \quad \frac{\text{Observed density}}{\text{Ideal density}} \quad 100$$

$$\frac{2.4 \quad 10^3}{2.5 \quad 10^3} \quad 100 \quad 96\%$$

23. (d) Octahedral void present at the centre of cube and tetrahedral void is present at $(1/4)^{th}$ of the distance along each body diagonal.

$\frac{\sqrt{3}a}{2}$ 2 distance between octahedral and tetrahedral void.

24. (d) Vol. of unit cell $a^2 \sin 60 \quad b$

$$173.2 \quad 10^{-24} \text{ cm}^3$$

a 5 Å

Mass of unit cell

$$173.2 \quad 10^{-24} \quad 5 \quad 6 \quad 10^{23}$$

519.6 g

No. of molecules present in given unit cell

$$\frac{519.6}{259.8} \quad 2$$

25. (b) In hcp arrangement total height

$$h \quad 4 \quad \sqrt{\frac{2}{3}}\, r$$

as per given

$$\frac{h}{2} \quad 2\sqrt{\frac{2}{3}}\, r \quad 3.35 \text{ Å} \quad 3.35 \quad 10^{-8} \text{ cm.}$$

Volume of unit cell $24\sqrt{2}r^3$

$$24\sqrt{2} \quad \sqrt{\frac{3}{2}} \quad \frac{1}{2} \quad 3.35 \quad 10^{-8} \quad ^3$$

$$2.93 \quad 10^{-22} \text{ cm}^3$$

Effective no. of atoms in hexagonal unit cell

6

$$\text{density} \quad \frac{6 \quad 12}{6.023 \quad 10^{23} \quad 2.93 \quad 10^{-22}}$$

0.41 g/cc

26. (a) No. of Na 4 1 3

No. of Cl $4 \quad 2 \quad \frac{1}{2} \quad 3$

27. (b) Along one body diagonal $2X$ atoms from 2 corners, one Y particle (at the centre of cube) will be removed.

So, effective no. of X particles in a unit cell

$$4 \quad 2 \quad \frac{1}{8} \quad \frac{15}{4}$$

and effective no. of Y particles in a unit cell

$$4 \quad 1 \quad 3$$

$$X : Y \quad \frac{15}{4} : 3$$

or 5 : 4

28. (b) Effective no. of Na $4 \quad 1 \quad 2 \quad \frac{1}{4} \quad \frac{5}{2}$

For Cl $4\sqrt{2}a \quad 4r$

P. F. $\frac{\text{Vol. of effective no. of cations and anions}}{\text{Vol. of unit cell}}$

29. (d) No. of X particles per unit cell 4

No. of Y particles 4

No. of Z particles 8

Along one body diagonal $2X$ atoms from 2 corners, one Y particle and $2Z$ particles will be removed.

So, effective no. of X particles in a unit cell

$$4 \quad \frac{1}{8} \quad 2 \quad \frac{15}{4}$$

Effective no. of Y particles in a unit cell

$$4 \quad 1 \quad 3$$

Effective no. of Z particles in a unit cell

$$8 \quad 2 \quad 6$$

$$X \quad : \quad Y \quad : \quad Z$$

$$\frac{15}{4} \quad : \quad 3 \quad : \quad 6$$

$$5 \quad : \quad 4 \quad : \quad 8$$

30. (a) Effective no. of A particles are removed

$$4 \quad \frac{1}{8} \quad 2 \quad \frac{1}{2} \quad \frac{3}{2}$$

Effective no. of A particles present in a unit cell

$$4 \quad \frac{3}{2} \quad \frac{5}{2}$$

Effective no. of B particles are removed

$$2 \quad \frac{1}{4} \quad 1 \quad \frac{3}{2}$$

For More Material Join: @JEEAdvanced_2024

Effective no. of B particles present in a unit cell

$$4 \quad \frac{3}{2} \quad \frac{5}{2}$$

$$A \quad : \quad B$$

$$\frac{5}{2} \quad : \quad \frac{5}{2} \quad \text{or} \quad 1 : 1$$

Level 3

Passage-1

1. (c) P.F. $\dfrac{2 \quad \frac{4}{3} \quad r^3}{\left(\frac{4r}{\sqrt{3}}\right)^3} \sim 0.68$

% of empty space $100 \quad 0.68 \quad 100 \quad 32$

2. (a) P.F. $\dfrac{4 \quad \frac{4}{3} \quad r^3}{\left(\frac{4r}{\sqrt{2}}\right)^3} \quad 0.7406$

Passage-2

1. (a) $10.6 \quad \dfrac{4 \quad 108}{a^3 \quad 6.023 \quad 10^{23}}$

$a \quad 40.7$ nm

2. (a) Mass of $12 \quad 10^{23}$ atoms $\quad 100$ gm

Mass of $6.022 \quad 10^{23}$ atom

$$\frac{100}{12 \quad 10^{23}} \quad 6.023 \quad 10^{23}$$

$$50.18$$

$$\frac{4 \quad 50.18 \quad 10}{6.023 \quad 10^{23} \quad (200 \quad 10^{-10})^3}$$

41.66 g/cm^3

3. (c) By Hit and Trial : For fcc $Z \quad 4$ (4 pairs of KBr)

$M \quad 39 \quad 80 \quad 119$

Passage-3

3. (a) 1 to 3

c.c.p. generated f.c.c., octahedral voids = effective number of atoms in a unit cell, tetrahedral voids $\quad 2 \quad$ effective no. of atoms in a unit cell.

4. (a) O^{2-} is replaced by $X^{8/3-}$ so formula of spinel is $MgAl_2X_3$ and deficiency by one anion.

Passage-6

1. (d) $D_{\text{Theo}} \quad \dfrac{ZM}{N_A \quad a^3}$

$$\frac{4 \quad 197}{6 \quad 10^{23} \quad (407 \quad 10^{-10})^3}$$

$$\frac{4 \quad 197}{6 \quad 10^{23} \quad 6.74 \quad 10^{-23}}$$

19.485 g/cm^3

2. (a) Distance between next neighbour $\quad a$

407 pm

Passage-7

1. (a) Effective no. of C-atoms per unit cell $\quad 8$

mass $\quad 12 \quad 8 \quad 96$ amu

2. (d) $\dfrac{\sqrt{3}a}{4} \quad B.L_{C-C}$

$D \quad \dfrac{4 \quad 154}{\sqrt{3}} \quad 355.66$ pm

3. (d) $D \quad \dfrac{ZM}{N_A a^3} \quad \dfrac{8 \quad 12}{6 \quad 10^{23} \quad (355.66 \quad 10^{-10})^3}$

3.56 g/cm^3

One or More Answers is/are Correct

4. (a, c)

$\dfrac{r_{Na}}{r_{Cl}} \quad \dfrac{95}{181} \quad 0.52$

Cl ion at F.C.C. lattice point

Na ion are at octahedral void

C.No. of NaCl $\quad 6 : 6$

$r_c \quad r_a \quad \dfrac{a}{2}$

7. (a, b)

Refer Q. no. 4

11. (a, c, d)

CsCl ionic solid is B.C.C.

12. (a, b)

On increasing temperature C.No.

C.No. change $8 : 8$ to $6 : 6$

On increasing pressure C. No.

14. (a, b, c)

Refer Q. no. 4

16. (a, b, c)

Refer Q. no. 90 (Level-1)

For More Material Join: @JEEAdvanced_2024

17. (a, c)

Refer Q. no. 97 (Level-1)

18. (a, b, c, d)

(A) 2*D*-hexagonal packing

Packing efficiency $\frac{R^2 \quad 6 \quad \frac{R^2}{3}}{6 \quad \frac{\sqrt{3}(2R)^3}{4}}$

(B) Simple cubic $\frac{1 \quad \frac{4}{3} \quad R^3}{a^3}$

$a \quad 2R$

H.C.P. $(3D) \quad \frac{6 \quad \frac{4}{3} \quad R^3}{24\sqrt{2}R^3}$

F.C.C. $(3D) \quad \frac{4 \quad \frac{4}{3} \quad R^3}{(2\sqrt{2}R)^3}$

20. (a, c)

$\frac{r_{A^2}}{r_{B^2}} \quad \frac{100}{300} \quad 0.333 \qquad \frac{r_C}{r_D} \quad \frac{240}{480} \quad 0.5$

B^2 are at F.C.C. D at F.C.C. lattice point lattice point

and A^2 ions are C in all octahedral void alternate

tetrahedral void

C.No. of AB 4 : 4 C.No. of CD 6 : 6

Subjective Problems

1. Cubic 90

Tetragonal 90

Orthorhombic 90

Rombohedral 90

Hexagonal 90

2. Cubic, tetragonal, orthorhombic, monoclinic

3. $Z \quad 8 \quad \frac{1}{8} \quad 1 \quad 2$

4. No. of x-atoms $\quad 8 \quad \frac{1}{8} \quad 2 \quad \frac{1}{2} \quad 2$

5. Distance b/w body centred and corner atom $\frac{\sqrt{3}a}{2}$

6. Mg^2 is in octahedral void

So its C.No. 6

7. No. Ca^2 per unit cell 4

No. of F per unit cell 8

No. of CaF_2 unit per unit cell 4

8. Na ions are present in tetrahedral void in Na_2O, So C.No. of Na 4

9. $D \quad \frac{ZM}{N_A a^3}$

$Z \quad 2$

10. Cl ions are at corners and Cs at body centre

So C.No. of Cl 8 Cs

11. Simple cubic

body centred cubic

face centred cubic

12. $\frac{r_A}{r_B} \quad \frac{1.7}{1.8} \quad 0.944$

B are at corner of cube A at body centre

C.No. of A 8

13. Fraction of edge unoccupied $\frac{a \quad 2R}{a}$

$a \quad 2\sqrt{2}R \qquad X \quad \frac{2(\sqrt{2} \quad 1)}{2\sqrt{2}}$

$X \quad \frac{0.414}{1.414} \quad 0.293$

$Z \quad \frac{X}{0.097} \quad \frac{0.293}{0.097} \quad 3$

14. NaA is salt of SBWA

pH $\frac{pK_w}{2} \quad \frac{1}{2}(pK_a \quad \log c)$

$8 \quad 6.5 \quad \frac{1}{2}(5 \quad \log c)$

$1.5 \quad 2 \quad 5 \quad \log c$

$\log c \quad 2$

$c \quad 0.01 \quad \frac{2.592}{M \quad 2}$

$M \quad \frac{2.592}{2 \quad 0.01} \quad 129.6$


$D \quad \frac{ZM}{N_A a^3} \qquad Z \quad 4$

$4 \text{ g/cm}^3 \qquad a \quad 2 \quad 300 \quad 600 \text{ pm}$

15. $Z \quad 4 \quad 6 \quad 6 \quad 12 \quad 12 \quad 40$

$\frac{Z}{10} \quad \frac{40}{10} \quad 4$

For More Material Join: @JEEAdvanced_2024



SURFACE CHEMISTRY

Terms Related to Adsorption and Absorption

- **Adsorption :** The accumulation of molecular species at the surface rather than bulk of solid or liquid is known as adsorption.
- **Adsorbate :** The molecular species or substance which accumulates at the surface is known as adsorbate.
- **Adsorbent :** The surface at which accumulation of adsorbate takes place is known as adsorbent.

  Ex : Water vapours (adsorbate) adsorb at the surface of silica gel (adsorbent).
- **Desorption :** The process of removing an adsorbed substance from adsorbent is known as desorption. When equilibrium is established then rate of adsorption and rate of desorption are equal.
- **Absorption :** When atoms, molecules or ions enter in bulk phase of solid or liquid is known as absorption.

  Ex. Water vapours absorbed by anhydrous $CaCl_2$.
- **Sorption :** When adsorption and absorption takes place simultaneously then phenomena is known as sorption.

  Ex. Dyeing of fabric

Characteristics of Adsorption

- Adsorption is surface phenomena.
- In adsorption enthalpy decreases.
- In adsorption entropy decreases.
- When decrease in free energy takes place then adsorption takes place.

For More Material Join: @JEEAdvanced_2024

Types of Adsorption

(1) Based on concentration :

- Positive adsorption : If the concentration of adsorbate is more at the surface as compared to its concentration in the bulk phase then it is called positive adsorption
- Negative adsorption : If concentration of adsorbate is less at the surface as compared to its concentration in the bulk phase then it is called negative adsorption

(2) Based on forces existing between adsorbate molecule and adsorbent :

- **Physical adsorption :** If the forces of attraction existing between adsorbate and adsorbent are vander Waal's forces, then adsorption is called physical adsorption.
- **Chemical adsorption :** If the forces of attraction existing between adsorbate particles and adsorbent are almost of the same strength as chemical bonds then adsorption is called chemical adsorption.

Comparison between physisorption and chemisorption

S.No.	Physisorption	Chemisorption
1.	Low heat of adsorption usually in range of 20 40 kJ/mol.	High heat of adsorption in the range of 80 240 kJ/mol.
2.	Forces of attraction are vander Waal's forces.	Forces of attraction are chemical bond forces.
3.	It is reversible in nature.	It is irreversible in nature.
4.	It usually takes place at low temp and it decreases with increasing temp.	It takes place at high temperature and it increase with increasing temperature.
5.	More easily liquefiable gases are adsorbed readily.	Gases which can react with the adsorbent show chemisorption.
6.	It forms multimolecular layers.	It forms monomolecular layer.
7.	It does not require any activation energy.	It requires activation energy.
8.	It is not specific.	It is highly specific.
9.	It increases with increase the surface area of adsorbent.	It also increases with increase the surface area of adsorbent.

Factors Affecting Adsorption

- **Nature of the gas :** Easily liquefiable gases adsorb to greater extent.
- **Effect of nature of adsorbent :** When adsorbent is activated then extent of adsorption increases.
- **Specific area of the solid :** Greater the specific area of the solid, greater would be its adsorbent power.
- **Effect of pressure of the gas :** The adsorption causes a net decrease in pressure as the gas gets adsorbed and thus the increase in pressure favours the process of adsorption.


For More Material Join: @JEEAdvanced_2024

Adsorption Isotherm

Graph between extent of the adsorption $\frac{x}{m}$ & pressure (P) at a given temperature is called adsorption isotherm. There are two types of adsorption isotherms

(i) Freundlich Adsorption isotherm :


Freundlich's Equation $\frac{x}{m}$ k $P^{1/n}$



where x mass of adsorbate adsorbed, m mass of adsorbent, p pressure

- **At low pressure :** $\frac{x}{m}$ k P
- **At intermediate pressure :** $\frac{x}{m}$ k $P^{1/n}$, where n 1
- **At high pressure :** $\frac{x}{m}$ constant

$\log \frac{x}{m} \log k$ $\frac{1}{n} \log p$




(ii) Langmuir Adsorption isotherm : According to Langmuir

- There is adsorption of gas molecules on the surface of the solid.
- There is desorption of the adsorbed molecules from the surface of the solid.
- There is formation of unimolecular layer and so it is chemisorption
- A dynamic equilibrium is attained when rate of adsorption rate of desorption.

$$\frac{x}{m} \quad \frac{ap}{1 \quad bp}$$

Where a & b Langmuir parameters.



Case-I At very high pressure

$$bp \quad 1,$$
$$\frac{x}{m} \quad \frac{ap}{bp} \quad \frac{a}{b} \quad \text{constant}$$

Case-II At very low pressure


$$bp \quad 1$$
$$\frac{x}{m} \quad ap$$

Adsorption Isobar

Graph between extent of the adsorption $\frac{x}{m}$ T at a given pressure is called adsorption isobar.


For More Material Join: @JEEAdvanced_2024

A physical adsorption isobar shows a decrease in x/m as the temperature rises.



Physical adsorption isobar

The isobar of chemisorption shows an increase in the beginning and then decrease as the temperature rises.



Chemical adsorption isobar

Application of Adsorption

- Activated charcoal is used in gas masks to remove poisonous gases such as CH_4, CO etc.
- Animal charcoal is used as decolorizer in manufacture of sugar.
- Silica gel is used for removing moisture and controlling humidity.
- Some industrial processes take place in the presence of catalysts which are based upon adsorption.
- Chromatographic purification of compounds which is based upon adsorption.
- The ion exchange resins used for removing hardness of water are also based upon adsorption.

Catalyst

1. Catalyst : A substance which influence the rate of reaction is known as catalyst provided it remains unchanged in amount a well as in composition.

- **Positive catalyst :** The catalyst which increase the rate of chemical reaction is known as positive catalyst. Fe act as positive catalyst in following reaction.

$$N_2 \quad 3H_2 \quad \overset{Fe}{} \quad 2NH_3$$

- **Negative catalyst :** The catalyst which decrease the rate of chemical reaction is known as negative catalyst. C_2H_5OH act as negative catalyst in following reaction.

$$CHCl_3 \quad O \quad \overset{C_2H_5OH}{} \quad COCl_2 \quad HCl$$

- **Auto catalyst :** When one of product act as catalyst then such type of catalyst is known as auto catalyst. Mn^2 act as auto catalyst in following reactions

$$2MnO_4 \quad 5C_2O_4^2 \quad 16H \quad 2Mn^2 \quad 10\,CO_2 \quad 8H_2O$$

2. Catalyst promoters : Substance which increase the activity of catalyst is known as catalyst promoters. Mo act as catalyst promoters in following reaction.

For More Material Join: @JEEAdvanced_2024

$$N_2 \quad 3H_2 \xrightarrow[Mo]{Fe} 2NH_3 ;$$

3. Homogeneous catalytic reaction : If catalyst is present in same phase as reactant then it is called homogenous catalytic reaction.

$$2SO_2(g) \quad O_2(g) \xrightarrow{NO(g)} 2SO_3(g)$$

4. Heterogeneous catalytic reaction : If catalyst is present in different phase as reactant then it is called heterogenous catalytic reaction.

$$N_2(g) \quad 3H_2(g) \xrightarrow{Fe(s)} 2NH_3(g)$$

5. Adsorption theory of heterogeneous catalysis : The mechanism of heterogenous catalysis involves the following steps :

- Adsorption of reactant molecules on the catalyst surface.
- Occurrence of a chemical reaction through the formation of an intermediate.
- Desorption of products from the catalyst surface.
- Diffusion of products away from the catalyst surface.

Enzyme Catalyst

Complex nitrogenous compounds which are produced by living plants and animals are known as enzymes. The characteristic features of enzyme catalyst are :

- They are highly efficient.
- They are highly specific in nature.
- They are highly active under optimum temperature.
- They are highly active under optimum pH.
- There is increase in their activity in the presence of activators and coenzymes.
- They are influenced by inhibitors and poisons.

$$C_{12}H_{22}O_{11} \quad H_2O \xrightarrow{\text{Invertase}} \underset{\text{Glucose}}{C_6H_{12}O_6} \quad \underset{\text{Fructose}}{C_6H_{12}O_6}$$

$$C_{12}H_{22}O_{11} \quad H_2O \xrightarrow{\text{Maltase}} 2\underset{\text{Glucose}}{C_6H_{12}O_6}$$

$$\underset{\text{Starch}}{(C_6H_{10}O_5)_n} \quad nH_2O \xrightarrow{\text{Diastase}} \underset{\text{Maltose}}{C_{12}H_{22}O_{11}}$$

$$\underset{\text{Glucose}}{C_6H_{12}O_6} \xrightarrow{\text{Zymase}} 2C_2H_5OH \quad 2CO_2$$

Mechanism of Enzyme Catalysis

Step 1 : Binding of enzyme to substrate to form an activated complex $E \quad S \quad ES$

Step 2 : Decomposition of the activated complex to form products $ES \quad E \quad P$

Colloids

Depending on size of particles mixture are divided into three classes :

- **True solution :** In true solution size of particles is less than 1 nm.
- **Suspension :** In suspension size of particles is greater than 1000 nm.
- **Colloidal solution :** In colloidal solution size of particles lies from 1 nm to 1000 nm.

It consists two phase *i.e.,* dispersed phase and dispersion medium.

For More Material Join: @JEEAdvanced_2024

Dispersed Phase : It is the component present in small proportion and is just like a solute in a true solution. For example, in the colloidal state of sulphur in water, the former acts as a dispersed phase.

Dispersion Medium : It is normally the component present in excess and is just like a solvent in a solution.

The particles of the dispersed phase are scattered in the dispersion medium in a colloidal system.

Classification of Colloids

1. Based on Physical State of Dispersed Phase & Dispersion Medium :

Dispersed phase	Dispersion medium	Colloidal system	Examples
Gas	Liquid	Foam or froth	Soap sols, lemonade froth, whipped cream.
Gas	Solid	Solid foam/Sol	Pumice stone, styrene, foam, foam rubber.
Liquid	Gas	Aerosols of Liquids	Fog, clouds, fine insecticide sprays.
Liquid	Liquid	Emulsions	Milk
Liquid	Solid	Gels	Cheese, butter, boot polish, table jellies.
Solid	Gas	Aerosols of Solid	Smoke, dust
Solid	Liquid	Sols	Most paint starch dispersed in water, gold sol, muddy water, inks.
Solid	Solid	Solid sols	Ruby glass, some gem stones.

2. Based on Interaction or Affinity of Phases :

(a) Lyophilic Colloids : The colloidal system in which the particle of dispersed phase have great affinity for the dispersion medium, are called lyophilic

(b) Lyophobic colloids : The colloidal system in which the dispersed phase have no affinity for the dispersion medium are called lyophobic.

	Property	Lyophilic sols	Lyophobic sols
1.	Nature	Reversible	Irreversible
2.	Preparation	They are prepared very easily by shaking or warming the substance with dispersion medium.	They are difficult to prepare, Special methods are used.
3.	Stability	They are very stable and are not easily coagulated by electrolytes.	They are generally unstable and get easily coagulated on addition of electrolytes.
4.	Charge	Particles carry no or very little charge depending upon the pH of the medium.	Colloidal particles have characteristic charge (positive or negative)
5.	Viscosity	Viscosity is much higher than that of the medium.	Viscosity is nearly the same as that of the medium
6.	Surface Tension	Surface tension is usually less than that of the medium.	Surface tension is nearly the same as that of the medium.

For More Material Join: @JEEAdvanced_2024

7.	Migration in electric field	The particles may or may not migrate in an electric field.	The colloidal particles migrate either towards cathode or anode in an electric field.
8.	Solvation	Particles are heavily solvated.	Particles are not solvated.
9.	Visibility	The particles can not be seen under ultra microscope.	The particles though invisible, can be seen under ultra microscope.
10.	Tyndall effect	Less distinct.	More distinct.
11.	Action of electrolyte	Large amount of electrolyte is required to cause coagulation.	Small amount of electrolyte is sufficient to cause coagulation.
12.	Examples	Mostly organic substances *e.g.*, starch, gums, proteins, gelatin etc.	Generally inorganic substance *e.g.*, metal sols, sulphides and oxides sols.

3. Based on Type of Particles of the Dispersed Phase

- **Multimolecular colloids :** The multimolecular colloidal particles consists of aggregate of atoms of small molecules with diameters less than 10^{-9} m or 1 nm.

  For example, a sol of gold contains particles of various sizes having several atoms.

- **Macromolecular colloids :** The macromolecular colloidal particles themselves are large molecules. They have very high molecular mass varying from thousands to millions. These substances are generally polymers.

  For example, starch, cellulose and proteins, polyethene, nylon etc.

- **Associated colloids :** Certain substances behave as strong electrolytes at low concentration but at higher concentrations these substances exhibit colloidal characteristics due to the formation of aggregated particles. These aggregated particles are called micelles, which are also known as associated colloids.

Kraft Temperature : The formation of micelles takes place only above a particular temperature called Kraft temperature.

Critical Micelle Concentration : Formation of micelle takes place above kraft temperature and above a particular concentration called the critical micelle concentration (CMC).

For soaps, the CMC is about $10^{-4} M$ to $10^{-3} M$.

4. Method of Preparation of Colloids :

- **Mechanical Dispersion :** In this method the substance is first finely powdered and a suspension is made by shaking the powdered substance with the dispersion medium. This suspension is then passed through a colloidal mill where suspension is converted into colloids. The colloidal solutions of rubber, ink, paints and varnishes are prepared by this method.
- **Electrical Dispersion (Bredig's arc method) :** In this method an electric arc is struck between two metallic rods and kept under the dispersion medium. The particles of the metal break away from the rods and disperse in the liquid. Colloidal solutions of Au, Pt, Ag, Cu and such other metals can be prepared by this method.
- **Peptization :** The process in which freshly formed precipitate converts into a sol by the addition of suitable electrolyte is known as peptization.

(a) Added electrolyte is known as peptizing agent.

(b) The peptization action is due to preferential adsorption of common ion.

For More Material Join: @JEEAdvanced_2024

Ex. $\underset{\text{ppt}}{AgI}$ $\underset{\text{Peptising agent}}{AgNO_3}$ $[AgI]\ Ag^+ : NO_3^-$

$\underset{\text{ppt}}{AgI}$ $\underset{\text{Peptising agent}}{KI}$ $[AgI]\ I^- : K^+$

❖ **Chemical methods :**

Double decomposition : $As_2O_3 \quad 3H_2S \qquad \underset{\text{Yellow sol}}{As_2S_3} \quad 3H_2O$

Oxidation : $2H_2S \quad SO_2 \qquad 2H_2O \quad \underset{\text{Sol}}{3S}$

Reduction : $2AuCl_3 \quad 3SnCl_2 \qquad 3SnCl_4 \quad \underset{\text{Sol}}{2Au}$

Hydrolysis : $FeCl_3 \quad 3H_2O \qquad \underset{\text{Red sol}}{Fe(OH)_3} \quad 3HCl$

5. Purification of Colloidal Solution :

❖ **Dialysis :** The process of separating the particles of colloid from those of crystalloid, by means of diffusion through a suitable membrane is called dialysis. Its principle is based upon the fact that colloidal particles can not pass through a parchment or cellophane membrane, while the ions of the electrolyte can pass through it.

❖ **Electrodialysis :** The ordinary process of dialysis is slow. To increase the process of purification, the dialysis is carried out by applying electric field. This process is called electrodialysis. The artificial kidney machine works on the principle of dialysis.

6. Properties of Colloidal Solution :

(i) Physical Properties :

❖ **Heterogeneity :** Colloidal solutions are heterogeneous in nature consisting of two phases viz, the dispersed phase and the dispersion medium.

❖ **Filterability :** Colloidal particles readily pass through ordinary filter papers. It is because the size of the pores of the filter paper is larger than that of the colloidal particles.

❖ **Non-settling nature :** Colloidal solutions are quite stable as the colloidal particles remain suspended in the dispersion medium indefinitely. Thus there is no effect of gravity on the colloidal particles.

❖ **Colour :** The colour of the colloidal solution is not always the same as the colour of the substances in the bulk. The colour of the colloidal solution depends upon the following factors :

(i) Size and shape of colloidal particles.

(ii) Wavelength of the source of light.

(iii) Method of preparation of the colloidal solution.

(iv) Nature of the colloidal solution.

(v) The way an observer receives the light.

❖ **Stability :** Colloidal solutions are quite stable. Only a few solutions of larger particles may settle but very slowly.

For More Material Join: @JEEAdvanced_2024

(ii) Mechanical Properties :

- **Brownian movement :** It involves motion of colloidal particles in zig-zag paths. It is due to the moving molecules of the dispersion medium which are constantly colliding with colloidal particles.
  - (i) It is not observed in ordinary suspension.
  - (ii) It offers a visible proof of the random kinetic motion of molecules in a liquid.
  - (iii) It provides a direct demonstration of ceaseless motion of molecule as postulated by kinetic theory.
  - (iv) It provide stability to colloidal sol by not allowing them to settle down.
- **Sedimentation :** Under the influence of gravity, colloids tends to settle down very slowly. This phenomena is known as sedimentation. It can be accelerated by ultracentrifugation.

(iii) Optical properties :

- **Tyndall Effect :** The scattering of light by colloidal particles is known as Tyndall effect.

The colloidal particles first absorb light and then a part of the absorbed light is scattered from the surface of the colloidal particles. Maximum scattered intensity is being in a plane at right angles to the plane of incident light. The path becomes visible when seen from that direction.

(iv) Electrical properties : Colloidal particles carry an electric charge which is equal and opposite to dispersion medium.

- **Electrophoresis :** The movement of colloidal particles towards a particular electrode in an electric field is known as electrophoresis. It provides information about the nature of the electrical charge on the colloidal particle.
- **Electro-osmosis :** When electrophoresis of dispersed particles in a colloidal system is prevented by suitable means, it is observed that the dispersion medium itself begins to move in an electric field. This phenomenon is called electro-osmosis.

Electrically charged sols

	Positively charged sols	Negatively charged sols
1.	Ferric hydroxide, aluminium hydroxide	Metals such as Pt, Cu, Au, Ag, Sb_2,S_3, CdS Metals sulphides, *e.g.*, arsenic sulphide
2.	Basic dyes such as methylene blue	Starch, clay, silicic acid, gum, gelatin, charcoal
3.	Haemoglobin	Acid dyes, such as eosin, congo Red

- **Electrical double layer :** The surface of a colloidal particle acquires a positive or a negative charge by selective adsorption of ions carrying +ve or –ve charges respectively. The charged layer attracts counter ions from the medium which forms a second layer. Thus, an electrical double layer is formed on the surface of the particles.

For More Material Join: @JEEAdvanced_2024



Electro kinetic potential : The potential difference between the fixed charge layer and diffused layer of opposite charge is called electro kinetic potential or zeta potential.

- **Coagulation :** The stability of the colloidal system is due to the existence of charged particles. If the charges are neutralised or destroyed, then the colloidal solution gets precipitated. This is called Coagulation or Flocculation. It can be done by
  (i) adding electrolyte
  (ii) mutual action of sols
  (iii) persistent dialysis
  (iv) By cooling
  (v) By electrophoresis

Coagulation value : The minimum concentration of the electrolyte required in millimoles per litre of solution to cause coagulation or flocculation is called coagulation value.

Hardy Schulze Rule : According to this rule the coagulating power of the active ion increases with the valency of the active ion.

Coagulating power of cations : Al^{3} Ba^{2} or Mg^{2} Na or K .

Coagulating power of anion : $[Fe(CN)_6]^{4}$ PO_4^{3} SO_4^{2} Cl

- **Isoelectric Point of Colloid :** The hydrogen ion concentration at which the colloidal particles are neither positively charged nor negatively charged (*i.e.*, uncharged) is known as isoelectric point of the colloid.
- **Protective colloids :** In presence of lyophilic colloids like starch no coagulation of lyophobic colloids takes places and lyophilic colloids act as protective colloids. Greater protecting power of a lyophilic colloid lesser would be gold number.

Gold Number : It is defined as the minimum amount of the protective colloid in milligrams which must be added to 10 mL solution of standard gold sol, which prevents coagulation on adding one mL of 10% NaCl solution.

- **Emulsions :** An emulsion is a colloidal solution of a liquid. It may be defined as a heterogeneous system consisting of more than one immiscible liquids dispersed in one another in the form of droplets

Oil in water type emulsion (*O/W*) : In this emulsion, oil is the dispersed phase and water is the dispersion medium. It is denoted by *O*/*W* or *O* in *W*. For example, milk (liquid fat dispersed in water), vanishing cream, etc.

Water in oil type : In this emulsion, water is the dispersed phase and oil is the dispersion medium. It is denoted by *W*/*O* or *W* in *O*. For example, butter, cod liver oil, cold cream, etc.

For More Material Join: @JEEAdvanced_2024

Distinction between Two Types of Emulsions

Dye test : It involves the addition of oil soluble dye to the emulsion under experiment. If the emulsion acquires the colour of the dye readily, it is water-in-oil type emulsion and if the emulsion remains colourless, it is oil-in-water type emulsion.

Dilution test : As a general rule, an emulsion can be diluted with the dispersion medium while the addition of the dispersed phase forms a separate layer. Thus, if an emulsion can be diluted with oil, it is water-in-oil type.

Preparation of emulsion (Emulsification) : Emulsification is the process which involves the preparation of emulsion. Generally, an emulsion is prepared by subjecting a mixture of the immiscible liquid to a distinct layers upon standing. This substance which stabilizes the emulsion is called emulsifier or emulsifying agent. The commonly used emulsifying agents are soaps, detergents and lyophilic colloids, casein.

Function of emulsifier : The main function of emulsifier or emulsifying agents is to lower the interfacial tension between oil and water and thus helps the intermixing of two liquids.

Demulsification : The process which involves the breaking of an emulsion into two separate liquid layers is called demulsification.

- **Gels :** Colloidal system in which liquids are the dispersed phase and solid act as the dispersion medium are called gels. The common examples are : boot polishes, jelly, gum arabic, agar agar, processed cheese and silicic acid.

When the gels are allowed to stand for a long time, they give out small quantities of trapped liquids with accumulate on its surface. This action of gels is known as Synresis or Weeping.

- **Uses of Colloids**
  - (i) Medicines
  - (ii) Dyes
  - (iii) Rubber industry
  - (iv) Formation of delta
  - (v) Purification of water
  - (vi) Artificial rain
  - (vii) Smoke precipitation
  - (viii) Sewage disposal
  - (ix) Cleansing action of soap and detergent
  - (x) In Photography

For More Material Join: @JEEAdvanced_2024



Level 1

1. The size of particles in suspension, true solution and colloidal solution varies in the order :
(a) suspension > colloidal > true solution
(b) true solution > suspension > colloidal
(c) suspension > colloidal true solution
(d) none of these

2. A colloidal system has what size of particles?
(a) 10^{-4} m to 10^{-10} m
(b) 10^{-5} m to 10^{-7} m
(c) 10^{-9} m to 10^{-12} m
(d) 10^{-6} m to 10^{-9} m

3. Which are not purely surface phenomena?
(a) Adsorption, surface tension
(b) Surface tension, viscosity
(c) Adsorption, viscosity
(d) Absorption, viscosity

4. Adsorbed acetic acid on activated carbon is :
(a) adsorber
(b) absorber
(c) adsorbent
(d) adsorbate

5. Adsorption due to strong chemical forces is called :
(a) chemisorption
(b) physisorption
(c) both (a) and (b)
(d) none of these

6. Adsorption of gases on solid surface is exothermic because :
(a) free energy increases
(b) entropy decreases
(c) entropy increases
(d) interaction developed between gas and solid particles

7. The nature of bonding forces in adsorption are :
(a) purely physical such as van der Waals' forces
(b) purely chemical
(c) both chemical and physical are possible
(d) none of these

8. Which one of the following is not applicable to chemisorption?
(a) Heat of adsorption is negative
(b) It takes place at high temperature
(c) It is reversible
(d) It forms mono-molecular layer

9. Which one of the following characteristics is correct for physical adsorption?
(a) It is very specific
(b) Adsorption on solids is irreversible
(c) Adsorption decreases with decrease in temperature
(d) Generally both enthalpy and entropy of adsorption are negative

10. Which of the following statement is not correct?
(a) Physical adsorption is due to van der Waals' forces
(b) Physical adsorption is irreversible

For More Material Join: @JEEAdvanced_2024

(c) Chemical adsorption increases with increase in temperature upto certain limit than decreases
(d) Enthalpy of adsorption ($|H|$) for a chemical adsorption is greater than that of physical adsorption

11. Which gas will be adsorbed on a solid to greater extent?
(a) Having non-polar molecule
(b) Having highest critical temperature
(c) Having lowest critical temperature
(d) Having lowest critical pressure

12. Which of the following factors affects the adsorption of a gas on solid?
(a) Critical temperature (T_c) (b) Temperature of gas
(c) Pressure of gas (d) All of these

13. Which gas is adsorbed to maximum amount by activated carbon?
(a) $H_2(g)$ (b) He(g) (c) CO(g) (d) $CO_2(g)$

14. The volume of gases NH_3, CO_2 and H_2 adsorbed by one gram of charcoal at 300 K are in order of:
(a) $H_2 > CO_2 > NH_3$ (b) $NH_3 > H_2 > CO_2$
(c) $NH_3 > CO_2 > H_2$ (d) $CO_2 > NH_3 > H_2$

15. Which of the following is used to adsorb water?
(a) Silica gel (b) Calcium acetate
(c) Hair gel (d) Anhydrous $CaCl_2$

16. Absorption and adsorptions are respectively:
(a) surface phenomena, bulk phenomena
(b) bulk phenomena, surface phenomena
(c) both are bulk phenomena
(d) both are surface phenomena

17. Adsorption is multilayer in case of:
(a) physical adsorption (b) chemisorption
(c) both (a) and (b) (d) none of these


18. Reversible adsorption is:
(a) chemical adsorption (b) physical adsorption
(c) both (a) and (b) (d) none of these

19. The effect of pressure on adsorption is high if:
(a) temperature is low (b) temperature is high
(c) temperature is very high (d) larger charcoal piece is taken

20. Sorption is the phenomenon:
(a) reverse of adsorption
(b) reverse of absorption
(c) when adsorption and absorption takes place simultaneously
(d) none of these

21. Which of the following adsorption isotherms represents the adsorption of a gas by a solid involving multilayers of formation? (P_s saturation pressure)

For More Material Join: @JEEAdvanced_2024



22. A plot of $\log \frac{x}{m}$ against $\log P$ for the adsorption of a gas on a solid gives a straight line with slope equal to:
(a) $\frac{1}{n}$ (b) n (c) log K (d) K

23. The heat evolved in chemisorption lies in the range (in kJ/mol) of :
(a) 80 to 240 (b) 20 to 40
(c) 40 to 80 (d) 20 to 100

24. The heat evolved in physisorption lies in the range (in kJ/mol) of :
(a) 20–40 (b) 40–100
(c) 100–200 (d) 200–400

25. According to the adsorption theory of catalysis, the speed of the reaction increases because :
(a) in the process of adsorption, the activation energy of the molecules becomes large
(b) adsorption produces heat which increases the speed of the reaction
(c) adsorption lowers the activation energy of the reaction
(d) adsorption increases the activation energy of the reaction

26. 3.6 gram of oxygen is adsorbed on 1.2 g of metal powder. What volume of oxygen adsorbed per gram of the adsorbent at 1 atm and 273 K?
(A) $0.19\,L\,g^{-1}$ (b) $1\,L\,g^{-1}$
(C) $2.1\,L\,g^{-1}$ (d) None of these

27. A catalytic poison renders the catalyst ineffective because:
(a) It is preferentially adsorbed on the catalyst
(b) It adsorbs the molecules of the reactants
(c) It combines chemically with the catalyst
(d) It combines chemically with one of the reactants

28. The catalyst used in the hydrogenation of oils is:
(a) Fe (b) Ni (c) Pt (d) V_2O_5

29. The function of zymase is to:
(a) change starch into sugar
(b) ferment glucose to alcohol and CO_2
(c) change malt sugar into glucose
(d) change starch into malt sugar and dextrin

30. The conversion of maltose to glucose is possible by the enzyme:
(a) zymase (b) lactase
(c) maltase (d) diastase

31. Shape selective catalysis is a reaction catalysed by:
(a) enzymes (b) ziegler-Natta Catalyst
(c) zeolites (d) platinum

For More Material Join: @JEEAdvanced_2024

32. The process which is catalysed by one of the product is called:
(a) acid-base catalysis (b) autocatalysis
(c) negative catalysis (d) homogeneous catalysis

33. An inhibitor is essentially:
(a) a negative catalyst (b) a heterogeneous catalyst
(c) an auto catalyst (d) a homogeneous catalyst

34. A catalyst in the finely divided form is most effective because:
(a) less surface area is available
(b) more active sites are formed
(c) more energy gets stored in the catalyst
(d) none of these

35. Identify the correct statement regarding enzymes:
(a) Enzymes are specific biological catalysts that normally works at high temperature
(b) Enzymes are normally heterogeneous catalysts which decrease the reaction rate
(c) Enzymes are specific biological catalysts with low molar masses
(d) Enzymes are specific biological catalysts that are very specific in nature

36. A liquid leaves no residue when passed through the ultra-filter paper. The liquid is:
(a) a suspension (b) oil (c) a colloidal sol (d) a true solution

37. Crystalloids differ from colloids mainly in respect of :
(a) electrical behaviour (b) particle nature
(c) particle size (d) solubility

38. Surface tension of lyophilic sols is:
(a) lower than water (b) more than water
(c) equal to water (d) none of these

39. Which one of the following is not used for preparing lyophilic sols?
(a) Starch (b) Gum
(c) Gelatin (d) Metal sulphide

40. Which one of the sols acts as protective colloid?
(a) As_2S_3 (b) Gelatin (c) Au (d) $Fe(OH)_3$

41. Which one of the following is lyophilic colloid?
(a) Pt (b) Gum (c) Fog (d) Blood

42. Small liquid droplets dispersed in another liquid is called:
(a) suspension (b) emulsion
(c) gel (d) true solution

43. Which of the following is an example of associated colloid?
(a) Protein in water (b) Soap in water
(c) Rubber in benzene (d) $FeCl_3$ in H_2O

44. Among the following, select the properties of lyophilic colloidal sols:
(a) viscosity same as that of the medium
(b) extensive hydration takes place
(c) particles migrate either towards cathode or anode in an electric field
(d) particle can be readily detected under ultramicroscope

45. Fog is an example of colloidal system of:
(a) liquid in a gas (b) gas in a liquid
(c) gas in a solid (d) solid in a liquid

For More Material Join: @JEEAdvanced_2024

46. Colloidal systems are not classified on the basis of :
(a) molecular size
(b) nature of the particles
(c) surface tension value
(d) interaction between disperse phase and dispersion medium

47. All colloids:
(a) are suspensions of one phase in another
(b) are two-phase systems
(c) contain only water-soluble particles
(d) are true solutions

48. Which of the following is a lyophobic colloid?
(a) Gelatin (b) Sulphur
(c) Starch (d) Gum arabic

49. Which of the following is not a gel?
(a) Cheese (b) Jellies (c) Curd (d) Milk

50. Cleaning action of soap occurs because:
(a) non-polar tails of soap molecules dissolve in grease
(b) oil and grease dissolved into hydrophilic centres of soap micelles acid washed away
(c) hydrophilic heads dissolves in grease
(d) grease dissolve in soap solution

51. Arsenous sulphide sol is prepared by passing H_2S through arsenous oxide solution. The charge developed on the particles is due to adsorption of:
(a) H^+ (b) S^{2-} (c) OH^- (d) O^{2-}

52. Bredig's arc method cannot be used for the preparation of colloidal sol of :
(a) copper (b) gold (c) silver (d) sodium

53. As_2S_3 sol is:
(a) positive colloid (b) negative colloid
(c) neutral colloid (d) none of the above

54. Which of the following electrolyte will be most effective in coagulation of negative sol ?
(A) KNO_3 (b) $K_4[Fe(CN)_6]$
(C) Na_3PO_4 (d) $MgCl_2$

55. The minimum amount of an electrolyte required to cause coagulation of a sol is called :
(a) Coagulation value (b) Gold number
(c) Protective value (d) None of these

56. The electrical charge on a colloidal particle is indicated by:
(a) Brownian movement (b) electrophoresis
(c) ultramicroscope (d) molecular sieves

57. Colloidal particles in a sol can be coagulated by :
(a) heating (b) adding an electrolyte
(c) adding oppositely charged sol (d) any of the above methods

58. Peptization involves:
(a) precipitation of colloidal particles
(b) disintegration of colloidal aggregates
(c) purification of colloids
(d) impact of molecules of the dispersion medium on the colloidal particles

For More Material Join: @JEEAdvanced_2024

59. Hardy-Schulze law states that:
(a) solution must have higher gold number
(b) disperse phase and dispersion medium must be of the same sign
(c) micelles coagulate in presence of surfactants
(d) the ions carrying more opposite charge to that of sol particle are effective in coagulation

60. Given below are a few electrolytes, indicate which one among them will bring about the coagulation of a gold sol quickest and in the least of concentration?
(a) NaCl (b) $MgSO_4$
(C) $Al_2(SO_4)_3$ (d) $K_4[Fe(CN)_6]$

61. The ability of an ion to bring about coagulation of a given colloid depends upon:
(a) its size
(b) the magnitude of its charge only
(c) the sign of its charge alone
(d) both magnitude and sign of its charge

62. An emulsifying agent is a substance which
(a) stabilizes the emulsion
(b) de-stabilizes the emulsion
(c) coagulates the emulsion
(d) break the interfacial film between suspended particle and medium

63. Colloidal solution of gold is prepared by:
(a) colloidal mill
(b) double decomposition method
(c) Bredig's method
(d) peptization

64. The formation of colloid from suspension is called:
(a) peptisation (b) condensation
(c) sedimentation (d) fragmentation

65. The method usually employed for the destruction of a colloidal solution is:
(a) dialysis (b) addition of electrolytes
(c) diffusion through animal membrane (d) condensation

66. Which of the following has minimum flocculation value for positively charged sol?
(A) Cl^- (b) SO_4^{2-}
(C) PO_4^{3-} (d) $[Fe(CN)_6]^{4-}$

67. Which of the following will have the highest coagulating power for $Fe(OH)_3$ colloid?
(a) PO_4^{3-} (b) SO_4^{2-} (c) Ca^{2+} (d) Al^{3+}

68. Artificial rain is caused by spraying:
(a) opposite charged collidal dust over a cloud
(b) same charged collidal dust over a cloud
(c) both
(d) none of these

69. Colloids can be purified by :
(a) condensation (b) peptization
(c) coagulation (d) dialysis

For More Material Join: @JEEAdvanced_2024

70. At what pH range the rate of enzyme catalyzed reaction is maximum ?
(a) 5 to 7 (b) 6 to 8
(c) 7 to 9 (d) 8 to 10

71. Protective sols are:
(a) lyophilic (b) lyophobic
(c) both (a) and (b) (d) none of these

72. Electro-osmosis is observed when:
(a) dispersion medium particles begins to move in an electric field
(b) dispersed phase begins to move in an electric field
(c) both (a) and (b)
(d) none of the above

73. On adding $AgNO_3$ solution into KI solution, a negatively charged colloidal sol is obtained when they are in:
(a) 50 mL of 0.1 *M* $AgNO_3$ + 50 mL of 0.1 *M* KI
(b) 50 mL of 0.1 *M* $AgNO_3$ + 50 mL of 0.2 *M* KI
(c) 50 mL of 0.2 *M* $AgNO_3$ + 50 mL of 0.1 *M* KI
(d) None of these

74. A sol is prepared by addition of excess of $AgNO_3$ solution in KI solution. The charge likely to develop on colloidal particles is:
(a) positive (b) negative
(c) no charge (d) both charges

75. The gold numbers of protective colloids A, B, C and D are 0.04, 0.004, 10 and 40 respectively. The protective powers of A, B, C and D are in the order:
(A) $A > B > C > D$ (b) $B > A > C > D$
(C) $D > C > A > B$ (d) $D > C > B > A$

76. In Brownian motion, the paths of the particles are:
(a) linear (b) curved
(c) zig-zag (d) uncertain

77. Which one of the following is not represented by sols?
(a) Adsorption (b) Tyndall effect
(c) Flocculation (d) Paramagnetism

78. The Tyndall effect associated with colloidal particles is due to:
(a) presence of electrical charges (b) scattering of light
(c) absorption of light (d) reflection of light


79. Blue colour of the sky is due to:
(a) absorption of light by dust particles (b) reflection of light by dust particles
(c) scattering of light by dust particles (d) presence of clouds


80. The apparatus used to coagulate carbon particles from smoke is called:
(a) cottrel smoker (b) cottrell precipitator
(c) cottrell absorber (d) none of these


For More Material Join: @JEEAdvanced_2024


Level 2

1. Select correct adsorption isobars for chemisorption and physisorption respectively :
(where $\frac{x}{m}$ extent of adsorption, T temperature)

(a) 

(b) 

(c) 

(d) 

2. Which among the following statements is false?
(a) Increase of pressure increases the amount of adsorption
(b) Increase of temperature may decrease the amount of adsorption
(c) Adsorption may be monolayered or multilayered
(d) Particle size of the adsorbent will not affect the amount of adsorption

3. Select incorrect statement;
(a) Lyophilic sols are reversible
(b) Lyophilic sols are self stabilized
(c) Lyophobic sols are obtained from inorganic materials
(d) Lyophobic sols particles are hydrated

4. Which one of the following statements is not correct in respect of lyophilic sols?
(a) There is a considerable interaction between the dispersed phase and dispersion medium
(b) These are quite stable and are not easily coagulated
(c) They always carry significant characteristic charge
(d) The particles are hydrated

5. Alums purify muddy water by:
(a) dialysis (b) absorption (c) coagulation (d) ultrafiltration

6. Lyophilic sols are more stable than lyophobic sols because:
(a) the colloidal particles have positive charge
(b) the colloidal particles have negative charge
(c) the colloidal particles are solvated
(d) there is strong electrostatic repulsion between the colloidal particles

7. Arrange the following electrolytes in the increasing order of coagulation power for the coagulation of As_2S_3 sol:

$\underset{(I)}{K_2SO_4}$ $\underset{(II)}{CaCl_2}$ $\underset{(III)}{Na_3PO_4}$ $\underset{(IV)}{AlCl_3}$

For More Material Join: @JEEAdvanced_2024

(a) I < II < III < IV (b) I = III < II < IV
(c) II < IV < I < II (d) II < III < IV < I

8. Equal volume each of two sols of AgI, one obtained by adding $AgNO_3$ to slight excess of KI and another obtained by adding KI to slight excess of $AgNO_3$, are mixed together. Then:
(a) the two sols will stabilize each other
(b) the sol particles will acquire more electric charge
(c) the sols will coagulate each other mutually
(d) a true solution will be obtained

9. Which of the following statements is not true about the oil-in-water type emulsion?
(a) Addition of small amount of water, no separate layer of water is formed
(b) Addition of a small amount of oil soluble dye renders the entire emulsion coloured
(c) Addition of oil results in the formation of two layers
(d) Addition of a small amount of an electrolyte increases the conductivity of the emulsion

10. Under the influence of an electric field, the particles in a sol migrate towards cathode. The coagulation of the same sol is studied using NaCl, Na_2SO_4 and Na_3PO_4 solutions. Their coagulating values will be in maximum for:
(a) NaCl (b) Na_2SO_4 (c) Na_3PO_4 (d) same for all

11. Select incorrect statement:
(a) Micelles are associated colloids
(b) The electrical charge on a colloid particle is indicated by electrophoresis
(c) Formation of micelles takes place above Kraft temperature
(d) Formation of micelles takes place below CMC

12. Bredig's arc method involves:
(a) dispersion of metal (b) condensation of metal
(c) dispersion as well as condensation (d) neither dispersion nor condensation

13. A freshly prepared $Fe(OH)_3$ precipitate is peptized by adding $FeCl_3$ solution. The charge on the colloidal particle is due to preferential adsorption of:
(a) Cl^- ions (b) Fe^{3+} ions (c) OH^- ions (d) none of these

14. Select incorrect statement :
(a) Gold sol is multimolecular colloid
(b) Large number of particles of a substance aggregate together and formed multimolecular colloids
(c) Metal sulphides are lyophobic colloids
(d) Sulphur sol is multimolecular colloid and hydrophilic in nature

15. Associated colloids :
(a) raise both the surface tension and viscosity of water
(b) lower both the surface tension and viscosity of water
(c) lower the surface tension and raise the viscosity of water
(d) have greater concentration at the surface layer than the bulk of the solution

16. Below critical micelle concentration (CMC):
(a) the surfactant molecules of ions undergo association to form clusters
(b) the viscosity of solution increases abruptly
(c) substances like grease, fat, etc. dissolve colloidally
(d) salt behave as normal, strong electrolyte

For More Material Join: @JEEAdvanced_2024

17. During electro-osmosis of $Fe(OH)_3$ sol:
(a) sol particles move towards anode
(b) sol particles move towards cathode
(c) the dispersion medium moves towards anode
(d) the dispersion medium moves towards cathode


18. Select incorrect statement:
(a) Soap and detergent lower the interfacial surface tension between oil and water
(b) Basic principle of peptization is reverse of coagulation
(c) Soap and detergent used as emulsifiers
(d) Lyophilic sols need stabilizing agent

19. Smoke precipitator works on the principle of :
(a) centrifugation (b) neutralization of charge on colloids
(c) absorption (d) addition of electrolytes

20. What is the correct sequence of the decreasing coagulation value of the following electrolyte for the coagulation of ferric hydroxide sol?
(I) Na_3PO_4 (II) KCl (III) K_2SO_4 (IV) $K_4[Fe(CN)_6]$
(a) IV > I > III > II (b) II > III > I > IV (c) I > II > III > IV (d) IV > III > II > I

21. Which of the following is true in respect of chemical adsorption?
(a) H 0, S 0, G 0 (b) H 0, S 0, G 0
(c) H 0, S 0, G 0 (d) H 0, S 0, G 0

22. For the graph below, select correct order of temperature?



(a) T_1 T_2 T_3 (b) T_2 T_3 T_1 (c) T_3 T_2 T_1 (d) T_1 T_2 T_3

23. Although, nitrogen does not adsorb on a surface at room temperature, it adsorbs on the surface at 83 K. Which one of the following statements is correct ?
(a) At 83 K, there is formation of monolayer
(b) At 83 K, nitrogen is adsorbed as atoms
(c) At 83 K, nitrogen molecules are held by chemical bonds
(d) At 83 K, there is formation of multimolecular layers

24. For the coagulation of 50 mL of ferric hydroxide sol 10 mL of 0.5 M KCl is required. What is the coagulation value of KCl?
(a) 5 (b) 10 (c) 100 (d) None of these

25. 100 mL of 0.6 *M* acetic acid is shaken with 2 g activated carbon. The final concentration of the solution after adsorption is 0.5 *M*. What is the amount of acetic acid adsorbed per gram of carbon?
(a) 0.6 g (b) 0.3 g (c) 1.2 g (d) None of these

For More Material Join: @JEEAdvanced_2024

26. A detergent ($C_{12}H_{25}SO_4^-Na^+$) solution becomes a colloidal sol at a concentration of $10^{-3}M$. On an average 10^{13} colloidal particles are present in 1 mm^3. What is the average number of ions which are contained by one colloidal particle (micelle)? [Given: N_A 6 10^{23}]

(a) 6 10^7 (b) 10 (c) 60 (d) None of these

27. One gram of activated carbon has a surface area of 1000 m^2. Considering complete coverage as well as monomolecular adsorption, how much ammonia at 1 atm and 273 K would be adsorbed on the surface of $\frac{44}{7}$ g carbon if radius of a ammonia molecules is 10 ^{8} cm.

[Given: N_A 6 10^{23}]

(a) 7.46 L (b) 0.33 L (c) 44.8 L (d) 23.5 L

28. At 1 atm and 273 K the volume of nitrogen gas required to cover a sample of silica gel, assuming Langmuir monolayer adsorption, is found to be 1.30 cm^3g^{-1} of the gel. The area occupied by a nitrogen molecule is 0.16 nm^2. What is the surface area per gram of silica gel? [Given: N_A 6 10^{23}]

(a) 5.568 $m^2\ g^{-1}$ (b) 3.48 $m^2\ g^{-1}$ (c) 1.6 $m^2\ g^{-1}$ (d) None of these

29. 10% sites of catalyst bed have adsorbed by H_2. On heating H_2 gas is evolved from sites and collected at 0.03 atm and 300 K in a small vessel of 2.46 cm^3 .

No. of sites available is 5.4 10^{16} per cm^2 and surface area is 1000 cm^2. Find out the no. of surface sites occupied per molecule of H_2. (Given N_A 6 10^{23})

(a) 1 (b) 2 (c) 3 (d) None of these

30. A sample of 16 g charcoal was brought into contact with CH_4 gas contained in a vessel of 1 litre at 27°C. The pressure of gas was found to fall from 760 to 608 torr. The density of charcoal sample is 1.6 g/cm^3. What is the volume of the CH_4 gas adsorbed per gram of the adsorbent at 608 torr and 27°C?

(a) 125 mL/g (b) 16.25 mL/g (c) 26 mL/g (d) None of these

For More Material Join: @JEEAdvanced_2024

Level 3


Passage 1

A graph between x/m and the pressure P of the gas at a constant temperature is called adsorption isotherm. Where x is the no. of moles of the adsorbate and m is the mass of the adsorbent. Adsorption isotherms of different shapes have been experimentally observed. According to Frundlich adsorption isotherm.

$$x/m \quad kP^{1/n}$$

where k and n are constant parameters depending upon the nature of the solid and gas.

1. In the given isotherm select the incorrect statement :




(a) $\frac{x}{m}$ $P^{1/n}$ along OA

(b) $\frac{x}{m}$ P^{0} when point B is reached

(c) $\frac{x}{m}$ does not increase as rapidly with pressure along BC due to less surface area available for adsorption

(d) nature of isotherm is different for two gases for same adsorbent

2. Adsorption isotherm of log $\frac{x}{m}$ and log P was found of the type :




This is true when :

(a) P 0 (b) P 1

(c) $\frac{1}{n}$ 1 (d) P 1

3. Graph between log $\frac{x}{m}$ and log P is a straight line at angle 45° with intercept OA as shown.



Hence, $\frac{x}{m}$ at a pressure of 2 atm is :

(a) 2 (b) 4

(c) 8 (d) 1

Passage 2

The protective power of the lyophilic colloids is expressed in terms of gold number, a term introduced by Zsigmondy. Gold number is the number of milli-gram of the protective colloid which prevent the coagulation of 10 mL of red gold sol, when 1 mL of a 10 per cent solution of sodium chloride is added to it. Thus, smaller the gold number of lyophilic colloid, the greater is the protective power.

For More Material Join: @JEEAdvanced_2024

1. On addition of one mL of solution of 10% NaCl to 10 mL of red gold sol in presence of 0.025 g of starch, the coagulation is just prevented. The gold number of starch is:
(a) 0.025 (b) 0.25 (c) 2.5 (d) 25

2. Which of the following statement(s) is/are correct?
(a) Higher the gold number, more protective power of colloid
(b) Lower the gold number, more the protective power
(c) Higher the coagulation value, more the coagulation power
(d) Lower the coagulation value, higher the coagulation power

3. Gold number gives an indication of:
(a) protective nature of colloids
(b) purity of gold in suspension
(c) the charge on a colloidal solution of gold
(d) gram of gold per litre of solution



Passage 3

Coagulation is the process by which the dispersed phase of a colloid is made to aggregate and thereby separate from the continuous phase. The minimum concentration of an electrolyte in milli-moles per litre of the electrolyte solution which is required to cause the coagulation of colloidal sol is called coagulation value.

Therefore higher is the coagulating power of effective ion, smaller will be the coagulation value of the electrolyte.

$$\text{Coagulation value} \quad \frac{1}{\text{coagulating power}}$$

The coagulation values of different electrolytes are different. This behaviour can be easily understood by Hardy-Schulze rule which states.

"The greater is the valency of the effective ion greater is its coagulating power."

1. Which one has the highest coagulating power?
(a) K^+ (b) Ca^{2+} (c) Al^{3+} (d) Sn^{4+}

2. As_2S_3 sol is negatively charged. Capacity to precipitate it is highest in:
(a) K_2SO_4 (b) Na_3PO_4 (c) $AlCl_3$ (d) $CaCl_3$

3. The ability of an ion to bring coagulation of a given colloid depends upon:
(a) the sign of its charge (b) magnitude of its charge
(c) both magnitude and sign (d) none of these

4. The coagulation of colloidal particles of the sol can be caused by:
(a) heating
(b) adding electrolyte
(c) adding oppositely charged sol
(d) all of these

For More Material Join: @JEEAdvanced_2024



Passage 4

Emulsions are normally prepared by shaking the two components together vigorously although some kind of emulsifying agent usually has to added to stabilize the product. This emulsifying agent may be a soap or other surfactant (surface active) species or a lyophilic sol that forms a protective film around the dispersed phase.

Emulsions broadly classified into two types :

(i) Oil in water emulsions (O/W) : Oil acts as dispersed phase and water acts as dispersion medium.

(ii) Water in oil emulsions (W/O) : Water acts as dispersed phase and oil acts as dispersion medium. Dye test, dilution test may be employed for identification of emulsions.

1. Read two statements:
(1) Milk is an example of oil in water (O/W) type emulsion
(2) Cold cream is an example of water in oil (W/O) type emulsion
(a) Only statement 1 is correct (b) Only statement 2 is correct
(c) Both are correct (d) None of these

2. Select correct statement:
(a) Water in oil emulsions are less viscous than the aqueous emulsions
(b) Electrical conductance of aqueous emulsions is less than that of oil emulsions
(c) Deemulsification can be done by soap or detergent
(d) An emulsion can be diluted with H_2O then it is oil in water (O/W) type

One or More Answers is/are Correct



1. Select the correct statement(s) :
(a) Physical adsorption is specific in nature
(b) Chemical adsorption highly specific in nature
(c) Physical adsorption is due to free valence of atoms
(d) Chemical adsorption is due to stronger interaction or bond formation

2. Select the correct statement(s):
(a) Adsorption is a non-spontaneous process
(b) Surface energy decreases during the process of adsorption
(c) Adsorption takes place with decrease of entropy
(d) In general adsorption is exothermic process

3. Select the correct statement(s):
(a) Physisorption is favoured by low temperature
(b) Chemisorption is favoured by very high temperature because the process is endothermic
(c) Chemisorption increases with increase in temperature owing to high activation energy
(d) Oxygen adsorbed by charcoal can be desorbed by lowering pressure and temperature

For More Material Join: @JEEAdvanced_2024

4. If adsorption of a gas on a solid is limited to monolayer formation, then which of the following statements are true?

(a) At low pressures, $\frac{x}{m}$ varies proportionately with p

(b) At moderate pressures, $\frac{x}{m}$ varies less than proportionately with p

(c) At high pressures, $\frac{x}{m}$ becomes independent of p

(d) At high pressures, $\frac{x}{m}$ varies more than proportionately with p

5. Which of the following are colloids?

(a) Sulphur sol (b) Starch (c) Gold sol (d) Soap solution

6. Which of the following is not ?

(a) Gelatin sol (b) Silver sol

(c) Sulphur sol (d) As_2S_3 sol

7. Which of the following is/are correct for sols?

(a) Its surface tension is lower than that of water

(b) Its viscosity is higher than that of water

(c) Its surface tension is higher than that of water

(d) Its viscosity is equal to that of water

8. Select the correct statement(s):

(a) Benzene is dispersed phase in benzosol

(b) sols are irreversible and not so stable

(c) sol can be produced by double decomposition

(d) When a solution of sulphur in alcohol is added in excess of water a sol of alcohol is formed

9. When negatively charged colloid like As_2S_3 sol is added to positively charged $Fe(OH)_3$ sol in stoichiometric amounts

(a) both the are precipitated simultaneously

(b) this process is called mutual coagulation

(c) they becomes positively charged colloid

(d) they become negatively charged colloid

10. Colloidal gold can be prepared by:

(a) Bredig's arc method (b) reduction of $AuCl_3$

(c) hydrolysis (d) peptization

11. The coagulation of sol particles may be brought about by:

(a) boiling (b) persistent dialysis

(c) adding electrolyte (d) adding oppositely charged sol

12. Select the correct statement(s) :

(a) A sol is prepared by addition of excess of $AgNO_3$ solution in KI solution. The charge likely to develop on colloidal particle is positive

(b) The effects of pressure on physical adsorption is high if temperature is low

(c) Ultracentrifugation process is used for preparation of colloids

(d) Gold number is the index for extent of gold plating done

For More Material Join: @JEEAdvanced_2024

13. Colloidal solution can be purified by:
(a) dialysis (b) electrodialysis (c) electrophoresis (d) ultrafiltration

14. Coagulation of colloids can be achieved by:
(a) centrifugation (b) adding electrolyte (c) change in pH (d) adding water

15. Which are the properties of colloidal solution ?
(a) Adsorption (b) Tyndall effect
(c) Flocculation (d) Depression of freezing point

16. In the aqueous solution of soaps above :
(a) the cations associate to form the aggregates
(b) the anions associate to form the clusters of colloidal dimension
(c) the polar ends of the ions forming the clusters are directed towards water
(d) the non-polar (hydrocarbon) ends are directed towards water

17. Amongst the following which is/are correct statement about the metal sulphide sols?
(a) The sol particles are positively charged due to preferential adsorption of metal ions
(b) The sol particles are negatively charged due to preferential adsorption of sulphide ions
(c) The cations of added electrolytes are effective in causing the coagulation of the sol
(d) The sol is due to both the electric charge and hydration of the particles

18. Emulsion can be destroyed by:
(a) the addition of an emulsifier (b) electrophoresis with a high potential
(c) freezing (d) all of these

19. Which of the following statement is/are correct for electrophoresis?
(a) Colloids are uncharged particles and do not migrate towards the electrodes when electric field is applied
(b) In electrophoresis, sol migrates either to the anode or to the cathode depending on the positively or negatively charged sol
(c) Electrophoresis is useful for finding the charge on a sol
(d) All of these

20. Select the false statement(s):
(a) Brownian motion and Tyndall effect are shown by true solutions
(b) Sorption process is combinations of adsorption and adsorption process
(c) Law Hardy Schulze is related with coagulation of a sol
(d) Higher is the gold number greater will be the protective power of a colloid

21. Select correct statement(s)
(a) colloids are used to protect lyophilic colloids
(b) Dehydrating agent is used to coagulate sols
(c) Rubber is obtained by coagulation of latex
(d) Sometimes, the rainfall occurs when two oppositely charged clouds meet

22. In which of the followings, effect is/are not observed?
(a) Sugar solution (b) Emulsions
(c) Urea solution (d) Solution of proteins

23. Which of the following reactions are examples for heterogeneous catalysis?
(a) $2H_2O_2(l) \xrightarrow{MnO_2(s)} 2H_2O(l) \quad O_2(g)$
(b) $2SO_2(g) \quad O_2(g) \xrightarrow{NO(g)} 2SO_3(g)$
(c) $H_2(g) \quad C_2H_4(g) \xrightarrow{Ni(s)} C_2H_6(g)$
(d) $CH_3COOC_2H_5(aq) \quad H_2O(l) \xrightarrow{HCl(aq)} CH_3COOH(aq.) \quad C_2H_5OH(aq.)$

For More Material Join: @JEEAdvanced_2024

24. Select correct statement(s)
(a) The role of a catalyst in a reversible reaction is to allow the equilibrium to be achieved quickly
(b) Diffusion process is involved in mechanism of heterogeneous catalysis process
(c) Hydrolysis of cane sugar is catalysed by H
(d) Promotors enhance the activity of a catalyst

25. Select correct statement(s)
(a) Blood is a colloidal solution
(b) Alum is used in water purification
(c) River water is a colloidal solution of clay
(d) Colloidal medicines are more effective due to large surface area

Match the Column



Column-I and **Column-II** contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

1.

Column-I	Column-II
(A) Chemisorption	(P) Exothermic
(B) Physical adsorption	(Q) Endothermic
(C) Desorption	(R) Removal of adsorbed material
(D) Activation of adsorbent	(S) Highly specific in nature

2.

Column-I	Column-II
(A) Chemisorption	(P) Not specific and decreases with temperature
(B) Physisorption	(Q) Specific and increases with temperature
(C) Desorption of a solute on liquid surface	(R) Increases the surface tension of the liquid
(D) Adsorption of a solute on a liquid surface	(S) Decreases the surface tension of the liquid

3.

Column-I	Column-II
(A) Milk	(P) Aerosol
(B) Dust	(Q) Emulsion
(C) Cheese	(R) Gel
(D) Froth	(S) Foam

For More Material Join: @JEEAdvanced_2024

4.

Column-I	Column-II
(A) Liquid dispersed in gas	(P) Foam
(B) Gas dispersed in liquid	(Q) Emulsion
(C) Liquid dispersed in solid	(R) Aerosol
(D) Liquid dispersed in liquid	(S) Gel

5.

Column-I	Column-II
(A) As_2S_3 sol	(P) Lyophobic colloid
(B) Sulphur sol	(Q) Macromolecular colloid
(C) Starch	(R) Multimolecular colloid
(D) Soap	(S) Associated colloid

6.

Column-I	Column-II
(A) Coagulation	(P) Due to presence of charge
(B) Electrophoresis	(Q) Due to scattering of light
(C) Tyndall effect	(R) Due to neutralization of charge
(D) Brownian movement	(S) Due to impact of the molecules of the dispersion medium with colloidal particles

7.

Column-I	Column-II
(A) Peptization	(P) Preparation of sols
(B) Ultra centrifugation	(Q) Purification of sols
(C) Electrodialysis	(R) Preparation of metal sols
(D) Bredig's arc method	(S) Movement of ions across the membrane in presence of electric field

For More Material Join: @JEEAdvanced_2024

Assertion-Reason Type Questions



Each questions contains STATEMENT-1 (Assertion) and STATEMENT-2 (Reason).

Examine the statement carefully and mark the correct answer according to the instructions given below:

(A) If both the statement are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1

(B) If both the statements are TRUE and STATEMENT-2 is NOT the correct explanation of STATEMENT-1

(C) If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

(D) If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

1. STATEMENT-1 : For adsorption G, H, S all have –ve values.
STATEMENT-2 : Adsorption is a exothermic process in which randomness decreases due to force of attraction between adsorbent and adsorbate.

2. STATEMENT-1 : The extent of adsorption of CO_2 is much more higher than of H_2.
STATEMENT-2 : $CO_2(g)$ has higher critical temperature and more van der Waals' force of attraction as compared to $H_2(g)$.

3. STATEMENT-1 : In absorption, the molecules of a substance are uniformly distributed throughout the body of other substance.
STATEMENT-2 : In some cases, both absorption and adsorption takes place simultaneously.

4. STATEMENT-1 : More heat evolved in physisorption than in chemisorption.
STATEMENT-2 : Molecules of adsorbate and adsorbent are held by van der Waals' forces in physisorption and by chemical bonds in chemisorption.

5. STATEMENT-1 : Colloidal solution is electrically neutral.
STATEMENT-2 : Due to similar nature of the charge carried by the dispersed phase particles, they repel each other and do not combine to form bigger particles.

6. STATEMENT-1 : Soap and detergent are macro-molecular colloids.
STATEMENT-2 : Soap and detergent are molecules of large size.

7. STATEMENT-1 : Micelles are formed by surfactant molecules above the critical micelle concentration (CMC).
STATEMENT-2 : The conductivity of a solution having surfactant molecules decrease sharply at the CMC.

8. STATEMENT-1 : The micelle formed by sodium stearate in water has COO^- groups at the surface.
STATEMENT-2 : Surface tension of water is reduced by the addition of stearate.

9. STATEMENT-1 : Protein, starch are lyophilic colloids.
STATEMENT-2 : They have strong interaction with the dispersion medium.

10. STATEMENT-1 : Colloidal AgI is prepared by adding KI in slight excess to $AgNO_3$ solution, the sol particles migrate toward cathode under electric field.
STATEMENT-2 : Colloidal particles adsorb ions and thus becomes electrically neutral.

For More Material Join: @JEEAdvanced_2024

11. STATEMENT-1 : When SnO_2 is reacted with NaOH, then its sol particles are attracted towards cathode.

STATEMENT-2 : When SnO_2 is reacted with NaOH, then it gives SnO_3^{2} which is adsorbed by SnO_2, so it is negatively charged sol.

12. STATEMENT-1 : For coagulation of positively charged sols, $[Fe(CN)_6]^{4-}$ ion has higher coagulating power than that of PO_4^{3-}, SO_4^{2-}, Cl^-.

STATEMENT-2 : Because according to Hardy Schulze rule, higher is the valency of ions for the oppositely charged sol particles, better will be the precipitation.

13. STATEMENT-1 : Dispersed phase particles of colloidal solution cannot pass through ultra-filter paper.

STATEMENT-2 : The size of colloidal particles are larger than the size of true solution particles.

14. STATEMENT-1 : When a finely divided active carbon is stirred into a dilute solution of a dye, the intensity of colour in the solution is decreased.

STATEMENT-2 : The dye is adsorbed on the solid surface.

15. STATEMENT-1 : Silica gel is used for drying air.

STATEMENT-2 : Silica gel adsorbs moisture from air.

16. STATEMENT-1 : ZSM-5 is a type of zeolites used as a catalyst in petrochemical industries.

STATEMENT-2 : Zeolites are microporous aluminosilicates three dimensional network silicates in which some silicon atoms are replaced by aluminium atoms.

17. STATEMENT-1 : Lyophilic colloids are called as reversible sols.

STATEMENT-2 : Lyophilic sols are extensively hydrated.

18. STATEMENT-1 : A catalyst is more effective in finely divided form.

STATEMENT-2 : Finely divided form has more surface area.

19. STATEMENT-1 : Sky appears blue in colour.

STATEMENT-2 : Colloidal particles of dust along with water suspended in air scatter blue light.

20. STATEMENT-1 : The conversion of fresh precipitate to colloidal state is called peptization.

STATEMENT-2 : It is caused by addition of common ions.

21. STATEMENT-1 : Colloidal solutions are stable but colloidal particles do not settle down.

STATEMENT-2 : Brownian movement counters the force of gravity act on colloidal particles.

22. STATEMENT-1 : A colloid gets coagulated by addition of an electrolyte.

STATEMENT-2 : Coagulation depends on the valance and sign of the charge of the coagulant ion.

23. STATEMENT-1 : Suspensions are visible to naked eye.

STATEMENT-1 : Size of suspension particles are smaller than collidal particles.

24. STATEMENT-1 : At pH of isoelectric point, the sol particles of amino acids neither move towards anode nor towards cathode.

STATEMENT-2 : Because at the isoelectric point, the concentration of conjugate base and conjugate acid of the Zwitter ions becomes equal and so one's charge is counterbalanced by other.

For More Material Join: @JEEAdvanced_2024

25. **STATEMENT-1 :** Gold sol is multimolecular and hydrophobic in nature.
STATEMENT-2 : Gold sol is prepared by Bredig's arc method.

Subjective Problems



1. How many colloidal systems exist in nature ?
2. How many colloidal systems exist in nature in which liquid is dispersed phase ?
3. How many colloidal systems exist in nature in which gas is dispersed phase ?
4. What is the minimum diameter (in nm) of colloidal particles ?
5. For soaps critical micelle concentration (CMC) is 10^{x} (min.) to 10^{y} (max.) mol/L. What is the value of x ?

For More Material Join: @JEEAdvanced_2024

ANSWERS

Level 1

1.	(a)	**2.**	(d)	**3.**	(d)	**4.**	(d)	**5.**	(a)	**6.**	(d)	**7.**	(c)	**8.**	(c)	**9.**	(d)	**10.**	(b)
11.	(b)	**12.**	(d)	**13.**	(d)	**14.**	(c)	**15.**	(a)	**16.**	(b)	**17.**	(a)	**18.**	(b)	**19.**	(a)	**20.**	(c)
21.	(a)	**22.**	(a)	**23.**	(a)	**24.**	(a)	**25.**	(c)	**26.**	(c)	**27.**	(a)	**28.**	(b)	**29.**	(b)	**30.**	(c)
31.	(c)	**32.**	(b)	**33.**	(a)	**34.**	(b)	**35.**	(d)	**36.**	(d)	**37.**	(c)	**38.**	(a)	**39.**	(d)	**40.**	(b)
41.	(b)	**42.**	(b)	**43.**	(b)	**44.**	(b)	**45.**	(a)	**46.**	(c)	**47.**	(b)	**48.**	(b)	**49.**	(d)	**50.**	(a)
51.	(b)	**52.**	(d)	**53.**	(b)	**54.**	(d)	**55.**	(a)	**56.**	(b)	**57.**	(d)	**58.**	(b)	**59.**	(d)	**60.**	(c)
61.	(d)	**62.**	(a)	**63.**	(c)	**64.**	(a)	**65.**	(b)	**66.**	(d)	**67.**	(a)	**68.**	(a)	**69.**	(d)	**70.**	(a)
71.	(a)	**72.**	(a)	**73.**	(b)	**74.**	(a)	**75.**	(b)	**76.**	(c)	**77.**	(d)	**78.**	(b)	**79.**	(c)	**80.**	(b)

Level 2

1.	(c)	**2.**	(d)	**3.**	(d)	**4.**	(c)	**5.**	(c)	**6.**	(c)	**7.**	(b)	**8.**	(c)	**9.**	(b)	**10.**	(a)
11.	(d)	**12.**	(c)	**13.**	(b)	**14.**	(d)	**15.**	(c)	**16.**	(d)	**17.**	(c)	**18.**	(d)	**19.**	(b)	**20.**	(b)
21.	(b)	**22.**	(c)	**23.**	(d)	**24.**	(c)	**25.**	(b)	**26.**	(c)	**27.**	(a)	**28.**	(a)	**29.**	(c)	**30.**	(b)

Level 3

Passage–1	**1.**	(b)	**2.**	(c)	**3.**	(b)			
Passage–2	**1.**	(d)	**2.**	(b)	**3.**	(a)			
Passage–3	**1.**	(d)	**2.**	(c)	**3.**	(c)	**4.**	(d)	
Passage–4	**1.**	(c)	**2.**	(d)					

For More Material Join: @JEEAdvanced_2024

One or More Answers is/are Correct

1. (b,d)	**2.** (b,c,d)	**3.** (a,c, d)	**4.** (a,b,c)	**5.** (a,c)	**6.** (b,c,d)	**7.** (a,b)	**8.** (b,c)		
9. (a,b)	**10.** (a,b)	**11.** (a,b,c,d)	**12.** (a,b)	**13.** (a,b,d)	**14.** (a,b,c)	**15.** (a,b,c,d)	**16.** (b,c)		
17. (b,c)	**18.** (b,c)	**19.** (c)	**20.** (a,d)	**21.** (b,c,d)	**22.** (a,c)	**23.** (a,c)	**24.** (a,b,c,d)		
25. (a,b,c,d)									

Match the Column

1.	A	P,S;	B	P;	C	Q,R;	D	Q
2.	A	Q;	B	P;	C	R;	D	S
3.	A	Q;	B	P;	C	R;	D	S
4.	A	R;	B	P;	C	S;	D	Q
5.	A	P, R;	B	P, R;	C	Q;	D	S
6.	A	R;	B	P;	C	Q;	D	S
7.	A	P;	B	Q;	C	Q,S;	D	P,R

Assertion-Reason Type Questions

1. (A)	**2.** (A)	**3.** (B)	**4.** (D)	**5.** (B)	**6.** (D)	**7.** (A)	**8.** (B)	**9.** (A)	**10.** (C)
11. (D)	**12.** (A)	**13.** (A)	**14.** (A)	**15.** (A)	**16.** (B)	**17.** (B)	**18.** (A)	**19.** (A)	**20.** (B)
21. (A)	**22.** (B)	**23.** (C)	**24.** (A)	**25.** (B)					

Subjective Problems

1. 8	**2.** 3	**3.** 2	**4.** 1	**5.** 4

For More Material Join: @JEEAdvanced_2024

Hints and Solutions

Level 1

1. (a) Solution Particle size
True solution Size < 1 nm
Colloidal solution 1 nm < size < 1000 nm
Suspension Size > 1000 nm

2. (d) 10^{-6} m to 10^{-9} m

3. (d) Absorption is not surface phenomena.

4. (d) Adsorbate adsorb at the surface of adsorbent.

5. (a) Chemical adsorption.

6. (d) Attractive force developed at the surface between adsorbent and adsorbate.

7. (c) Physical adsorption and chemical adsorption.

8. (c) Chemical adsorption is irreversible in nature.

9. (d) H O S O

10. (b) Physical adsorption is reversible in nature.

11. (b) Critical temp. Attractive force
Extent of adsorption .

12. (d) Adsorption T_c
Adsorption T
Adsorption P

13. (d) Van der Waal force adsorption

14. (c) Van der Waal force adsorption


15. (a) Silica gel adsorb water.

16. (b) Adsorption : surface phenomena
Absorption : bulk phenomena

17. (a) Physical adsorption is multilayer adsorption.


18. (b) Physical adsorption is reversible adsorption.

19. (a) At low temperature, pressure have greater effect on adsorption.



20. (c) In sorption, adsorption and absorption takes place simultaneously.

21. (a)



22. (a) $\frac{x}{m}$ $KP^{\frac{1}{n}}$

$\log \frac{x}{m}$ $\log K$ $\frac{1}{n}$ $\log P$

Slope $\frac{1}{n}$

23. (a) H : 80 to 240 kJ mol^{-1}

24. (a) H : 20 to 40 kJ mol^{-1}

25. (c) Adsorption lowers the activation energy of reaction because surface energy released during adsorption.

26. (c) Mass of O_2 per gram of adsorbent $\frac{3.6}{1.2}$ 3

No. of moles of O_2 per gram of adsorbent

$$\frac{3}{32}$$

Volume of O_2 per gram of adsorbent

$$\frac{3}{32} \quad \frac{0.0821 \quad 273}{1}$$

$= 2.10$

27. (a) It is preferentially adsorb on the catalyst.

28. (b) Ni used as catalyst in hydrogenation of oil.

29. (b) $C_6H_{12}O_6 \xrightarrow{\text{Zymase}} CO_2 \quad C_2H_5OH$

30. (c) Maltose $\xrightarrow{\text{Maltase}}$ glucose

31. (c) Zeolites is shape selective catalyst.

32. (b) Autocatalyst

33. (a) Negative catalyst

34. (b) Surface area No of sites

35. (d) Enzyme are specific in nature.

36. (d) True solution.

37. (c) particle size

38. (a) Surface tension of lyophilic solution is lower than water.

39. (d) Metal sulphide are lyophobic solution.


For More Material Join: @JEEAdvanced_2024

40. (b) Gelatin act as protective colloids.
41. (b) Gum is lyophilic solution.
42. (b) Emulsion : D.P. liquid
D.M. liquid
43. (b) Soap formed associated colloids in water.
44. (b) D.P. have greater affinity towards D.M.
45. (a) fog : liquid in gas.
46. (c) Colloids divided on the basis of molecular size, affinity of D.P. towards D.M. and particle nature.
47. (b) Colloids have D.P. and D.M.
48. (b) Sulphur is lyophobic colloids.
49. (d) Milk is emulsion.
50. (a) Non-polar tails of soap molecules dissolve grease.
51. (b) As_2S_3 H_2S As_2S_3 : S^2 : H


Preferential adsorption of S^2 takes place. So negative charge developed.
52. (d) Bredig's arc method is used to prepare metal sol.
53. (b) Metal sulphides is negative colloids.
54. (d) Positive charge Coagulating power
$MgCl_2$
55. (a) Millimoles of electrolyte required to coagulates one litre colloidal solution is called coagulation value.
56. (b) By electrophoresis, we can determine charge of colloidal solution.
57. (d) Coagulation takes place by heating, adding electrolyte and addition of opposite charge colloids.
58. (b) In peptisation, suspended state converts into colloidal state.
59. (d) Coagulating power charge.
60. (c) Gold sol. have negative charge. So $Al_2(SO_4)_3$ is most effective for coagulation.
61. (d) Coagulating power depend upon sign of charge and magnitude of charge.
62. (a) Emulsifying agent stablise emulsion.
63. (c) Gold solution is prepared by Bredig's arc method.
64. (a) In peptisation, colloid formed from suspension.
65. (b) By addition of electrolyte.
66. (d) For positive charge colloids coagulating power coagulation value . $[Fe(CN)_6]^4$.
67. (a) $Fe(OH)_3$ is a positive sol. Hence, greater the charge on anion more is the coagulation power.
68. (a) Mutual coagulation
69. (d) Dialysis is a method of purification of colloids.
70. (a) 5 to 7
71. (a) Lyophilic solution are protective colloids.
72. (a) In electro osmosis D.M. move towards opposite charge electrode.
73. (b) $AgNO_3$ KI AgI(*s*) KNO_3
5 10^{3} 10 10^{3}
0 5 10^{3} 5 10^{3}
AgI(*s*) I AgI : I
Negative charge colloids.
74. (a) $AgNO_3$ KI AgI KNO_3
excess
AgI Ag AgI Ag
75. (b) Smaller is the 'gold number' of protective colloid, greater is its protective power.
76. (c) *Zig-zag* path.
77. (d) paramagnetism.
78. (b) Tyndall effect takes place due to scattered of light.
79. (c) Light scattered by dust particles.
80. (b) Cottrell precipitator.

Level 2

1. (c)



Chemical adsorption



Physical adsorption

2. (d) Surface area rate of adsorption
3. (d) In lyophobic sol D.P. have less affinity towards D.M.
4. (c) In lyophilic sol no charge present.

For More Material Join: @JEEAdvanced_2024

5. (c) Alum, purified to muddy water by coagulation.

6. (c) D.P. particles have great affinity towards D.M. in lyophilic colloids.

7. (b) Charge coagulation power

8. (c) The sols obtained in the two cases will be oppositely charged so coagulate each other.

9. (b) Colour of dye appear in patches in *o/w* emulsion.

10. (a) The sol particles migrate towards cathode. So they are positively charged. Hence, anions would be effective in coagulation. Greater is the valence of effective ion, smaller will be its coagulating value.

11. (d) Formation of micelles takes place above CMC.

12. (c) Bredig's arc method involve dispersion and condensation.

13. (b) Preferential adsorption of Fe^3 takes place. So positive charge develop.

14. (d) Sulphur solution is multimolecular by hydrophobic in nature.

15. (c) Associate colloids decrease surface tension and increase in viscosity.

16. (d) Below CMC, salt behave as normal electrolyte.

17. (c) In electro-osmosis, the sol particles are prevented from migration whereas the dispersion medium migrates in the direction opposite to those of particles. Here, the medium is negatively charged.

18. (d) No stabilising agent required for the preparation of lyophilic sol.

19. (b) Neutralisation of charge.

20. (b) Ferric hydroxide solution have positive charge. So order of coagulation value.

KCl K_2SO_4 Na_3PO_4 $K_4[Fe(CN)_6]$

Charge coagulation value

21. (b) H 0, S 0, G 0

22. (c) Temp Extent of adsorption

23. (d) At 83 K physical adsorption takes place effectively so multimolecular layers formed.

24. (c) Coagulation value

$$\frac{\text{Number of millimoles of electrolyte required}}{\text{Volume of colloidal solution (in litre)}}$$

$$\frac{10 \quad 0.5}{50} \quad 1000 \quad 100$$

25. (b) Mass of acetic acid adsorbed by 2 g charcoal

$$100 \quad 10^{\ 3} \quad (0.6 - 0.5) \quad 60$$

$$0.6; \frac{x}{m} \quad \frac{0.6}{2} \quad 0.3$$

26. (c) No. of sodium lauryl sulphate molecules $(CH_3(CH_2)_{11}SO_4^-Na^+)$ in 1 litre solution

$$10^{\ 3} \quad 6 \quad 10^{23}$$

$$6 \quad 10^{20}$$

No. of sodium lauryl sulphate molecules per $mm^3 = 6 \quad 10^{14}$

No. of colloidal particles per $mm^3 = 10^{13}$

No. of molecules per colloidal particle

$$\frac{6 \quad 10^{14}}{10^{13}}$$

$$= 60$$

27. (a) Total surface area of carbon $\frac{44}{7} \quad 10^7 \ cm^2$

$$r \quad 10^{\ 8} \ cm$$

Surface area of $NH_3 = \quad r^2$

$$\frac{22}{7} \quad 10^{\ 16} \ cm^2$$

No. of NH_3 molecules adsorbed $\frac{\frac{44}{7} \quad 10^7}{\frac{22}{7} \quad 10^{\ 16}}$

$$2 \quad 10^{23}$$

Vol. of NH_3 adsorbed at 1 atm 273 K

$$\frac{2 \quad 10^{23}}{6 \quad 10^{23}} \quad 22.4 = 7.46 \text{ L}$$

28. (a) No. of molecules per gram of N_2 in monolayer

$$\frac{6 \quad 10^{23}}{22{,}400} \quad 1.30$$

$$= 3.48 \quad 10^{19}$$

Cross-sectional area of a molecule

$$1.6 \quad 10^{-19} \ m^2$$

Area covered by molecules per gram

$$3.48 \quad 10^{19} \quad 1.6 \quad 10^{-19} \quad 5.568 \ m^2$$

Surface area $5.568 \ m^2 g^{-1}$

29. (c) Adsorbed moles of $H_2 = \frac{0.03 \quad 2.46 \quad 10^{\ 3}}{0.0821 \quad 300}$

$$3 \quad 10^{\ 6}$$

No. of absorbed molecules of

$$H_2 \quad 3 \quad 10^{\ 6} \quad 6 \quad 10^{23}$$

$$18 \quad 10^{17}$$

For More Material Join: @JEEAdvanced_2024

Total no. of surface sites available

$5.4 \quad 10^{16} \quad 1000$

$5.4 \quad 10^{19} \text{ cm}^2$

No. of surface sites that is occupied by adsorption of

$H_2 \quad \frac{10}{100} \quad 5.4 \quad 10^{19}$

$5.4 \quad 10^{18}$

No. of surface sites occupied by one molecule

$H_2 \quad \frac{5.4 \quad 10^{18}}{18 \quad 10^{17}} \quad 3$

30. (b) Final volume of gas, at 608 torr pressure

$V_2 \quad \frac{P_1 V_1}{P_2}$

$\frac{760 \quad 1}{608}$

1.25 L or $V_2 \quad 1250$ mL

Volume occupied by gas Volume of vessel

Volume occupied by charcoal

$1000 \quad \frac{16}{1.6} \quad 990$ mL

Difference of volume is due to adsorption of gas by charcoal.

Volume of gas adsorbed by charcoal

$1250 \quad 990$

260 mL

Volume of the gas adsorbed per gram of charcoal

$\frac{260}{16}$ 16.25 mL/g at 608 torr and 27 C.

Level 3

Passage-1

1. (b) At point $B \; \frac{x}{m} \quad P^{1/n}$

2. (c) $\log \frac{x}{m} \quad \log K \quad \frac{1}{n} \quad \log P$

3. (b) $\frac{x}{m} \quad K \quad P^{1/n}$

$\frac{x}{m} \quad 2 \quad 2^1 \quad 4$

Passage-2

1. (d) Gold no. $0.025 \quad 10^3 \quad 25$

2. (b) Gold no protecting power

3. (a) Protective nature of colloids.

Passage-3

1. (d) Charge coagulating power

2. (c) Al^3 charge coagulating power

3. (c) Coagulation depend upon charge and magnitude.

4. (d) Coagulation takes place by heating, adding electrolyte and mutual coagulation.

Passage-4

1. (c) Milk : o/w emulsion

Cold cream : w/o emulsion

2. (d) o/w emulsion diluted by addition of water.

One or More Answers is/are Correct

1. (b, d)

Physical adsorption are not specific in nature due to van der Waal's forces and chemical adsorption are specific in nature and due to chemical bond.

2. (b,c,d)

Adsorption is spontaneous process.

Enthalpy of adsorption is –ve


Change in entropy = –ve

3. (a, c, d)

Low temperature is favourable for physical adsorption.

Chemical adsorption required activation energy.

4. (a, b, c)



5. (a, c)

Sulphur and gold are multimolecular colloids.

6. (b, c, d)

Gelatin is lyophilic colloids.

7. (a, b)

Surface tension is less than water and viscosity is greater than water in case of lyophilic sol.

For More Material Join: @JEEAdvanced_2024

8. (b, c)

Lyophobic sol are ir-reversible and less stable and they produced by double decomposition.

9. (a, b)

Mutual coagulation.

10. (a, b)

Gold solution prepared by Bredig's arc method and by reduction of $AuCl_3$.

11. (a, b, c, d)

Coagulation takes place by boiling,

persistent dialysis

add of electrolyte

mutual coagulation.

12. (a, b)

$AgNO_3$ KI AgI KNO_3

AgI $AgNO_3$ AgIAg $:NO_3$

At low temperature, extent of physical adsorption is greater due to increase in pressure.

13. (a, b, d)

Colloidal solution can be purified by dialysis, electrolysis and ultrafiltration.

14. (a, b, c)

Coagulation occurs due to centrifugation, add of electrolyte and change in pH.

15. (a, b, c, d)

Sol shows Tyndall effect adsorption coagulation sol have colligative properties

16. (b, c)

Above CMC, associate colloids formed.

Non polar part directed towards water.

17. (b, c)

Charge develop due to preferential adsorption of ion addition of electrolyte cause coagulation.

18. (b, c)

Coagulation occurs due to electrophoresis and freezing.

19. (c)

In electrophoresis colloidal particle migrate towards opposite charge electrode.

20. (a,d)

Tyndall effect does not shown by true solution due to small size of particles.

Sorption involve adsorption and absorption coagulating power charge, smaller the gold number greater would be the protecting power.

21. (b, c, d)

Lyophilic colloids used as protecting colloids electrolyte used for coagulation. Rubber is obtained by coagulation of latex. By mutual coagulation rain fall occurs.

22. (a, c)

Due to size of particle

23. (a, c)

MnO_2, Ni

24. (a,b,c,d)

In presence of catalyst equilibrium obtained quickly.

Mechanism of hetrogenous catalyst is based on adsorption.

H act as auto catalyst

Catalyst promotor increase the efficiency of catalyst.

25. (a,b, c,d)

Blood is colloidal solution, water purification by alum is based on coagulation. River water is colloidal solution of clay surface area preferential adsorption .

Subjective Problems

1. Eight gas colloidal system does not exists.

2. Aerosol, Emulsion, Gels.

3. Foam, Solid foam.

4. Size 1nm

5. CMC of soap

$10^{-4}\ M$ to $10^{-3}\ M$

x 4

For More Material Join: @JEEAdvanced_2024