



Fabio  
Strozzi

Fisica Teorica

Studente esterno  
Scuola Normale Superiore  
Pisa, a.a. 2021-2022

Rel.: Prof. Dario Buttazzo  
Tesi di Laurea Magistrale



# UNIVERSITÀ DI PISA



Dipartimento di Fisica Enrico Fermi  
Corso di Laurea in Fisica

## **Path along different solutions for the B-Physics Anomalies**

### **Introduction of one or two mediators Beyond Standard Model to accomodate those Anomalies**

**Relatori:**  
**Prof. Dario Buttazzo**

**Presentata da:**  
**Fabio Strozzi**

**Sessione autunnale**  
**Anno Accademico 2021/2022**

# Contents

Index	2
Introduction	3
1 B-Physics Anomalies	5
1.1 Main Processes for NC transitions . . . . .	5
1.1.1 $B_s \rightarrow \mu\mu$ . . . . .	7
1.2 Main Processes for CC transitions . . . . .	7
2 Effective Field Theory approach	9
2.1 Extended Fermi Theory . . . . .	9
2.1.1 Flavour Changing Neutral Currents . . . . .	10
2.1.2 4-fermions operators . . . . .	11
2.2 Fierz identities . . . . .	12
2.2.1 Fierzing in the scalar lagrangian . . . . .	14
2.3 Effective Semileptonic Lagrangian . . . . .	15
3 Some of the possible heavy bosons	18
3.1 Colour-less bosons . . . . .	19
3.1.1 $B' \sim (\mathbf{1}, \mathbf{1})_0$ . . . . .	19
3.1.2 $W' \sim (\mathbf{1}, \mathbf{3})_0$ . . . . .	20
3.2 Leptoquarks . . . . .	21
3.2.1 Proton decay . . . . .	22
3.3 Vector Leptoquarks . . . . .	23
3.3.1 $U_1 \sim (\mathbf{3}, \mathbf{1})_{2/3}$ . . . . .	23
3.3.2 $U_3 \sim (\mathbf{3}, \mathbf{3})_{2/3}$ . . . . .	25
3.4 Scalar Leptoquarks . . . . .	26
3.4.1 $S_1 \sim (3^*, 1)_{1/3}$ . . . . .	26
3.4.2 $R_2 \sim (3, 2)_{7/6}$ . . . . .	27
3.4.3 $S_3 \sim (3^*, 3)_{1/3}$ . . . . .	28
3.5 Summary . . . . .	29
Appendix	31
A Flavour Physics	32

# Introduction

In this work we are trying to explore the huge world of Beyond Standard Model (BSM) theories using as gate some experimental observations seen recently, that bring clues of the breaking of an accidental symmetry. In the Standard Model (SM) as is known today we have an accidental symmetry which is called Lepton Flavour Universality (LFU) which tells us that all the gauge interactions are flavour blind; in other words the Electro-Weak (EW) processes have the same strenght for the electron, the muon, the tau and same for the different types of neutrinos. Since the only difference between the different lepton families is the mass, there is the phase space factor which affect the rate or the cross section of a process (for instance the decay of charged  $\pi$  produces mostly muons for this reasons). Nevertheless there is also a dynamical effect which is not mass-blind, which is the coupling with the Higgs Boson but since leptons are very light (the  $\tau$ , which is the heaviest, weighs less than 50 times the Higgs' vacuum expectation value) we neglect the Higgs' coupling and say that LFU is a well approximated symmetry of the SM.

If we have accepted that LFU is a good symmetry we can also understand why testing it is a good place to find clues for BSM theories. Recently in different experiments were found different hints of LFU Violation (LFUV) in semileptonic decays of the B mesons (mesons with non vacuum difference of  $b$  and  $\bar{b}$  as valence quark). All the deviations from the SM appearing in these decays go under the name of *B-Physics Anomalies*.

Since, as we will discuss, the B-Physics Anomalies appear mostly at the hadronic scale (order few GeV) the most natural approach is the Effective Field Theory (EFT) approach, as is usually done with the Fermi Theory. In this approach finding New Physics (NP) basically means to find deviation from the Lagrangian's coefficients of the operators (or Wilson's coefficients).

So our purpose is to find the right heavy mediators which, once integrated out from the Lagrangian (again as we do with W's and Z bosons to get the Fermi theory), give us the appropriate contribute to accomodate the B-Physics Anomalies. We also have to be very careful about the processes that are already tested, because introducing NP can affect also processes that don't concern B or hadrons at all.

This is one of the reason because we like to introduce LeptoQuarks (LQs): coloured bosons which can be absorbed from a quark to become a lepton (+h.c.) and try in this way to affect just semileptonic processes without disturbing others SM constraints. We will take count also of the possible colour-less bosons (heavier version of W and Z) and in both cases we will try to find the best flavour structure to accomodate the Anomalies.

We also would like to see those bosons, that's why apart from the low energy processes we want to take in consideration the direct searches in which we try to see if this bosons are produced at LHC smashing proton against proton. In this point we will find interesting constraints on the masses and some reasons to improve the colliders we already have.

A lot of papers are already written about B-Anomalies and LeptoQuarks; the main purpose of this work is to offer a catalogue as much clear and complete we can do of solution with single NP mediator and with a pair of them.

In the end we have to mention that adding bosons to SM is not enough to say that we have a NP theory, we need a theory in which with few assumptions and few input values all the particles and processes come out naturally (as in SM). These complete theories are known as UV Completions and we will mention some of them. Most of them belong to the family of Great Unification Theories (GUTs) which are the natural enviroment to get LQs.


Hope you readers will find the work interesting and light to read, for now we have done enough of introductions; let's begin with B-Physics Anomalies.

# 1

## B-Physics Anomalies


The two fundamental processes in which we are interested in are: the neutral current (NC) decay

$$b \rightarrow sl^+l^- \text{ or } b \rightarrow s\nu\bar{\nu}$$



and the charged current (CC) decay

$$b \rightarrow cl\bar{\nu}$$



### 1.1 Main Processes for NC transitions

For what concern the NC decays the first process we want to discuss is the decay:

$$B \rightarrow K^*l^+l^-.$$

Table 1.1: Table with the most important anomalies in  $b \rightarrow sl^+l^-$  transition.  $R_{K^*}^{[q_1^2, q_2^2]}$  means the ratio  $R_{K^*}$  in which the momenta of the pair lepton-antilepton has energy at rest  $q^2$  included between  $q_1^2$  and  $q_2^2$ . Fonte: [1]

<i>Observable</i>	<i>Experiment</i>	<i>SM</i>
$R_{K^*}^{[0.045, 1.1]}$	$0.66_{-0.07}^{+0.11} \pm 0.03$	$0.906 \pm 0.028$
$R_{K^*}^{[1.1, 6.0]}$	$0.69_{-0.07}^{+0.11} \pm 0.05$	$1.00 \pm 0.01$
$R_K^{[1.1, 6.0]}$	$0.846_{-0.039-0.012}^{+0.042+0.013}$	$1.00 \pm 0.01$
$BR(B_s \rightarrow \mu^- \mu^+)$	$2.85_{-0.31}^{+0.32} \cdot 10^{-9}$	$(3.66 \pm 0.14) \cdot 10^{-9}$

We know that in the SM this process is loop-induced (as all the Flavour Changing Neutral Current (FCNC)) and should have the same rate for the charged lepton being electron or muon because of LFU. Given the masses of the particle involved:

$$m_B \simeq 5.3\text{GeV} \quad m_K \simeq 430\text{MeV} \quad m_\mu \simeq 106\text{MeV} \quad m_e \simeq 500\text{keV}$$

the phase space factor is neglectable; and the ratio

$$R_{K^*}^{\mu e} \equiv \frac{\mathcal{B}(B \rightarrow K^* \mu^- \mu^+)}{\mathcal{B}(B \rightarrow K^* e^- e^+)}$$

is equal to 1 in the SM.

The reasons why we use ratios of Branching Ratios are mainly three:

- To reduce the dependence from hadronic form factors
- To reduce the dependence from CKM matrix elements
- To reduce the systematic error in general

In the Table 1.1 we can finally see the first B-Physics Anomaly; in fact the lack of muons among the decay's products is a clear hint of LFU violation. The formal definition for the ratio  $R_{K^*}^{[q_1^2, q_2^2]}$  is:

$$R_{K^*}^{[q_1^2, q_2^2]} = \frac{\int_{q_1^2}^{q_2^2} dq^2 \frac{d\mathcal{B}}{dq^2}(B \rightarrow K^* \mu^+ \mu^-)}{\int_{q_1^2}^{q_2^2} dq^2 \frac{d\mathcal{B}}{dq^2}(B \rightarrow K^* e^+ e^-)}$$

and same for the pseudoscalar  $K$ .






Figure 1.1: Diagram of the fundamental transition  $b \rightarrow s\mu^-\mu^+$  in the SM.

### 1.1.1 $B_s \rightarrow \mu\mu$

:

Also in Table 1.1 we can see the data referred to the  $B_s \rightarrow \mu\mu$  decay we can easily see that the 4-fermion vertex factor involved is exactly the same of the  $B \rightarrow K^*\mu\mu$  decay because of the so called *crossing symmetry*.

With this decay indeed we need to be careful to the possibility that perturbativity condition is not satisfied because of  $c\bar{c}$  resonances. The diagram describing this decay in SM is the one in Figure 1.1.1 which is one of the so called *penguin diagrams*.

If the energy of the virtual photon is few GeV, there is a no-neglectable contribution due to the fact that the photon can produce a pair  $c\bar{c}$  nearby of the known resonances that have those quark as valence quark :  $J/\psi, \psi(1S), \psi(2S), \dots$

So we can have issues due to QCD non-perturbativity but on the other hand we have a final state which is *clean* theoretically, since is a two lepton state with no hadronic form factor. That's why we will treat this kind of process separately.

## 1.2 Main Processes for CC transitions

The main CC decay in which we are interested in is:

$$B \rightarrow D^* l \nu_l$$

We define the ratio analogously:

$$R_{D^*}^{tl} \equiv \frac{\mathcal{B}(B \rightarrow D^* \tau \nu_\tau)}{\frac{1}{2} \sum_{l=e,\mu} \mathcal{B}(B \rightarrow D^* l \nu_l)}.$$


In Table 1.2 we see the most important CC anomalies.

The main differences with the NC can be summarized as follows:

- From a first look to the Table 1.2 we can already notice that the significance of the anomaly is less important than the NC case;
- In SM this process is tree level generated (no penguins):

Table 1.2: Table with the most important anomalies in  $b \rightarrow sl\nu$  transition.  $\rho$  stays for the correlation between the ratio with  $D$  and  $D^*$ . Fonte:[1]

<i>Observable</i>	<i>Experiment</i>	<i>SM</i>
$\{R_D, R_{D^*}\}$	$\{0.337(30), 0.298(14)\}$	$\{0.299(3), 0.258(5)\}$
$\rho$	$-0.42$	$-$
$BR(B^- \rightarrow \tau^- \bar{\nu}_\tau)$	$1.09(24) \cdot 10^{-4}$	$0.812(54) \cdot 10^{-4}$



and that's why the SM prediction is more accurated than the FCNC process.

- in the  $B^- \rightarrow \tau \nu$  decay we have no issues linked to the charmonic resonances but the final state is not clean as the  $\mu\mu$  pair (the neutrino is invisible and the  $\tau$  decays briefly in hadrons). Also the rate is way bigger than the  $B_s \rightarrow \mu\mu$ .

If we begin to wonder about the NP's shape we notice that the NP couple mostly at the third generation for what concern quarks. A naive approach would be to guess a coupling growing with the mass (*Higgs-like*) but the NC decays suggest a smaller coupling for muons that the one for electrons. In the CC case, instead, we could imagine that the new physics couples more to the heaviest lepton: the  $\tau$ .

All this kind of considerations (and way more of them) will be the ones that allow us to build a low energy model independent Field Theory: the Effective Field Theory; which later will give us the shape of the BSM particles we need.

## 2

# Effective Field Theory approach

The first step to see how the low energy deviations from the SM is to parametrize these effects in a Quantum Field Theory which is independent from the physics at higher energies.

The usual example of an EFT is the Fermi Theory which is used to describe the weak processes of the SM when the energy is way below  $M_W = 80 \text{ GeV}$ . This theory can describe the weak processes at low energy ignoring the details of the UV Physics like the coupling between fermions and vectors that mediate the force, the mass of these mediators nor the theoretical nature of all the particles heavier than  $80 \text{ GeV}$ , which is called the *matching scale*.

In an EFT all the physics beyond the matching scale is contained in the numerical coefficient in front of the Lagrangian's operators. In fact the condition for the EFT to be a low energy version of an UV theory is for those coefficient to satisfy the *matching condition* that consists in impose the coefficients of the two theories, which normally run with the energy scale of the process via Renormalization Group Equations (RGE), to recreate the same transition amplitudes at an energy equal to the matching scale.

In the Fermi Theory, as is normally intended, we find the description of the 4-fermion weak processes that are tree-level generated in SM without flavour suppression. So, for first, we introduce the *Extended Fermi Theory* which takes in consideration basically all the allowed 4-fermion process.

Once this is done we can take just the semileptonic operators which concern the  $b$  quark and then try to find the proper flavour structure to explain the Anomalies without bothering the SM.

## 2.1 Extended Fermi Theory

The Fermi Theory was historically the first theory that described the weak processes as the beta decay or the muon decay.

When the EW theory was formulated from Glashow-Weinberg-Salam they of course needed to predict the low energy decays with same rates tested by the Fermi Theory. Basically the EW theory had to satisfy the matching condition with the Fermi Theory.





Figure 2.1: Left: Feynman diagram for muon decay in SM, the lagrangian terms responsible for this interaction is of the form  $\sim g_2 \bar{\psi}_{1L} \gamma^\mu \psi_{2L} W_\mu$ . Right: Feynman diagram for muon decay in Fermi Theory, the lagrangian terms responsible for this interaction is of the form  $\sim G_F \bar{\psi}_{1L} \gamma^\mu \psi_{2L} \bar{\psi}_{3L} \gamma_\mu \psi_{4L}$ .

In Figure 2.1 we can see how Feynman diagrams change passing from the SM to the Fermi Theory or, in other words, integrating out all the particles heavier than  $80 \text{ GeV}$ . Basically, if we study processes in which the energy available to produce particles is less than the rest energy of the  $W$  boson, we can ignore it in the spectrum of our theory and take count of its presence as virtual particle through a form factor which happens to be constant at low energy: the Fermi's constant  $G_F^1$ .

### 2.1.1 Flavour Changing Neutral Currents


A complete and synthetic discussion on Flavour Physics can be found in Appendix. Here we report just the main parts that make us understand how we get tree level operators in the EFT that can't appear at tree level in SM.

The most natural example consists is the so called Flavour Changing Neutral Currents (FCNC) which are not allowed at a classical level in SM. For example the vertex:




is not allowed from SM because the neutral interaction is diagonal in mass's basis (see Appendix). On the other hand that doesn't mean that we can't have the  $b \rightarrow sZ/\gamma$  at all, in fact at one loop level we can have:

<sup>1</sup> $G_F = 1.166376(7) \cdot 10^{-5} \text{ GeV}^{-2}$ , [2]



Now if we integrate out W's and Z bosons we get:



Here we don't have  $G_F$  as vertex factor, in fact there is to multiply the loop factor and the Cabibbo-Kobayashi-Maskawa (CKM) matrix elements and both those factors make the transition suppressed. This is why we say that the SM allows FCNCs but they are a suppressed effect and so a good place to look for NP.

In the previous diagram we neglected the Z boson because in Fermi Theory it doesn't belong to the spectrum. Indeed if the  $Z/\gamma$  were virtual in the SM diagram, perhaps producing a pair  $l^+l^-$ , we would have exactly the effective vertex mentioned in section 1 that allows the decay  $b \rightarrow sl^-l^+$  and so the decay  $B \rightarrow K^*l^-l^+$  which is one of the most interesting for us.

In SM we have CKM, loop factor, and gauge coupling that give me a prediction for the coefficient in front of the operator that mediate this decay: the Wilson coefficient. So if the measured Wilson coefficient is different from the theoretical one, it means that probably to build the EFT we need to integrate out other heavy particles which we don't know yet.

### 2.1.2 4-fermions operators

:

Once we have included all the 4-fermions operators in the effective lagrangian we can choose the ones we need to calculate the amplitudes of the processes of interest. Neglecting for the moment the Barion Number Violating (BNV) operators we have just three types of operators allowed by the gauge symmetry of the SM:

- Purely quark operators, which can mediate for instance the Kaon's decay in pions' channels. Nevertheless those operators are quite hard to match with the SM because we can have loop of gluons between the two currents which both couple with gluons. Since at the meson scale of energies the strong interaction is non-perturbative we have to consider a lot of contributions that are not easy to parametrize.
- Purely leptonic operators which can, for instance mediate the muon's decay. Those operators are used to the processes that have just leptons in the initial and final

state avoiding all the QCD's mess for both of them. In fact the cited muon decay is predicted so much accurately that it is the main process used to measure the Fermi's constant  $G_F$ .

- Semileptonic operators, which can mediate the charged pion's decay, but also all the processes to which we can address the B-Physics Anomalies. Of course the prediction are not clean as the purely leptonic case, nevertheless we have no gluon loop between the two currents and this tells us that we can see how the quark current renormalize just using global symmetries of QCD. In fact, in QCD, the vector current is conserved in the massless limit and so the quark current in a semileptonic decay is not affected by renormalization group of QCD. This fact reduces the theoretical uncertainty to the hadronic form factor.

The Lorentz and the  $SU(2)_L$  structure of these operators depend from the heavy mediators we integrate out to build the effective Lagrangian. Nevertheless all the possible operators we can generate are, a priori, non independent.

That's why we will choose a proper basis of effective semileptonic operators and, once we have generated the operators from UV couplings, we will project the operators we obtained on that basis.

Before to show the basis chosen we will explain what is the way to pass from a basis of 4-fermion operators to another through *Fierz identities*.

## 2.2 Fierz identities

Is possible to prove that, since any hermitian matrix  $N \times N$  can be written as

$$H = c_0 \mathbf{I} + \sum_{i=1}^{N^2-1} c_i T_i$$

where  $T_i$  are the generators of the fundamental representation of  $SU(N)$ , these generators satisfy the completeness condition:

$$\sum_{a=1}^{N^2-1} T_{ij}^a T_{kl}^a = \frac{1}{2} (\delta_{il} \delta_{jk} - \frac{1}{N} \delta_{ij} \delta_{kl}).$$

In the case  $N = 2$  since  $\sigma_a = 2T_a$  we find (neglecting the sum on  $a$ ):

$$\sigma_{a\ ij} \sigma_{kl}^a = 2\delta_{il} \delta_{kj} - \delta_{ij} \delta_{kl} \quad (2.1)$$

this relation can help us to write the EW structure to have all the currents contracted in the operators listed in 2.3, i.e. to have every single current transforming with an irreducible representation of  $SU(2)_L$ .

Comparing two terms with the indexes sorted at the way we can rewrite the relation 2.1:

$$\sigma_{ij}^a \sigma_{a\ kl} = \frac{1}{2} (3\delta_{il} \delta_{jk} - \sigma_{il}^a \sigma_{a\ kj}). \quad (2.2)$$

For what concern the Lorentz group, since the group is  $SU(2) \times SU(2)^*$ , the situation is

analogous.

Having written fermions as Weyl spinors we have explicit the relation between the Lorentz structure of the currents and Pauli matrices:

$$\sigma^\mu = (\mathbf{I}, \vec{\sigma}) \quad \bar{\sigma}^\mu = (\mathbf{I}, -\vec{\sigma})$$

These matrices allow us to write a current which transform as a vector under Lorentz group, combining two spinors transforming both as the  $(\frac{1}{2}, 0)$  or the  $(0, \frac{1}{2})$ , and so the current with explicit lorentz index is written with the *dotted notation*<sup>2</sup>

$$J_R^\mu = (\psi_R^\dagger)^\alpha (\sigma^\mu)_{\alpha\dot{\alpha}} \chi_R^{\dot{\alpha}} \sim (0, 1) \quad , \quad J_L^\mu = (\psi_L^\dagger)^{\dot{\alpha}} (\bar{\sigma}^\mu)_{\dot{\alpha}\alpha} \chi_L^\alpha \sim (1, 0)$$

In which the dots distinguish the two fundamental representation of the Lorentz group. The scalar bilinear instead is the product of two spinors transforming one as  $(\frac{1}{2}, 0)$  and the other as  $(0, \frac{1}{2})$ :

$$S = (\psi_R^\dagger)^\alpha \chi_{L\alpha} \sim (0, 0)$$

and same with all the indexes dotted.

It is useful to introduce the charge conjugated spinor:

$$(\psi_R)_\alpha^c \equiv \varepsilon_{\alpha\beta} (\psi_R^*)^\beta \sim (\frac{1}{2}, 0)$$

where the  $\varepsilon$  is the totally antisymmetric tensor in two dimensions (fixed  $\varepsilon^{12} = -\varepsilon^{21} = -\varepsilon_{12} = \varepsilon_{21} = 1$ ). This allow us to build a scalar with two spinors belonging to the same representation of the Lorentz group, eventually even two copies of the same spinor. In that case the  $\varepsilon$  tensor guarantees the antisymmetry typical of the singlet.

Now that we described the structure of the vector and the scalar identities we can take back 2.1 and, mindful that  $\sigma_{\alpha\dot{\alpha}}^0 = \bar{\sigma}_{\alpha\dot{\alpha}}^0 = \delta_{\alpha\dot{\alpha}}$  and including the Minkowski's metric, write:

$$\sigma_{\mu\alpha\dot{\alpha}} \sigma_{\beta\dot{\beta}}^\mu = 2(\delta_{\alpha\dot{\alpha}} \delta_{\beta\dot{\beta}} - \delta_{\alpha\dot{\beta}} \delta_{\beta\dot{\alpha}}) \quad (2.3)$$

now using that  $\varepsilon_{\alpha\beta} \varepsilon_{\gamma\delta} = \delta_{\alpha\gamma} \delta_{\beta\delta} - \delta_{\alpha\delta} \delta_{\beta\gamma}$  we find:

$$\sigma_{\alpha\dot{\alpha}}^\mu \sigma_{\mu\beta\dot{\beta}} = 2\varepsilon_{\alpha\beta} \varepsilon_{\dot{\alpha}\dot{\beta}} = -\sigma_{\alpha\dot{\beta}}^\mu \sigma_{\mu\beta\dot{\alpha}} \quad (2.4)$$

Now, generalizing at the overlined matrices:

$$\bar{\sigma}_\mu^{\dot{\alpha}\alpha} \bar{\sigma}^{\mu\dot{\beta}\beta} = 2\varepsilon^{\alpha\beta} \varepsilon^{\dot{\alpha}\dot{\beta}} \quad (2.5)$$

$$\sigma_{\alpha\dot{\alpha}}^\mu \bar{\sigma}_\mu^{\dot{\beta}\beta} = 2\delta_\alpha^\beta \delta_{\dot{\alpha}}^{\dot{\beta}}. \quad (2.6)$$

Despite to the similar aspect of the equations it is clear that the relations for  $SU(2)_L$  and for  $SU(2) \times SU(2)^*$  has to be used independently.

In some cases we have to use both to write the operators generated in the basis presented in section 2.3. In these cases the tensor structure could be complicated and eventually confusing, so we are going to present the useful results now to have them ready when we will handle the physics.

---

<sup>2</sup>The dotted indexes indicate  $SU(2)^*$ , i.e. the RH fermions.

### 2.2.1 Fierzing in the scalar lagrangian

When we introduce to the theory scalar Leptoquarks, integrating them out from the lagrangian we could generate four-fermion operators of this shape:

$$\overline{l^1 c} \epsilon q^1 \overline{q^2} \epsilon l^{2c} = l_{\alpha a}^1 q_{\beta b}^1 q_{\gamma c}^{2*} l_{\delta d}^{2*} \epsilon^{ab} \epsilon^{cd} \epsilon^{\alpha\beta} \epsilon^{\gamma\delta}$$

where  $i, j, k, l$  are EW indexes and  $\alpha, \beta, \gamma, \delta$  are Lorentz indexes and the  $\epsilon$  of the Lorentz's group comes from the charge coniugation. Since all the fermions involved happens to be LH we have to write them as a linear combination of  $O_S$  and  $O_T$ . To do that we need to write

$$\epsilon^{ab} \epsilon^{cd} \epsilon^{\alpha\beta} \epsilon^{\gamma\delta} = c_1 \delta^{ad} \delta^{bc} \overline{\sigma}_\mu^{\alpha\delta} \overline{\sigma}^\mu_{\beta\gamma} + c_2 \sigma^{ad} \sigma^{bc} \overline{\sigma}_\mu^{\alpha\delta} \overline{\sigma}^\mu_{\beta\gamma}$$

basically switching the position of  $q^1 \longleftrightarrow l^2$ .

Now we use that

$$\epsilon^{ab} \epsilon^{cd} = \delta^{ac} \delta^{bd} - \delta^{ad} \delta^{bc} = \frac{1}{2} \sigma_a^{ad} \sigma^{bc a} - \frac{1}{2} \delta^{ad} \delta^{bc}$$

Where we used 2.1 in the second equality.

Then, acting on the 2.5,

$$\epsilon^{\alpha\beta} \epsilon^{\gamma\delta} = \frac{1}{2} \overline{\sigma}_\mu^{\alpha\gamma} \overline{\sigma}^\mu_{\beta\delta} = -\frac{1}{2} \overline{\sigma}_\mu^{\alpha\delta} \overline{\sigma}^\mu_{\beta\gamma}$$

And multiplying is straightforward to obtain:

$$\overline{l^1 c} \epsilon q^1 \overline{q^2} \epsilon l^{2c} = \frac{1}{4} [q^{2\dagger} \overline{\sigma}^\mu q^1 l^{2\dagger} \overline{\sigma}_\mu l^1 - q^{2\dagger} \overline{\sigma}^\mu \sigma^a q^1 l^{2\dagger} \overline{\sigma}_\mu \sigma_a l^1] \quad (2.7)$$

Once obtained that result is easy to derive the same in the case of the contraction of two triplet scalar currents:

$$\overline{l^1 c} \epsilon \sigma_a q^1 \overline{q^2} \sigma^a \epsilon l^{2c} = l_{\alpha a}^1 q_{\beta b}^1 q_{\gamma c}^{2*} l_{\delta d}^{2*} (\sigma^a \epsilon)^{ab} (\epsilon \sigma_a)^{cd} \epsilon^{\alpha\beta} \epsilon^{\gamma\delta}$$

We already know how to treat the lorentz structure that will bring a factor  $-\frac{1}{2} \overline{\sigma}_\mu^{\alpha\delta} \overline{\sigma}^\mu_{\beta\gamma}$ . The EW structure instead is different:

$$\begin{aligned} \epsilon^{ae} \epsilon^{fd} \sigma_a^{eb} \sigma^{ce a} &= (\delta^{af} \delta^{ed} - \delta^{ad} \delta^{fe}) (2\delta^{ej} \delta^{kj} - \delta^{mj} \delta^{kn}) = \\ &= 2\delta^{ad} \delta^{bc} - 4\delta^{ad} \delta^{bc} + \delta^{ad} \delta^{bc} - \delta^{ac} \delta^{bd} = -(\delta^{ad} \delta^{bc} + \delta^{ac} \delta^{bd}) \end{aligned}$$

then recalling the 2.1

$$\begin{aligned} \sigma_{da}^a \sigma_{cb a} &= 2\delta^{ac} \delta^{bd} - \delta^{ad} \delta^{bc} \rightarrow \\ \epsilon^{ae} \epsilon^{fd} \sigma_a^{eb} \sigma^{ce a} &= -\frac{1}{2} (\sigma_a^{ad} \sigma^{bc a} + 3\delta^{da} \delta^{cb}) \end{aligned}$$

multiplying the factors coming from EW and Lorentz group:

$$\overline{l^1 c} \epsilon \sigma_a q^1 \overline{q^2} \sigma^a \epsilon l^{2c} = \frac{1}{4} [3q^{2\dagger} \overline{\sigma}^\mu q^1 l^{2\dagger} \overline{\sigma}_\mu l^1 + q^{2\dagger} \overline{\sigma}^\mu \sigma^a q^1 l^{2\dagger} \overline{\sigma}_\mu \sigma_a l^1] \quad (2.8)$$



## 2.3 Effective Semileptonic Lagrangian

As we mentioned many times so far we want to find the coefficients that parametrize the B-Physics Anomalies which appear just in semileptonic  $B$  mesons decays.

Since every four fermion operator has to be made of two spinor bilinears both being vector or scalar<sup>3</sup>. The choice we make is to avoid charge's conjugated spinors and, if possible, to have every bilinear being a colour singlet. This choice is due to the fact that in hadronic transitions is comfortable to have few shapes of quark operators in order to simplify them, when it's possible, in the computation of experimental observables.

Our first step is to collect all the effective operators that can contribute to those processes. For first we will list five operators that can be written as contraction of two vector currents.

If we assume for NP to be coupled just to LH fermions, as done by [3], we would need just two operators:

$$\begin{aligned}\mathcal{O}_S &= q_L^\dagger \bar{\sigma}^\mu q_L l_L^\dagger \bar{\sigma}_\mu l_L = (V^* u_L)^\dagger \bar{\sigma}^\mu (V^* u_L) e_L^\dagger \bar{\sigma}_\mu e_L + d_L^\dagger \bar{\sigma}^\mu d_L e_L^\dagger \bar{\sigma}_\mu e_L + \\ &\quad (V^* u_L)^\dagger \bar{\sigma}^\mu (V^* u_L) \nu_L^\dagger \bar{\sigma}_\mu \nu_L + d_L^\dagger \bar{\sigma}^\mu d_L \nu_L^\dagger \bar{\sigma}_\mu \nu_L\end{aligned}$$

$$\mathcal{O}_T = q_L^\dagger \bar{\sigma}^\mu \sigma_a q_L l_L^\dagger \bar{\sigma}_\mu \sigma^a l_L = 2(V^* u_L)^\dagger \bar{\sigma}^\mu d_L l_L^\dagger \bar{\sigma}_\mu \nu_L + 2d_L^\dagger \bar{\sigma}^\mu (V^* u_L) \nu_L^\dagger \bar{\sigma}_\mu l_L +$$

$$(V^* u_L)^\dagger \bar{\sigma}^\mu (V^* u_L) \nu_L^\dagger \bar{\sigma}_\mu \nu_L + d_L^\dagger \bar{\sigma}^\mu d_L l_L^\dagger \bar{\sigma}_\mu l_L - d_L^\dagger \bar{\sigma}^\mu d_L \nu_L^\dagger \bar{\sigma}_\mu \nu_L - (V^* u_L)^\dagger \bar{\sigma}^\mu (V^* u_L) l_L^\dagger \bar{\sigma}_\mu l_L$$

where we neglect the flavour indexes. Writing them explicitly we have, for instance, the singlet operator equal to  $\mathcal{O} = \mathcal{O}^{abcd} = q_L^a \bar{\sigma}^\mu q_L^b l_L^c \bar{\sigma}_\mu l_L^d$  with

$$q_L^i = \begin{pmatrix} V^{*ij} u_{Lj} \\ d_L^i \end{pmatrix}, \quad l_L^i = \begin{pmatrix} \nu_L^i \\ e_L^i \end{pmatrix}$$

Also the EW indexes are neglected and here that means that both the vector currents composing the operator are irreducible representation of  $SU(2)_L$  (two triplets for  $\mathcal{O}_T$  and two singlets for any other). Besides we want to point out that  $\mathcal{O}_T$  can mediate CC transitions as well as NC ones.


Including the RH fermions, we find other two operators compatible with the gauge symmetry  $SU(3)_c \times SU(2)_L \times U(1)_Y$  already present in SM:

$$\mathcal{O}_{LR1} = q_L^\dagger \bar{\sigma}^\mu q_L e_R^\dagger \bar{\sigma}_\mu e_R = (V^* u_L)^\dagger \bar{\sigma}^\mu (V^* u_L) e_R^\dagger \bar{\sigma}_\mu e_R + d_L^\dagger \bar{\sigma}^\mu d_L e_R^\dagger \bar{\sigma}_\mu e_R$$

$$\mathcal{O}_{LR2}^{u/d} = q_R^\dagger \sigma^\mu q_R l_L^\dagger \bar{\sigma}_\mu l_L = q_R^\dagger \sigma^\mu q_R \nu_L^\dagger \bar{\sigma}_\mu \nu_L + q_R^\dagger \sigma^\mu q_R e_L^\dagger \bar{\sigma}_\mu e_L$$

when the u/d means that we have two independent versions of the  $\mathcal{O}_{LR2}$  for  $q_R$  equal to the up or down quark's flavour triplet. These two operators can describe just NC transitions and so the flavour diagonal part is not much interesting. The flavour breaking contributes are as well generated in SM but suppressed for different reasons: the FCNC contribute given by  $\mathcal{O}_{LR1}$  described by the diagram:

<sup>3</sup>we could have even 2 rank tensor if we used Dirac representation for the fields



is accidentally suppressed because of the smallness of the *Weinberg's angle*  $\theta_W$  that suppresses the coupling between the Z boson and RH fermions;  $O_{LR2}$  is even more suppressed because, even if the Z this time couple to its favourite LH fermions, to have the FCNC for RH quarks we need to flip the chirality twice because of the Flavour theorem which states that flavour breaking couplings of the SM are allowed just for LH fermions, and so the contribute is suppressed due to the light mass of the quarks involved.

The last vector-vector operator is the one that, in SM, takes both the suppressions described above:

$$\mathcal{O}_R^{u/d} = q_R^\dagger \sigma^\mu q_R e_R^\dagger \sigma_\mu e_R$$

again in the  $u/d$  versions according to the flavour triplet involved.

Now we have three operators made up the contraction of two scalar currents. For first:

$$\begin{aligned}\mathcal{O}_S^u &= q_L^\dagger u_R \varepsilon l_L^\dagger e_R = (V^* u)_L^\dagger u_R e_L^\dagger e_R - d_L^\dagger u_R v_L^\dagger e_R \\ \mathcal{O}_S^d &= q_L^\dagger d_R e_R^\dagger l_L = (V^* u)_L^\dagger d_R e_R^\dagger v_L + d_L^\dagger d_R e_R^\dagger e_L\end{aligned}$$

in which  $\mathcal{O}_S^d$  could indeed be written as vector currents contraction through Fierz identities, but renouncing to the request of having currents transforming as  $SU(3)_c$  singlets. The  $\varepsilon$  tensor is the one introduced in section 2.2, a part that acts on  $SU(2)_L$  to select the singlet.

In the end we have the *LeptoQuark* operator:

$$\mathcal{O}_{LQ} = e_R^\dagger q_L \varepsilon u_R^\dagger l_L = e_R^\dagger (V^* u)_L u_R^\dagger e_L - e_R^\dagger d_L u_R^\dagger v_L$$

that can't be written as contraction of two colour singlets but just of two LQs scalar currents, unless we involve charge's conjugate fields, in fact

$$\varepsilon_{ij}\varepsilon_{kl} = \delta_{ik}\delta_{jl} - \delta_{il}\delta_{jk} \rightarrow q_L^c \varepsilon l_L e_R^\dagger u_R^c = \mathcal{O}_{LQ} - \mathcal{O}_S^u$$

that anyway wouldn't help the simplification of hadronic form factors, since we would have anyway a current which appears just in this operator.

It is straightforward to see, from the electric charge's eigenstates form, that all the scalar operators contribute both to CC and NC transitions,

We choose to separate the SM Lagrangian from the operators defined above, i.e. to use a Lagrangian that recreate the SM predictions in the limit  $C_S = C_T = C_{LR1} = C_{LR2}^{u/d} =$

$$C_R^{u/d} = C_{LQ} = C_S^u = C_S^d = 0:$$

$$\begin{aligned} \mathcal{L}_{EFT} = & \mathcal{L}_{SM} + C_S \mathcal{O}_S + C_T \mathcal{O}_T + C_{LQ} \mathcal{O}_{LQ} + C_{LR1} \mathcal{O}_{LR1} + \\ & \sum_{q=u,d} [C_R^q \mathcal{O}_R^q + C_{LR2}^q \mathcal{O}_{LR2}^q + C_S^q \mathcal{O}_S^q] \end{aligned} \quad (2.9)$$

in which, expliciting flavour indexes  $C\mathcal{O} = C_{abcd}\mathcal{O}^{abcd}$  and hence we define  $c_i$  as

$$C_i^{q_1 q_2 l_1 l_2} \equiv c_i \Lambda_i^{q_1 q_2 l_1 l_2}$$

that seem to be redundant but will be useful to describe separately the flavour structure separately from the rest. In our convention  $q_1$  and  $l_1$  are the ones appearing in the operator *daggered*.

From the experiments we acknowledge what are the observables of interest to explore the clues of NP. Plus they suggest us that NP, at leading level, doesn't concern the lightest families of quark and leptons and couples preferly to the heaviest.

### 3

## Some of the possible heavy bosons

As we previously said one possible way to modify the Wilson coefficients in an EFT is to introduce heavy particles to the theory which couple with the fermions involved in the processes we want to accomodate.

When we introduce new particle interacting with the SM particles we need to look carefully at the experimental constraints. In particular the particle has to satisfy two bounds:

- The contribution to the observable given by the diagrams in which the new particles appears as virtual particles has to show agreement with the experiments, both the ones who seem anomalous to the SM and the ones that has tested the SM.
- If the new particles interact with the particles that are smashing at the colliders, so the mass range allowed for those particles introduced has to not contain the energies explored at that colliders so far.

To begin we will see how we can accomodate the B-Physics Anomalies introducing different type of heavy vector and scalar bosons.

We will begin with the most familiar case to the ones who know the SM: colour-less vector charged under  $SU(2)_L$ . Indicating the quantum number as  $(SU(3)_c, SU(2)_L)_Y$  we can address to these particles  $B' \sim (\mathbf{1}, \mathbf{1})_0$  and  $W' \sim (\mathbf{1}, \mathbf{3})_0$ , where the names already suggest the connection with their SM's lighter sisters  $W, B$ <sup>1</sup>.

Then we will describe the vector Leptoquarks  $U_1 \sim (\mathbf{3}, \mathbf{1})_{2/3}$  and  $U_3 \sim (\mathbf{3}, \mathbf{3})_{2/3}$ .

In the end we will describe the behavior of some scalar Leptoquarks  $S_1 \sim (\mathbf{3}^*, \mathbf{1})_{1/3}$ ,  $S_3 \sim (\mathbf{3}^*, \mathbf{3})_{1/3}$  and  $R_2 \sim (\mathbf{3}, \mathbf{2})_{7/6}$ .

Once we have acknowledged what contributions are taken from the different bosons we will be ready to see what mediator, or what combination of mediators, are proper to accomodate the B-Physics anomalies without contradicting the Flavour tests done so far.

---

<sup>1</sup>That phenomenological are better known as  $W^\pm$ ,  $Z$  and the photon  $\gamma$  because of the so known EWSB.

### 3.1 Colour-less bosons

#### 3.1.1 $B' \sim (\mathbf{1}, \mathbf{1})_0$

The first candidate is the heaviest version of the EW singlet of the SM  $B$ :  $B' \sim (\mathbf{1}, \mathbf{1})_0$ . Apart from the bigger mass the main difference between these two bosons is the flavour structure, which is diagonal in SM because of the gauge symmetry and is free a priori instead for the BSM version.

Once introduced this boson the most general UV Lagrangian contains:

$$\begin{aligned} \mathcal{L}_{UV} \supset & \frac{M_{B'}^2}{2} B_\mu B^\mu - \frac{1}{4} B_{\mu\nu} B^{\mu\nu} + g_{B'} l_L^\dagger \bar{\sigma}^\mu \lambda_B^l l_L B_\mu + g_{B'} q_L^\dagger \bar{\sigma}^\mu \lambda_B^q q_L B_\mu + \\ & g_{B'} e_R^\dagger \sigma^\mu \lambda_B^e e_R B_\mu + g_{B'} u_R^\dagger \sigma^\mu \lambda_B^u u_R B_\mu + g_{B'} d_R^\dagger \sigma^\mu \lambda_B^d d_R B_\mu \end{aligned} \quad (3.1)$$

where the  $\lambda$  matrices are flavour matrices that are real since  $B'$  belongs to a real representation of the gauge group.

When we go down to energies  $\ll M_{B'}$   $B'$  can't be produced on shell anymore, and so it can't appear in a process but as virtual particle. In this condition is possible to *integrate it out* from the lagrangian, i.e. replacing  $B'^\mu$  with the solution of the equation of motion:

$$\begin{aligned} \frac{\delta \mathcal{L}}{\delta B_\mu} = 0 = & \frac{M_{B'}^2}{2} B^\mu + g_{B'} l_L^\dagger \bar{\sigma}^\mu \lambda_B^l l_L + g_{B'} q_L^\dagger \bar{\sigma}^\mu \lambda_B^q q_L \\ & g_{B'} e_R^\dagger \sigma^\mu \lambda_B^e e_R + g_{B'} u_R^\dagger \sigma^\mu \lambda_B^u u_R + g_{B'} d_R^\dagger \sigma^\mu \lambda_B^d d_R \rightarrow \\ B^\mu = & -\frac{2g_{B'}}{M_{B'}^2} [l_L^\dagger \bar{\sigma}^\mu \lambda_B^l l_L + q_L^\dagger \bar{\sigma}^\mu \lambda_B^q q_L + e_R^\dagger \sigma^\mu \lambda_B^e e_R + u_R^\dagger \sigma^\mu \lambda_B^u u_R + d_R^\dagger \sigma^\mu \lambda_B^d d_R]. \end{aligned}$$

obtaining an effective lagrangian:

$$\mathcal{L}_{EFT} \supset -2G_{B'} \left[ \left( \sum_{\psi} \bar{\psi} \lambda_B^\psi \sigma_\mu \psi \right) \left( \sum_{\chi} \bar{\chi} \lambda_B^\chi \sigma_\mu \chi \right) \right]$$

where  $G_{B'} \equiv \frac{g_{B'}^2}{M_{B'}^2}$  and the sums on  $\psi$  and  $\chi$  are done over every SM fermion.

Among the 25 four-fermion operators, made of all the possible combination of these five neutral currents we recognize some semileptonic operators, precisely all the ones constructed with two vectorcurrents  $SU(2)_L$ -singlet inside the list shown in section 2.3. The coefficients of these operators are equal to:


$$\begin{aligned} C_S = -2G_{B'} \lambda_B^q \lambda_B^l, \quad C_R^{u/d} = -2G_{B'} \lambda_B^e \lambda_B^{u/d} \\ C_{LR1} = -2G_{B'} \lambda_B^q \lambda_B^e, \quad C_{LR2}^{u/d} = -2G_{B'} \lambda_B^{u/d} \lambda_B^l \end{aligned}$$

Plus integrating out  $B'$  we generate lots of four-quarks and four-leptons operators that, according to the parameters, could generate uncomfortable contributes to observables tested by EW Precision Tests (EWPT) or to the meson mixing. In particular, since to accomodate the anomaly of the  $b \rightarrow sl^- l^+$  transition i need a non vacuum  $(\lambda_B^{q/u/d})^{bs} =$

$(\lambda_B^{q/u/d})^{sb}$  (where the / in this case means that has to be non vacuum at least in one of those three flavour matrices), we could produce an unpleasant tree-level contribute to  $B_s \leftrightarrow \bar{B}_s$  mixing which is one-loop induced in the SM. Also we need to point out that we haven't generate any operator which concerns CC transition, so  $B'$  can't be used to accomodate the anomaly of the  $b \rightarrow cl\nu$  decays.

### 3.1.2 $W' \sim (\mathbf{1}, \mathbf{3})_0$

Since  $B'$  can't accomodate CC anomalies, we expect that to be done by an  $SU(2)_L$  triplet as happens in SM, where  $b \rightarrow c\bar{l}\nu$  is mediated by  $W^\pm$  bosons which are the non  $SU(2)_L$  diagonal components of the gauge boson  $W_a$ . The amplitude for the fundamental process in the SM takes the leading contribute from



which takes a flavour suppression from the quark vertex equal to  $V_{cb}$  and is diagonal and flavour universal (in one word the flavour matrix is the  $\mathbf{1}_{3 \times 3}$ ) for what concern the lepton current.

Since the anomaly consists in a preference for the decay channel with the  $\tau$  as charged lepton, the most easy try to accomodate it is another  $SU(2)_L$  triplet (much heavier than the  $W^\pm$  to avoid collider constraints) with a non universal flavour structure in the lepton sector, that gets involved just when  $b$  quark belongs to the process.

Adding to SM  $W' \sim (\mathbf{1}, \mathbf{3})_0$  the most general additional terms to high energy Lagrangian are:

$$\mathcal{L}_{UV} \supset \frac{M_{W'}^2}{2} W_\mu'^a W_a'^\mu - \frac{1}{4} W_{\mu\nu}'^a W_a'^{\mu\nu} + g_{W'} l_L^\dagger \bar{\sigma}^\mu \lambda_W^l \sigma_a l_L W_\mu'^a + g_{W'} q_L^\dagger \bar{\sigma}^\mu \lambda_W^q \sigma_a q_L W_\mu'^a \quad (3.2)$$

To see how this new terms contribute to processes at hadronic scale we need again to integrate out  $W'$  to see what effective semileptonic operators are generated in the EFT.

$$\frac{\delta \mathcal{L}}{\delta W_\mu'^a} = 0 = \frac{M_{W'}^2}{2} W_a'^\mu + g_{W'} l_L^\dagger \bar{\sigma}^\mu \sigma_a \lambda_W^l l_L + g_{W'} q_L^\dagger \bar{\sigma}^\mu \sigma_a \lambda_W^q q_L \rightarrow$$

$$W_a'^\mu = -\frac{2g_{W'}}{M_{W'}^2} [l_L^\dagger \bar{\sigma}^\mu \sigma_a \lambda_W^l l_L + q_L^\dagger \bar{\sigma}^\mu \sigma_a \lambda_W^q q_L].$$

Then inserting it in the lagrangian we obtain:

$$\mathcal{L}_{EFT} \supset -2G_{W'} [l_L^\dagger \lambda_W^l \bar{\sigma}^\mu \sigma_a l_L l_L^\dagger \lambda_W^l \bar{\sigma}_\mu \sigma^a l_L + q_L^\dagger \lambda_W^q \bar{\sigma}^\mu \sigma_a q_L q_L^\dagger \lambda_W^q \bar{\sigma}_\mu \sigma^a q_L + 2q_L^\dagger \lambda_W^q \bar{\sigma}^\mu \sigma_a q_L l_L^\dagger \lambda_W^l \bar{\sigma}_\mu \sigma^a l_L] \quad (3.3)$$

in which we recognize the triplet operator  $\mathcal{O}_T$  in the last term implying

$$C_T = -4G_{W'}\lambda_W^l\lambda_W^q$$

with  $G_{W'} \equiv \frac{g_{W'}^2}{M_{W'}^2}$ . Notice that  $C_T \neq 0$  could accomodate CC anomalies.


Again we generate in addition four-leptons and four-quark operators that are unpleasant because of the tension they create between the anomalies and EWPT or meson mixing.

In conclusion colour-less vectors could together, a priori, accomodate both CC and NC anomalies but we can already see, without numerical analysis, that they generate a lot of unpleasant terms that can't be tuned easily once we included precision observables. In fact we will see, after a proper analysis, that colourless vectors don't constitute much more than an exercise to exemplify our *modus operandi*.

## 3.2 Leptoquarks

A possible way to avoid the tension between the anomalies and the precisely tested sector of the SM is to introduce bosons with quantum numbers different from the Higgs and gauge's bosons.

Since our purpose is to generate semileptonic operators, a good guess is to introduce bosons that connect quark and lepton sector at classical level, allowing the vertex



Bosons like that are known as *Leptoquark* and there could be different shape of them differing from the mass and the representation under their fields transform under Lorentz group and SM gauge group.

The only features common to all the LQs is that, to link a colour triplet (or antitriplet) and a colour singlet, it has to transform under the fundamental (or antifundamental) representation of  $SU(3)_c$ .


Being the LQs coloured they have to be confined and that protects the test on Baryon Number conservation at a phenomenological level. That's true in the limit in which the only vertex allowed is a quark-lepton vertex.

Also, in that limit, we would generate at tree level just semileptonic operators.

Nevertheless, because of  $SU(3)$  structure we have that:


$$\mathbf{3}^* \otimes \mathbf{3}^* \sim \mathbf{6}^* \oplus \mathbf{3}$$

and so, if the Lorentz and  $SU(2)_L \times U(1)_Y$  group allow, is possible to have



at tree level.

The presence of this vertex would generate the already mentioned meson mixing because of four-quarks tree level contributions. Also with both of the vertexes above allowed is possible to generate the amplitude



where the lepton is conjugated because of the  $U(1)_Y$  symmetry and that would violate the Barion Number even at a phenomenological level.


The easiest solution at the level of simplified model would be just to switch off the quark-quark vertex. But when we want to implement a properly said UV completion, kill that contribute could need a non natural fine tuning.

We won't concern about naturalness in our analysis but it is still worthy to discuss the BNV issue in the Leptoquarks frame.


### 3.2.1 Proton decay

Before B-Physics Anomalies came up there where different attempt to go BSM making LQs arise. This is because LQs arise naturally as gauge bosons of most of the GUT groups.

The most common reason to not believe in LQs (and as consequence in GUT as well) has always been that they could mediate proton decay  $p \rightarrow e^+ \pi^0$



or even in t-channel





Now the available experimental limit for the proton lifetime is  $\tau_p > 10^{34}$ ys and so if LQs arise the contribute sketched above should be very suppressed and that can happen if:

- $M_{LQ} \simeq \Lambda_{GUT} \simeq 10^{16}$  GeV to have a decoupling at low energies,
- a flavour structure that kills the contribute for the lightest families.

As we will see later  $M_{LQ}$  is supposed to be  $O(TeV)$  to accomodate the anomalies so clearly, to implement a complete theory, we would fall in the second case; in fact, as we said many times so far, NP is requested to not couple lightest families.

This could be not enough to say that our LQs don't concern proton decay in a naturalness'scenario if we go beyond tree level.

### 3.3 Vector Leptoquarks

For first we are going to analyze Leptoquark vectors named  $U_1 \sim (\mathbf{3}, \mathbf{1})_{2/3}$  and  $U_3 \sim (\mathbf{3}, \mathbf{3})_{2/3}$ . We will see that they are a pleasant option because, among the rest, since they couple to vector currents the value of the hypercharge  $Y = 2/3$  doesn't allow them to couple with a quark-quark current and then they can never mediate proton decay.

#### 3.3.1 $U_1 \sim (\mathbf{3}, \mathbf{1})_{2/3}$

The first LQ we take in consideration is the  $SU(2)_L$  singlet  $U_1$  which is one of the most popular in the available literature for reasons that will be clear at the end.

Including  $U_1$  in our theory we get an UV lagrangian that reads:

$$\mathcal{L}_{UV} \supset -\frac{1}{2}U_{1\mu\nu}^\dagger U_1^{\mu\nu} + M_{U_1}^2 U_{1\mu}^\dagger U_1^\mu + [g_U U_{1\mu} (q_L^\dagger \beta_{U_1}^L \bar{\sigma}^\mu l_L + d_R^\dagger \beta_{U_1}^R \sigma^\mu e_R) + h.c.] \quad (3.4)$$

One difference that can be seen immediately is that belonging to the fundamental representation of  $SU(3)_c$ , which is complex, the current coupled to it is complex too and so, in general, the flavour matrices  $\beta$  are complex in general.

Treating the physics at the hadronic scale we can integrate out  $U_1$  from the Lagrangian through the equation of motion:

$$\frac{\delta \mathcal{L}}{\delta U_{1\mu}^\dagger} = 0 = M_{U_1}^2 U_1^\mu + g_{U_1} [l_L^\dagger \beta_{U_1}^{*L} \bar{\sigma}^\mu q_L + e_R^\dagger \beta_{U_1}^{*R} \sigma^\mu d_R] \rightarrow$$

$$U_1^\mu = -\frac{g_{U_1}}{M_{U_1}^2} [l_L^\dagger \beta_{U_1}^{*L} \bar{\sigma}^\mu q_L + e_R^\dagger \beta_{U_1}^{*R} \sigma^\mu d_R]$$

Obtaining

$$\mathcal{L}_{EFT} \supset -G_{U_1} [l_L^\dagger \beta_{U_1}^L \bar{\sigma}^\mu q_L q_L^\dagger \beta_{U_1}^{*L} \bar{\sigma}_\mu l_L + 2l_L^\dagger \beta_{U_1}^L \bar{\sigma}^\mu q_L d_R^\dagger \beta_{U_1}^{*R} \sigma_\mu e_R + e_R^\dagger \beta_{U_1}^R \sigma^\mu d_R d_R^\dagger \beta_{U_1}^{*R} \sigma^\mu e_R + h.c.]$$

where,  $G_{U_1} \equiv \frac{g_{U_1}^2}{M_{U_1}^2}$ .

The effective lagrangian is made of a sum of semileptonic operators, now we need to project them on the basis listed in section 2.3.

The second term is proportional to  $\mathcal{O}_S^d$  because of the relation  $\bar{\sigma}_{\alpha\alpha}^\mu \sigma_{\mu}^{\beta\dot{\beta}} = 2\delta_{\alpha}^{\beta} \delta_{\dot{\alpha}}^{\dot{\beta}}$  it's precisely equal to  $2\mathcal{O}_S^d$ .

The lorentz structure of the first and the secon get us a minus sign in the first and the third term because of the relations  $\sigma_{\alpha\dot{\alpha}}^\mu \sigma_{\mu\beta\dot{\beta}} = 2\varepsilon_{\dot{\alpha}\dot{\beta}} \varepsilon_{\alpha\beta} = -\sigma_{\alpha\dot{\beta}}^\mu \sigma_{\mu\beta\dot{\alpha}}$  and  $\sigma_{\alpha\dot{\alpha}}^\mu \sigma_{\mu\beta\dot{\beta}} = 2\varepsilon_{\dot{\alpha}\dot{\beta}} \varepsilon_{\alpha\beta} = -\sigma_{\alpha\dot{\beta}}^\mu \sigma_{\mu\beta\dot{\alpha}}$ .

Using these relations we see that the third term is equal to  $-\mathcal{O}_R$  while the first term seems equal to  $-\mathcal{O}_S$  apart from the fact that the quark and the lepton current aren't  $SU(2)_L$  singlets. In fact both quarks contract the  $SU(2)_L$  index with a lepton, to project that operator on our basis we need to recall 2.1

$$\sigma_{ad}^i \sigma_{cbi} = 2\delta_{ab}\delta_{cd} - \delta_{ad}\delta_{bc} \rightarrow \delta_{ab}\delta_{cd} = \frac{\sigma_{ad}^i \sigma_{cbi} + \delta_{ad}\delta_{cb}}{2}$$

to rewrite the first term as  $-\frac{1}{2}(\mathcal{O}_S + \mathcal{O}_T)$ . Adding up everything we obtain:

$$\mathcal{L}_{EFT} \supset G_{U_1} [Re[\beta_{U_1}^L \beta_{U_1}^{*L}](\mathcal{O}_S + \mathcal{O}_T) + (\beta_{U_1}^R \beta_{U_1}^{*R} \mathcal{O}_R^d - 4\beta_{U_1}^R \beta_{U_1}^{*L} \mathcal{O}_S^d + h.c.)] \quad (3.5)$$

that means in other words

$$C_S = C_T \equiv G_{U_1} Re[\beta_{U_1}^{L*} \beta_{U_1}^L] \quad C_R^d \equiv G_{U_1} \beta_{U_1}^{R*} \beta_{U_1}^R \quad C_S^d \equiv -4G_{U_1} \beta_{U_1}^{R*} \beta_{U_1}^L$$

The first feature we notice is that, even though we take all the possible terms in the UV lagrangian, just semileptonic contribute are generated. This means that when we accomodate the anomalies we need to be mindful of constraints that come just from semileptonic observables.

Without considering the details of the computation we can also see how one of this constraints is avoided from the fact that  $C_S = C_T$ . In fact  $C_S = C_T \neq 0$  gives us a way to accomodate the anomaly on  $b \rightarrow sl^+l^-$  where the amplitude due to NP effects is (considering just purely LH contributions)  $\propto (C_S + C_T)$  as can be seen from the electric charge eigenstates form of  $\mathcal{O}_S$  and  $\mathcal{O}_T$  shown in section 2.3.

From that form we can see also that the transition  $b \rightarrow s\bar{\nu}$  is linked to  $b \rightarrow sl^+l^-$  by the gauge symmetry and results to be  $\propto C_S - C_T$  which means no contribution coming from the integration of  $U_1$  as shown by [3]. This is very pleasant since  $\mathcal{B}(B \rightarrow K^*\bar{\nu})$  is a rare decay extremely suppressed in SM and very constrained by experimental data. Nevertheless  $U_1$  doesn't contribute to that decay even taking count of the coupling to RH fermions which can as well help to accomodate the anomalies.

Another reason to support  $U_1$  is that it comes naturally as gauge boson in the UV completion proposed by Pati and Salam [4] that will be briefly discussed later.

### 3.3.2 $U_3 \sim (\mathbf{3}, \mathbf{3})_{2/3}$

The vector  $SU(2)_L$  triplet LQ couples just to the triplet LQ current  $J_T^a = q_L^\dagger \bar{\sigma}^\mu \sigma^a \beta_{U_3} l_L$  i.e. just to LH fermions so, for what said in section 2.3, we expect to generate at low energy a combination of  $\mathcal{O}_S$  and  $\mathcal{O}_T$ .

The UV lagrangian gains the following terms:

$$\mathcal{L}_{UV} \supset -\frac{1}{2} U_{3\mu\nu}^\dagger U_{3a}^{\mu\nu} + M_{U_3}^2 U_{3a}^\dagger U_{3\mu}^a + g_{U_3} [U_{3\mu}^a q_L^\dagger \beta_{U_3} \bar{\sigma}^\mu \sigma_a l_L + h.c.] \quad (3.6)$$

To build the effective lagrangian at  $m_b$  scale we set as always

$$\frac{\delta \mathcal{L}}{\delta U_{3\mu}^a} = 0 = M_{U_3}^2 U_{3a}^{mu} + g_{U_3} l_L^\dagger \beta_{U_3}^* \bar{\sigma}^\mu \sigma_a q_L \rightarrow$$

$$U_{3a}^\mu = -\frac{g_{U_3}}{M_{U_3}^2} l_L^\dagger \beta_{U_3}^* \bar{\sigma}^\mu \sigma_a q_L$$

generating

$$\mathcal{L}_{EFT} \supset -G_{U_3} [q_L^\dagger \beta_{U_3} \bar{\sigma}^\mu \sigma_a l_L l_L^\dagger \beta_{U_3}^* \bar{\sigma}_\mu \sigma^a q_L + h.c.]$$

Now we get a minus sign as in the  $U_1$  case to make colour singlet currents, meanwhile the EW structure recalls 2.2

$$\sigma_{ij}^a \sigma_{a\ kl} = \frac{1}{2} (3\delta_{il}\delta_{jk} - \sigma_{il}^a \sigma_{a\ kj}) \rightarrow$$

$$\mathcal{L}_{EFT} \supset G_{U_3} \text{Re}[\beta_{U_3} \beta_{U_3}^*] [3\mathcal{O}_S - \mathcal{O}_T]$$

in which we explicited the (+h.c.) because both operators are hermitian. The Wilson coefficients are equal to

$$C_S = -3C_T \equiv 3G_{U_3} \text{Re}[\beta_{U_3} \beta_{U_3}^*]$$

that can accomodate both CC and NC anomalies, nevertheless since we mentioned for  $U_1$  the constraint from  $B \rightarrow K \bar{\nu}$  we need to point out that this time the constraint is not passed automatically and could need a fine tuning.

To summarize the features of vector LQs we would say that they are a good possibility since they affect at tree level just semileptonic processes even if we include all possible terms allowed by SM symmetries.

Particularly  $U_1$  is mentioned many times in the literature as the only one that can accomodate both anomalies alone passing most of the constraints, while  $U_3$  is not that appreciated as solution of the anomalies but possibly it can be combined with other mmediators to build a relevant model.

### 3.4 Scalar Leptoquarks

In the end of this list we will show what are the scalar LQs that can be helpful for our purposes.

By convention the ones belonging to real representation of  $SU(2)_L$  are defined in the antifundamental of  $SU(3)_c$  i.e. we have  $S_1 \sim (3^*, 1)_{1/3}$ ,  $S_3 \sim (3^*, 3)_{1/3}$  and  $R_2 \sim (3, 2)_{7/6}$ . In literature sometimes another scalar LQ is mentioned :  $\tilde{R}_2 \sim (3, 2)_{1/6}$ ; nevertheless it doesn't serve the purpose properly while it has more than one quark-quark coupling, that's why we have chosen to neglect it.

We will see that the scalar currents that couple LQs can be of many shapes for two reasons. For first the values of the hypercharge allow  $S_1$  and  $S_3$  to couple a quark-quark current, secondly the scalar current has to be made by  $(\psi_{L/R})^\dagger \chi_{R/L}$  and it is possible to use the charge conjugation to flip the chirality since  $(\psi_L^c) \sim (0, \frac{1}{2})$  and viceversa. That was possible also for vector LQ but there the hypercharge wasn't the proper to a current  $(\psi_{L/R})^c \sigma^\mu \chi_{R/L}$  for any  $\psi_L, \chi_R \in \{q_L, l_L, u_R, d_R, e_R\}$ .

So, to see what current couples to a given LQ, we have to see all the bilinears, according to  $SU(2)_L$ , with the right hypercharge, mindful that the charge conjugation brings a minus sign to it.

#### 3.4.1 $S_1 \sim (3^*, 1)_{1/3}$

A scalar current which couples with a  $SU(2)_L$  singlet has to involve charge's conjugation. Mindful of this point we find that, this time, the hypercharge allows a quark-quark coupling

Precisely the UV lagrangian that includes  $S_1$  is:

$$\begin{aligned} \mathcal{L}_{UV} \supset (D_\mu S_1)^\dagger D^\mu S_1 + M^2 S_1^\dagger S_1 + \\ + g_{S_1} S_1^i [q_{Li}^{\dagger c} \beta_{S_1}^L \epsilon l_L + u_{Ri}^{\dagger c} \beta_{S_1}^R e_R + \overline{q_L^j} \lambda_{S_1} \epsilon q_L^c \epsilon_{ijk}] + h.c. \end{aligned} \quad (3.7)$$

Where the  $\epsilon$  guarantees the current to be a EW singlet.

Integrating out  $S_1$  we need to explicit the colour index, since this time is not contracted trivially in all the terms:

$$\frac{\delta \mathcal{L}_{UV}}{\delta S_1^{\dagger i}} = 0 = M_{S_1}^2 S_{1a} + g_{S_1} [l_L^{\dagger} \epsilon \beta_{S_1}^{L*} q_{Li}^c + e_R^{\dagger} \beta_{S_1}^{R*} u_{Ri}^c + q_L^{\dagger c j} \lambda_{S_1}^* \epsilon q_L^k \epsilon_{ijk}] \rightarrow$$

$$S_{1i} = -\frac{g_{S_1}}{M_{S_1}^2} [\overline{l_L} \beta_{S_1}^* \epsilon q_{Li}^c + e_R^{\dagger} \beta_{S_1}^{R*} u_{Ri}^c + \overline{q_L^j} \lambda_{S_1}^* \epsilon q_L^k \epsilon_{ijk}]$$

Inserting this term in the UV lagrangian we would obtain nine terms, for simplicity we here neglect all the BNV terms,

$$\begin{aligned} \mathcal{L}_{EFT} \supset -G_{S_1} [\overline{l_L} \beta_{S_1}^* \epsilon q_{Li}^c + e_R^{\dagger} \beta_{S_1}^{R*} u_{Ri}^c + \overline{q_L^j} \lambda_{S_1}^* \epsilon q_L^k \epsilon_{ijk}] [q_{Li}^{\dagger c} \beta_{S_1}^L \epsilon l_L + u_{Ri}^{\dagger c} \beta_{S_1}^R e_R + \overline{q_L^j} \lambda_{S_1} \epsilon q_L^c \epsilon_{ijk}] + h.c. = \\ -G_{S_1} [l_L^{\dagger} \beta_{S_1}^{L*} \epsilon q_{Li}^c q_{Li}^{\dagger c} \beta_{S_1}^L \epsilon l_L + e_R^{\dagger} \beta_{S_1}^{R*} u_{Ri}^c q_{Li}^{\dagger c} \beta_{S_1}^L \epsilon l_L + q_L^{\dagger c j} \lambda_{S_1}^* \epsilon q_L^k \epsilon_{ijk} \overline{q_L^j} \lambda_{S_1} \epsilon q_L^c \epsilon_{ilm} + \end{aligned}$$

$$l_L^\dagger \beta_{S_1}^{L*} \varepsilon q_{Li}^c u_{Ri}^\dagger \beta_{S_1}^R e_R + e_R^\dagger \beta_{S_1}^{R*} u_{Ri}^c u_{Ri}^\dagger \beta_{S_1}^R e_R + (\text{BNV terms})]$$

Here we can see three terms describing semileptonic contributes and a four-quark term that can generate amplitudes contributing to meson mixing.

To project these terms on our operators basis we need to recall the relations obtained using Fierz Identities of EW group together with the ones of the Lorentz group, in particular we need 2.7:

$$\overline{l^1}^c \varepsilon q^1 \overline{q^2} \varepsilon l^{2c} = \frac{1}{4} [q^{2\dagger} \overline{\sigma}^\mu q^1 l^{2\dagger} \overline{\sigma}_\mu l^1 - q^{2\dagger} \overline{\sigma}^\mu \sigma^a q^1 l^{2\dagger} \overline{\sigma}_\mu \sigma_a l^1]$$

that projects the semileptonic term with LH fermions. Then we notice that the left-right semileptonic term is generated in pair with his hermitian conjugate. Then, since we add (+h.c.) it gains a 2 factor in front. To project that term we need to use that:

$$\varepsilon_{ij} \varepsilon_{kl} = \delta_{ik} \delta_{jl} - \delta_{il} \delta_{jk}$$

obtaining two scalar operators that we recognize to be  $\mathcal{O}_{LQ}$  and  $-\mathcal{O}_{S^\dagger}^u$ .

In the end, the completely RH term turns to a vector-vector operator using

$$\varepsilon_{\alpha\beta} \varepsilon_{\dot{\alpha}\dot{\beta}} = -\frac{1}{2} \sigma_{\alpha\dot{\alpha}}^\mu \sigma_{\mu\dot{\beta}}^\nu$$

obtaining  $-\frac{1}{2} \mathcal{O}_R$ .

Adding up all the terms we obtain the following effective lagrangian:

$$\begin{aligned} \mathcal{L}_{EFT} \supset -G_{S_1} & \left[ \frac{1}{4} \beta_{S_1}^L \beta_{S_1}^{L*} \mathcal{O}_S - \frac{1}{4} \beta_{S_1}^L \beta_{S_1}^{L*} \mathcal{O}_T + 2 \beta_{S_1}^{R*} \beta_{S_1}^L \mathcal{O}_{LQ} - 2 \beta_{S_1}^{R*} \beta_{S_1}^L \mathcal{O}_S^{u\dagger} - \frac{1}{2} \beta_{S_1}^{R*} \beta_{S_1}^R \mathcal{O}_R^u \right. \\ & \left. + \overline{q^j} \lambda_{S_1}^* \varepsilon q_L^k \varepsilon_{ijk} \overline{q_L} \lambda_{S_1} \varepsilon q_{Lm}^c \varepsilon^{ilm} + (\text{BNV terms}) \right] + h.c. \end{aligned} \quad (3.8)$$

Here we have generated different semileptonic contributes that contribute both to CC and NC processes and a term that contributes to tree level amplitude of meson mixing, plus some BNV terms that we neglected.

There is one substantial difference with the colour-less case, in fact here we can set  $\lambda_{S_1} = 0$  without affecting semileptonic contributes and turn off all four-quark and BNV terms.

The Wilson coefficients of semileptonic operators result to be

$$C_S = -C_T \equiv -\frac{G_{S_1}}{4} \beta_{S_1}^{L*} \beta_{S_1}^L, \quad C_{LQ} = -C_S^{u*} = 2G_{S_1} \beta_{S_1}^{R*} \beta_{S_1}^L, \quad C_R = -\frac{1}{2} G_{S_1} \beta_{S_1}^{R*} \beta_{S_1}^R$$

For what we said about  $U_1$  is easy to see that the purely LH tree level contribute to  $b \rightarrow sl^+ l^-$ , which is proportional to  $C_S + C_T$ , turns out to be null and that takes out from our toolbox a possible weapon to accomodate NC anomalies.

Nevertheless as we will see  $S_1$  is a good candidate which appears in many UV scenarios.

### 3.4.2 $R_2 \sim (3, 2)_{7/6}$

The scalar doublet with  $Y = 7/6$  couples to a left-right current to contract the  $SU(2)_L$  index.

That means that it can't generate the purely LH operators  $\mathcal{O}_S, \mathcal{O}_T$  and this is the reason why [3] neglects every LQ EW doublet.

The lagrangian terms allowed are:

$$\mathcal{L}_{UV} \supset (D_\mu R_2)^\dagger D^\mu R_2 + M_{R_2}^2 R_2^\dagger R_2 + g_{R_2} R_2 [u_R^\dagger \beta_{R_2}^l l_L \varepsilon + q_L^\dagger \beta_{R_2}^q e_R] + h.c. \quad (3.9)$$

where replacing at low energies the equation of motion

$$\frac{\delta \mathcal{L}_{UV}}{\delta R_2^\dagger} = 0 = M_{R_2}^2 R_2 + g_{R_2} [\varepsilon l_L^\dagger \beta_{R_2}^{l*} u_R + e_R^\dagger \beta_{R_2}^{q*} q_L] \rightarrow$$

$$R_2 = -\frac{g_{R_2}}{M_{R_2}^2} [\varepsilon l_L^\dagger \beta_{R_2}^{l*} u_R + e_R^\dagger \beta_{R_2}^{q*} q_L]$$

we find

$$\mathcal{L}_{EFT} \supset -G_{R_2} [-u_R^\dagger \beta_{R_2}^l l_L l_L^\dagger \beta_{R_2}^{l*} u_R + u_R^\dagger \beta_{R_2}^l l_L e_R^\dagger \beta_{R_2}^{q*} q_L + q_L^\dagger \beta_{R_2}^{q*} e_R l_L^\dagger \beta_{R_2}^{l*} u_R + q_L^\dagger \beta_{R_2}^q e_R e_R^\dagger \beta_{R_2}^{q*} q_L] + h.c.$$

where we used  $\varepsilon \cdot \varepsilon = -\mathbf{1}$ . Apart from the flavour structure, the second and the third term are  $\mathcal{O}_{LQ}$  and its hermitian conjugate, then using the (+h.c.) we can write the sum of them as  $2\text{Re}[\beta_{R_2}^{l*} \beta_{R_2}^q] \mathcal{O}_{LQ}$ .

The first and the fourth terms can be fierzed through

$$\delta_a^b \delta_c^d = \frac{1}{2} \sigma_{ac}^\mu \overline{\sigma}_{\mu}^{bc}$$

to obtain  $\frac{1}{2} \mathcal{O}_{LR1}$  and  $\frac{1}{2} \mathcal{O}_{LR2}^u$ .

Adding up we obtain

$$\mathcal{L}_{EFT} \supset -\frac{G_{R_2}}{2} [-\beta_{R_2}^l \beta_{R_2}^{l*} \mathcal{O}_{LR2}^u + 4\text{Re}[\beta_{R_2}^l \beta_{R_2}^{q*}] \mathcal{O}_{LQ} + \beta_{R_2}^q \beta_{R_2}^{q*} \mathcal{O}_{LR1}] + h.c. \quad (3.10)$$

Again we generated just semileptonic terms even including all possible terms.

The Wilson coefficients result to be

$$C_{LR2}^u = \frac{1}{2} G_{R_2} \beta_{R_2}^l \beta_{R_2}^{l*}, \quad C_{LQ} = -\frac{1}{2} G_{R_2} \beta_{R_2}^q \beta_{R_2}^{q*}, \quad C_{LR1} = -2 G_{R_2} \text{Re}[\beta_{R_2}^l \beta_{R_2}^{q*}]$$

### 3.4.3 $S_3 \sim (3^*, 3)_{1/3}$

The last heavy boson we want to compute the affection at low energy is the scalar EW triplet.

As  $U_3$  it couples with a triplet current, hence it doesn't interact with RH fermions. The main difference with  $U_3$  is that it allows quark-quark coupling, in fact the UV lagrangian turns out to be:

$$\mathcal{L}_{UV} \supset (D_\mu S_{3a})^\dagger D^\mu S_{3a}^a + M_{S_3}^2 S_{3a}^\dagger S_{3a}^a + g_{S_3} S_{3a}^{ia} (\overline{q_L}^j \lambda_{S_3} \varepsilon \sigma_a q_L^k \varepsilon_{ijk} + \overline{q_L^c} \beta_{S_3} \varepsilon \sigma_a l_L) + h.c. \quad (3.11)$$

in which  $i = 1, 2, 3$  is the colour index and  $a = 1, 2, 3$  is the EW index. Again the annoying part could be turned off without affecting semileptonic operators, as for  $S_1$ . Integrating it out

$$\frac{\delta \mathcal{L}_{UV}}{\delta S_3^{ia\dagger}} = 0 = M_{S_3}^2 S_{3ia} + g_{S_3} [\bar{q}_L^c \lambda_{S_3}^* \varepsilon \sigma_a q_{Lk} \varepsilon^{ijk} + \bar{l}_L \beta_{S_3}^* \varepsilon \sigma_a q_L^c] \rightarrow$$

$$S_{3ia} = -\frac{g_{S_3}}{M_{S_3}^2} [\bar{q}_L^c \lambda_{S_3}^* \varepsilon \sigma_a q_{Lk} \varepsilon^{ijk} + \bar{l}_L \beta_{S_3}^* \varepsilon \sigma_a q_L^c]$$

hence the EFT lagrangian results

$$\mathcal{L}_{EFT} \supset -G_{S_3} [\bar{q}_{Li} \beta_{S_3} \varepsilon \sigma_a l_L \bar{l}_L \beta_{S_3}^* \varepsilon \sigma_a q_L^c + \bar{q}_{Li} \beta_{S_3} \varepsilon \sigma_a l_L \bar{q}_{Lj} \lambda_{S_3}^* \varepsilon \sigma_a q_{Lk} \varepsilon^{ijk} +$$

$$\bar{q}_L^j \lambda_{S_3} \varepsilon \sigma_a q_L^k \varepsilon_{ijk} \bar{l}_L \beta_{S_3}^* \varepsilon \sigma_a q_L^c + \bar{q}_L^j \lambda_{S_3} \varepsilon \sigma_a q_L^k \varepsilon_{ijk} \bar{q}_{Ll} \lambda_{S_3}^* \varepsilon \sigma_a q_{Lm} \varepsilon^{ilm}]$$

The first term is the only one affecting semileptonic processes at tree level. To project it on our basis we remember that we obtained 2.8

$$\bar{l}^c \varepsilon \sigma_a q^1 \bar{q}^2 \sigma^a \varepsilon l^{2c} = \frac{1}{4} [3q^{2\dagger} \bar{\sigma}^\mu q^1 l^{2\dagger} \bar{\sigma}_\mu l^1 + q^{2\dagger} \bar{\sigma}^\mu \sigma^a q^1 l^{2\dagger} \bar{\sigma}_\mu \sigma_a l^1]$$

which means

$$\mathcal{L}_{EFT} \supset -G_{S_3} [\frac{3}{4} \beta_{S_3}^* \beta_{S_3} \mathcal{O}_S + \frac{1}{4} \beta_{S_3}^* \beta_{S_3} \mathcal{O}_T + \bar{q}_{Li} \beta_{S_3} \varepsilon \sigma_a l_L \bar{q}_{Lj} \lambda_{S_3}^* \varepsilon \sigma_a q_{Lk} \varepsilon^{ijk} +$$

$$\bar{q}_L^j \lambda_{S_3} \varepsilon \sigma_a q_L^k \varepsilon_{ijk} \bar{l}_L \beta_{S_3}^* \varepsilon \sigma_a q_L^c + \bar{q}_L^j \lambda_{S_3} \varepsilon \sigma_a q_L^k \varepsilon_{ijk} \bar{q}_{Ll} \lambda_{S_3}^* \varepsilon \sigma_a q_{Lm} \varepsilon^{ilm}] + h.c. \quad (3.12)$$

As already mentioned, meson mixing and BNV amplitudes would arise at tree level just for non vacuum values of the flavour matrix  $\lambda_{S_3}$  that we can turn off for our purposes. The Wilson coefficients of semileptonic terms are

$$C_S = 3C_T \equiv -\frac{3}{4} G_{S_3} \beta_{S_3}^* \beta_{S_3}$$

that means, a priori, contributions to both NC and CC processes.

## 3.5 Summary

The last thing we need to decide what recipes are suitable to accomodate the anomalies is to compute what values of Wilson coefficients we need to accomodate the B-Physics anomalies and pass the constraint given by the tests to Flavour Physics done so far.

Nevertheless since we have introduced many parameters in section 2.3 to fit the observables to effective parameters we will need to do some assumptions on the UV scenario to reduce the number of parameters, these could be suggested by looking at the contributions generated by the boson listed above. So it is useful to see some features that can be already pointed out at this point. In tables 3.1 we have summarized all the

Figure 3.1: Wilson coefficients corresponding to effective generators generated by the integration of heavy vector bosons (above) and scalar bosons (below).

	$B'$	$W'$	$U_1$	$U_3$
$C_S$	$-2G_{B'}\lambda_B^q\lambda_B^l$	$\emptyset$	$G_{U_1}\text{Re}[\beta_{U_1}^{L*}\beta_{U_1}^L]$	$3G_{U_3}\beta_{U_3}\beta_{U_3}^*$
$C_T$	$\emptyset$	$-4G_{W'}\lambda_W^q\lambda_W^l$	$G_{U_1}\text{Re}[\beta_{U_1}^{L*}\beta_{U_1}^L]$	$-G_{U_3}\beta_{U_3}\beta_{U_3}^*$
$C_{LR1}$	$-2G_{B'}\lambda_B^q\lambda_B^e$	$\emptyset$	$\emptyset$	$\emptyset$
$C_{LR2}^u$	$-2G_{B'}\lambda_B^u\lambda_B^l$	$\emptyset$	$\emptyset$	$\emptyset$
$C_{LR2}^d$	$-2G_{B'}\lambda_B^d\lambda_B^l$	$\emptyset$	$\emptyset$	$\emptyset$
$C_R^u$	$-2G_{B'}\lambda_B^u\lambda_B^e$	$\emptyset$	$\emptyset$	$\emptyset$
$C_R^d$	$-2G_{B'}\lambda_B^d\lambda_B^e$	$\emptyset$	$G_{U_1}\beta_{U_1}^{R*}\beta_{U_1}^R$	$\emptyset$
$C_{LQ}$	$\emptyset$	$\emptyset$	$\emptyset$	$\emptyset$
$C_S^u$	$\emptyset$	$\emptyset$	$\emptyset$	$\emptyset$
$C_S^d$	$\emptyset$	$\emptyset$	$-4G_{U_1}\beta_{U_1}^{R*}\beta_{U_1}^R$	$\emptyset$

	$S_1$	$R_2$	$S_3$
$C_S$	$-G_{S_1}\beta_{S_1}^L\beta_{S_1}^{L*}$	$\emptyset$	$-\frac{3}{4}G_{S_3}\beta_{S_3}^*\beta_{S_3}$
$C_T$	$-G_{S_1}\beta_{S_1}^{L*}\beta_{S_1}^L$	$\emptyset$	$-G_{S_3}\beta_{S_3}^*\beta_{S_3}$
$C_{LR1}$	$\emptyset$	$-2G_{R_2}\text{Re}[\beta_{R_2}^l\beta_{R_2}^{q*}]$	$\emptyset$
$C_{LR2}^u$	$\emptyset$	$\frac{1}{2}G_{R_2}\beta_{R_2}^{l*}\beta_{R_2}^l$	$\emptyset$
$C_{LR2}^d$	$\emptyset$	$\emptyset$	$\emptyset$
$C_R^u$	$-\frac{1}{2}G_{S_1}\beta_{S_1}^{R*}\beta_{S_1}^R$	$\emptyset$	$\emptyset$
$C_R^d$	$\emptyset$	$\emptyset$	$\emptyset$
$C_{LQ}$	$2G_{S_1}\beta_{S_1}^{R*}\beta_{S_1}^L$	$-\frac{1}{2}G_{R_2}\beta_{R_2}^{q*}\beta_{R_2}^q$	$\emptyset$
$C_S^u$	$-2G_{S_1}\beta_{S_1}^{R*}\beta_{S_1}^L$	$\emptyset$	$\emptyset$
$C_S^d$	$\emptyset$	$\emptyset$	$\emptyset$



contributions generated by the corresponding BSM particle.

As previously said we should ignore  $W'$  and  $B'$  since we can't have non vacuum contributions to semileptonic processes without affecting with tree level amplitudes  $B_s$  meson mixing.

First thing we notice is that, apart from  $R_2$ , all LQs generate  $\mathcal{O}_S$  and  $\mathcal{O}_T$  plus, when other contributions arise is possible to turn them off independently from  $C_S$  and  $C_T$  just set equal to zero the flavour matrices that involves RH fields. In this way we would implemented a scenario in which the NP is purely LH as suggested by [3] and [5]. In this framework the purpose is to accomodate both CC and NC anomalies since  $C_S$  and  $C_T$  are forced to be generated together and they are always linked by some algebraic relation.

**A**

# **Flavour Physics**

ggggg

# Bibliography

- <sup>1</sup>C. Cornella, D. A. Faroughy, J. Fuentes-Martin, G. Isidori, and M. Neubert, “Reading the footprints of the b-meson flavor anomalies”, (2021).
- <sup>2</sup>*Particle data group* (2021).
- <sup>3</sup>D. Buttazzo, A. Greljo, G. Isidori, and D. Marzocca, “B-physics anomalies: a guide to combined explanations”, JHEP **11**, 044 (2017).
- <sup>4</sup>J. C. Pati and A. Salam, “Unified Lepton-Hadron Symmetry and a Gauge Theory of the Basic Interactions”, Phys. Rev. D **8**, 1240–1251 (1973).
- <sup>5</sup>A. Angelescu, D. Bečirević, D. A. Faroughy, and O. Sumensari, “Closing the window on single leptoquark solutions to the  $B$ -physics anomalies”, JHEP **10**, 183 (2018).