Chapter Exercises

Federico

Exercise 1-1

> *Show that $a + b$ can be less than $\min(a, b)$.*

Let $a = -1$ and $b = -1$.

Then $a + b = -2$ and $\min(a, b) = -1$.

Since $-2 < -1$, the claim that $a + b$ can be less than $\min(a, b)$ holds.

Exercise 1-2

> *Show that $a \times b$ can be less than $\min(a, b)$.*

Let $a = -1$ and $b = 2$.

Then $a \times b = -2$ and $\min(a, b) = -1$.

Since $-2 < -1$, the claim that $a \times b$ can be less than $\min(a, b)$ holds.

Exercise 1-3

> *Design/draw a road network with two points a and b such that the fastest route between a and b is not the shortest route.*

We can realize such a network by using speed 'limits'– which should instead be treated as fixed speed specifications.



1. **Route 1 (direct):**
   - By the Pythagorean thourem, its length is $d_1 = \sqrt{2}l$.
   - Let its speed limit (and minimum speed) be $s_1 = v$.
   - The time to traverse it is $t_1 = \frac{d_1}{s_1} = \frac{\sqrt{2}}{v}l$

2. **Route 2 (through c):**
   - Its lenght is the sums of the lengths of paths a to c and c to b, whicth totals $d_2 = 2l$.
   - Let its speed limit (and minimum speed) be $s_2 = 2v$.
   - The time to traverse it is $t_2 = \frac{d_2}{s_2} = 2\frac{l}{2}v = \frac{l}{v}$

Conclusion:
While $d_1 < d_2$, it was shown that $t_1 > t_2$.

■

Exercise 1-4

> *Design/draw a road network with two points a and b such that the shortest route between a and b is not the route with the fewest turns.*

We can go from a to b by taking small deviations, which add turns but amount to little additional distance. This will yield a short path with a high number of turns. It is easy to create a second path with a small number of turns that deviates more from the straight-line path, yielding the desired network.

$$\begin{tikzcd}
a \arrow[r, "\text{Route 1}"] \arrow[ddr, "\text{Route 2}"'] & c_1 \arrow[d] & c_4 \arrow[r] & b \\
& c_2 \arrow[r] & c_3 \arrow[u] & \\
& & d \arrow[uur] &
\end{tikzcd}$$

1. **Route 1:**
   - It's easy to see that the total route length is $d_1 = 2l + l + 2l + l + 2l = 8l$.
   - Number of turns $b_1 = 4$.
2. **Route 2:**
   - By the Pythagorean thourem, its length is $d_2 = 2 \cdot \sqrt{2} \cdot 3l \approx 8.485l$.
   - Number of turns $b_2 = 1$.

Conclusion:
While $d_1 < d_2$, it was shown that $b_1 > b_2$.

■

Exercise 1-5

> *The* knapsack problem *is defined as follows: given a set of integers $S = \{s_1, s_2, ..., s_n\}$ and a target number T, find a subset of S that adds up exactly to T. For example, there exists a subset within $S = \{1, 2, 5, 9, 10\}$, that adds up to $T = 22$ but not $T = 23$.*
>
> *Provide counterexamples for the following algorithms for the knapsack problem. That is, give an S and T where the algorithm does not find a solution that leves the knapsack completely full, even though a full-knapsack solution exists.*
>
> *(a) Put the elements of S in the knapsack in left to right order if they fit, that is, the first-fit algorithm.*
> *(b) Put the elements of S in the knapsack from smallest to largest, that is, the best-fit algorithm.*
> *(c) Put the elements of S in the knapsack from largest to smallest.*

Solutions

These are all greedy heuristics, which can be tricked by constructing sets in ways that they make them choose elements which are not part of any solution for the problem.

(a) Left-to-right

Let $S = \{1, 2\}$ and $T = 2$. Picking the first element, 1, will leave no room for the second element, which would be the solution for this problem.

(b) Smallest to largest

Let $S = \{1, 2\}$ and $T = 2$. Picking the smallest element, 1, will leave no room for element 2, which would be the solution for this problem.

(c) Largest to smallest

A counter example for this case requires three elements. For example, let $S = \{2, 3, 4\}$ and $T = 5$. Picking the largest element, 4, will be outright wrong, as it is not part of the solution, $\{2, 3\}$.

■