



Università di Pisa
Physics Department

PhD PreThesis

Quantum phase transition behavior in unitary and dissipation time evolution

Analysis of out-of-equilibrium regimes: round trip and dissipation protocols

Author:
Francesco Tarantelli

Supervisor:
Prof. Ettore Vicari

Session 2023/2024

Contents

Introduction	ii
1 Out-of-equilibrium dynamics	1
1.1 Introduction	1
1.2 Equilibrium Quantum Transitions	1
1.2.1 Continuous Quantum Transition (CQT)	1
1.2.2 First Order Quantum Transition ()	2
1.2.3 Models	2
2 Round Trip	3
3 Dissipation	4
Conclusions	5
References	6

Introduction

Chapter 1

Out-of-equilibrium dynamics

1.1 Introduction

In the following work, we expose the analysis of the many-body systems in the presence of time-dependent perturbations. These interferences send

$$H = H_c + w(t)H_p ; \quad (1.1)$$

$$\frac{d\rho}{dt} = -i[H, \rho] + \mathbb{D}(\rho) . \quad (1.2)$$

[1] 1.2

1.2 Equilibrium Quantum Transitions

1.2.1 Continuous Quantum Transition (CQT)

In the thermodynamic limit, i.e. infinite volume limit, same specific models could undergo a continuous phase transition. With the word phase, we intend the physical properties of the ground state associated with the Hamiltonian and characterized by the values of the Hamiltonian parameters. If this phase changes driving one of this parameter, we are in the presence of a quantum transition and the point, in which that happens, is called transition point.

In the particular case of the CQT, the quantum ground-state properties are continuous close to the transition point, called also critical point. In this special point, the system develops a long-distance correlations and its microscopic behavior becomes negligible. Independent from the local details, the global properties determine a notable universal critical behavior in which we collect different physical systems in universality class. In terms of the Renormalization Group (RG) theory, the critical point and its universal behavior are associated the fixed points of the RG flux. Hence, if we call b the characteristic unit length of the system and we define the RG transformation as the parameter rescaling respect b , the physical observables satisfy general scaling law unchanged along all the RG flux. From a physics point of view, we can interpret the factor b like the spatial correlation length ξ which diverges approaching the critical point.

According to the renormalization group (RG) theory of critical phenomena, these global properties may be the spatial dimension-ality, the nature of the

order parameter, the symmetry and the symmetry-breaking pattern. Moreover, at CQTs the systems develop an equilibrium and dynamic scaling behavior in the thermodynamic limit and their quantum functions satisfy scaling power laws characterized by universal critical exponents.

1.2.2 First Order Quantum Transition ()

1.2.3 Models

Chapter 2

Round Trip

Chapter 3

Dissipation

Conclusions

References

- [1] Francesco Tarantelli and Ettore Vicari, “Quantum critical systems with dissipative boundaries”, [Phys. Rev. B **104**, 075140 \(2021\)](#).