

# CSEN3001: DESIGN AND ANALYSIS OF ALGORITHMS

## UNIT-I: INTRODUCTION TO ALGORITHMS Strassen's Matrix Multiplication

# Divide and conquer – general idea

- **Divide a problem into subprograms of the same kind**
- **Solve subprograms using the same approach**
- **Combine partial solution (if necessary)**

# Matrix multiplication

Multiply two matrices  $A$  and  $B$ , each of size  $[n \times n]$

$$\begin{bmatrix} A \\ \phantom{A} \end{bmatrix}_{n \times n} \cdot \begin{bmatrix} B \\ \phantom{B} \end{bmatrix}_{n \times n} = \begin{bmatrix} C \\ \phantom{C} \end{bmatrix}_{n \times n}$$

# Matrix multiplication – General Method

$$\text{Mat C} = \text{Mat A} * \text{Mat B}$$

$$\begin{bmatrix} c_{11} & c_{12} & c_{13} & c_{14} \\ c_{21} & c_{22} & c_{23} & c_{24} \\ c_{31} & c_{32} & c_{33} & c_{34} \\ c_{41} & c_{42} & c_{43} & c_{44} \end{bmatrix} \quad \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{bmatrix} \quad \begin{bmatrix} b_{11} & b_{12} & b_{13} & b_{14} \\ b_{21} & b_{22} & b_{23} & b_{24} \\ b_{31} & b_{32} & b_{33} & b_{34} \\ b_{41} & b_{42} & b_{43} & b_{44} \end{bmatrix}$$

Where,

$$c_{11}=a_{11}*b_{11}+a_{12}*b_{21}+a_{13}*b_{31}+a_{14}*b_{41}$$

$$c_{12}=a_{11}*b_{12}+a_{12}*b_{22}+a_{13}*b_{32}+a_{14}*b_{42}$$

$$c_{21}=a_{21}*b_{11}+a_{22}*b_{21}+a_{23}*b_{31}+a_{24}*b_{41}$$

$$c_{22}=a_{21}*b_{12}+a_{22}*b_{22}+a_{23}*b_{32}+a_{24}*b_{42}$$

So no....

$$C_{ij} = \sum_{k=1}^n A_{ik} \times B_{kj}$$

$$\therefore T(n) = O(n^3)$$

# Matrix multiplication – Divide and Conquer Method

Transform the problem of multiplying  $A$  and  $B$ , each of size  $[n \times n]$  into subproblems, each of size  $\left[\frac{n}{2} \times \frac{n}{2}\right]$

$$\begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix} = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \cdot \begin{bmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{bmatrix}$$

$$\begin{aligned} \therefore T(n) &= 8 \cdot T\left(\frac{n}{2}\right) + an^2 \\ &= O(n^3) \end{aligned}$$

$$C_{11} = \underline{A_{11} \cdot B_{11}} + \underline{A_{12} \cdot B_{21}}$$

$$C_{12} = \underline{A_{11} \cdot B_{12}} + \underline{A_{12} \cdot B_{22}}$$

$$C_{21} = \underline{A_{21} \cdot B_{11}} + \underline{A_{22} \cdot B_{21}}$$

$$C_{22} = \underline{A_{21} \cdot B_{12}} + \underline{A_{22} \cdot B_{22}}$$

where  $an^2$  is for addition.  
So, it is no improvement compared with the traditional way

# Matrix multiplication – Divide and Conquer Method

$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix} \cdot \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \cdot \begin{bmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 3 & 3 & 3 & 0 \\ 3 & 3 & 3 & 0 \\ 3 & 3 & 3 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$C_{11} = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \cdot \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix} \cdot \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 3 & 3 \\ 3 & 3 \end{bmatrix}$$

$$C_{12} = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \cdot \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix} \cdot \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 3 & 0 \\ 3 & 0 \end{bmatrix}$$

$$C_{21} = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} \cdot \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \cdot \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 3 & 3 \\ 0 & 0 \end{bmatrix}$$

$$C_{22} = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} \cdot \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \cdot \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 3 & 0 \\ 0 & 0 \end{bmatrix}$$

# Matrix multiplication – Strassen's Matrix Multiplication

- Discover a way to compute the  $C_{ij}$ 's using 7 multiplications and 18 additions or subtractions

$$P = (A_{11} + A_{22})(B_{11} + B_{22})$$

$$Q = (A_{21} + A_{22})B_{11}$$

$$R = A_{11}(B_{12} - B_{22})$$

$$S = A_{22}(B_{21} - B_{11})$$

$$T = (A_{11} + A_{12})B_{22}$$

$$U = (A_{21} - A_{11})(B_{11} + B_{12})$$


$$V = (A_{12} - A_{22})(B_{21} + B_{22})$$

$$C_{11} = P + S - T + V$$

$$C_{12} = R + T$$

$$C_{21} = Q + S$$

$$C_{22} = P + R - Q + U$$



# Algorithm: Matrix Multiplication Strassen's Method

Algorithm Strassen ( $n, A, B, C$ )

//  $n$  is size,  $A, B$  the input matrices,  $C$  output matrix

begin

    if  $n == 2$

    {

$$C_{11} = a_{11} \cdot b_{11} + a_{12} \cdot b_{21}$$

$$C_{12} = a_{11} \cdot b_{12} + a_{12} \cdot b_{22}$$

$$C_{21} = a_{21} \cdot b_{11} + a_{22} \cdot b_{21}$$

$$C_{22} = a_{21} \cdot b_{12} + a_{22} \cdot b_{22}$$

    }

    else

    {

        Partition Matrix A into 4 submatrices  
( $A_{11}, A_{12}, A_{12}, A_{22}$ )

        Partition Matrix B into 4 submatrices  
( $B_{11}, B_{12}, B_{21}, B_{22}$ )

        call Strassen( $n/2, A_{11}+A_{22}, B_{11}+B_{22}, P$ )

        call Strassen( $n/2, A_{21}+A_{22}, B_{11}, Q$ )

        call Strassen( $n/2, A_{11}, B_{12}-B_{22}, R$ )

        call Strassen( $n/2, A_{22}, B_{21}-B_{11}, S$ )

        call Strassen( $n/2, A_{11}+A_{12}, B_{22}, T$ )

        call Strassen( $n/2, A_{21}+A_{22}, B_{11}, Q$ )

        call Strassen( $n/2, A_{21}-A_{11}, B_{11}+B_{12}, U$ )

        call Strassen( $n/2, A_{11}-A_{22}, B_{21}+B_{22}, V$ )

    }

end;



# Algorithm: Matrix Multiplication Strassen's Method

$$C_{11} = P + S - T + V;$$

$$C_{12} = R + T;$$

$$C_{21} = Q + S;$$

$$C_{22} = P + R - Q + U;$$

# Matrix Multiplication Strassen's Method: Time Complexity

$$T(n) = \begin{cases} 7 \cdot T(\frac{n}{2}) + an^2 & n > 2 \\ b & n \leq 2 \end{cases}$$

$$\begin{aligned} T(n) &= an^2[1 + 7/4 + (7/4)^2 + \cdots + (7/4)^{k-1}] + 7^k T(1) \\ &\leq cn^2(7/4)^{\log_2 n} + 7^{\log_2 n}, \text{ } c \text{ a constant} \\ &= cn^{\log_2 4 + \log_2 7 - \log_2 4} + n^{\log_2 7} \\ &= O(n^{\log_2 7}) \approx O(n^{2.81}) \end{aligned}$$

# Matrix Multiplication Strassen's Method: Time Complexity

$$T(n) = \begin{cases} 7 \cdot T\left(\frac{n}{2}\right) + an^2 & n > 2 \\ b & n \leq 2 \end{cases}$$


$$\begin{aligned} T(n) &= 7 \cdot T\left(\frac{n}{2}\right) + an^2 \\ &= 7^2 \cdot T\left(\frac{n}{2^2}\right) + \left(\frac{7}{4}\right)an^2 + an^2 \\ &= 7^3 \cdot T\left(\frac{n}{2^3}\right) + \left(\frac{7}{4}\right)^2 \cdot an^2 + \left(\frac{7}{4}\right) \cdot an^2 + an^2 \\ &\quad \dots \end{aligned}$$

# Matrix Multiplication Strassen's Method: Time Complexity

Assume  $n = 2^k$  for some integer  $k$

$$\begin{aligned} &= 7^{k-1} \cdot T\left(\frac{n}{2^{k-1}}\right) + an^2 \cdot \left[ \left(\frac{7}{4}\right)^{k-2} + \dots + 1 \right] \\ &= 7^{k-1} \cdot b + an^2 \left[ \frac{\left(\frac{7}{4}\right)^{k-1} - 1}{\frac{7}{4} - 1} \right] \\ &\leq b \cdot 7^k + c \cdot n^2 \cdot \left(\frac{7}{4}\right)^k \\ &= b \cdot 7^{\lg n} + cn^2 \cdot \left(\frac{7}{4}\right)^{\lg n} = b \cdot 7^{\lg n} + cn^2 (n)^{\lg \frac{7}{4}} \\ &= b \cdot n^{\lg 7} + cn^{\lg 7} = (b + c) \cdot n^{\lg 7} \\ &= O(n^{\lg 7}) = O(n^{2.81}) \end{aligned}$$

# Comparison : Straight method vs. Strassen's Method



# Improvements and other methods

1978	<b>V. Y. Pan.</b> Strassen's algorithm is not optimal.	$O(n^{2.796})$
1979	<b>D. Bini et al.</b> $O(n^{2.7799})$ complexity for $n \times n$ approximate matrix multiplication.	$O(n^{2.7799})$
1981	<b>A. Schönhage.</b> Partial and total matrix multiplication.	$O(n^{2.522})$

## First to break the 2.5 barrier:

1981	<b>Copper Smith and Winograd</b> On the asymptotic complexity of matrix multiplication.	$O(n^{2.496})$
1986	<b>Volker Strassen</b>	$O(n^{2.479})$
1989	<b>CopperSmith and Winograd</b> Matrix multiplication via arithmetic progressions.	$O(n^{2.376})$
2011	<b>Virginia Vassilevska Williams:</b> Breaking the Coppersmith-Winograd barrier	$O(n^{2.3727})$

THANK YOU