

Contents

1	Simplicial complexes and polyhedra	2
1.1	Simplicial complexes geometrically and combinatorially	2
1.2	The face poset and nerves.	4
1.3	Subdivisions	4
2	Definability of compact polyhedra	6
2.1	Disjoint union	6
2.2	generated subframes	7
2.3	bounded morphic images	7
2.4	Compact Goldblatt-Thomason for polyhedra	9
3	Homology	10
3.1	simplicial homology	10
3.2	Modally undefinable homology	11
4	Future Work	12

1 Simplicial complexes and polyhedra

Our aim is to create a good environment to prove an analogon of the Goldblatt-Thomason theorem for polyhedral logic.

Let's begin by setting the objects we are going to find interplay between:

1.1 Simplicial complexes geometrically and combinatorially

Intuitively, any polygon can be obtained by gluing together triangles on their edges or vertices. Similarly we can describe a polyhedron as a gluing of simplices (higher dimensional triangles):

Definition 1.1 (Geometric Simplex).

The **standard n -simplex** in $\mathbb{R}^{n+1}$ is the set

$$\Delta^n = \{x \in \mathbb{R}^{n+1} : x_0 + \dots + x_n = 1, x_i \geq 0 \text{ for } i = 0 \dots n\}$$

endowed with the topology induced by $\mathbb{R}^{n+1}$.

Note that every standard n -simplex comes equipped with $n+1$ 'canonical' subsimplices, those given by the intersection with the canonical hyperplanes:

$$\Delta^n \geq \Delta_i^{n-1} = \{x \in \mathbb{R}^{n+1} : x_0 + \dots + \hat{x}_i + \dots + x_n = 1, x_i \geq 0 \text{ for } i = 0 \dots n\}^1$$

We call such subsimplices and all their such subsimplices **Faces** of Δ^n

We call an **n -simplex** in $\mathbb{R}^m$ an affine immersion $\Delta^n \hookrightarrow \mathbb{R}^m$.

With this definition, we have what we expect:

a 0-simplex is a point, a 1-simplex is a segment, a 2-simplex is a triangle, etc.

We can obtain a polyhedra by gluing together simplices:

Definition 1.2 (Polyhedron).

A **(Geometric) Simplicial Complex** is a collection K of simplices in $\mathbb{R}^m$ such that

1. Every face of a simplex in K is in K ,
2. The intersection of two simplices of K is a face of both simplices.

We call a **Polyhedron** in $\mathbb{R}^m$ the union of all elements of a geometric simplicial complex.

If K is a simplicial complex we call $|K|$ its **realization**, i.e. its associated polyhedron, and given a polyhedron P , any simplicial complex K such that $|K| = P$ is called a **triangulation** of P .

Lemma 1.3. If K is finite, then $|K|$ is compact

Proof. It's a closed and bounded subset of $\mathbb{R}^n$. □

We make polyhedra into a category via **PL**-homeomorphisms: We consider maps that are affine when restricted to any face. More details can be found at [Col72] or (closer to the aim of the paper [Bez+21][tGS09])

We may notice that a simplex is uniquely determined by its vertices, and a simplicial complex is uniquely determined by its simplices;

This means that we can give an abstract definition

¹here I use what I hope is standard notation, the hat means 'that component is removed'.

Definition 1.4 (Simplicial complex).

A **(Combinatorial) Simplicial complex** is a collection of finite sets K closed under taking subsets, or equivalently, is a set of vertices V together with a collection $K \subseteq \mathcal{P}(V)$ closed under taking subsets.

From now on we refer to Combinatorial simplicial complexes as just Simplicial complexes.

We make Simplicial complexes into a category by saying that a map of simplicial complexes is a function $V \rightarrow V'$ such that the image of a simplex is a simplex.

Lemma 1.5. *Every geometric simplicial complex uniquely determines a combinatorial simplicial complex*

Proof. Let K be a geometric simplicial complex. Construct K^v by sending each simplex in K to the set of vertices of K .

This is a combinatorial simplicial complex, since if $(v_1, \dots, v_n) \in K^v$ and $(w_1, \dots, w_k) \subseteq (v_1, \dots, v_n)$, then $(w_1, \dots, w_k)$ are the vertices of a face of K , thus an element of K^v . $\square$

Lemma 1.6. *Every finite combinatorial simplicial complex determines a geometric simplicial complex up to PL-isomorphism.*

Proof. We can realize every simplex to the standard simplex of the correct dimension, then glue them according to the data of the simplicial complex:

$$\tilde{K} = \frac{\bigsqcup_{\sigma \in K} \Delta^{|\sigma|}}{\sim}$$

Where $\Delta_i \sim \Delta_j$ if they correspond to the same face in K .

The realization of this complex only depends on the immersion of its faces, thus any two will be isomorphic, and the total map will be affine when restricted to the faces, thus *PL*. $\square$

This means that we have a correspondence

$$\{\text{Simplicial complexes}\} \rightarrow \{\text{Polyhedra}\}$$

We can extend it to a functor by noticing that

Lemma 1.7. *Every simplicial map $K \rightarrow K'$ induces a PL map $|K| \rightarrow |K'|$*

sketch of proof. Notice that every morphism of simplices can be realized as an affine map between their realizations. $\square$

Thus we have what is commonly known as the realization:

Definition 1.8 (Realization functor). Call **Scplx** the category of simplicial complexes and **Pol** the category of polyhedra described above, We call the functor realizing a simplicial complex into a polyhedron

$$|-| : \text{Scplx} \rightarrow \text{Pol}$$

The **realization functor**

1.2 The face poset and nerves.

Notice that any simplicial complex comes equipped with a 'face' order:

Definition 1.9 (Face poset).

Let Σ be a simplicial complex. Then $(\Sigma, \subseteq)$ is a poset, that we call the **Face poset of Σ** and we denote $[\Sigma]$

Note that every morphism of simplicial complexes induces a morphism of the posets (by taking the image of the function), it's easy to check that we have a -sort of forgetful- functor

$$[-] : \mathbf{Scplx} \rightarrow \mathbf{Pos}$$

As a 'forgetful' we expect a way to 'freely' express the information of the poset through a simplicial complex; indeed

Definition 1.10 (Nerve of a poset).

Let $(P, \leq)$ be a poset. The **Nerve $\mathcal{N}(P)$** is a simplicial complex constructed in the following way:

- The **vertices** are the points of P .
- The **sides** are the 1-chains $p_0 \leq p_1$ in P .
- The **Triangles** are the 2-chains $p_0 \leq p_1 \leq p_2$ in P .
- $\vdots$
- The **n -simplices** are the n -chains $p_0 \leq p_1 \leq \dots \leq p_n$ in P .
- $\vdots$

This is clearly a simplicial complex as for any $m < n$, any sub- m -chain on an n -chain is part of the complex.

This is clearly not an inverse: $\mathcal{N}[\Sigma]$ generally has more points than Σ .

Let's talk of a way we can recover information from the nerve process.

1.3 Subdivisions

Definition 1.11 (Stellar moves).

Given two disjoint simplicial complexes Σ and Σ' we define:

Their **join** $\Sigma \vee \Sigma'$ to be

$$\{\sigma \cup \sigma' : \sigma \in \Sigma, \sigma' \in \Sigma'\}$$

The **Star** of a face σ to be

$$\star_{\Sigma}(\sigma) := \{\tau \in \Sigma : \tau \cup \sigma \in \Sigma\}$$

The **link** of a face σ to be

$$\ell_{\Sigma}(\sigma) := \{\tau \in \star_{\Sigma}(\sigma) : \tau \cap \sigma = \emptyset\}$$

Lastly, the **deletion** of a face σ as

$$\Sigma \setminus \sigma := \{\tau \in \Sigma : \sigma \not\subseteq \tau\}$$

and the **boundary** of a face σ as

$$\partial\sigma = \{\tau \in \Sigma : \tau \not\subseteq \sigma\}$$

A **stellar subdivision** of a complex Σ at σ is a simplicial complex

$$(\Sigma \setminus \sigma) \cup (\{x\} \vee \partial\sigma \vee \ell_\Sigma(\sigma))$$

where x is a new vertex we are introducing.

We call a **stellar move** of a complex either a stellar subdivision or its inverse, and we say that two complexes are **stellarly equivalent** $\Sigma \sim_* \Sigma'$ if there exist a complex Σ'' that can be obtained by Σ or Σ' via a sequence of stellar moves.

Theorem 1.12 (Alexander).

Two simplicial complexes are stellarly equivalent if and only if their geometric realizations are **PL**-homeomorphic.

Proof. [Col72] □

Also, a theorem that is of interest for us:

Theorem 1.13.

Stellarly equivalent simplicial complexes have bisimilar face posets.

Proof. A complete account can be found in [Dek23]: The idea is that given a triangulation of a face $\text{Subd}(\Sigma, \sigma)$, we can find a p-morphism $\text{Subd}(\Sigma, \sigma) \rightarrow \Sigma$ sending every new subdivided piece to σ .

This means that we have a span of p-morphisms $\Sigma \leftarrow \Sigma'' \rightarrow \Sigma'$, thus $\Sigma \rightleftharpoons \Sigma'^2$ □

²placeholder because I cannot find the bisimulation symbol.

2 Definability of compact polyhedra


[BRV01] in chapter 3 gives us a definability result for classes of finite transitive frames

Theorem 2.1. *A class of finite transitive frame is modally definable if and only if it's closed under finite disjoint union, bounded morphic images and generated subframes.*

Suppose we have a class of simplicial complexes, it's modally definable if and only if the corresponding class of posets is.

We want to translate these conditions in terms of simplicial complexes and then realize them in terms of polyhedra.

The picture we want to have in mind when building classes is the following:



2.1 Disjoint union

Let us start with disjoint unions: these are the easiest construction since they are the same in all three settings

Proposition 2.2. *For a pair of simplicial complexes Σ, Σ' , $[\Sigma \sqcup \Sigma'] = [\Sigma] \sqcup [\Sigma']$*

Proof. Let σ be a simplex of $[\Sigma \sqcup \Sigma']$, then -as the union is disjoint- it must be a simplex of either $[\Sigma]$ or $[\Sigma']$, and vice versa. $\square$

Similarly

Proposition 2.3. *For a pair of simplicial complexes Σ, Σ' , $|\Sigma \sqcup \Sigma'| = |\Sigma| \sqcup |\Sigma'|$*

Proof. There are direct ways to show it but we appeal to abstract nonsense and say that the geometric realization is the coend

$$\int^{n \in \mathbb{N}} \Delta^n \cdot \Sigma_n$$

thus the constructions commutes with colimits.

This is not as clear, but theoretically better than a direct proof in $\mathbb{R}^n$ since it eventually will allow us to work with arbitrarily-sized polyhedra too. $\square$

This means that

Lemma 2.4. *A class of simplicial complex C is closed under disjoint union if and only if its corresponding class of posets $[C]$ is closed under disjoint union if and only if its corresponding class of polyhedra $|C|$ is closed under disjoint union.*

2.2 generated subframes

Now suppose we have a simplicial complex Σ : What corresponds to generated subframes of $[\Sigma]$? Recall that

Definition 2.5 (Generated subframe). A **Generated subframe** of a kripke frame (W, R) , is a subset $W' \subseteq W$ closed upwards under R .

Recall that the posets we are considering are the 'coface' posets, i.e. the posets whose elements are the faces of Σ and whose relation is the opposite of the inclusion.

This means that a generated subframe is $\uparrow$ -closed under $\supseteq$, thus $\downarrow$ -closed under $\subseteq$.
in other words

Lemma 2.6.

Let C be a class of simplicial complexes, then $[C]$ is closed under generated subframes if and only if C is closed under subcomplexes.

Proof. Let Σ be a simplicial complex and Σ' a subcomplex. As Σ' is a complex as well, then it's downward-closed.

Conversely, if $S' \leq [\Sigma]$ is a downward-closed subposet, then by definition it's closed under taking subsets, thus it's a simplicial complex. $\square$

Lemma 2.7. Let C be a class of simplicial complexes, then $|C|$ is closed under subpolyhedra if and only if C is closed under subcomplexes and triangulations.

Proof. if $\Sigma' \leq \Sigma$ is a subcomplex then $|\Sigma'| \subseteq |\Sigma|$ is a closed subpolyhedron.

Conversely, if $S \subseteq |\Sigma|$ is a closed subpolyhedron, then there exists a triangulation of $|\Sigma|$ that triangulates S , meaning that there exist two simplicial complexes Σ', Σ'' such that $|\Sigma'| = S$, $|\Sigma''| = |\Sigma|$ $\square$

2.3 bounded morphic images

Lastly we need to interpret bounded morphism in our triad. Recall

Definition 2.8.

Let (W, R) and (W', R') be kripke frames and $f : W \rightarrow W'$ a function.
 f is a **bounded morphism** if and only if

- wRv implies $f(w)R'f(v)$
- $f(w)R'v'$ implies that there exists v such that wRv and $f(v) = v'$.

Interpreting this as the coface relation this means

- $\sigma \subseteq \tau$ implies $f(\sigma) \subseteq f(\tau)$
- $\sigma' \subseteq f(\tau)$ implies that there exists $\sigma \subseteq \tau$ such that $f(\sigma) = \sigma'$

Let's define a class of simplicial maps:

Definition 2.9 (Bisimplicial map).

Let $(V, \Sigma), (V', \Sigma')$ be simplicial complexes. $f : V \rightarrow V'$ is called a **bisimplicial map** if

$$\forall s \in \mathcal{P}(V), f(s) \in \Sigma' \iff s \in \Sigma$$

Lemma 2.10.

A map of simplicial complexes $f : \Sigma \rightarrow \Sigma'$ induces a bounded morphism $[\Sigma] \rightarrow [\Sigma']$ if and only if it's bisimplicial.

Proof. Suppose f is a bisimplicial map, then suppose $\sigma \subseteq \tau$, then $f(\sigma)$ and $f(\tau)$ are simplices and $f(\sigma) \subseteq f(\tau)$ as it's an image map.

Now suppose $\sigma' = f(\tau)$. suppose $\tau = v_1, \dots, v_n$, thus $f(\tau) = f(v_1), \dots, f(v_n)$. up to reordering, $\sigma' = f(v_1), \dots, f(v_k)$. As f is bisimplicial, $f(v_1), \dots, f(v_k) = f(v_1, \dots, v_k)$ is a simplex, thus $\sigma := v_1, \dots, v_k$ is a simplex and - by structure- a subsimplex of $v_1, \dots, v_n$.

Now conversely, suppose that f is a bounded morphism $[\Sigma] \rightarrow [\Sigma']$, the (co)induced $f : \Sigma \rightarrow \Sigma'$

First we need to check that image of a vertex is a vertex. This is because vertices are simplices, and minimal ones at that: suppose we have a vertex v such that $f(v) = \sigma'$. then if σ' is not a vertex, there exists $w' \subsetneq \sigma' = f(v)$.

This means that there must exist $w \subseteq v$ such that $f(w) = w'$, but such w does not exist as v is minimal. $\rightarrow \leftarrow$.

Image of a simplex is a simplex (by definition), and suppose

$f(v_1, \dots, v_n)$ is a simplex of Σ' but $v_1, \dots, v_n$ is not a simplex of Σ . but $f(v_1, \dots, v_n) = f(v_1, \dots, v_n)$ (thus in particular $\subseteq$) But then there exist a simplex $v_1, \dots, v_k \subseteq v_1, \dots, v_n$ such that $f(v_1, \dots, v_k) = f(v_1, \dots, v_n)$. But then for a question of cardinality it must be the case that $v_1, \dots, v_k = v_1, \dots, v_n$, thus $v_1, \dots, v_n$ is a simplex of Σ . $\square$

Now to interpret bisimplicial maps as maps of polyhedra:

Suppose we have the map $|f| : |\Sigma| \rightarrow |\Sigma'|$ that is the realization of a map $f : \Sigma \rightarrow \Sigma'$.

If f is bisimplicial, then this results in the conditions

- if $\text{Hull}(v_1, \dots, v_k) \subseteq |\Sigma|$, then $f(\text{Hull}(v_1, \dots, v_k)) = \text{Hull}(f(v_1), \dots, f(v_k)) \subseteq |\Sigma'|$, and
- if $\text{Hull}(f(v_1), \dots, f(v_k)) \subseteq |\Sigma'|$, then $\text{Hull}(v_1, \dots, v_k) \subseteq |\Sigma|$

This is equivalent to it being a retraction onto the image:

Theorem 2.11. Suppose f if a map $\Sigma \rightarrow \Sigma'$, then it's a bisimplicial map if and only if $f : \Sigma \rightarrow \text{im}(f)$ is a retraction.

Proof.

Suppose that f is bisimplicial.

First, note that f is surjective onto the vertices of $\text{im}(f)$. Moreover if $\sigma' = v'_1, \dots, v'_k$ is a simplex in $\text{im}(f)$, then there exist $v_1, \dots, v_k$ such that $f(v_i) = v'_i$ and $v_1, \dots, v_k$ is a simplex in Σ .

This means that $f : \Sigma \rightarrow \text{im } f$ is epimorphic at the level of simplices as well. Now let $g : \text{im}(f) \rightarrow \Sigma$ be defined as $g(v'_i) = v_i$ for some v_i such that $f(v_i) = v'_i$

This is a well-defined map, sending vertices to vertices, and $f(g(v'_1, \dots, v'_n)) = f(v_1, \dots, v_n) = v_1', \dots, v_n'$ thus $fg = 1_{\text{im}(f)}$

Conversely, suppose that f is a section, i.e. there exists a $g : \text{im}(f) \rightarrow \Sigma$ such that $fg = 1_{\text{im}(f)}$ Then suppose $f(v_1), \dots, f(v_n) = v'_1, \dots, v'_n$ is a simplex. then $g(v'_1), \dots, g(v'_n) = v_1, \dots, v_n$ is a simplex since g is simplicial. $\square$

Corollary 2.12. Let C be a class of simplicial complexes, then $|C|$ is closed under retracts if and only if C is closed under bisimplicial images and triangulations.

Proof. **PL** –retracts of $|\Sigma|$ are retracts of a triangulation of Σ which are bisimplicial images by the lemma above. $\square$

2.4 Compact Goldblatt-Thomason for polyhedra

Recall that the frames we defined are models for a modal logic (described extensively in [Bez+21]):

Essentially propositions are interpreted in unions of open subpolyhedra, $[\Diamond\phi]$ is the least closed subpolyhedron containing $[\phi]$.

We connect those as simplicial models by associating every simplex of a complex Σ to its relative interior in $|\Sigma|$, thus obtaining a neighborhood-description:

Corollary 2.13. *If Σ is a simplicial complex, for each point $x \in |\Sigma|$ there exists a unique face $\sigma \in \Sigma$ such that $x \in \sigma$, and such Σ can be interpreted as a poset through the face (or coface) relation.*

Here we can finally express the main result of this paper:

Theorem 2.14 (Compact GTT for Polyhedra).

A class of compact polyhedra is modally definable if and only if it's closed under disjoint unions, closed subpolyhedra and retractions.

Proof. There's a direct correspondence between classes of polyhedra and class of simplicial complexes closed under triangulations: Indeed we showed that two simplicial complexes are **PL**-homeomorphic if and only if they have a common triangulation, thus the triangulation-equivalence forms the kernel of the realization functor.

As the constructions we built are stable under triangulation (indeed, up to triangulation equivalence, $\mathbb{N}$ and $[-]$ are inverses!), we have an equivalence between modally definable classes of polyhedra and of simplicial complexes modulo triangulation. $\square$

3 Homology

Homology is a hot topic when talking about polyhedra. Here is a synthetic account

3.1 simplicial homology

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

Nulla malesuada porttitor diam. Donec felis erat, congue non, volutpat at, tincidunt tristique, libero. Vivamus viverra fermentum felis. Donec nonummy pellentesque ante. Phasellus adipiscing semper elit. Proin fermentum massa ac quam. Sed diam turpis, molestie vitae, placerat a, molestie nec, leo. Maecenas lacinia. Nam ipsum ligula, eleifend at, accumsan nec, suscipit a, ipsum. Morbi blandit ligula feugiat magna. Nunc eleifend consequat lorem. Sed lacinia nulla vitae enim. Pellentesque tincidunt purus vel magna. Integer non enim. Praesent euismod nunc eu purus. Donec bibendum quam in tellus. Nullam cursus pulvinar lectus. Donec et mi. Nam vulputate metus eu enim. Vestibulum pellentesque felis eu massa.

Quisque ullamcorper placerat ipsum. Cras nibh. Morbi vel justo vitae lacus tincidunt ultrices. Lorem ipsum dolor sit amet, consectetur adipiscing elit. In hac habitasse platea dictumst. Integer tempus convallis augue. Etiam facilisis. Nunc elementum fermentum wisi. Aenean placerat. Ut imperdiet, enim sed gravida sollicitudin, felis odio placerat quam, ac pulvinar elit purus eget enim. Nunc vitae tortor. Proin tempus nibh sit amet nisl. Vivamus quis tortor vitae risus porta vehicula.

Fusce mauris. Vestibulum luctus nibh at lectus. Sed bibendum, nulla a faucibus semper, leo velit ultricies tellus, ac venenatis arcu wisi vel nisl. Vestibulum diam. Aliquam pellentesque, augue quis sagittis posuere, turpis lacus congue quam, in hendrerit risus eros eget felis. Maecenas eget erat in sapien mattis porttitor. Vestibulum porttitor. Nulla facilisi. Sed a turpis eu lacus commodo facilisis. Morbi fringilla, wisi in dignissim interdum, justo lectus sagittis dui, et vehicula libero dui cursus dui. Mauris tempor ligula sed lacus. Duis cursus enim ut augue. Cras ac magna. Cras nulla. Nulla egestas. Curabitur a leo. Quisque egestas wisi eget nunc. Nam feugiat lacus vel est. Curabitur consectetur.

Suspendisse vel felis. Ut lorem lorem, interdum eu, tincidunt sit amet, laoreet vitae, arcu. Aenean faucibus pede eu ante. Praesent enim elit, rutrum at, molestie non, nonummy vel, nisl. Ut lectus eros, malesuada sit amet, fermentum eu, sodales cursus, magna. Donec eu purus. Quisque vehicula, urna sed ultricies auctor, pede lorem egestas dui, et convallis elit erat sed nulla. Donec luctus. Curabitur et nunc. Aliquam dolor odio, commodo pretium, ultricies non, pharetra in, velit. Integer arcu est, nonummy in, fermentum faucibus, egestas vel, odio.

Sed commodo posuere pede. Mauris ut est. Ut quis purus. Sed ac odio. Sed vehicula hendrerit sem. Duis non odio. Morbi ut dui. Sed accumsan risus eget odio. In hac habitasse platea dictumst.

Pellentesque non elit. Fusce sed justo eu urna porta tincidunt. Mauris felis odio, sollicitudin sed, volutpat a, ornare ac, erat. Morbi quis dolor. Donec pellentesque, erat ac sagittis semper, nunc dui lobortis purus, quis congue purus metus ultricies tellus. Proin et quam. Class aptent taciti sociosqu ad litora torquent per conubia nostra, per inceptos hymenaeos. Praesent sapien turpis, fermentum vel, eleifend faucibus, vehicula eu, lacus.

3.2 Modally undefinable homology

While there's little hope to give much information about homology through a modal language we shall note that our constructions map neatly onto the homology chain. In particular

Proposition 3.1. $H_{\bullet}(\Sigma \sqcup \Sigma') = H_{\bullet}(\Sigma) \oplus H_{\bullet}(\Sigma')$

Proof. There's a proof of this somewhere for sure, can't be bothered now □

Proposition 3.2. *If $\Sigma \leq \Sigma'$ is a closed subcomplex, then there's a map $H_{\bullet}(\Sigma) \rightarrow H_{\bullet}(\Sigma')$*

Proof. Same here □

Proposition 3.3. *If $f : \Sigma \rightarrow \Sigma'$ is a simplicial map, then there's a bisimplicial map $f_{\bullet} : H_{\bullet}(\Sigma) \rightarrow H_{\bullet}(\Sigma')$*

Proof. and here □

This means

Lemma 3.4. *The homologies of a modally definable class of polyhedra must be closed under direct sum and images.*

Or, in the more useful direction:

*If the homology groups of a class of polyhedra is **not** closed under direct sums and images, then the class is not modally definable.*

Proof. Follows from the three translations. □

Corollary 3.5. *The class of genus n polyhedra is not modally for any positive n*

Proof. $H_1(\Sigma) \simeq \mathbb{Z}^n$ for all $\Sigma \in C$. As $\mathbb{Z}^n \oplus \mathbb{Z}^n \neq \mathbb{Z}^n$, the class is not closed under direct sums, thus not modally definable. □

4 Future Work

An interesting question is what happens when we allow for arbitrarily sized simplicial complexes, both in dimension and number of simplices. Then the class is not of transitive finite frames but arbitrary ones.

- What is the geometric realization of an arbitrarily big simplicial complex?
- What is the ultrafilter extension of “
- How do we describe such ultrafilter extension geometrically?
- Simplicial complexes allow for a coalgebraic description: They are a class of $\mathcal{P} \circ \mathcal{P}$ coalgebras over (finite) sets (the map $V \rightarrow \mathcal{P} \mathcal{P} V$ is the one associating to every point the up-set of simplices containing it). Do completeness and definability results coincide to the usual ones in that setting?

References

- [Col72] Brian J. Sanderson Colin P. Rourke. *Introduction to Piecewise-Linear Topology*. 1972. ISBN: 978-3-642-81735-9. DOI: <https://doi.org/10.1007/978-3-642-81735-9>.
- [CK90] C. C. Chang and H. J. Keisler. *Model Theory*. en. Google-Books-ID: uiHq0EmaFp0C. Elsevier, June 1990. ISBN: 978-0-08-088007-5.
- [Lic99] W B R Lickorish. “Simplicial moves on complexes and manifolds”. en. In: *Proceedings of the Kirbyfest*. MSRI, Berkeley, California, USA: Mathematical Sciences Publishers, Nov. 1999, pp. 299–320. DOI: 10.2140/gtm.1999.2.299. URL: <http://www.msp.org/gtm/1999/02/p016.xhtml>.
- [BRV01] Patrick Blackburn, Maarten de Rijke, and Yde Venema. “Frames”. In: *Modal Logic*. Cambridge Tracts in Theoretical Computer Science. Cambridge University Press, 2001, pp. 123–187.
- [tGS09] Balder ten Cate, David Gabelaia, and Dmitry Sustretov. “Modal languages for topology: Expressivity and definability”. In: *Annals of Pure and Applied Logic* 159.1 (2009), pp. 146–170. ISSN: 0168-0072. DOI: <https://doi.org/10.1016/j.apal.2008.11.001>. URL: <https://www.sciencedirect.com/science/article/pii/S016800720800184X>.
- [IS10] Ivan Izestiev and Jean-Marc Schlenker. “On the infinitesimal rigidity of polyhedra with vertices in convex position”. In: *Pacific Journal of Mathematics* 248.1 (Oct. 2010). arXiv:0711.1981 [math], pp. 171–190. ISSN: 0030-8730, 0030-8730. DOI: 10.2140/pjm.2010.248.171.
- [IKN17] Ivan Izestiev, Steven Klee, and Isabella Novik. “Simplicial moves on balanced complexes”. en. In: *Advances in Mathematics* 320 (Nov. 2017), pp. 82–114. ISSN: 00018708. DOI: 10.1016/j.aim.2017.08.036.
- [Fri20] Greg Friedman. *Singular Intersection Homology*. en. 1st ed. Cambridge University Press, Sept. 2020. ISBN: 978-1-316-58444-6. DOI: 10.1017/9781316584446. URL: <https://www.cambridge.org/core/product/identifier/9781316584446/type/book>.
- [Bez+21] Nick Bezhanishvili et al. “Geometric Model Checking of Continuous Space”. In: *CoRR* abs/2105.06194 (2021). arXiv: 2105.06194. URL: <https://arxiv.org/abs/2105.06194>.
- [Dek23] P.M. Dekker. “polyhedral semantics of modal logic”. PhD thesis. UVA Amsterdam, 2023.
- [FH24] Stefan Friedl and Julian Hannes. “Pachner’s Theorem”. en. In: *2021-2022 MATRIX Annals*. Ed. by David R. Wood, Jan De Gier, and Cheryl E. Praeger. Vol. 5. MATRIX Book Series. Cham: Springer Nature Switzerland, 2024, pp. 777–832. ISBN: 978-3-031-47416-3. DOI: 10.1007/978-3-031-47417-0_34. URL: https://link.springer.com/10.1007/978-3-031-47417-0_34.
- [Sol25] Sławomir Solecki. “Simplicial complexes, stellar moves, and projective amalgamation”. In: arXiv:2503.14825 (Mar. 2025). arXiv:2503.14825 [math]. DOI: 10.48550/arXiv.2503.14825. URL: <http://arxiv.org/abs/2503.14825>.
- [AP] Karim A Adiprasito and Igor Pak. “ALL TRIANGULATIONS HAVE A COMMON STELLAR SUBDIVISION”. en. In: ().