



**RAJARATA UNIVERSITY OF SRI LANKA  
FACULTY OF APPLIED SCIENCES**

B.Sc. (General) Degree Examination  
Second Year – Semester II Examination – March / April 2014

**PHY 2204 - PHYSICAL OPTICS**

Answer **FOUR** questions only.

Time allowed: 2 Hours

1.

a. Derive equation,  $n_2 \frac{1}{V} - n_1 \frac{1}{U} = (n_2 - n_1) \frac{1}{R}$  for the refraction at a spherical surface. All the symbols have their usual meanings.

b. Show that the first and the second focal lengths of the spherical surface are given by,

$$f_1 = -\frac{Rn_1}{(n_2 - n_1)} \text{ and } f_2 = \frac{Rn_2}{(n_2 - n_1)} \text{ respectively.}$$

c. The incident face of a glass block is bounded by a concave surface of radius 3 cm. A thin object of 2 cm height is situated in air and on the axis 10 cm from the curved face. If the refractive index of glass is 1.5, find,

- i. the first and second focal lengths.
- ii. the image distance.
- iii. the lateral magnification and
- iv. image height.

2.

- a. Two thin lenses of focal lengths  $f_1$  and  $f_2$  are separated in air by a distance  $d$ . Prove that the equivalent focal length  $F$  of the two lens system is given by  $F = \frac{f_1 f_2}{f_1 + f_2 - d}$ .
- b. Derive Newton's relationship for the two lens combination.
- c. Two thin convex lenses of focal lengths 15.0 cm and 20.0 cm are placed coaxially 10.0 cm apart.
  - i. Find the equivalent focal length of the system.
  - ii. Locate the principal points and focal points of the system.
  - iii. Locate the image of an object placed 60 cm away from the first lens.

3.

- a. Derive equations for the distance to bright and dark fringes from central fringe of a double slit interference experiment.
  - b. In Young's double-slit experiment, an electron beam is used. What will happen to the interference pattern, if the velocity of the electrons is increased?
  - c. In another double-slit experiment sources with slit separation 0.2 mm is located 1.5 m from a screen. The distance between successive bright fringes on the screen is measured to be 3.30 mm. What is the wavelength of the light?
  - d. What is the color of the light?
4. A beam of, initially unpolarized, light is incident normally on a piece of polarizing sheet (polaroid)  $A$ , the transmitted light then falling on a second similar sheet  $B$ .  $A$  and  $B$  are set with their axes at right angles. A third polarizing sheet  $C$  is placed between the first two with its axis at an angle  $\phi$  to that of sheet  $A$ .
- a. If the amplitude of incident light is  $E$ , what are the amplitudes of transmitted light before and after insertion of polarizer  $C$ ?
  - b. What are the intensities of transmitted light before and after insertion of polarizer  $C$ ?

5.

- a. How do you employ Newton's rings phenomenon for measuring the wavelength of light? Give necessary theory.
- b. Light of 589 nm is used to produce Newton rings using a plano-convex lens and a flat glass side. The radius of curvature of the lens is 1 m. A liquid is inserted between lens and the glass. Find the refractive index of the liquid if the diameter of the 10<sup>th</sup> dark ring is 4.0 mm.
- c. Are light waves longitudinal waves or transverse waves? How do you prove this?
- d. What are the major differences between *Fresnel* and *Fraunhofer* diffraction?

6.

- a. Derive the conditions for dark and bright regions in single slit *Fraunhofer* diffraction pattern.
- b. Monochromatic blue light has been used to obtain the diffraction pattern of a narrow slit on a screen. Keeping the experimental set up unchanged if blue light is replaced by red monochromatic light, what will happen to the diffraction pattern?
- c. Two coherent monochromatic light beams of intensities  $I$  and  $4I$  are superimposed. Calculate the maximum and minimum possible intensities of the resulting beam.
- d. Obtain an equation for the resolving power of an optical instrument having a slit width  $a$ .
- e. Light of wavelength 550 nm is used to observe an object under a microscope. If the aperture of the objective lens has a diameter of 0.8 cm, what is the limiting angle of resolution?