

RAJARATA UNIVERSITY OF SRI LANKA FACULTY OF APPLIED SCIENCES

B.Sc. in Information Technology First Year - Semester I Examination - May 2022

ICT 1404 - Mathematics and Statistics for Computing

Time: Three (03) hours

Instructions

- Answer ANY FIVE (05) questions.
- Calculators are allowed.
- Each question carries equal marks.
- Statistical tables and formulae are provided as attachments.
- 1. a) How many different words can be made using four letters of the word 'STATISTICS'?

(08 Marks)

- b) A group consists of 4 girls and 7 boys. In how many ways can a team of 5 members be selected if the team has
 - (i) no girls
 - (ii) three girls
 - (iii) at least three girls

(12 Marks)

2. a) If $A = \begin{pmatrix} 1 & 2 \\ 5 & 4 \end{pmatrix}$ and $B = \begin{pmatrix} 1 & -2 \\ 3 & 5 \end{pmatrix}$.

Find (i) A+B (ii) AB (iii) A-1

(06 Marks)

b) Using matrices or Cramer's rule solve the following linear equations. x+3y+z=5, 3x+4y+3z=5, 9x+y+7z=-3

(14 Marks)

3. a) A basket contains 4 red, 5 blue and 3 green marbles. If 2 marbles are drawn at random from the basket, what is the probability that both are red?

(08 Marks)

b) A student travels to college by three routes A, B and C. The probabilities he choose those routes are respectively 0.6, 0.3 and 0.1. The probabilities that he is late for college if he go via route A, B and C respectively are 0.02, 0.04 and 0.05. Find the probability he is late for the college.

(12 Marks)

- 4. a) At a fuel filling station, 80% of customers pay by credit cards. Find the probability that in a randomly selected sample of five customers,
 - a) Exactly two customers pay by credit cards,
 - b) More than two customers pay by credit cards.

(10 Marks)

- b) On average a machine breaks down four times during a month. Assuming that the number of breaks down can be modeled by a Poisson distribution, find the probability that it breaks down
 - a) Exactly once in a given month,
 - b) At most two times a given month,.

(10 Marks)

- 5. The weight of a sugar packet which is normally distributed with a mean 400g and standard deviation 10g. Find the probability that the weight of a randomly selected sugar packet is
 - a) more than 410g
 - b) less than 390g.
 - c) between 380g and 420g
 - d) between 395g and 400g

(20 Marks)

6. The information of age and value of a car are given by the following table.

Age, x (years)	Value, Rs. y (million)				
2.0	6.1				
2.5	5.5				
3.0	5.0				
3.5	4.6				
4.5	3.8				
5.0	3.5				
6.0	3.3				
7.0	1.0				

- a) Calculate the equation of line of regression of y on x using (10 Marks) least square method.
- b) Estimate the value of the car when the age is 4 years.

(04 Marks)

c) Compute the coefficient of Determination.

(06 Marks)

--END--

Statistical Formulae

Probability Distributions

$$Z = \frac{X - \mu}{\pi}$$

 $Z = \frac{X - \mu}{\sigma}$, here $\mu = \text{mean and } \sigma = \text{standard deviation}$

$$P(x) = {}^{n}C_{x}p^{x}q^{n-x};$$
 here, $q=1-p$, $x=0,1,2,...n$

$$P(x) = \frac{e^{-\lambda} \lambda^x}{x!}$$
; here $\lambda = \text{mean} = \text{variance}$

$$P(x < T) = 1 - e^{\frac{T}{\beta}}$$
 or $P(x \ge T) = e^{\frac{T}{\beta}}$ here $\beta = \text{mean}$

"no-memory property".
$$P(x>a+b \mid x>a)=P(x>b)$$

$$P(x>a+b\mid x>a)=P(x>b)$$

Uniform distribution

$$P(x) = \frac{1}{b-a} \quad \text{for} \quad a < x < b$$
$$= 0 \quad \text{elsewhere}$$

Correlation and Regression Analysis

$$r = \frac{n\sum xy - \sum x\sum y}{\sqrt{[n\sum x^2 - (\sum x)^2][n\sum y^2 - (\sum y)^2]}}$$

Line of regression,
$$y=a+bx$$

Line of regression,
$$y=a+bx$$
 where $b = \frac{n\sum xy - \sum x\sum y}{n\sum x^2 - (\sum x)^2}$ $a = y - bx$


$$a = \overline{y} - b\overline{x}$$

Coefficient of Determination, R
$$R^2 = b^2 \left(\frac{\sum x^2 - n\overline{x}^2}{\sum y^2 - n\overline{y}^2} \right)$$
 or $R^2 = b^2 \left(\frac{n\sum x^2 - (\sum x)^2}{n\sum y^2 - (\sum y)^2} \right)$

$$R^{2} = b^{2} \left(\frac{n \sum x^{2} - (\sum x)^{2}}{n \sum y^{2} - (\sum y)^{2}} \right)$$

$$R^2=r^2$$


සෘණ Z අගයයන් සඳහා සම්මත පුමත වගුව Standard Normal Table for negative Z values



වගු අගය = වර්ගඵලය Table value =Area

2	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
-3.4	.0003	.0003	.0003	.0003	,0003	.0003	.0003	.0003	.0003	.0002
-3.3	.0005	.0005	.0005	.0004	.0004	.0004	.0004	.0004	.0004	.0003
-3.2	.0007	.0007	.0006	.0006	.0006	.0006	.0006	.0005	.0005	.0005
-3.1	.0010	.0009	.0009	.0009	.0008	.0008	.0008	.0008	.0007	.0007
-3.0	.0013	.0013	.0013	.0012	.0012	.0011	.0011	.0011	.0010	.0010
-2.9	.0019	.0018	.0018	- 2017 -	WI6	.0016		,0015	, and	
2.3	.0026	.0025	797797	.0023		.0022	(0021	1824		,0019
227	.0035	.0034	.0033		(00)	.0030	.0029	.0028		
-2.6	.0047	-0.45	(64)	.0043		,0040	.0039	- 100018	.0037	
医肾盂	0062	(000)	.0059	.0057	2005	10.56	.0052	10051	.000	OX.
-2.4	.0082	.0080	.0078	.0075	.0073	.0071	.0069	.0068	.0066	.0064
-2.3	.0107	.0104	.0102	.0099	.0096	.0094	.0091	.0089	.0087	.0084
-2.2	.0139	.0136	.0132	.0129	.0125	.0122	.0119	.0116	.0113	.0110
-2.1	.0179	.0174	.0170	.0166	.0162	.0158	.0154	.0150	.0146	.0143
-2.0	.0228	.0222	.0217	.0212	.0207	.0202	.0197	.0192	.0188	.0183
210	10.31	0281	20274		11112				MAN TO SERVICE	100
	1039	.0351		.0336	.0329	3022	40.14	100	400	.024
37.2		6436			.0409	.0001			.0375	.0367
87	4841	05.17	1624	.0516	.0505	.0495		.aris		(4.5)
-1.5	.0668	.0e55	.0643	.0630	.0618	.060e	4504		(0574	4,5%
-1.4	.0808	.0793	.0778	.0764	.0749	.0735	.0721	.0708	.0694	.0681
-1.3	.0968	.0951	.0934	.0918	.0901	.0885	.0869	.0853	.0838	.0823
-1.2	.1151	.1131	.1112	.1093	.1075	.1056	.1038	.1020	.1003	.0985
-1.1	.1357	.1335	.1314	.1292	.1271	.1251	.1230	.1210	.1190	.1170
-1.0	.1587	.1562	.1539	.1515	.1492	.1469	.1446	.1423	.1401	.1379
_0.9		.3814	.1789	.1762	and the second	.1711	1/015	.1040		
	2119		-2861	2003	1005	.1977	.1949	1924	.1894	الطال
-0.7	2420	2389		2127	21%	2266	62.0	22.6	alve.	214
-0.6	2743	.2709	.2676			2576	2546	2514		245)
	.3025	3050	.3015			.2912	200		2.10	11.
-0.4	3446	.3409	.3372	.3336	.3300	.3264	_3228	3192	.3156	.3121
-0.3	.3821	.3783	.3745	.3707	3669	.3632	.3594	.3557	.3520	.3483
-0.2	.4207	.4168	.4129	.4090	A052	A013	3974	.3936	.3897	.3859
-0.1	.4602	A562	.A522	.4483	,4443	.4404	.4364	.4325	A286	.4247
0.0	.5000	.4960	.4920	.4880	.4840	.4801	A761	.4721	.4681	.4641

ධන Z අගයයන් සඳහා සම්මත පුමත වගුව Standard Normal Table for positive Z values



වගු අගය = වර්ගඵලය

Table value =Area

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	.5000	.5438	.5080	.5120 .5517	.5160 .5357	.5199 .5596	.5239	.5279 .5675	.5319 .5714	.539 .575
0.1	5793	5832	.5478 .5871	.5910	.5948	.5987	.5636 .6026	.6064	.5/14	.614
4	.5179	2.40 1						.6443		
0.4	.6554	.6217 .6591	,6255 ,6628	.6664	.6331 .6700	.6368 .6736	.6406	.6808	.6480	.651
0.3										
o e										1
才團	750				374		472			
						8023				
0.9										
1.0	.8413	.8438	.8461				8554			.862
1.1	.8643	.8665	.2686	£708	.8729	.8749	.8770	.8790	.8810	.883
1.2	8849	.8869	.8888	.8907	.8925	.8944	.8962	.8980	.8997	.901
1.3	.9032	9049	.9066	9082	9099	.9115	.9131	.9147	.9162	.917
1.4	.9192	.9207	9222	9236	.9251	9265	.9279	9292	.9306	.931
15	9302	2345			0.02	. 9594	4000	SUIT		944
1.6		to ki	1.777		200	9305	.9515			
1.7			.9573		1989]	.0599	9.00	1		and the second
					4491		10.04		9009	. 7
10			9726			.9744			9761	
2.0	.9772	.9778	.9783	.9788	.9793	.9798	.9803	.9808	.9812	,981
2.1	.9821	.9826	.9830	.9834	.9838	.9842	.9846	.9850	.9854	.985
2.2	.9861	.9864	.9868	.9871	.9875	.9878	.9881	.9884	.9887	.989
2.3	.9893	.9896	.9898	.9901	.9904	.9906	.9909	.9911	.9913	.991
2.4	.9918	.9920	.9922	.9925	.9927	.9929	.9931	.9932	.9934	.993
2.5			.9941			.9946		39989	Mal	995
2.6	.9833	3983	.9956		1.5	4.250	.9961	.9962		
2.7	9965	996	100 yr	Light	.9969	.9970	3971	.0012	Jase District	
2.3	9074	9916	.9936		. am.	9978			1910	
					.5984				. 5486	998
3.0	.9987	.9987	.9987	.9988	.9988	.9989	.9989	.9989	.9990	.999
3.1	.9990	.9991	,9991	.9991	.9992	,9992	.9992	.9992	.9993	.999
3.2	.9993	.9993	.9994	.9994	.9994	.9994	.9994	.9995	.9995	.999
3.3	.9995	.9995	.9995	.9996	.9996	.9996	.9996	.9996	.9996	.999
3.4	.9997	.9997	.9997	.9997	.9997	,9997	.9997	.9997	.9997	,9991