



RAJARATA UNIVERSITY OF SRI LANKA
FACULTY OF APPLIED SCIENCES

B.Sc (General) Degree in Applied Sciences
 First Year - Semester II Examination – October/November 2017

CHE 1302 – PHYSICAL CHEMISTRY I

Answer **FIVE** questions only

Time: Three (3) hours

$$R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}, 1 \text{ bar} = 10^5 \text{ Pa}, 1 \text{ atm} = 1.01325 \times 10^5 \text{ Pa}$$

1.

- a) Assuming ideal gas behavior, calculate the root mean square speed of a helium atom (relative atomic mass 4.0) at 298 K.
- b) Show that the mean molar translational kinetic energy of an ideal gas is $3/2 RT$.
- c) Use the laws put forward by Boyles, Charles and Avogadro to show that the Boltzmann constant (k) and the universal gas constant (R) are independent of the nature of gas.
- d) Indicate briefly the advantage of using the reduced equation of state instead of the van der Waal's equation of state to describe the behavior of gases.
- e) Explain why real gaseous CO_2 and NH_3 can be liquefied very easily at room temperature while it is impossible to liquefy gaseous H_2 and O_2 under similar conditions.
- f) Calculate the three critical constants for NH_3 given that the van der Waals constants for this gas are $a = 0.366 \text{ Nm}^4 \text{ mol}^{-2}$ and $b = 4.3 \times 10^{-5} \text{ m}^3 \text{ mol}^{-1}$

2.

(100 marks)

a)

- i. What are extensive and intensive properties?
- ii. Classify the following into extensive and intensive properties

Enthalpy, density, pressure, heat capacity, molar volume, free energy, kinetic energy, entropy, molecular weight, volume

(20 marks)

- b) 5 mol of an ideal gas at 400 K and 4 atm pressure undergo expansion isothermally to half the initial pressure. If this expansion take place;

- i. Reversibly
- ii. Irreversible against zero pressure
- iii. Irreversible against 5 atm external pressure

Calculate the work done on the gas by surrounding in each case. (30 marks)

- c) 2 mol of an ideal gas is allowed to expand reversibly at a constant temperature of 0 °C from an initial pressure of 25 atm to final pressure of 0.2 atm.

Calculate

- i. The work done w by the system
- ii. Internal energy change ΔU of the system
- iii. Enthalpy change ΔH of the system

(30 marks)

- d) What is the ΔH when 10 mol of H_2O at 20 atm is heated from 30 °C to 40 °C?

$$C_{p,m}(\text{liquid, } H_2O) = 75 \text{ J K}^{-1} \text{ mol}^{-1}$$

$$\Delta H_{\text{Vapourization}} = 47.3 \text{ kJ mol}^{-1} \text{ at } 373 \text{ K}$$

$$C_{p,m}(\text{gaseuos, } H_2O) = 35.4 \text{ J K}^{-1} \text{ mol}^{-1}$$

(20 marks)

3.

- a) Define the compressibility factor (z) of a gas and show how the deviation of the behavior of a real gas from ideality can be expressed in terms of the compressibility factor for

- i) A given gas at various temperatures
- ii) A series of gases at a temperature

(25 marks)

- b) Explain what do you understand by the term 'critical temperature' of a gas

(15 marks)


c)

- i. Briefly explain the Gibbs free energy criterion for equilibrium and spontaneity in a system
- ii. Ammonium chloride dissolves in water at 300 K with a molar increase in enthalpy of 243 k J mol⁻¹. The corresponding molar entropy change for the dissolution is 168.5 kJ mol⁻¹. Calculate the value of ΔG

- iii. Deduce whether ammonium chloride will spontaneously dissolve in water at constant pressure and temperature
- iv. If this conversion take place reversibly (hypothetically) under condition of constant pressure and constant temperature, calculate the value of ΔG

(60 marks)

4. The galvanic cell represented below consists of a hydrogen half-cell and a magnesium half-cell at standard conditions.



$$E^0 (\text{Mg}^{2+}/\text{Mg}) = -2.37 \text{ V}$$

- a) Write down the balanced equation with the help of half cell reactions, for the net (overall) cell reaction taking place in the above cell.
- b) Write down the cell notation for the above cell.
- c) Calculate the cell potential using Nernst equation and Gibbs free energy change.
- d) How will the initial voltmeter reading be affected if the concentration of the electrolyte in the $\text{Mg(s)}/\text{Mg}^{2+}(\text{aq})$ half-cell is decreased? Explain your answer.
- e) What is 'X' in the diagram and its role in this cell?

(40 marks)

- f) Write short notes on the following:
 - (i) Helmholtz electrical double layer
 - (ii) Liquid junction potential
 - (iii) $\text{Ag}/\text{AgCl}/\text{Cl}^-$ electrode

(60 marks)

5.

- a) The molar conductivity of a solution containing 2.54 g of CuSO_4 /L is $91.0 \text{ S cm}^2 \text{ mol}^{-1}$. What is the resistance of a 100 cm^3 of this solution when placed between two electrodes 1.00 cm apart, each having an area of 1.00 cm^2 . (atomic weight: Cu = 63.5, S = 32.0, O = 16.0 g mol^{-1}) (30 marks)

- b) Sketch and explain the variation of molar conductivity of KCl and CH_3COOH against concentration. (30 marks)

- c) Give detailed experimental method for the determination of the solubility product of PbSO_4 (40 marks)

6.

- a) One mole of an ideal gas is compressed isothermally but irreversibly at 130°C from 2.5 to 6.5 bar in a piston/cylinder device. The work required is 30% greater than the work of a reversible isothermal compression. The heat transferred from the gas during the compression flows to a heat reservoir at 25°C . Calculate the entropy changes of the gas, the heat reservoir, and ΔS_{total} . (30 marks)

- b) Derive an equation for the determination of pH of an acid solution by quinhydrone electrode. (40 marks)

- c) The limiting molar conductivities of KCl, KNO_3 , and AgNO_3 are $14.99 \text{ mS m}^2 \text{ mol}^{-1}$, $14.50 \text{ mS m}^2 \text{ mol}^{-1}$ and $13.34 \text{ mS m}^2 \text{ mol}^{-1}$ respectively at 25°C . Calculate the limiting molar conductivity of AgCl at this temperature?

(Hint: The basis for the solution is Kohlrausch's law of independent of ions)

(30 marks)