



RAJARATA UNIVERSITY OF SRI LANKA
FACULTY OF APPLIED SCIENCES

B.Sc. (Four Year) Degree in Applied Sciences
Fourth Year - Semester I Examination – October / November 2017

PHY 4312 - STATISTICAL THERMODYNAMICS

Time: Three (3) hours

Answer all **five** questions

The use of a non-programmable calculator is permitted.

1.

- a) Derive the binomial distribution function, $W_N(n_1) = [N!/(n_1!n_2!)]p^{n_1}q^{n_2}$ using approximations in one dimensional random walk where, p and q are probabilities of stepping to the right and left respectively, n_1 is number of steps to the right and n_2 is number of steps to the left out of total N steps.

(40 marks)

b) Prove that;

- i) the mean value of n_1 is given by $\bar{n}_1 = Np$,
- ii) the mean value of n_2 is given by $\bar{n}_2 = Nq$ and
- iii) the mean displacement is given by $\bar{m} = N(p - q)$.
- iv) Mean dispersion $\overline{\Delta n_1^2} = Npq$

(60 marks)

2.

- a) A particle of mass m is confined to a one dimensional box with length L . Prove that the number of states accessible to the system is $\frac{L\sqrt{2m}}{2\pi\hbar\sqrt{E}}\delta E$ (symbols have their usual meaning).

(50 marks)

- b) A particle of mass m is free to move in one dimension and its position and momentum coordinates are x and p respectively. Suppose that this particle is confined within $x = 0$ and $x = L$ and its energy lies between E and $E + \delta E$. Draw classical phase space for this particle, indicating the regions accessible to the particle.

(50 marks)

3.

- a) Probability $W_N(n)$ of an event characterized with a probability p occurs n times in N trials is given by the binomial distribution, $W_N(n) = [N!/(n!(N-n)!)]p^n(1-p)^{N-n}$. Consider a situation where n is very small compared to N ($n \ll N$). Using the result $\ln(1-p) \approx -p$ for very small p , show that $(1-p)^{N-n} \approx e^{-Np}$.

(25 marks)

- b) Prove that $\frac{N!}{(N-n)!} \approx N^n$

(25 marks)

- c) Hence, prove that $W_N(n)$ reduces to $W_N(n) = \frac{\lambda^n}{n!} e^{-\lambda}$ where λ is the mean number of events.

(25 marks)

- d) Pure Na sample prepared in a lab contains K as an impurity. The probability of finding K atom is 0.01 and Na is 99.99. What is the probability of containing exactly five K atoms in a sample of 300 g ?

(25 marks)

4.

- a) Define purely thermal interaction and quasi-static process.

(30 marks)

- b) An ideal gas has a temperature-independent molar specific heat capacity c_v at constant volume. Let $\gamma = \frac{c_p}{c_v}$ denotes the ratio of its specific heats. The gas is thermally insulated and allowed to expand quasi-statically from an initial volume V_i at temperature T_i to the final volume V_f .

i) Use the relation $pV^\gamma = \text{Constant}$, to find the final temperature.

ii) Use the fact that the entropy remains constant in this process to find the final temperature, T_f .

(70 marks)

5.

- a) Show that $\beta(\tilde{E})$ is a constant for a purely thermal quasi static macroscopic interaction of two systems where, $\Omega(E)$ is number of accessible states to the system, \tilde{E} is most probable energy of the system and $\beta(E) = \frac{\partial}{\partial E} \ln \Omega(E)$.

(60 marks)

- b) Prove that, when the system is at the equilibrium,

i) the entropy change is given by $\frac{dE}{T}$,

ii) the entropy change in the combined system is maximum and

iii) the temperature difference between two systems is zero.

(40 marks)

-END-