

QCQI Chapter 4 Exercises

Felix Tripier

October 20, 2018

4.2: Let x be a real number and A a matrix such that $A^2 = I$. Show that

$$\exp(iAx) = \cos(x)I + i \sin(x)A$$

Answer:

$$\exp(iAx) = \sum_{n=1}^{\infty} \frac{1}{n!} (iAx)^n \quad (1a)$$

$$= \sum_{n=1}^{\infty} \frac{1}{(2n)!} (iAx)^{2n} + \frac{1}{(2n+1)!} (iAx)^{2n+1} \quad (1b)$$

$$= I \sum_{n=1}^{\infty} \frac{(-1)^n}{(2n)!} x^n + \frac{1}{(2n+1)!} (iAx)^{2n+1} \quad (1c)$$

$$= I \sum_{n=1}^{\infty} \frac{(-1)^n}{(2n)!} x^n + iA \frac{(-1)^n}{(2n+1)!} x^n \quad (1d)$$

$$= \cos(x)I + i \sin(x)A \quad (1e)$$

4.3: Show that, up to a global phase, the $\pi/8$ gate satisfies $T = R_z(\pi/4)$

Answer:

$$\pi/8 \text{ gate} = \exp(i\pi/8) \begin{bmatrix} e^{-i\pi/8} & 0 \\ 0 & e^{-i\pi/8} \end{bmatrix}$$

and

$$R_z(\pi/4) = \begin{bmatrix} e^{-i\pi/8} & 0 \\ 0 & e^{-i\pi/8} \end{bmatrix}$$

$\exp(i\pi/8)$ is just a global phase factor dude.

4.4: Express the Hadamard gate H as a product of R_x and R_z rotations and $e^{i\varphi}$

Answer:

$$\text{Hadamard} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \quad (2a)$$

$$R_z(\theta) = \begin{bmatrix} e^{-i\theta/2} & 0 \\ 0 & e^{i\theta/2} \end{bmatrix} \quad (2b)$$

$$R_x(\theta) = \begin{bmatrix} \cos \frac{\theta}{2} & -i \sin \frac{\theta}{2} \\ -i \sin \frac{\theta}{2} & \cos \frac{\theta}{2} \end{bmatrix} \quad (2c)$$

$$R_z(-2\pi) = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \quad (2d)$$

$$R_x(\pi) = \begin{bmatrix} 0 & -i \\ -i & 0 \end{bmatrix} \quad (2e)$$

$$e^{i \cdot \frac{5\pi}{2}} \cdot R_x(\pi/2) \cdot R_z(\pi/2) \cdot R_x(\pi/2) = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \quad (2f)$$

The insight here is that this series of rotations is equivalent (within a global phase factor) to the Hadamard's more obvious set of rotations: 90 degrees around the y axis followed by 180 degrees around the x axis.

4.5 Prove that $(\hat{n} \cdot \sigma)^2 = I$ and use this to verify this equation:

$$R_{\hat{n}}(\theta) \equiv \exp(-i\theta \hat{n} \cdot \sigma/2) = \cos\left(\frac{\theta}{2}\right)I - i \sin\left(\frac{\theta}{2}\right)(n_x X + n_y Y + n_z Z)$$

Okay, proof by equational reasoning:

\hat{n} is a unit vector so $n_z^2 + n_x^2 + n_y^2 = 1$.

$$(\hat{n} \cdot \sigma)^2 = \left(\begin{bmatrix} 0 & n_x \\ n_x & 0 \end{bmatrix} + \begin{bmatrix} 0 & -in_y \\ -in_y & 0 \end{bmatrix} + \begin{bmatrix} n_z & 0 \\ 0 & -n_z \end{bmatrix} \right)^2 \quad (3a)$$

$$= \left(\begin{bmatrix} n_z & n_x - in_y \\ n_x + in_y & n_z \end{bmatrix} \right)^2 = \begin{bmatrix} n_z^2 + n_x^2 + n_y^2 & 2n_z(n_x - in_y) \\ 2n_z(n_x + in_y) & n_z^2 + n_x^2 + n_y^2 \end{bmatrix} \quad (3b)$$

$$= \begin{bmatrix} 1 & 2n_z(n_x - in_y) \\ 2n_z(n_x + in_y) & 1 \end{bmatrix} \quad (3c)$$

Hermitian operators must be equal to their adjoint and so

$$2n_z(n_x + in_y) = 2n_z(n_x - in_y)$$

We'll call this term "scooby-doo".

The determinant of a rotation matrix must be 1, and so we have

$$1 - \text{scooby-doo}^2 = 1 \tag{4a}$$

$$-\text{scooby-doo}^2 = 0 \tag{4b}$$

$$\text{scooby-doo} = 0 \tag{4c}$$

and therefore

$$\begin{bmatrix} 1 & \text{scooby-doo} \\ \text{scooby-doo} & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I$$