

SURGICAL PATHOLOGY REPORT

UUID:33C8B2EA-980D-4CDC-B8A9-B8583E048CCA
TCGA-XF-A8HH-01A-PR

Redacted



Collected
Ordered by

FINAL DIAGNOSIS:

RADICAL CYSTECTOMY WITH BILATERAL PELVIC LYMPH NODE DISSECTION, BOWEL RESECTION FOR STUDDOR POUCH TO URETHRA AND GASTROSTOMY TUBE PLACEMENT:

RIGHT DISTAL URETER (A):

FROZEN SECTION DIAGNOSIS:

- NO HIGH GRADE UROTHELIAL ATYPIA OR CARCINOMA IDENTIFIED

FINAL DIAGNOSIS:

- BENIGN URETER
- NO EVIDENCE OF MALIGNANCY

LEFT DISTAL URETER (B):

FROZEN SECTION DIAGNOSIS:

- NO HIGH GRADE UROTHELIAL ATYPIA OR CARCINOMA IDENTIFIED

FINAL DIAGNOSIS:

- BENIGN URETER
- NO EVIDENCE OF MALIGNANCY

APICAL URETHRAL MARGIN (C):

FROZEN SECTION DIAGNOSIS:

- NO HIGH GRADE UROTHELIAL ATYPIA OR CARCINOMA IDENTIFIED

FINAL DIAGNOSIS:

- NO HIGH GRADE UROTHELIAL ATYPIA OR CARCINOMA IDENTIFIED

URINARY BLADDER, UTERUS, BILATERAL FALLOPIAN TUBES AND OVARIES (D):

URINARY BLADDER:

- TRANSITIONAL CELL CARCINOMA, GRADE 3-4/4, SHOWING FULL THICKNESS INVASION OF THE BLADDER WALL TO INVOLVE THE PERIVESICAL FAT
- MULTIFOCAL TRANSITIONAL CELL CARCINOMA IN SITU (FLAT LESION) IDENTIFIED
- FOCAL VASCULAR SPACE INVASION IDENTIFIED
- METASTATIC TRANSITIONAL CELL CARCINOMA IDENTIFIED IN ONE POSTERIOR PERIVESICAL LYMPH NODE (1/1)
- ALL MARGINS ARE NEGATIVE FOR TUMOR

UTERUS:

- THE UTERINE CERVIX IS UNREMARKABLE
- UTERUS SHOWING ADENOMYOSIS
- A SINGLE INTRAMURAL LEIOMYOMA
- PROLIFERATIVE ENDOMETRIUM

BILATERAL FALLOPIAN TUBES AND OVARIES:

- BILATERAL FALLOPIAN TUBES ARE UNREMARKABLE AND FREE OF TUMOR
- BILATERAL OVARIES SHOW NORMAL PHYSIOLOGIC CHANGES AND ARE FREE OF TUMOR

LEFT PERIAORTIC LYMPH NODES (E):

- NO EVIDENCE OF MALIGNANCY IN 12 LYMPH NODES (0/12)

LEFT COMMON ILIAC LYMPH NODES (F):

- NO EVIDENCE OF MALIGNANCY IN FIVE LYMPH NODES (0/5)

RIGHT PARACAVAL LYMPH NODES (G):

- NO EVIDENCE OF MALIGNANCY IN FIVE LYMPH NODES (0/5)

RIGHT COMMON ILIAC LYMPH NODES (H):

- NO EVIDENCE OF MALIGNANCY IN SIX LYMPH NODES (0/6)

RIGHT NODE OF CLOQUET (I):

ICD-O-3

Carcinoma, transitional cell NOS
8120/3

Site: Bladder NOS
C67.9

JW 1/23/14

SURGICAL PATHOLOGY REPORT

Collected:
Ordered by:

- NO EVIDENCE OF MALIGNANCY IN ONE LYMPH NODE (0/1)

RIGHT EXTERNAL ILIAC LYMPH NODES (J):

- METASTATIC TRANSITIONAL CELL CARCINOMA IDENTIFIED IN ONE OF TWO LYMPH NODES (1/2)

RIGHT OBTURATOR / HYPOGASTRIC LYMPH NODES (K):

- NO EVIDENCE OF MALIGNANCY IN 21 LYMPH NODES (0/21)

LEFT NODE OF CLOQUET (L):

- NO EVIDENCE OF MALIGNANCY IN ONE LYMPH NODE (0/1)

LEFT EXTERNAL ILIAC LYMPH NODES (M):

- METASTATIC TRANSITIONAL CELL CARCINOMA IDENTIFIED IN ONE OF THREE LYMPH NODES (1/3)

LEFT OBTURATOR / HYPOGASTRIC LYMPH NODES (N):

- NO EVIDENCE OF MALIGNANCY IN 12 LYMPH NODES (0/12)

RIGHT PRESACRAL LYMPH NODES (O):

- NO LYMPHOID TISSUE IDENTIFIED
- NO EVIDENCE OF MALIGNANCY

PRESACRAL LYMPH NODES (P):

- NO EVIDENCE OF MALIGNANCY IN FOUR LYMPH NODES (0/4)

LEFT PRESACRAL LYMPH NODES (Q):

- NO LYMPHOID TISSUE IDENTIFIED
- NO EVIDENCE OF MALIGNANCY

LEFT PROXIMAL URETER (R):

- BENIGN URETER
- NO EVIDENCE OF UROTHELIAL ATYPIA OR MALIGNANCY IDENTIFIED

RIGHT PROXIMAL URETER (S):

- BENIGN URETER
- NO EVIDENCE OF UROTHELIAL ATYPIA OR MALIGNANCY IDENTIFIED

LYMPH NODE SUMMARY: METASTATIC TRANSITIONAL CELL CARCINOMA
IDENTIFIED IN THREE OUT OF 73 LYMPH NODES
EXAMINED (3/73)

INTRAOPERATIVE DIAGNOSIS:

FROZEN SECTION DIAGNOSIS:

AFS: Right distal ureter:

- no high-grade urothelial atypia or carcinoma identified

BFS: Left distal ureter:

- no high-grade urothelial atypia or carcinoma identified

CFS: Apical urethral margin:

- no high-grade urothelial atypia or carcinoma identified

GROSS DESCRIPTION:

A: The specimen is received fresh from the O.R. and labeled "Right distal ureter."
It consists of a 1 x 0.5 x 0.5 cm fragment of pink-tan tissue. Entirely submitted

SURGICAL PATHOLOGY REPORT

Collected: [REDACTED]

Ordered by: [REDACTED]

for frozen section.

B: The specimen is received fresh from the O.R. and labeled "Left distal ureter." It consists of a 1 x 0.5 x 0.3 cm fragment of red-tan tissue. Entirely submitted for frozen section.

C: The specimen is received fresh from the O.R. and labeled "Apical urethral margin." It consists of a 1.5 x 0.3 x 0.2 cm fragment of red-tan tissue. Entirely submitted for frozen section.

D: The specimen is received fresh from the O.R. and labeled "Urinary bladder, Uterus, Bilateral fallopian tubes & ovaries." It consists of a 15 x 12 x 5.5 cm cystectomy, total abdominal hysterectomy and bilateral salpingo-oophorectomy. The bladder measures 6.5 x 7 x 4 cm. The uterus measures 9 x 6.5 x 2.5 cm. The right ovary measures 2 x 1.5 x 1 cm, and the left ovary measures 2.5 x 1.5 x 1 cm. The right fallopian tube measures 7.5 cm in length and 0.4 cm in diameter. The left fallopian tube measures 6.5 cm in length and 0.5 cm in diameter. There is no serosal involvement. The specimen is inked blue and the urethral margin is submitted for frozen section. The bladder is then opened along its anterior surface to reveal a 4 x 3.5 x 1 cm endophytic, ulcerated mass in the posterior portion of the bladder. Both ureterovesical junctions are grossly negative. Tumor extends to within 0.2 cm of each ureterovesical junction. The remaining bladder mucosa is otherwise unremarkable. The ureters are opened to reveal grossly normal mucosa. The left ureter measures 4.5 cm in length and 1.2 cm in circumference. The right ureter measures 4.6 cm in length and 1.1 cm in circumference. The right ovary and fallopian tube are grossly unremarkable on external examination and on sectioning. The left ovary and fallopian tube are unremarkable on external examination and sectioning. The ectocervix measures 4 x 3 cm and is lined by smooth tan epithelium. The endocervical canal measures 2.9 cm in length and 0.7 cm in diameter and is lined by smooth, tan epithelium. The endometrial canal measures 3 x 1.7 x 0.1 cm and is lined by smooth, tan epithelium. Serial sectioning of the uterus reveals an unremarkable myometrium and serosa with the exception of a single nodule 0.5 cm in greatest dimension, located intramurally. Careful palpation of the perivesical fat pad reveals no lymphadenopathy. Representative sections are submitted in 22 cassettes.

E: The specimen is received in formalin and labeled "Left peri aortic lymph nodes." It consists of a 4.5 x 2.2 x 1.2 cm of fatty tissue. Entirely submitted in two cassettes.

F: The specimen is received in formalin and labeled "Left common iliac lymph nodes." It consists of a 5 x 2 x 0.5 cm of greasy yellow tissue. Entirely submitted in two cassettes.

G: The specimen is received in formalin and labeled "Right para caval lymph nodes." It consists of a 4.2 x 2.7 x 1.5 cm of greasy yellow tissue. Entirely submitted in two cassettes.

H: The specimen is received in formalin and labeled "Right common iliac lymph nodes." It consists of a 4 x 2 x 1.5 cm greasy yellow tissue. Entirely submitted in two cassettes.

I: The specimen is received in formalin and labeled "Right node of cloquet." It consists of a 2 x 1 x 1 cm of greasy yellow tissue. Entirely submitted in one cassette.

J: The specimen is received in formalin and labeled "Right external iliac lymph nodes." It consists of a 4 x 3 x 0.7 cm greasy yellow tissue. Entirely submitted in two cassettes.

SURGICAL PATHOLOGY REPORT

Collected

Ordered by

K: The specimen is received in formalin and labeled "Right obturator / hypogastric lymph nodes." It consists of a 5 x 5 x 1 cm of greasy yellow tissue fragment. Entirely submitted in five cassettes.

L: The specimen is received in formalin and labeled "Left node of cloquet." It consists of a 1 x 0.7 x 0.5 cm yellow-tan tissue fragment. Entirely submitted in one cassette.

M: The specimen is received in formalin and labeled "Left external iliac lymph nodes." It consists of a 3 x 2 x 1 cm greasy yellow tissue. Entirely submitted in two cassettes.

N: The specimen is received in formalin and labeled "Left obturator / hypogastric lymph nodes." It consists of a 5 x 5 x 1 cm of greasy yellow tissue fragment. Entirely submitted in five cassettes.

O: The specimen is received in formalin and labeled "Right presciatic lymph nodes." It consists of a 2 x 1 x 0.5 cm greasy yellow tissue fragment. Entirely submitted in one cassette.

P: The specimen is received in formalin and labeled "Presacral lymph nodes." It consists of a 3 x 2 x 1 cm of greasy yellow tissue. Entirely submitted in one cassette.

Q: The specimen is received in formalin and labeled "Left presciatic lymph nodes." It consists of a 1 x 1 x 0.5 cm of greasy yellow tissue fragment. Entirely submitted in one cassette.

R: The specimen is received in formalin and labeled "Left proximal ureter." It consists of a 1.5 cm length of ureter, 0.8 cm in diameter. Entirely submitted in one cassette.

S: The specimen is received in formalin and labeled "Right proximal ureter." It consists of a 1.5 cm length of ureter, 0.7 cm in diameter. Entirely submitted in one cassette.

SECTIONS:

AFS: frozen section, right distal ureter
 BFS: frozen section, left distal ureter
 CFS: frozen section, apical urethral margin
 D1,2: tumor with cervix
 D3-5: tumor with perivesical fat
 D6: right ureterovesical junction
 D7: left ureterovesical junction
 D8: right ureter random sections
 D9: left ureter random sections
 D10: bladder, random section, right
 D11: bladder, random section, left
 D12: right ovary and fallopian tube
 D13: left ovary and fallopian tube
 D14: posterior cervix
 D15: anterior cervix
 D16: posterior endomyometrium and serosa (includes intramural nodule)
 D17: anterior endomyometrium with serosa
 D18-22: perivesical fat pad
 E1,2: left periaortic lymph nodes
 F1,2: left common iliac lymph nodes
 G1,2: right paracaval lymph nodes
 H1,2: right common iliac lymph nodes
 I: right node of Cloquet

SURGICAL PATHOLOGY REPORT

Collected:
Ordered by:

J1,2: right external iliac lymph nodes
K1-5: right obturator/hypogastric lymph nodes
L: left node of Cloquet
M1,2: left external iliac lymph nodes
N1-5: left obturator/hypogastric lymph nodes
O: right presciatic lymph nodes
P: presacral lymph nodes
Q: left presciatic nodes
R: left proximal ureter
S: right proximal ureter

MICROSCOPIC DESCRIPTION:

A-C: See final microscopic-diagnosis.

D: Sections show invasive transitional cell carcinoma, nuclear grade 3-4/4. The cells are medium to large with high nuclear to cytoplasmic ratio prominent nucleoli, vesicular chromatin and coarse chromatin clumping. The mitotic rate is high. The tumor cells grow in sheets and trabeculae and penetrate the muscularis propria extending into the perivesical fat. The tumor does not extend into the vaginal wall. Lymphovascular space invasion is identified (D5). Multifocal carcinoma in situ is identified (A5,11). The cervix is uninvolved. Both ovary and fallopian tubes are unremarkable. The ovaries show normal physiologic changes. The uterus shows proliferative endometrium, adenomyosis and a single intramural leiomyoma. Metastatic transitional cell carcinoma is identified in a single posterior perivesical lymph node (D18).

E-S: See final microscopic-diagnosis.

Signed by:

ELECTRONIC SIGNATURE

POST-OPERATIVE DIAGNOSIS:

Same

PRE-OPERATIVE DIAGNOSIS:

Transitional cell carcinoma of the bladder

ORDERING PHYSICIAN:

Ordering Physician:

Criteria	Yes	No
Diagnosis Discrepancy		✓
Primary Tumor Site Discrepancy		✓
HPV Discrepancy		✓
Prior Malignancy History		✓
Real/Syn. Chronic Primary Note		✓
See Is (C, C, C)		
Reviewer Initials	Date Reviewed: 10/21/2013	