


UUID:7F4BAF33-CF00-4819-914A-5AA8CFB3D797
TCGA-XF-A9T8-01A-PR

Redacted



Collected
Ordered by

Location

DIAGNOSIS:

RADICAL CYSTECTOMY WITH URINARY DIVISION (ILEAL CONDUIT) AND PELVIC LYMPH NODE
DISSECTION :

RIGHT URETER (AFS):

FROZEN SECTION DIAGNOSIS:

- NO TUMOR IDENTIFIED

FINAL DIAGNOSIS:

- BENIGN URETER

- NO DYSPLASIA OR MALIGNANCY IDENTIFIED

LEFT URETER (BFS):

FROZEN SECTION DIAGNOSIS:

- NO TUMOR IDENTIFIED

FINAL DIAGNOSIS:

- BENIGN URETER

- NO DYSPLASIA OR MALIGNANCY IDENTIFIED

BLADDER (C):

- INVASIVE POORLY DIFFERENTIATED UROTHELIAL CARCINOMA WITH SQUAMOID FEATURES (12.0 X 9.5 X 3.2 CM), GRADE 3-4/4, GROSSLY AND MICROSCOPICALLY EXTENDING THROUGH BLADDER WALL INTO PERIVESICLE SOFT TISSUE WITH IMPINGEMENT OF RIGHT DISTAL URETER AND RIGHT ANTERIOR PROXIMAL PROSTATE
- EXTENSIVE UROTHELIAL CARCINOMA IN SITU (FLAT LESION) THROUGHOUT BLADDER
- LYMPHOVASCULAR SPACE INVASION IDENTIFIED
- UNINVOLVED BLADDER SHOWS SQUAMOUS METAPLASIA AND EDEMATOUS CHANGE
- URETHRAL, URETERAL AND PERIVESICLE SOFT TISSUE RESECTION MARGINS, FREE OF DYSPLASIA AND MALIGNANCY

PROSTATE (C):

- BILATERAL PROSTATIC ADENOCARCINOMA, GLEASON'S SCORE 6 (3+3), CONFINED TO PROSTATE WITHOUT TRANSGRESSION OF CAPSULE
- BILATERAL SEMINAL VESICLES, FREE OF PROSTATIC AND UROTHELIAL CARCINOMA
- UNINVOLVED PROSTATE SHOWS GLANDULAR HYPERPLASIA
- APICAL AND RADIAL RESECTION MARGINS, FREE OF DYSPLASIA AND MALIGNANCY

LEFT EXTERNAL ILIAC LYMPH NODES (D):

- NO MALIGNANCY IDENTIFIED IN THREE LYMPH NODES EXAMINED (0/3)

LEFT LYMPH NODE OF CLOQUET (E):

- NO MALIGNANCY IDENTIFIED IN ONE LYMPH NODE EXAMINED (0/1)

LEFT OBTURATOR/HYPOGASTRIC LYMPH NODE (F):

- NO MALIGNANCY IDENTIFIED IN SEVEN LYMPH NODES EXAMINED (0/7)

RIGHT LYMPH NODE OF CLOQUET (G):

- NO MALIGNANCY IDENTIFIED IN ONE LYMPH NODE EXAMINED (0/1)

RIGHT EXTERNAL ILIAC LYMPH NODE (H):

- NO MALIGNANCY IDENTIFIED IN ONE LYMPH NODE EXAMINED (0/1)

RIGHT OBTURATOR/HYPOGASTRIC LYMPH NODE (I):

- NO MALIGNANCY IDENTIFIED IN FOUR LYMPH NODES EXAMINED (0/4)

LEFT PROXIMAL URETER (J):

- BENIGN URETER

- NO DYSPLASIA OR MALIGNANCY IDENTIFIED

*ICD O-3
Carcinoma, urothelial NOS 8120/3
Site: Bladder NOS C67.9
Bladder wall NOS C67.9
JDS 3/26/14*

Per TSS, 10% squamous.

Collected
Entered by

Location

RIGHT PROXIMAL URETER (K):

- BENIGN URETER
- NO DYSPLASIA OR MALIGNANCY IDENTIFIED

LYMPH NODE SUMMARY: NO MALIGNANCY IDENTIFIED IN A TOTAL OF 17 LYMPH NODES EXAMINED (0/17).

PATHOLOGIC STAGING (AJCC 1997):

BLADDER: pT3bNOMX

PROSTATE: pT2bNOMX

SPECIMEN SOURCE:

AFS: "Right ureter"

BFS: "Left ureter"

C: "Prostate + bladder"

D: "Left external iliac LN"

E: "Left LN of cloquet"

F: "Left obturator/hypogastric LN"

G: "Right LN of cloquet"

H: "Right external iliac LN"

I: "Right obturator/hypogastric LN"

J: "Left proximal ureter"

K: "Rt proximal ureter"

CLINICAL INFORMATION:

Pre-Op Dx: Bladder carcinoma

Post-Op Dx: Same

GROSS EXAMINATION:

A: The specimen is received fresh from the O.R. and labeled "Right ureter". It consists of a 0.3 cm length of ureter measuring 0.4 cm in diameter. The outer surface is inked black and the specimen is entirely submitted in cassette AFS, following frozen section.

B: The specimen is received fresh from the O.R. and labeled "Left ureter". It consists of a 0.3 cm length of ureter measuring 0.4 cm in diameter. Entirely submitted in cassette BFS, following frozen section.

C: The specimen is received fresh from the O.R. and labeled "Prostate + bladder". It consists of a 590-gram cystoprostatectomy specimen consisting of bladder (10 x 6.5 x 3.2 cm), attached peritoneum (14 x 10 x < 0.1 cm), attached right ureter (4 x 0.4 cm), attached left ureter (4.5 x 0.4 cm), prostate (4 x 4 x 3.5 cm), right seminal vesicle (2.5 x 1.3 x 0.8 cm) and left seminal vesicle (2.5 x 1 x 0.6 cm). Opening the prostate and bladder anteriorly reveals a ulcerated and centrally necrotic tan-white tumor (12 x 9.5 x 3.2 cm) involving the entire right half of the bladder wall and extending to the dome, posterior wall, and inferiorly to the right anterior proximal prostate. Tumor diffusely replaces the bladder wall and extends into the perivesical fat with thickness of 3.2 cm. Tumor comes within 1 mm of the inked right resection margin. The mucosa beyond the central ulceration is granular and the uninvolved left lateral wall mucosa is edematous and smooth. The right ureter is impinged upon by tumor for a length of 2 cm from the ureterovesical junction. The mucosa of the bilateral ureters is otherwise unremarkable. The

Collected
Ordered by

Location

tumor causes a large area of retraction at the peritoneal surface but does not grossly transgress the peritoneum. The prostatic urethra measures 3.5 x 1.3 cm and the verumontanum measures 0.6 x 0.4 x 0.3 cm. The urethral mucosa is unremarkable.

Serial sectioning through the prostate reveals bilateral periurethral tan-white nodularities. Tumor is again noted extending to the right anterior proximal prostate. The seminal vesicles are uninvolved by tumor. Representative sections are submitted in 20 cassettes.

D: The specimen is received in formalin and labeled "Left external iliac LN". It consists of a 5 x 3.5 x 1 cm aggregate of tan-yellow fat and soft tissue. Three lymph nodes are identified ranging in size from 1 cm to 2.3 cm. Entirely submitted in three cassettes.

E: The specimen is received in formalin and labeled "Left LN of cloquet". It consists of a 0.7 x 0.6 x 0.3 cm tan-white apparent lymph node. Entirely submitted in one cassette.

F: The specimen is received in formalin and labeled "Left obturator/hypogastric LN". It consists of a 3.5 x 2.5 x 1 cm aggregate of tan-yellow fat and soft tissue. Multiple lymph nodes are identified ranging in size from 0.2 cm to 0.7 cm. Entirely submitted in three cassettes.

G: The specimen is received in formalin and labeled "Right LN of cloquet". It consists of a 0.8 x 0.8 x 0.6 cm lymph node which is bisected and entirely submitted in one cassette.

H: The specimen is received in formalin and labeled "Right external iliac LN". It consists of a 2.5 x 1.5 x 0.6 cm aggregate of tan-yellow fat and soft tissue. A single 1.2 cm lymph node is identified. The lymph node is bisected and submitted in cassette H1. The remaining tissue is submitted in cassette H2..

I: The specimen is received in formalin and labeled "Right obturator/hypogastric LN". It consists of a 3 x 2.5 x 1 cm aggregate of tan-yellow fat and soft tissue. Multiple lymph nodes are identified ranging in size from 0.3 cm to 2.5 cm. The largest lymph node is bisected and submitted in cassettes I1 and I2. The remaining tissue is submitted in cassette I3.

J: The specimen is received in formalin and labeled "Left proximal ureter". It consists of a 1.5 cm length of ureter measuring 0.5 cm in diameter. Serially sectioned and entirely submitted in one cassette.

K: The specimen is received in formalin and labeled "Rt proximal ureter". It consists of a 0.8 cm length of ureter measuring 0.5 cm in diameter. Serially sectioned and entirely submitted in one cassette.

SECTIONS:

- AFS: frozen section, right ureter
- BFS: frozen section, left ureter
- C1: prostate + bladder, right lateral bladder wall and tumor
- C2,3: tumor with closest right surgical resection margin
- C4: right ureterovesical junction
- C5: anterior wall and tumor
- C6: dome and tumor
- C7: posterior wall and tumor
- C8: left lateral wall
- C9: left ureterovesical junction
- C10: bladder neck/trigone
- C11: right perivesical fat and possible lymph nodes
- C12: left perivesical fat
- C13: right posterior proximal prostate

Collected:
Ordered by:

Location:

C14: right anterior proximal prostate
 C15: left proximal prostate
 C16: right mid prostate
 C17: left mid prostate
 C18: right distal prostate
 C19: left distal prostate
 C20: apical margin
 D1: left external iliac lymph node; one bisected lymph node
 D2: one bisected lymph node
 D3: remaining lymph node and soft tissue
 E: left lymph node of Cloquet - all embedded
 F1-3: left obturator/hypogastric lymph node - all embedded
 G: right lymph node of Cloquet - all embedded
 H1: right external iliac lymph node; one lymph node, bisected
 H2: remaining tissue
 I1,2: right obturator/hypogastric lymph node; largest lymph node
 I3: remaining tissue
 J: left proximal ureter - all embedded
 K: right proximal ureter - all embedded

INTRAOPERATIVE FROZEN CONSULTATION:

AFS: Right ureter:
- no tumor identified

BFS: Left ureter:
- no tumor identified

Criteria	Yes	No
Diagnosis Discrepancy		✓
Primary Tumor Site Discrepancy		✓
HIPAA Discrepancy		✓
Prior Malignancy History		✓
Dual/Synchronous Primary Noted		✓
Case Is (circle):	QUALIFIED	DISQUALIFIED
Reviewer Initials	gac	1/2/2014