



SPECIMEN

1. Bladder tumour.
2. Bladder tumour base.

CLINICAL DETAILS

Papillary tumour (solid bladder resected).

MACROSCOPY

1. 3 g of friable tissue chippings.
2. 3 pieces of tan coloured tissue the largest measuring 12 mm in maximum dimension.

MICROSCOPY

Parts 1&2.

Grade:

3 poorly differentiated (WHO 1973) high grade (WHO 2004).

Growth pattern:

Mixed papillary and solid.

Type:

Transitional cell carcinoma with some signet ring cell differentiation and some sarcomatoid (spindle cell) differentiation. No heterologous elements are seen. Extensive necrosis is present.

Muscularis propria present:

Present.

Stromal invasion:

Invasion of muscularis propria (pT2) (at least) (part 2).

Vascular channel invasion:

Not identified.

Background urothelium:

A small amount of flat urothelium is present which is not dysplastic.

SUMMARY

Bladder; G3 pT2 (at least).

U6E
ICD-O-3
Carcinoma, urothelial NOS 8120/3
path
Carcinoma, transitional cell NOS 8120/3
Site: Bladder NOS C67.9
J 5/2/14

0% spindle
0% signet ring > per TSS.

Criteria	Yes	No
Diagnosis Discrepancy		
Primary Tumor Site Discrepancy		
HIPAA Discrepancy		
Prior Malignancy History		
Dual/Synchronous Primary Noted		
Case is (circle): QUALIFIED / DISQUALIFIED		
Reviewer Initials	Date Reviewed: 1/3/14	