

FINAL DIAGNOSIS:

PART 1: RIGHT URETER, DISTAL, SHAVE MARGIN, BIOPSY-

- A. BENIGN URETER.
- B. NEGATIVE FOR NEOPLASIA.

PART 2: LEFT, URETER, DISTAL, SHAVE MARGIN, BIOPSY-

- A. BENIGN URETER.
- B. NEGATIVE FOR NEOPLASIA.

UUID: 8716FEB2-A4DD-4EFF-B1A6-D679A5C75C76
TCGA-BT-A42C-01A-PR

Redacted



PART 3: URINARY BLADDER, DISTAL URETERS, PROSTATE, BILATERAL SEMINAL VESICLES, AND BILATERAL DISTAL VASA DEFERENTIA, RADICAL CYSTOPROSTATECTOMY -

- A. INVASIVE UROTHELIAL CARCINOMA, HIGH-GRADE (2004 WHO/ISUP) (SLIDES 3E, 3F, 3L, 3R, 3S, 3T, 3U, 3V), ASSOCIATED WITH FOCAL HIGH-GRADE PAPILLARY UROTHELIAL CARCINOMA (SLIDES 3H, 3O, 3S). TUMOR MEASURES 3.5 x 1.5 x 0.4 CM, AND IS PRESENT IN THE LEFT LATERAL WALL AND TRIGONE. CARCINOMA INFILTRATES INTO THE SUPERFICIAL DETRUSOR MUSCLE (INNER HALF).
- B. UROTHELIAL DYSPLASIA THROUGH CARCINOMA IN SITU IS PRESENT IN THE RIGHT URETERAL OSTIUM (SLIDE 3G), LEFT URETERAL OSTIUM (SLIDE 3H), AND ANTERIOR WALL / DOME (SLIDE 3N).
- C. NO DEFINITIVE ANGIOLYMPHATIC INVASION IS IDENTIFIED (SLIDE 3E).
- D. ALL EXAMINED SURGICAL RESECTION MARGINS ARE FREE OF UROTHELIAL NEOPLASIA.
- E. TNM PATHOLOGIC STAGE: pT2a N0 MX (SEE SYNOPSIS).
- F. BACKGROUND BLADDER WITH GRANULOMATOUS INFLAMMATION (HISTORY OF BCG THERAPY) AND BIOPSY SITE CHANGES.
- G. BENIGN PROSTATE WITH MULTIFOCAL NECROTIZING GRANULOMAS (HISTORY OF BCG THERAPY), NODULAR HYPERPLASIA, FOCAL GLANDULAR ATROPHY, AND CHRONIC INFLAMMATION.
- H. BENIGN BILATERAL SEMINAL VESICLES.

PART 4: LYMPH NODES, RIGHT PELVIC, EXCISION

- A. NO EVIDENCE OF MALIGNANCY IN SEVENTEEN LYMPH NODES (0/17).
- B. ONE LYMPH NODE WITH FEW NON-NECROTIZING GRANULOMAS (SLIDE 4B).

PART 5: LYMPH NODES, LEFT PELVIC, EXCISION

- A. NO EVIDENCE OF NEOPLASIA IN SIX LYMPH NODES (0/6).
- B. ONE LYMPH NODE WITH A SMALL NON-NECROTIZING GRANULOMA (SLIDE 4K).

CASE SYNOPSIS:

SYNOPTIC DATA - PRIMARY URINARY BLADDER TUMORS

SPECIMEN TYPE:	Radical cystoprostatectomy
TUMOR SITE:	Trigone, Left lateral wall
TUMOR SIZE:	Greatest dimension: 3.5 cm Additional dimensions: 1.5 x 0.4 cm
HISTOLOGIC TYPE:	Urothelial (transitional cell) carcinoma
ASSOCIATED EPITHELIAL LESIONS:	None identified
HISTOLOGIC GRADE:	Urothelial carcinoma - High-grade
TUMOR CONFIGURATION:	Papillary, Flat, Ulcerated
PATHOLOGIC STAGING (pTNM):	pT2a pN0 Number of nodes examined: 23 Number of nodes involved: 0
MARGINS:	pMX Margins uninvolved by invasive carcinoma Margin(s) uninvolved by carcinoma in situ
VENOUS/LYMPHATIC (LARGE/SMALL) VESSEL INVASION (V/L):	Absent
DIRECT EXTENSION OF INVASIVE TUMOR:	None identified
ADDITIONAL PATHOLOGIC FINDINGS:	Urothelial carcinoma in situ Therapy-related changes

ICD-O-3

urothelial, carcinoma, papillary
8130/3

Sites:

path: bladder, NOS

C67.9

CQCF: bladder, wall, lateral

C67.2

9-1-12 RD

Criteria	Yes	No
Diagnosis Discrepancy		
Primary Tumor Site Discrepancy		
HIPAA Discrepancy		
Prior Malignancy History		
Dual/Synchronous Primary Noted		
Case is (circle):	QUALIFIED	DISQUALIFIED
Reviewer Initials	7/24/12	