

SURGICAL PATHOLOGY REPORT

UID:07718E82-778A-4797-8BAD-4494AAE3CAA2
TCGA-XF-AAN8-01A-PR

Redacted



Collected
Ordered by

Location

DIAGNOSIS:

RADICAL CYSTECTOMY WITH ILEAL CONDUIT AND BILATERAL PELVIC LYMPH NODE DISSECTION:

RIGHT DISTAL URETER (A):

FROZEN SECTION DIAGNOSIS:

- NO TUMOR IDENTIFIED

FINAL DIAGNOSIS:

- BENIGN URETER

- NO UROTHELIAL DYSPLASIA OR MALIGNANCY IDENTIFIED

LEFT DISTAL STENTED URETER (B):

FROZEN SECTION DIAGNOSIS:

- HYDROURETER, NO TUMOR IDENTIFIED

FINAL DIAGNOSIS:

- BENIGN URETER WITH HYDROURETER

- NO UROTHELIAL DYSPLASIA OR MALIGNANCY IDENTIFIED

LEFT STENTED URETER (C):

FROZEN SECTION DIAGNOSIS:

- REACTIVE ATYPIA

- NO HIGH GRADE DYSPLASIA IDENTIFIED

FINAL DIAGNOSIS:

- BENIGN URETER WITH REACTIVE UROTHELIAL ATYPIA

- NO UROTHELIAL DYSPLASIA OR MALIGNANCY IDENTIFIED

OLD SCAR PERIVESICAL NODE, BLADDER, URETHRAL MARGIN, VAGINA STRIP (D):

BLADDER:

GROSS DIAGNOSIS:

- TUMOR CLOSELY APPROACHES DEEP RADIAL MARGIN

FINAL DIAGNOSIS:

- UROTHELIAL CARCINOMA, HIGH GRADE, NUCLEAR GRADE 4/4 WITH SQUAMOUS DIFFERENTIATION, INVOLVING POSTERIOR BLADDER WALL AND EXTENDING INTO LEFT URETER-VESICLE JUNCTION AND LEFT URETER

- TUMOR EXTENDS THROUGHOUT THE ENTIRE BLADDER WALL AND DEEPLY INFILTRATES INTO PERIVESICAL SOFT TISSUE

- FOCAL CARCINOMA IN SITU (FLAT LESION) ASSOCIATED WITH INVASIVE TUMOR

- TUMOR CLOSELY COMES WITHIN 1 MM OF THE POSTERIOR SOFT TISSUE RESECTION MARGIN, FINAL SOFT TISSUE MARGIN (SEE PART E) NEGATIVE FOR TUMOR

- ALL OTHER MARGINS, BILATERAL URETERS (2 LEFT AND 2 RIGHT), URETHRAL, AND VAGINAL MUCOSA MARGIN NEGATIVE FOR TUMOR

- EXTENSIVE FOLLICULAR CYSTITIS PRESENT

- SKIN WITH ASSOCIATED SCAR, NO TUMOR IDENTIFIED

RIGHT AND LEFT PERIVESICAL LYMPH NODES (D):

- NO TUMOR IDENTIFIED IN SIX PERIVESICAL LYMPH NODES (0/6)

LEFT PELVIC FLOOR MARGIN (E):

- BENIGN FIBRO CONNECTIVE TISSUE AND MUSCLE

- NO TUMOR IDENTIFIED

RIGHT PARACAVAL LYMPH NODES (F):

- NO MALIGNANCY IDENTIFIED IN FIVE LYMPH NODES EXAMINED (0/5)

RIGHT COMMON ILIAC LYMPH NODES (G):

- NO MALIGNANCY IDENTIFIED IN THREE LYMPH NODES EXAMINED (0/3)

RIGHT LYMPH NODE OF CLOQUET (H):

- BENIGN FIBRO CONNECTIVE TISSUE AND ADIPOSE TISSUE

- NO LYMPHOID TISSUE IDENTIFIED (0/0)

ICD O-3

Carcinoma, urothelial NOS 8120/3

Site: Bladder, posterior wall
C67.4

JW 3/26/14

20% squamous, per TSS.

SURGICAL PATHOLOGY REPORT

Collected
Ordered by

Location

LEFT PARA AORTIC LYMPH NODES (I):

- BENIGN FIBRO CONNECTIVE TISSUE AND ADIPOSE TISSUE
- NO LYMPHOID TISSUE IDENTIFIED (0/0)

LEFT COMMON ILIAC LYMPH NODES (J):

- NO MALIGNANCY IDENTIFIED IN TEN LYMPH NODES EXAMINED (0/10)

RIGHT OBTURATOR/HYPOGASTRIC LYMPH NODES (K):

- NO MALIGNANCY IDENTIFIED IN 16 LYMPH NODES EXAMINED (0/16)

RIGHT EXTERNAL ILIAC LYMPH NODES (L):

- NO MALIGNANCY IDENTIFIED IN THREE LYMPH NODES EXAMINED (0/3)

PRE SACRAL LYMPH NODES (M):

- NO MALIGNANCY IDENTIFIED IN TWO LYMPH NODES EXAMINED (0/2)

LEFT LYMPH NODE OF CLOQUET (N):

- NO MALIGNANCY IDENTIFIED IN FIVE LYMPH NODES EXAMINED (0/5)

LEFT OBTURATOR/HYPOGASTRIC LYMPH NODES (O):

- NO MALIGNANCY IDENTIFIED IN TWO LYMPH NODES EXAMINED (0/2)

LEFT EXTERNAL ILIAC LYMPH NODES (P):

- NO MALIGNANCY IDENTIFIED IN ONE LYMPH NODES EXAMINED (0/1)

LEFT PROXIMAL URETER (Q):

- HIGH GRADE URETER AND BENIGN URETER
- NO UROTHELIAL DYSPLASIA OR MALIGNANCY IDENTIFIED

RIGHT PROXIMAL URETER (R):

- BENIGN URETER
- NO UROTHELIAL DYSPLASIA OR MALIGNANCY IDENTIFIED

LEFT PRE SCIATIC LYMPH NODES (S):

- BENIGN FIBRO CONNECTIVE AND ADIPOSE TISSUE
- NO LYMPHOID TISSUE IDENTIFIED (0/0)

RIGHT PRE SCIATIC LYMPH NODES (T):

- NO MALIGNANCY IDENTIFIED IN ONE LYMPH NODES EXAMINED (0/1)

LEFT URETER STENT [GROSS ONLY] (U):

- MEDICAL DEVICE, URETER STENT

LYMPH NODE SUMMARY: NO METASTATIC CARCINOMA IDENTIFIED IN TOTAL OF
54 LYMPH NODES EXAMINED (0/54)

BLADDER TNM STAGE: pt3bn0MX

Electronically signed by
Verified:

SPECIMEN SOURCE:

- A: "Rt distal ureter FS"
- B: "Lt distal stented ureter"
- C: "Lt unstented ureter"
- D: "Old scar perivesicle node, bladder, urethral margin, vagina strip"

SURGICAL PATHOLOGY REPORT

Collected

Ordered by

Location

E: "Left pelvic floor margin"
 F: "Right para caval lymph nodes"
 G: "Right common iliac lymph nodes"
 H: "Right lymph node of cloquet"
 I: "Left para aortic lymph nodes"
 J: "Left common iliac lymph node"
 K: "Right obturator/hypogastric lymph node"
 L: "Right external iliac lymph node"
 M: "Pre sacral lymph nodes"
 N: "Left lymph node of cloquet"
 O: "Left obturator/hypogastric lymph nodes"
 P: "Left external iliac lymph nodes"
 Q: "Left proximal ureter"
 R: "Right proximal ureter"
 S: "Left pre sciatic lymph nodes"
 T: "Right pre sciatic lymph nodes"
 U: "Left ureter stent"

CLINICAL INFORMATION:

Pre-op Dx: Bladder cancer

Post-op Dx: Same

GROSS EXAMINATION:

A: The specimen is received fresh from the O.R. and labeled "Rt distal ureter FS". It consists of two segments of ureter measuring in aggregate 1.5 x 0.5 x 0.3 cm. The specimen is entirely submitted for frozen section diagnosis in AFS.

B: The specimen is received fresh from the O.R. and labeled "Lt distal stented ureter". It consists of a segment of pink-tan ureter measuring 1 x 0.8 x 0.3 cm. The specimen is entirely submitted for frozen section diagnosis in BFS.

C: The specimen is received fresh from the O.R. and labeled "Lt unstented ureter". It consists of a segment of pink ureter measuring 0.6 x 0.3 x 0.2 cm. The specimen is entirely submitted for frozen section diagnosis in CFS.

D: The specimen is received fresh from the O.R. and labeled "Old scar perivesicle node, bladder, urethral margin, vaginal strip". It consists of a radical cystectomy specimen measuring overall 12 x 15 x 5 cm. A flap of peritoneum is present on the posterior surface measuring 23 x 9 x 1 cm. The peritoneum is smooth, mobile and glistening. The bladder itself measures 9 x 4 x 3 cm. There is an area marked by silk on the left wall. Opening the bladder reveals a large exophytic mass at the left ureterovesical junction measuring 3 x 2.5 x 4 cm in depth. Both ureterovesical junctions are identified and are patent. The attached ureters are identified and are opened. There are two ureters on either side. The right attached ureter (A) measures 5 cm in length x 0.6 cm in circumference and second right ureter (B) measures 4.5 cm in length x 0.6 cm in circumference. The left ureter (A) measures 6 cm in length x 1.2 cm in circumference and the second left ureter (B) measures 6 cm in length x 1.5 cm in circumference. The urethra measures 2.5 cm in length x 2.5 cm in circumference. The vaginal wall measures 4 x 3 x 0.2 cm. There is a skin ellipse measuring 15 x 1 x 0.3 cm. On the surface of the skin, there is a healed surgical scar measuring 10 cm in length. Representative sections are submitted in 29 cassettes.

E: The specimen is received in formalin and labeled "Left pelvic floor margin". It consists of segment of yellow lobulated fatty tissue measuring 3 x 2.4 x 0.2 cm. The specimen is entirely submitted in three cassettes.

F: The specimen is received in formalin and labeled "Right para caval lymph nodes". It consists of a segment of pale yellow lobulated fatty tissue measuring 2.9 x 2 x 0.2 cm. The specimen is entirely submitted in one cassette.

Collected
Ordered by

G: The specimen is received in formalin and labeled "Right common iliac lymph nodes". It consists of multiple pieces of pale yellow lobulated fatty tissue measuring in aggregate 2 x 1.7 x 0.6 cm. The specimen is entirely submitted in one cassette.

H: The specimen is received in formalin and labeled "Right lymph node of cloquet". It consists of multiple pieces of pale yellow lobulated fatty tissue measuring in aggregate 2.5 x 1.9 x 0.3 cm. The specimen is entirely submitted in two cassettes.

I: The specimen is received in formalin and labeled "Left para aortic lymph nodes". It consists of multiple pieces of pale yellow lobulated fatty tissue measuring in aggregate 1.5 x 0.9 x 0.2 cm. The specimen is entirely submitted in one cassette.

J: The specimen is received in formalin and labeled "Left common iliac lymph node". It consists of multiple pieces of pale yellow lobulated fatty tissue measuring in aggregate 2.5 x 1.6 x 1 cm. The specimen is entirely submitted in three cassettes.

K: The specimen is received in formalin and labeled "Right obturator/hypogastric lymph node". It consists of multiple pieces of pale yellow lobulated fatty tissue measuring in aggregate 6.5 x 4 x 1 cm. The specimen is entirely submitted in five cassettes.

L: The specimen is received in formalin and labeled "Right external iliac lymph node". It consists of multiple pieces of pale yellow lobulated fatty tissue measuring in aggregate 4.5 x 3 x 1 cm. The specimen is entirely submitted in one cassette.

M: The specimen is received in formalin and labeled "Pre sacral lymph nodes". It consists of a segment of yellow lobulated fatty tissue measuring 4.5 x 3 x 0.2 cm. The specimen is entirely submitted in three cassettes.

N: The specimen is received in formalin and labeled "Left lymph node of cloquet". It consists of multiple pieces of pale yellow lobulated fatty tissue measuring in aggregate 2.6 x 2.5 x 0.5 cm. The specimen is entirely submitted in two cassettes.

O: The specimen is received in formalin and labeled "Left obturator/hypogastric lymph nodes". It consists of multiple pieces of pale yellow lobulated fatty tissue measuring in aggregate 2.5 x 2 x 0.2 cm. The specimen is entirely submitted in two cassettes.

P: The specimen is received in formalin and labeled "Left external iliac lymph nodes". It consists of multiple pieces of pale yellow lobulated fatty tissue measuring in aggregate 3 x 3 x 0.5 cm. The specimen is entirely submitted in three cassettes.

Q: The specimen is received in formalin and labeled "Left proximal ureter". It consists of two segments of ureter measuring 0.7 cm in diameter x 0.7 cm in length and 0.5 cm in diameter x 0.5 cm in length. The largest segment of ureter has a black suture around it. The larger segment is inked black. Specimen is entirely submitted in two cassettes.

R: The specimen is received in formalin and labeled "Right proximal ureter". It consists of two segments of ureter, each measuring 0.2 cm in diameter x 0.5 cm in length. The specimen is entirely submitted in one cassette.

S: The specimen is received in formalin and labeled "Left pre sciatic lymph nodes". It consists of multiple pieces of pale yellow lobulated fatty tissue

SURGICAL PATHOLOGY REPORT

Collected
Ordered by

Location

measuring in aggregate 2.5 x 1.5 x 0.3 cm. The specimen is entirely submitted in one cassette.

T: The specimen is received in formalin and labeled "Right pre sciatic lymph nodes". It consists of multiple pieces of pale yellow lobulated fatty tissue measuring in aggregate 2 x 2.2 x 0.5 cm. The specimen is entirely submitted in two cassettes.

U: The specimen is received in a plastic container and labeled "Left ureter stent". It consists of a plastic tubing measuring 35 cm in length with an average diameter of 0.2 cm. Gross description only.

SECTIONS:

AFS: frozen section, right distal ureter
 BFS: frozen section, left distal stented ureter
 CFS: frozen section, left unstented ureter
 D1: urethral margin
 D2: right vaginal margin
 D3: distal vaginal margin
 D4: left vaginal margin
 D5: right proximal ureter margin of specimen
 D6: left proximal ureter margin of specimen
 D7: right ureterovesical junction (A)
 D8: right ureterovesical junction (B)
 D9: cross section of ureters (A and B)
 D10: left ureterovesical junction with tumor involvement (left A)
 D11: left ureter (B) junction opening into left ureter (A)
 D12: mid section of left ureter
 D13: cross section of proximal left ureter
 D14-17: deep margin of tumor
 D18: trigone
 D19: bladder neck, proximal urethra
 D20: posterior wall of bladder
 D21: dome of bladder
 D22: surface of tumor with bladder junction
 D23, 24: right perivesical fat for probable lymph nodes
 D25-28: left perivesical fat for probable lymph nodes
 D29: representative of skin
 E1-3: left pelvic floor margin
 F: right para caval lymph nodes
 G: right common iliac lymph nodes
 H1-2: right lymph node of Cloquet
 I: left para aortic lymph nodes
 J1-3: left common iliac lymph node
 K1-5: right obturator/hypogastric lymph node
 L: right external iliac lymph node
 M1-3: pre sacral lymph nodes
 N1-2: left lymph node of Cloquet
 O1-2: left obturator/hypogastric lymph nodes
 P1-3: left external iliac lymph nodes
 Q1: left proximal ureter; larger segment
 Q2: smaller segment
 R: right proximal ureter
 S: left pre sciatic lymph nodes
 T: right pre sciatic lymph nodes

SURGICAL PATHOLOGY REPORT

Collected

Ordered by

Location

INTRAOPERATIVE FROZEN CONSULTATION:

- AFS: Right distal ureter
 - no tumor identified
- BFS: Left distal stented ureter
 - hydroureter
 - no tumor identified
- CFS: Left unstented ureter
 - reactive atypia
 - no high-grade dysplasia identified

INTRAOPERATIVE GROSS CONSULTATION

- Exophytic mass at left uretero-vesical junction. Dual ureters entering bladder on both sides.

MICROSCOPIC EXAMINATION:

A-U: See final microscopic-diagnosis.

Criteria	Yes	No
Diagnosis Discrepancy		✓
Primary Tumor Site Discrepancy		✓
HIPAA Discrepancy		✓
Prior Malignancy History	✓	✓
Dual/Synchronous Primary Noted		✓
Case is (circle):	QUALIFIED	DISQUALIFIED
Reviewer Initial	Date Reviewed: 1/2/14	