

Redacted

Pathology Review

ICD-0-3

Carcinoma, urothelial, NOS 8120/3

Site: bladder, anterior wall C67.3

W
6/1/11

DIAGNOSIS

(A) LEFT VAS:

Segment of vas deferens, no tumor present.

(B) LEFT URETER:

Segment of ureter with mild reactive atypia, no tumor present.

(C) RIGHT VAS:

Segment of vas deferens, no tumor present.

(D) RIGHT URETER:

Segment of ureter, no tumor present.

(E) BLADDER, PROSTATE AND SEMINAL VESICLES:

Bladder: UROTHELIAL CARCINOMA, NONPAPILLARY TYPE, GRADE 3.

TRANSMURAL INVASION OF BLADDER WALL WITH FOCAL
EXTENSION INTO PERIVESICAL ADIPOSE TISSUE.

UROTHELIAL CARCINOMA IN SITU IN A BACKGROUND OF
DIFFUSE LOW GRADE INTRAUROTHELIAL NEOPLASIA.

Serosal and resection margins, free of tumor.

Prostate: Prostate with microcystic adenomatous change. No evidence of
malignancy.

(F) RIGHT COMMON ILIAC LYMPH NODES:

Fibroadipose tissue, no lymph node or tumor present.

(G) RIGHT EXTERNAL ILIAC LYMPH NODE:

Four lymph nodes, no tumor present (0/4).

(H) RIGHT OBTURATOR LYMPH NODE:

Seven lymph nodes, no tumor present (0/7).

(I) LEFT COMMON ILIAC:

Five lymph nodes, no tumor present (0/5).

(J) LEFT EXTERNAL ILIAC:

Two lymph nodes, no tumor present (0/2).

(K) LEFT OBTURATOR:

Ten lymph nodes, no tumor present (0/10).

(L) LEFT URETER, FINAL MARGIN:

Wall of ureter, no tumor present.

(M) RIGHT URETER, FINAL MARGIN:

Wall of ureter, no tumor present.

GROSS DESCRIPTION

Criteria	Yes	No
Diagnosis Discrepancy		
Primary Tumor Site Discrepancy		
HIPAA Discrepancy		
Prior Malignancy History		
Dual/Synchronous Primary noted		
Case is (circle):	QUALIFIED	DISQUALIFIED
Reviewer Initials	MG	
Date Reviewed	5/21/11	

(A) LEFT VAS – A white-tan portion of vas deferens measuring 5 cm in length x 0.3 cm in diameter. A lumen is present on the cut surfaces. Specimen is representatively submitted in A.

(B) LEFT URETER – A segment of ureter 0.8 cm long and 0.1 cm in diameter with surrounding fibroadipose tissue. The resection margin is inked by the surgeon and entire specimen is submitted for frozen in block B (ink side down, en face).

*FS/DX: BENIGN URETER.

(C) RIGHT VAS – A segment white-tan vas deferens measuring 5.5 cm in length x 0.3 cm in diameter. A pinpoint lumen is visible in the cut surfaces. Specimen is representatively submitted in C.

(D) RIGHT URETER – A segment of ureter (1.0 x 0.7 cm). The margin, designated by the surgeon with ink, is reinked blue and submitted en face for frozen section in D.

*FS/DX: BENIGN URETER.

(E) BLADDER, PROSTATE AND SEMINAL VESICLES – A cystoprostatectomy measuring overall (14 x 7.5 x 7 cm). The specimen is opened longitudinally through the superior wall to reveal a fungating tumor (9 x 7 x 2.5 cm) involving the near entire anterior bladder wall and dome. The tumor appears to invade the muscular wall of the bladder but does not extend into the perivesical adipose tissue.

The remaining bladder mucosa shows hemorrhage, especially in the trigone, and is other unremarkable.

INK CODE: Black – outer surface of anterior wall.

SECTION CODE: E1-E48, bladder with tumor; E49-E57, prostate; E58, E59, urethra, en face.

(F) RIGHT COMMON ILIAC LYMPH NODES – A single red tan piece of tissue measuring 2.4 x 1.5 x 1 cm. Three potential lymph nodes are palpated, ranging in size from 0.4 to 0.6 cm in greatest dimension. No tumor is seen grossly. Specimen is entirely submitted.

SECTION CODE: F1, three potential lymph nodes; F2, remainder of specimen.

(G) RIGHT EXTERNAL ILIAC LYMPH NODE – Multiple red tan pieces of adipose tissue measuring 4.9 x 4 x 1 cm in aggregate. Five possible lymph nodes are found, ranging in size from 0.5 to 3.7 cm in greatest dimension. No tumor is seen grossly. Specimen is entirely submitted.

SECTION CODE: G1, three potential lymph nodes; G2, G3, single lymph node; G4, G5, single lymph node.

(H) RIGHT OBTURATOR LYMPH NODE – Multiple red tan pieces of adipose tissue measuring 5 x 2.5 x 2 cm in aggregate. Seven potential lymph nodes are found, ranging in size from 0.4 to 2.6 cm in greatest dimensions. No tumor is seen grossly.

SECTION CODE: H1, four potential lymph nodes; H2, one potential lymph node; H3, one potential lymph node; H4, H5, single lymph node, sectioned.

(I) LEFT COMMON ILIAC – Received is a 3.5 x 3.0 x 1.0 cm portion of fibroadipose tissue, which yields multiple possible lymph nodes, ranging from 0.6 cm to 1.5 x 0.8 x 0.6 cm. The lymph nodes are submitted entirely.

SECTION CODE: I1, three lymph nodes; I2, two lymph nodes.

(J) LEFT EXTERNAL ILIAC – Received is a 5.0 x 4.2 x 1.0 cm portion of fibroadipose tissue which yields two possible lymph nodes, measuring 2.0 x 0.8 x 0.5 cm and 3.0 x 1.5 x 0.5 cm. Both lymph nodes are submitted entirely.

SECTION CODE: J1, one lymph node; J2, J3, one lymph node, serially sectioned.

(K) LEFT OBTURATOR - Multiple fibroadipose tissue fragments (6.5 x 4.0 x 1.0 cm). Eleven possible lymph nodes, ranging from 0.5 to 2.5 cm in greatest dimension are identified and are submitted in toto.

SECTION CODE: K1, four lymph nodes; K2, three lymph nodes; K3, one lymph node, bisected; K4, one lymph node, bisected; K5, one lymph node, bisected; K6, one lymph node, bisected.

(L) LEFT URETER FINAL MARGIN – A segment of ureter (1.7 x 0.6 cm). The true margin, designated by the surgeon with ink, is reinked blue and submitted en face in L.

(M) RIGHT URETER FINAL MARGIN – A segment of ureter (0.8 x 0.8 cm). The margin, designated by the surgeon with ink, is reinked blue and submitted en face in M.

CLINICAL HISTORY

None given.