

Pathology Report:

Surgical Pathology Report

FINAL PATHOLOGIC DIAGNOSIS

A. Right pelvic lymph nodes; lymph node dissection:

-Three of eleven lymph nodes positive for metastatic urothelial carcinoma (3/11).

B. Left pelvic lymph nodes; lymph node dissection:

-Five of ten lymph nodes positive for metastatic urothelial carcinoma with extranodal extension (5/10).

C. Bladder, prostate; cystoprostatectomy:

Bladder: High grade invasive urothelial carcinoma, see pathologic parameter.

- Diverticulum

Prostate: Prostatic urethral ulcer and squamous metaplasia.

-High grade prostate intraepithelial neoplasia.

-Prostatic abscess.

-No involvement by urothelial carcinoma.

D. Distal right ureter; excision:

-Portion of ureter, no tumor.

Urothelial Carcinoma Pathologic Parameters

1. Tumor type: Invasive urothelial carcinoma.

2. Grade of tumor:

High grade

3. Depth of invasion:

Extra vesicular soft tissue or adjacent organ (specify)

4. Tumor distribution:

Solitary: 5 cm Size

Left lateral wall

Trigone

Posterior wall

5. Ureteral margins:

Negative

6. Distal urethral margin:

Negative

7. Soft tissue margin or serosa:

Negative

ICD-O-3
Carcinoma, urothelial NOS
8/20/13

Site: ^{path} Overlapping lesion of
bladder (trigone) posterior wall)
C67.8

~~C67.8~~ Trigone of bladder
C67.0
Jtu 4/16/13

UUID:41E61CCB-5006-4918-9487-BE81F8559D0E
TCGA-FD-A620-01A-PR

Redacted



8. Lymph nodes: Positive in 3 of 11 right and 5 of 10 left pelvic nodes (8/21)

9. pTNM: pT3a, N2, MX

Effective January 1, 2010 this Checklist utilizes the 7th edition TNM staging system for Bladder of the American Joint Committee on Cancer (AJCC) and the International Union Against Cancer (UICC).

Interpretation performed by the Attending Pathologist and reviewed with the

Clinical History:

The patient is a year-old male with high grade urothelial carcinoma undergoing cystoprostatectomy.

Specimens Received:

A: Right pelvic lymph nodes

B: Left pelvic lymph nodes

C: Bladder, prostate, vas deferens

D: Distal right ureter

Gross Description:

The specimens are received in four containers each labeled with the patient's name and medical record number.

A. The first container is additionally identified as, "1. Right pelvic lymph nodes". Received fresh and placed in formalin is a 6 x 5.5 x 1.5 cm aggregate of lobulated yellow fibrofatty tissue. The specimen is dissected for lymph nodes to reveal 15 apparent lymph nodes ranging from 0.5-4.5 cm in greatest dimension. All apparent lymph nodes are submitted as follows:

A1: 5 candidate lymph nodes

A2: 5 candidate lymph nodes A3: 3 candidate lymph nodes

A4-A5: One lymph node, serially sectioned

A6: One lymph node, serially sectioned

B. The second container is additionally identified as, "2. Left pelvic lymph nodes". Received fresh and placed in formalin is a 7.5 x 5 x 2 cm aggregate of lobulated yellow fibrofatty tissue. The specimen is dissected for lymph nodes to reveal 14 apparent lymph nodes ranging from 0.5-5.5 cm in greatest dimension. All apparent lymph nodes are submitted as follows:

B1: 6 candidate lymph nodes

B2: 3 candidate lymph nodes

B3: One candidate lymph node, bisected
B4: One candidate lymph node, bisected
B5: One candidate lymph node, bisected
B6: One candidate lymph node, bisected
B7-B9: One candidate lymph node, serially sectioned

C. The third container is additionally identified as, "3. Bladder, prostate, vas deferens". Received fresh and placed in formalin is a 334.3 g, 15 x 10.5 x 4.5 cm cystoprostatectomy specimen consisting of a 7.5 x 6.5 x 3.5 cm bladder with attached mesenteric fat and a 5 x 4 x 3.2 cm prostate. Also received is a 3.0 x 3.3 x 0.7 cm aggregate of yellow, lobulated adipose tissue and rubbery, tan, tubelike structures. The right seminal vesicles measure 3.5 x 2.2 x 0.7 cm and right vas deferens measures 8.5 x 0.6 cm. The left seminal vesicles measures 3.5 x 1.3 x 0.8 cm and left vas deferens measures 8 x 0.6 cm. The right ureteral stump measures 1.8 x 0.6 cm, the left measures 2 x 0.6 cm, and each demonstrates an intact lumen which is probe patent to the trigone.

The right half of the prostate is inked blue and the left half is inked black. The outside of the bladder is inked black. The bladder and prostate are opened anteriorly along the urethra. The outer surface of the prostate is shaggy and pink. Sectioning demonstrates rubbery, pink-tan, peri-urethral nodularity with a 1 x 0.5 x 0.5 cm area of tan-brown nodularity at the urethra proximally, which appears continuous with the bladder lesion.

The opened bladder reveals a 5 x 3.5 cm tan, plaque-like indurated, lobulated, ill-defined mass involving the posterior wall and entire trigone with extension into the prostatic urethra. Also present is a 3 x 2.5 x 1 cm smooth-walled cavity containing hemorrhagic fluid which is located subjacent to the right posterior-lateral wall. On cut section, the mass extends through the muscularis propria into the perivesical adipose tissue and is 0.1 cm from the deep margin. The mass also extends into the wall of the cystic lesion and the cystic lesion extends to 0.2 cm from the deep margin. The surrounding bladder mucosa is edematous and tan with a 1-2 cm wall thickness. Photographs are taken.

Representative sections are submitted as follows:

C1: Left ureter resection margin
C2: Right ureter resection margin
C3: Distal prostatic urethral margin
C4: Prostate apex
C5-C8: Representative right prostate from apex to base (nodular portion of urethra C7-C8)
C9-C12: Representative left prostate from apex to base (nodular portion of urethra C11-C12)
C13: Representative right vas deferens and seminal vesicle
C14: Representative left vas deferens and seminal vesicle
C15: Mass at deepest extent of invasion

C16: Mass extending into wall of cystic lesion with deepest extent of cystic lesion

C17: Mass and cystic lesion

C18: Additional sections of mass and adjacent mucosa

C19: Grossly uninvolved posterior bladder

C20: Grossly uninvolved right wall

C21: Grossly uninvolved left wall

C22: Grossly uninvolved bladder dome

C23: Grossly uninvolved anterior bladder

D. The fourth container is additionally identified as, "4. Distal right ureter". Received fresh and placed in formalin is a 4.5 x 1.5 x 0.7 cm tan-yellow, rubbery, tubelike structure. The specimen is oriented with a suture on the non-margin end. The margin end is shaved and submitted as D1.

Criteria	Yes	No
Diagnosis Discrepancy		✓
Primary Tumor Site Discrepancy		✓
HIPAA Discrepancy		✓
Prior Malignancy History		✓
Dual/Synchronous Primary Mx		✓
Cases (circle):	QUALIFIED	DISQUALIFIED
Reviewer initials	4/5/13	4/12/13