

Lyapunov Stability

Contents

- Motivation
- Lyapunov's Direct Method for Autonomous System
- The Invariance Principle
- LTI System and Exponential Stable

Recall the pendulum equation

$$\dot{x}_1 = x_2$$

$$\dot{x}_2 = -a \sin x_1 - bx_2$$

has two equilibrium points at

$$x_{e1} = (0, 0)^T$$

is (asymptotically) stable, and

$$x_{e2} = (\pi, 0)^T$$

is unstable.

The conclusion relies on explicit knowledge of solution or physical meaning of the system, which in general is NOT easy to obtain!

Recall the pendulum equation

$$\dot{x}_1 = x_2$$

$$\dot{x}_2 = -a \sin x_1 - bx_2$$

has two equilibrium points at

$$x_{e1} = (0, 0)^{\top}$$

is (asymptotically) stable, and

$$x_{e2} = (\pi, 0)^{\top}$$

is unstable.

The conclusion relies on explicit knowledge of solution or physical meaning of the system, which in general is NOT easy to obtain!

Motivation

However, the conclusion can also be reached by using the energy concepts. Define the energy of the pendulum as

$$\begin{aligned} E(x) &= \overbrace{mgl(1 - \cos x_1)}^{\text{potential energy}} + \overbrace{\frac{1}{2}m(l\dot{x}_2)^2}^{\text{kinetic energy}} \\ &= ml^2[a(1 - \cos x_1) + \frac{1}{2}\dot{x}_2^2] \end{aligned}$$

with the reference of the potential energy is chosen as $E(0) = 0$,
i.e. x_{e1} has no energy.

The derivative of the energy along the trajectory of the system

$$\frac{dE}{dt} = -ml^2 b \dot{x}_2^2 \leq 0$$

Motivation

However, the conclusion can also be reached by using the energy concepts. Define the energy of the pendulum as

$$\begin{aligned} E(x) &= \overbrace{mgl(1 - \cos x_1)}^{\text{potential energy}} + \overbrace{\frac{1}{2}m(\dot{x}_2)^2}^{\text{kinetic energy}} \\ &= ml^2[a(1 - \cos x_1) + \frac{1}{2}\dot{x}_2^2] \end{aligned}$$

with the reference of the potential energy is chosen as $E(0) = 0$,
i.e. x_{e1} has no energy.

The derivative of the energy along the trajectory of the system

$$\frac{dE}{dt} = -ml^2 b \dot{x}_2^2 \leq 0$$

Motivation

However, the conclusion can also be reached by using the energy concepts. Define the energy of the pendulum as

$$\begin{aligned} E(x) &= \overbrace{mgl(1 - \cos x_1)}^{\text{potential energy}} + \overbrace{\frac{1}{2}m(l\dot{x}_2)^2}^{\text{kinetic energy}} \\ &= ml^2[a(1 - \cos x_1) + \frac{1}{2}\dot{x}_2^2] \end{aligned}$$

with the reference of the potential energy is chosen as $E(0) = 0$,
i.e. x_{e1} has no energy.

The derivative of the energy along the trajectory of the system

$$\frac{dE}{dt} = -ml^2 b \dot{x}_2^2 \leq 0$$

Motivation

The derivative of the energy along the trajectory of the system

$$\dot{E} = -ml^2bx_2^2 \leq 0$$

- if $b = 0$, $E = \text{constant} \iff$ Stable equilibrium
- if $b \neq 0$, E keeps decreasing until it eventually reaches zero
 \iff Asymptotically stable equilibrium

In 1982, Lyapunov showed that certain energy-like function could be used to determine the stability of an equilibrium.

Motivation

The derivative of the energy along the trajectory of the system

$$\dot{E} = -ml^2bx_2^2 \leq 0$$

- if $b = 0$, $E = \text{constant} \iff$ Stable equilibrium
- if $b \neq 0$, E keeps decreasing until it eventually reaches zero
 \iff Asymptotically stable equilibrium

In 1982, Lyapunov showed that certain energy-like function could be used to determine the stability of an equilibrium.

Motivation

The derivative of the energy along the trajectory of the system

$$\dot{E} = -ml^2bx_2^2 \leq 0$$

- if $b = 0$, $E = \text{constant} \iff$ Stable equilibrium
- if $b \neq 0$, E keeps decreasing until it eventually reaches zero
 \iff Asymptotically stable equilibrium

In 1982, Lyapunov showed that certain energy-like function could be used to determine the stability of an equilibrium.

Contents

- Motivation
- Lyapunov's Direct Method for Autonomous System
- The Invariance Principle
- LTI System and Exponential Stable

Autonomous system

Recall: Consider the autonomous system

$$\dot{x} = f(x), \quad (1)$$

where $f(x) : \mathcal{D} \rightarrow \mathbb{R}^n$ is locally Lipschitz map from a domain $\mathcal{D} \subset \mathbb{R}^n$. Without loss of generality, we assume $x_e = \{0\} \in \mathcal{D}$.

Let function $V : \mathcal{D} \rightarrow \mathbb{R}$ be a \mathcal{C}^1 (continuous differentiable) function on x . $\dot{V}(x)$ is defined as

$$\dot{V}(x) = \frac{\partial V}{\partial x} \dot{x} = \left[\frac{\partial V}{\partial x_1}, \quad \frac{\partial V}{\partial x_2}, \quad \cdots, \quad \frac{\partial V}{\partial x_n} \right] \begin{bmatrix} f_1(x) \\ f_2(x) \\ \vdots \\ f_n(x) \end{bmatrix}$$

Autonomous system

Recall: Consider the autonomous system

$$\dot{x} = f(x), \quad (1)$$

where $f(x) : \mathcal{D} \rightarrow \mathbb{R}^n$ is locally Lipschitz map from a domain $\mathcal{D} \subset \mathbb{R}^n$. Without loss of generality, we assume $x_e = \{0\} \in \mathcal{D}$.

Let function $V : \mathcal{D} \rightarrow \mathbb{R}$ be a \mathcal{C}^1 (continuous differentiable) function on x . $\dot{V}(x)$ is defined as

$$\dot{V}(x) = \frac{\partial V}{\partial x} \dot{x} = \left[\frac{\partial V}{\partial x_1}, \quad \frac{\partial V}{\partial x_2}, \quad \cdots, \quad \frac{\partial V}{\partial x_n} \right] \begin{bmatrix} f_1(x) \\ f_2(x) \\ \vdots \\ f_n(x) \end{bmatrix}$$

Autonomous system

Recall: Consider the autonomous system

$$\dot{x} = f(x), \tag{1}$$

where $f(x) : \mathcal{D} \rightarrow \mathbb{R}^n$ is locally Lipschitz map from a domain $\mathcal{D} \subset \mathbb{R}^n$. Without loss of generality, we assume $x_e = \{0\} \in \mathcal{D}$.

Let function $V : \mathcal{D} \rightarrow \mathbb{R}$ be a \mathcal{C}^1 (continuous differentiable) function on x . $\dot{V}(x)$ is defined as

$$\dot{V}(x) = \frac{\partial V}{\partial x} \dot{x} = \left[\frac{\partial V}{\partial x_1}, \quad \frac{\partial V}{\partial x_2}, \quad \cdots, \quad \frac{\partial V}{\partial x_n} \right] \begin{bmatrix} f_1(x) \\ f_2(x) \\ \vdots \\ f_n(x) \end{bmatrix}$$

Lyapunov Stability Theorem

Theorem: For the autonomous system defined in (1), let

$V : \mathcal{D} \rightarrow \mathbb{R}$ be a \mathcal{C}^1 function such that

- i) $V(0) = 0$
- ii) $V(x) > 0$ for any $x \in \mathcal{D} - \{0\}$
- iii) $\dot{V} \leq 0$ for any $x \in \mathcal{D}$.

Then, the equilibrium at the origin is *stable* in the sense of Lyapunov. Moreover, origin is *asymptotically stable*, if

$$\dot{V} < 0$$

for any $x \in \mathcal{D} - \{0\}$.

Proof can be found in Khalil's *nonlinear systems* Theorem 4.1.

Lyapunov Stability Theorem

Theorem: For the autonomous system defined in (1), let

$V : \mathcal{D} \rightarrow \mathbb{R}$ be a \mathcal{C}^1 function such that

- i) $V(0) = 0$
- ii) $V(x) > 0$ for any $x \in \mathcal{D} - \{0\}$
- iii) $\dot{V} \leq 0$ for any $x \in \mathcal{D}$.

Then, the equilibrium at the origin is *stable* in the sense of Lyapunov. Moreover, origin is *asymptotically stable*, if

$$\dot{V} < 0$$

for any $x \in \mathcal{D} - \{0\}$.

Proof can be found in Khalil's *nonlinear systems* Theorem 4.1.

Lyapunov Stability Theorem

Theorem: For the autonomous system defined in (1), let

$V : \mathcal{D} \rightarrow \mathbb{R}$ be a \mathcal{C}^1 function such that

- i) $V(0) = 0$
- ii) $V(x) > 0$ for any $x \in \mathcal{D} - \{0\}$
- iii) $\dot{V} \leq 0$ for any $x \in \mathcal{D}$.

Then, the equilibrium at the origin is *stable* in the sense of Lyapunov. Moreover, origin is *asymptotically stable*, if

$$\dot{V} < 0$$

for any $x \in \mathcal{D} - \{0\}$.

Proof can be found in Khalil's *nonlinear systems* Theorem 4.1.

Lyapunov Stability Theorem

Theorem: For the autonomous system defined in (1), let

$V : \mathcal{D} \rightarrow \mathbb{R}$ be a \mathcal{C}^1 function such that

- i) $V(0) = 0$
- ii) $V(x) > 0$ for any $x \in \mathcal{D} - \{0\}$
- iii) $\dot{V} \leq 0$ for any $x \in \mathcal{D}$.

Then, the equilibrium at the origin is *stable* in the sense of Lyapunov. Moreover, origin is *asymptotically stable*, if

$$\dot{V} < 0$$

for any $x \in \mathcal{D} - \{0\}$.

Proof can be found in Khalil's *nonlinear systems* Theorem 4.1.

Example: Consider a first-order system

$$\dot{x} = -g(x)$$

where $g(x)$ is locally Lipschitz on $(-a, a)$ and satisfies

$$g(0) = 0, xg(x) > 0, \forall x \in (-a, 0) \cup (0, a)$$

The system has a isolated equilibrium at $x = 0$. Explicit solution is impossible to obtain. Consider a Lyapunov function

$$V(x) = x^2 \rightarrow \dot{V} = -2xg(x)$$

or

$$V(x) = \int_0^x g(y)dy \rightarrow \dot{V} = -g^2(x)$$

Note: The choice of Lyapunov function is not unique.

Example: Consider a first-order system

$$\dot{x} = -g(x)$$

where $g(x)$ is locally Lipschitz on $(-a, a)$ and satisfies

$$g(0) = 0, xg(x) > 0, \forall x \in (-a, 0) \cup (0, a)$$

The system has a isolated equilibrium at $x = 0$. Explicit solution is impossible to obtain. Consider a Lyapunov function

$$V(x) = x^2 \rightarrow \dot{V} = -2xg(x)$$

or

$$V(x) = \int_0^x g(y)dy \rightarrow \dot{V} = -g^2(x)$$

Note: The choice of Lyapunov function is not unique.

Example: Consider a first-order system

$$\dot{x} = -g(x)$$

where $g(x)$ is locally Lipschitz on $(-a, a)$ and satisfies

$$g(0) = 0, xg(x) > 0, \forall x \in (-a, 0) \cup (0, a)$$

The system has a isolated equilibrium at $x = 0$. Explicit solution is impossible to obtain. Consider a Lyapunov function

$$V(x) = x^2 \rightarrow \dot{V} = -2xg(x)$$

or

$$V(x) = \int_0^x g(y)dy \rightarrow \dot{V} = -g^2(x)$$

Note: The choice of Lyapunov function is not unique.

Example: Consider a first-order system

$$\dot{x} = -g(x)$$

where $g(x)$ is locally Lipschitz on $(-a, a)$ and satisfies

$$g(0) = 0, xg(x) > 0, \forall x \in (-a, 0) \cup (0, a)$$

The system has a isolated equilibrium at $x = 0$. Explicit solution is impossible to obtain. Consider a Lyapunov function

$$V(x) = x^2 \rightarrow \dot{V} = -2xg(x)$$

or

$$V(x) = \int_0^x g(y)dy \rightarrow \dot{V} = -g^2(x)$$

Note: The choice of Lyapunov function is not unique.

Example: Consider a first-order system

$$\dot{x} = -g(x)$$

where $g(x)$ is locally Lipschitz on $(-a, a)$ and satisfies

$$g(0) = 0, xg(x) > 0, \forall x \in (-a, 0) \cup (0, a)$$

The system has a isolated equilibrium at $x = 0$. Explicit solution is impossible to obtain. Consider a Lyapunov function

$$V(x) = x^2 \rightarrow \dot{V} = -2xg(x)$$

or

$$V(x) = \int_0^x g(y)dy \rightarrow \dot{V} = -g^2(x)$$

Note: The choice of Lyapunov function is not unique.

Lyapunov Stability Theorem

Definitions:

1) A function $V : \mathcal{D} \rightarrow \mathbb{R}$ is said to be *positive definite* if it satisfies

$$V(0) = 0 \text{ and } V(x) > 0 \text{ for any } x \in \mathcal{D} - \{0\}$$

2) A function $V(x)$ is said to be *positive semi-definite*, if it just satisfies the weaker condition

$$V(0) = 0 \text{ and } V(x) \geq 0 \text{ for any } x \in \mathcal{D} - \{0\}$$

3) A function $V(x)$ is said to be *negative definite (semi-definite)*, if $-V(x)$ is *positive definite (semi-definite)*.

Lyapunov Stability Theorem

Definitions:

1) A function $V : \mathcal{D} \rightarrow \mathbb{R}$ is said to be *positive definite* if it satisfies

$$V(0) = 0 \text{ and } V(x) > 0 \text{ for any } x \in \mathcal{D} - \{0\}$$

2) A function $V(x)$ is said to be *positive semi-definite*, if it just satisfies the weaker condition

$$V(0) = 0 \text{ and } V(x) \geq 0 \text{ for any } x \in \mathcal{D} - \{0\}$$

3) A function $V(x)$ is said to be *negative definite (semi-definite)*, if $-V(x)$ is *positive definite (semi-definite)*.

Lyapunov Stability Theorem

Definitions:

1) A function $V : \mathcal{D} \rightarrow \mathbb{R}$ is said to be *positive definite* if it satisfies

$$V(0) = 0 \text{ and } V(x) > 0 \text{ for any } x \in \mathcal{D} - \{0\}$$

2) A function $V(x)$ is said to be *positive semi-definite*, if it just satisfies the weaker condition

$$V(0) = 0 \text{ and } V(x) \geq 0 \text{ for any } x \in \mathcal{D} - \{0\}$$

3) A function $V(x)$ is said to be *negative definite (semi-definite)*, if $-V(x)$ is *positive definite (semi-definite)*.

Lyapunov Stability Theorem

Rephrase the Lyapunov's theorem as : *The origin is stable if there is a C^1 and positive definite function $V(x)$ so that $\dot{V}(x)$ is negative semi-definite, and the origin is a.s. if $\dot{V}(x)$ is negative definite.*

For the class of function of quadratic form

$$V(x) = x^\top P x, \quad x \in \mathbb{R}^n$$

where P is a real symmetric matrix. In this case,

$$V(x) \text{ is p.d.} \iff P \text{ is p.d.} \iff \text{eig}(P) > 0$$

Example: $V(x) = ax_1^2 + 2x_1x_3 + ax_2^2 + 4x_2x_3 + ax_3^2$ and

$\dot{V}(x) = bx_1^2 + 2x_1x_3 + bx_2^2 + 4x_2x_3 + bx_3^2$, under what condition

of a, b yields a stable origin? an a.s. origin?

Lyapunov Stability Theorem

Rephrase the Lyapunov's theorem as : *The origin is stable if there is a C^1 and positive definite function $V(x)$ so that $\dot{V}(x)$ is negative semi-definite, and the origin is a.s. if $\dot{V}(x)$ is negative definite.*

For the class of function of quadratic form

$$V(x) = x^\top P x, \quad x \in \mathbb{R}^n$$

where P is a real symmetric matrix. In this case,

$$V(x) \text{ is p.d.} \iff P \text{ is p.d.} \iff \text{eig}(P) > 0$$

Example: $V(x) = ax_1^2 + 2x_1x_3 + ax_2^2 + 4x_2x_3 + ax_3^2$ and

$\dot{V}(x) = bx_1^2 + 2x_1x_3 + bx_2^2 + 4x_2x_3 + bx_3^2$, under what condition

of a, b yields a stable origin? an a.s. origin?

Lyapunov Stability Theorem

Rephrase the Lyapunov's theorem as : *The origin is stable if there is a C^1 and positive definite function $V(x)$ so that $\dot{V}(x)$ is negative semi-definite, and the origin is a.s. if $\dot{V}(x)$ is negative definite.*

For the class of function of quadratic form

$$V(x) = x^\top P x, \quad x \in \mathbb{R}^n$$

where P is a real symmetric matrix. In this case,

$$V(x) \text{ is p.d.} \iff P \text{ is p.d.} \iff \text{eig}(P) > 0$$

Example: $V(x) = ax_1^2 + 2x_1x_3 + ax_2^2 + 4x_2x_3 + ax_3^2$ and

$\dot{V}(x) = bx_1^2 + 2x_1x_3 + bx_2^2 + 4x_2x_3 + bx_3^2$, under what condition

of a, b yields a stable origin? an a.s. origin?

Example: Recall the pendulum system

$$\dot{x}_1 = x_2$$

$$\dot{x}_2 = -a \sin x_1 - bx_2$$

with $\mathcal{D} := (-\pi, \pi) \times \mathbb{R}$, consider $x_e = (0, 0)^\top$ and candidate Lyapunov function is chosen as $V(x) = ml^2[a(1 - \cos x_1) + \frac{1}{2}x_2^2]$ satisfies $V(x)$ is p.d., and the time derivative of V is given by

$$\dot{V}(x) = -ml^2bx_2^2 \leq 0$$

is only n.s.d. One can conclude the origin is *stable*, **but NOT a.s.!!!!** However, according to Barbalat's Lemma, we have

$$x_2 \rightarrow 0 \quad \text{as} \quad t \rightarrow \infty$$

Example: Recall the pendulum system

$$\dot{x}_1 = x_2$$

$$\dot{x}_2 = -a \sin x_1 - bx_2$$

with $\mathcal{D} := (-\pi, \pi) \times \mathbb{R}$, consider $x_e = (0, 0)^\top$ and candidate Lyapunov function is chosen as $V(x) = ml^2[a(1 - \cos x_1) + \frac{1}{2}x_2^2]$ satisfies $V(x)$ is p.d., and the time derivative of V is given by

$$\dot{V}(x) = -ml^2bx_2^2 \leq 0$$

is only n.s.d. One can conclude the origin is *stable*, **but NOT a.s.!!!!** However, according to Barbalat's Lemma, we have

$$x_2 \rightarrow 0 \quad \text{as} \quad t \rightarrow \infty$$

Example: Recall the pendulum system

$$\dot{x}_1 = x_2$$

$$\dot{x}_2 = -a \sin x_1 - bx_2$$

with $\mathcal{D} := (-\pi, \pi) \times \mathbb{R}$, consider $x_e = (0, 0)^\top$ and candidate Lyapunov function is chosen as $V(x) = ml^2[a(1 - \cos x_1) + \frac{1}{2}x_2^2]$ satisfies $V(x)$ is p.d., and the time derivative of V is given by

$$\dot{V}(x) = -ml^2bx_2^2 \leq 0$$

is only n.s.d. One can conclude the origin is *stable*, **but NOT a.s.!!!!** However, according to Barbalat's Lemma, we have

$$x_2 \rightarrow 0 \quad \text{as} \quad t \rightarrow \infty$$

Example: Recall the pendulum system

$$\dot{x}_1 = x_2$$

$$\dot{x}_2 = -a \sin x_1 - bx_2$$

with $\mathcal{D} := (-\pi, \pi) \times \mathbb{R}$, consider $x_e = (0, 0)^\top$ and candidate Lyapunov function is chosen as $V(x) = ml^2[a(1 - \cos x_1) + \frac{1}{2}x_2^2]$ satisfies $V(x)$ is p.d., and the time derivative of V is given by

$$\dot{V}(x) = -ml^2bx_2^2 \leq 0$$

is only n.s.d. One can conclude the origin is *stable*, **but NOT a.s.!!!!** However, according to Barbalat's Lemma, we have

$$x_2 \rightarrow 0 \quad \text{as} \quad t \rightarrow \infty$$

Let's find a Lyapunov function $V(x)$ that would have a n.d. $\dot{V}(x)$.

Try a more general quadratic form

$$V(x) = x^T P x + a(1 - \cos x_1)$$

with

$$P = \begin{pmatrix} p_{11} & p_{12} \\ p_{12} & p_{22} \end{pmatrix}, \quad p_{11} > 0, \quad p_{11}p_{22} - p_{12}^2 > 0$$

The derivative is given by

$$\begin{aligned} \dot{V}(x) = & a(1 - p_{22})x_2 \sin x_1 + (p_{11} - p_{12}b)x_1x_2 - ap_{12}x_1 \sin x_1 \\ & + (p_{12} - p_{22}b)x_2^2 \end{aligned}$$

By taking $p_{22} = 1$ and $p_{11} = p_{12}b$, we have

Let's find a Lyapunov function $V(x)$ that would have a n.d. $\dot{V}(x)$.

Try a more general quadratic form

$$V(x) = x^\top P x + a(1 - \cos x_1)$$

with

$$P = \begin{pmatrix} p_{11} & p_{12} \\ p_{12} & p_{22} \end{pmatrix}, \quad p_{11} > 0, \quad p_{11}p_{22} - p_{12}^2 > 0$$

The derivative is given by

$$\begin{aligned} \dot{V}(x) = & a(1 - p_{22})x_2 \sin x_1 + (p_{11} - p_{12}b)x_1x_2 - ap_{12}x_1 \sin x_1 \\ & + (p_{12} - p_{22}b)x_2^2 \end{aligned}$$

By taking $p_{22} = 1$ and $p_{11} = p_{12}b$, we have

Let's find a Lyapunov function $V(x)$ that would have a n.d. $\dot{V}(x)$.

Try a more general quadratic form

$$V(x) = x^\top P x + a(1 - \cos x_1)$$

with

$$P = \begin{pmatrix} p_{11} & p_{12} \\ p_{12} & p_{22} \end{pmatrix}, \quad p_{11} > 0, \quad p_{11}p_{22} - p_{12}^2 > 0$$

The derivative is given by

$$\begin{aligned} \dot{V}(x) = & a(1 - p_{22})x_2 \sin x_1 + (p_{11} - p_{12}b)x_1x_2 - ap_{12}x_1 \sin x_1 \\ & + (p_{12} - p_{22}b)x_2^2 \end{aligned}$$

By taking $p_{22} = 1$ and $p_{11} = p_{12}b$, we have

$$\dot{V} = -ap_{12}x_1 \sin x_1 + (p_{12} - b) x_2^2$$

Summary the requirements of p.d. and n.d.

$$\begin{cases} p_{12}b > 0, & p_{12}b - p_{12}^2 > 0 \\ p_{12} > 0, & p_{12} - b < 0 \end{cases}$$

choose $p_{12} = 0.5b$, yields the time derivative

$$\dot{V} = -0.5abx_1 \sin x_1 - 0.5bx_2^2$$

is n.d. for any $x \in \mathcal{D} := \{x \in \mathbb{R}^2 | x_1 \in (-\pi, 0) \cup (0, \pi)\}$. Hence, the origin is *locally* a.s..

This example emphasis, the conditions in the Lyapunov stability theorem are only sufficient!

$$\dot{V} = -ap_{12}x_1 \sin x_1 + (p_{12} - b)x_2^2$$

Summary the requirements of p.d. and n.d.

$$\begin{cases} p_{12}b > 0, & p_{12}b - p_{12}^2 > 0 \\ p_{12} > 0, & p_{12} - b < 0 \end{cases}$$

choose $p_{12} = 0.5b$, yields the time derivative

$$\dot{V} = -0.5abx_1 \sin x_1 - 0.5bx_2^2$$

is n.d. for any $x \in \mathcal{D} := \{x \in \mathbb{R}^2 | x_1 \in (-\pi, 0) \cup (0, \pi)\}$. Hence, the origin is *locally* a.s..

This example emphasis, the conditions in the Lyapunov stability theorem are only sufficient!

Globally Asymptotically Stable

Theorem:(Barbashin-Krasovskii) For the autonomous system (1) that well-defined in \mathbb{R}^n , let $V : \mathbb{R}^n \rightarrow \mathbb{R}$ be a \mathcal{C}^1 function such that

- i) $V(x)$ is positive definite
- ii) $V(x)$ is *radially unbounded*, that is

$$\|x\| \rightarrow \infty \quad \text{implies} \quad V(x) \rightarrow \infty$$

- iii) $\dot{V}(x)$ is negative definite

then the equilibrium at origin is *globally* asymptotically stable.

Example: $V(x) = \frac{x_1^2}{1+x_1^2} + x_2^2$

Globally Asymptotically Stable

Theorem:(Barbashin-Krasovskii) For the autonomous system (1) that well-defined in \mathbb{R}^n , let $V : \mathbb{R}^n \rightarrow \mathbb{R}$ be a \mathcal{C}^1 function such that

- i) $V(x)$ is positive definite
- ii) $V(x)$ is *radially unbounded*, that is

$$\|x\| \rightarrow \infty \quad \text{implies} \quad V(x) \rightarrow \infty$$

- iii) $\dot{V}(x)$ is negative definite

then the equilibrium at origin is *globally* asymptotically stable.

Example: $V(x) = \frac{x_1^2}{1+x_1^2} + x_2^2$

Unstable

Lemma For the autonomous system (1), $V : \mathcal{D} \rightarrow \mathbb{R}$ be a \mathcal{C}^1 function, suppose the equilibrium at origin is contained by \mathcal{D} . If $V(0) = 0$ and there is a point at x_0 arbitrarily close to the origin such that $V(x_0) > 0$, then we can always construct a set

$$\mathcal{U} := \{x \in \mathcal{B}_r | V(x) > 0\}$$

such that

- \mathcal{U} is non-empty set contained in $\mathcal{B}_r = \{x \in \mathbb{R}^n | \|x\| < r\}$ for some $r > 0$.
- Its boundary is the surface $V(x) = 0$ or the sphere $\|x\| = r$

Example : $V(x) = x_1^2 - x_2^2$.

Unstable

Lemma For the autonomous system (1), $V : \mathcal{D} \rightarrow \mathbb{R}$ be a \mathcal{C}^1 function, suppose the equilibrium at origin is contained by \mathcal{D} . If $V(0) = 0$ and there is a point at x_0 arbitrarily close to the origin such that $V(x_0) > 0$, then we can always construct a set

$$\mathcal{U} := \{x \in \mathcal{B}_r | V(x) > 0\}$$

such that

- \mathcal{U} is non-empty set contained in $\mathcal{B}_r = \{x \in \mathbb{R}^n | \|x\| < r\}$ for some $r > 0$.
- Its boundary is the surface $V(x) = 0$ or the sphere $\|x\| = r$

Example : $V(x) = x_1^2 - x_2^2$.

Unstable

Lemma For the autonomous system (1), $V : \mathcal{D} \rightarrow \mathbb{R}$ be a \mathcal{C}^1 function, suppose the equilibrium at origin is contained by \mathcal{D} . If $V(0) = 0$ and there is a point at x_0 arbitrarily close to the origin such that $V(x_0) > 0$, then we can always construct a set

$$\mathcal{U} := \{x \in \mathcal{B}_r | V(x) > 0\}$$

such that

- \mathcal{U} is non-empty set contained in $\mathcal{B}_r = \{x \in \mathbb{R}^n | \|x\| < r\}$ for some $r > 0$.
- Its boundary is the surface $V(x) = 0$ or the sphere $\|x\| = r$

Example : $V(x) = x_1^2 - x_2^2$.

Unstable

Lemma For the autonomous system (1), $V : \mathcal{D} \rightarrow \mathbb{R}$ be a \mathcal{C}^1 function, suppose the equilibrium at origin is contained by \mathcal{D} . If $V(0) = 0$ and there is a point at x_0 arbitrarily close to the origin such that $V(x_0) > 0$, then we can always construct a set

$$\mathcal{U} := \{x \in \mathcal{B}_r | V(x) > 0\}$$

such that

- \mathcal{U} is non-empty set contained in $\mathcal{B}_r = \{x \in \mathbb{R}^n | \|x\| < r\}$ for some $r > 0$.
- Its boundary is the surface $V(x) = 0$ or the sphere $\|x\| = r$

Example : $V(x) = x_1^2 - x_2^2$.

Unstable

Lemma For the autonomous system (1), $V : \mathcal{D} \rightarrow \mathbb{R}$ be a \mathcal{C}^1 function, suppose the equilibrium at origin is contained by \mathcal{D} . If $V(0) = 0$ and there is a point at x_0 arbitrarily close to the origin such that $V(x_0) > 0$, then we can always construct a set

$$\mathcal{U} := \{x \in \mathcal{B}_r | V(x) > 0\}$$

such that

- \mathcal{U} is non-empty set contained in $\mathcal{B}_r = \{x \in \mathbb{R}^n | \|x\| < r\}$ for some $r > 0$.
- Its boundary is the surface $V(x) = 0$ or the sphere $\|x\| = r$

Example : $V(x) = x_1^2 - x_2^2$.

Unstable

Theorem:(Chetaev) For the autonomous system (1), $V : \mathcal{D} \rightarrow \mathbb{R}$ be a \mathcal{C}^1 function, suppose the equilibrium at origin is contained by \mathcal{D} , $V(0) = 0$ and there is a point at x_0 arbitrarily close to the origin such that $V(x_0) > 0$, if we can find a set aforementioned \mathcal{U} such that $\dot{V}(x) > 0$ for any $x \in \mathcal{U}$, then the origin is *unstable*.

Note: the simplest case is $\mathcal{U} = \mathcal{B}_r - \{0\}$.

Corollary: For the system (1), if a \mathcal{C}^1 function $V_1(x)$ can be found in a neighborhood of the origin such that $V_1(0) = 0$, and \dot{V}_1 is positive definite, but V_1 itself is NOT negative definite or negative semidefinite arbitrarily near the origin, then the origin is unstable.

Unstable

Theorem:(Chetaev) For the autonomous system (1), $V : \mathcal{D} \rightarrow \mathbb{R}$ be a \mathcal{C}^1 function, suppose the equilibrium at origin is contained by \mathcal{D} , $V(0) = 0$ and there is a point at x_0 arbitrarily close to the origin such that $V(x_0) > 0$, if we can find a set aforementioned \mathcal{U} such that $\dot{V}(x) > 0$ for any $x \in \mathcal{U}$, then the origin is *unstable*.

Note: the simplest case is $\mathcal{U} = \mathcal{B}_r - \{0\}$.

Corollary: For the system (1), if a \mathcal{C}^1 function $V_1(x)$ can be found in a neighborhood of the origin such that $V_1(0) = 0$, and \dot{V}_1 is positive definite, but V_1 itself is NOT negative definite or negative semidefinite arbitrarily near the origin, then the origin is unstable.

Contents

- Motivation
- Lyapunov's Direct Method for Autonomous System
- The Invariance Principle
- LTI System and Exponential Stable

Recall the energy-like Lyapunov function of pendulum system whose time derivative

$$\dot{V}(x) = -bx_2^2$$

According to Barbala's lemma, we have x_2 converges to zero.

Notice, however, \dot{V} is n.d. except on the line $x_2 = 0$. In other words, to maintain $\dot{V} = 0$, x_2 has to and *always* equal to 0,

$$x_2 \equiv 0 \Rightarrow \dot{x}_2 = 0 \Rightarrow -a \sin(x_1) = 0 \Rightarrow x_1 = 0$$

Hence, the system can maintain $\dot{V} = 0$ condition only at the origin, therefore, as $V(x)$ decreasing towards to 0, consequently, $x(t)$ converges to origin, i.e. origin is a.s.

A wrong derivation is $x_2 \rightarrow 0 \Rightarrow \dot{x}_2 = 0$!

Recall the energy-like Lyapunov function of pendulum system whose time derivative

$$\dot{V}(x) = -bx_2^2$$

According to Barbala's lemma, we have x_2 converges to zero.

Notice, however, \dot{V} is n.d. except on the line $x_2 = 0$. In other words, to maintain $\dot{V} = 0$, x_2 has to and *always* equal to 0,

$$x_2 \equiv 0 \Rightarrow \dot{x}_2 = 0 \Rightarrow -a \sin(x_1) = 0 \Rightarrow x_1 = 0$$

Hence, the system can maintain $\dot{V} = 0$ condition only at the origin, therefore, as $V(x)$ decreasing towards to 0, consequently, $x(t)$ converges to origin, i.e. origin is a.s.

A wrong derivation is $x_2 \rightarrow 0 \Rightarrow \dot{x}_2 = 0$!

Recall the energy-like Lyapunov function of pendulum system whose time derivative

$$\dot{V}(x) = -bx_2^2$$

According to Barbala's lemma, we have x_2 converges to zero.

Notice, however, \dot{V} is n.d. except on the line $x_2 = 0$. In other words, to maintain $\dot{V} = 0$, x_2 has to and *always* equal to 0,

$$x_2 \equiv 0 \Rightarrow \dot{x}_2 = 0 \Rightarrow -a \sin(x_1) = 0 \Rightarrow x_1 = 0$$

Hence, the system can maintain $\dot{V} = 0$ condition only at the origin, therefore, as $V(x)$ decreasing towards to 0, consequently, $x(t)$ converges to origin, i.e. origin is a.s.

A wrong derivation is $x_2 \rightarrow 0 \Rightarrow \dot{x}_2 = 0$!

Recall the energy-like Lyapunov function of pendulum system whose time derivative

$$\dot{V}(x) = -bx_2^2$$

According to Barbala's lemma, we have x_2 converges to zero.

Notice, however, \dot{V} is n.d. except on the line $x_2 = 0$. In other words, to maintain $\dot{V} = 0$, x_2 has to and *always* equal to 0,

$$x_2 \equiv 0 \Rightarrow \dot{x}_2 = 0 \Rightarrow -a \sin(x_1) = 0 \Rightarrow x_1 = 0$$

Hence, the system can maintain $\dot{V} = 0$ condition only at the origin, therefore, as $V(x)$ decreasing towards to 0, consequently, $x(t)$ converges to origin, i.e. origin is a.s.

A wrong derivation is $x_2 \rightarrow 0 \Rightarrow \dot{x}_2 = 0$!

LaSalle's Theorem

Definition: A set $\Omega \in \mathbb{R}^n$ is said to be an *invariant* set w.r.t the autonomous system (1), if $x(t_0) \in \Omega \Rightarrow x(t) \in \Omega$ for all $t \in \mathbb{R}$. That is, if a solution belongs to Ω at some instant, then it belongs to Ω for all future and past time, e.g. equilibrium.

Theorem: For the autonomous system (1), Let $V : \mathcal{D} \rightarrow \mathbb{R}$ be a p.d. \mathcal{C}^1 function, suppose $\dot{V}(x) \leq 0$ for all $x \in \mathcal{D}$. Let

$$E := \{x \in \mathcal{D} | \dot{V}(x) = 0\}$$

and Ω be the *largest invariant set* of E , then all solution starting in \mathcal{D} approaches Ω as $t \rightarrow \infty$.

LaSalle's Theorem

Definition: A set $\Omega \in \mathbb{R}^n$ is said to be an *invariant* set w.r.t the autonomous system (1), if $x(t_0) \in \Omega \Rightarrow x(t) \in \Omega$ for all $t \in \mathbb{R}$.

That is, if a solution belongs to Ω at some instant, then it belongs to Ω for all future and past time, e.g. equilibrium.

Theorem: For the autonomous system (1), Let $V : \mathcal{D} \rightarrow \mathbb{R}$ be a p.d. \mathcal{C}^1 function, suppose $\dot{V}(x) \leq 0$ for all $x \in \mathcal{D}$. Let

$$E := \{x \in \mathcal{D} | \dot{V}(x) = 0\}$$

and Ω be the *largest invariant set* of E , then all solution starting in \mathcal{D} approaches Ω as $t \rightarrow \infty$.

LaSalle's Theorem

For the autonomous system (1), Let $V : \mathcal{D} \rightarrow \mathbb{R}$ be a p.d. \mathcal{C}^1 function on domain \mathcal{D} containing the equilibrium at $x = 0$,

Corollary 1: suppose $\dot{V}(x) \leq -W(x) \leq 0$ for all $x \in \mathcal{D}$ and some positive-semidefinite $W(x)$, then it holds

$$W(x) \rightarrow 0 \text{ as } t \rightarrow \infty$$

Corollary 2: suppose $\dot{V}(x) \leq 0$ for all $x \in \mathcal{D}$. Let

$$E := \{x \in \mathcal{D} | \dot{V}(x) = 0\}$$

and suppose no other solution stay identically in E other than the trivial solution $x(t) \equiv 0$, then the origin is a.s.

LaSalle's Theorem

For the autonomous system (1), Let $V : \mathcal{D} \rightarrow \mathbb{R}$ be a p.d. \mathcal{C}^1 function on domain \mathcal{D} containing the equilibrium at $x = 0$,

Corollary 1: suppose $\dot{V}(x) \leq -W(x) \leq 0$ for all $x \in \mathcal{D}$ and some positive-semidefinite $W(x)$, then it holds

$$W(x) \rightarrow 0 \text{ as } t \rightarrow \infty$$

Corollary 2: suppose $\dot{V}(x) \leq 0$ for all $x \in \mathcal{D}$. Let

$$E := \{x \in \mathcal{D} | \dot{V}(x) = 0\}$$

and suppose no other solution stay identically in E other than the trivial solution $x(t) \equiv 0$, then the origin is a.s.

The Invariance Principle

Example: Now we are ready to analysis the motivating example

$$\dot{x} = \theta x + u$$

with the control law $u = -kx$ and k is given by the adaptive law

$$\dot{k} = \gamma x^2$$

Taking $x_1 = x$ and $x_2 = k$, the closed-loop system is

$$\dot{x}_1 = -(x_2 - \theta) x_1$$

$$\dot{x}_2 = \gamma x_1^2$$

Step I: Check the type of system and find the equilibrium

$$x_e = [0, c]^\top$$

Step II: Shift equilibrium to the origin by $z_1 = x_1, z_2 = x_2 - c$

$$\dot{z}_1 = -(z_2 + c - \theta) z_1$$

$$\dot{z}_2 = \gamma z_1^2$$

Step III: Consider the Lyapunov candidate function

$$V(z) = \frac{1}{2} z_1^2 + \frac{1}{2\gamma} z_2^2$$

and check the positive definiteness (and radially unboundedness)

Step IV: Calculate the time derivative

$$\dot{V} = -(z_2 + c - \theta)z_1^2 + z_2z_1^2 = -(c - \theta)z_1^2$$

Step V: Draw the conclusion about the equilibrium $x_e = [0, c]^\top$

- $c < \theta \Rightarrow \dot{V} \geq 0 \Rightarrow x_e$ is unstable
- $c = \theta \Rightarrow \dot{V} = 0 \Rightarrow x_e$ is stable
- $c > \theta \Rightarrow \dot{V} \leq 0 \Rightarrow x_e$ is stable.

According to the LaSalle's corollary, we have $x_1 \rightarrow 0$ as $t \rightarrow \infty$.

However, the origin is not the only solution of $\dot{V} = 0$, x_e is not g.a.s!

Step IV: Calculate the time derivative

$$\dot{V} = -(z_2 + c - \theta)z_1^2 + z_2z_1^2 = -(c - \theta)z_1^2$$

Step V: Draw the conclusion about the equilibrium $x_e = [0, c]^\top$

- $c < \theta \Rightarrow \dot{V} \geq 0 \Rightarrow x_e$ is unstable
- $c = \theta \Rightarrow \dot{V} = 0 \Rightarrow x_e$ is stable
- $c > \theta \Rightarrow \dot{V} \leq 0 \Rightarrow x_e$ is stable.

According to the LaSalle's corollary, we have $x_1 \rightarrow 0$ as $t \rightarrow \infty$.

However, the origin is not the only solution of $\dot{V} = 0$, x_e is not g.a.s!

Step IV: Calculate the time derivative

$$\dot{V} = -(z_2 + c - \theta)z_1^2 + z_2z_1^2 = -(c - \theta)z_1^2$$

Step V: Draw the conclusion about the equilibrium $x_e = [0, c]^\top$

- $c < \theta \Rightarrow \dot{V} \geq 0 \Rightarrow x_e$ is unstable
- $c = \theta \Rightarrow \dot{V} = 0 \Rightarrow x_e$ is stable
- $c > \theta \Rightarrow \dot{V} \leq 0 \Rightarrow x_e$ is stable.

According to the LaSalle's corollary, we have $x_1 \rightarrow 0$ as $t \rightarrow \infty$.

However, the origin is not the only solution of $\dot{V} = 0$, x_e is not g.a.s!

Step IV: Calculate the time derivative

$$\dot{V} = -(z_2 + c - \theta)z_1^2 + z_2z_1^2 = -(c - \theta)z_1^2$$

Step V: Draw the conclusion about the equilibrium $x_e = [0, c]^\top$

- $c < \theta \Rightarrow \dot{V} \geq 0 \Rightarrow x_e$ is unstable
- $c = \theta \Rightarrow \dot{V} = 0 \Rightarrow x_e$ is stable
- $c > \theta \Rightarrow \dot{V} \leq 0 \Rightarrow x_e$ is stable.

According to the LaSalle's corollary, we have $x_1 \rightarrow 0$ as $t \rightarrow \infty$.

However, the origin is not the only solution of $\dot{V} = 0$, x_e is not g.a.s!

Contents

- Motivation
- Lyapunov's Direct Method for Autonomous System
- The Invariance Principle
- LTI System and Exponential Stable

The LTI system

$$\dot{x} = Ax, \quad x(0) = x_0 \in \mathbb{R}^n$$

It has an equilibrium at origin. The equilibrium is isolated iff $\det(A) \neq 0$, otherwise every point in the null space of A

$$\text{null}(A) = \{x \in \mathbb{R}^n | Ax = 0\}$$

is an equilibrium of the system.

From the linear system theorem, we know that

- All $\text{Re}\lambda_i < 0$ or $\text{Re}\lambda_i = 0$ with associated Jordan block's rank is 1 \iff origin is "neutrally" stable or marginally stable
- All $\text{Re}\lambda_i < 0 \iff$ origin is stable
- Any $\text{Re}\lambda_i > 0 \iff$ origin is unstable

The LTI system

$$\dot{x} = Ax, \quad x(0) = x_0 \in \mathbb{R}^n$$

It has an equilibrium at origin. The equilibrium is isolated iff $\det(A) \neq 0$, otherwise every point in the null space of A

$$\text{null}(A) = \{x \in \mathbb{R}^n | Ax = 0\}$$

is an equilibrium of the system.

From the linear system theorem, we know that

- All $\text{Re}\lambda_i < 0$ or $\text{Re}\lambda_i = 0$ with associated Jordan block's rank is 1 \iff origin is "neutrally" stable or marginally stable
- All $\text{Re}\lambda_i < 0 \iff$ origin is stable
- Any $\text{Re}\lambda_i > 0 \iff$ origin is unstable

The LTI system

$$\dot{x} = Ax, \quad x(0) = x_0 \in \mathbb{R}^n$$

It has an equilibrium at origin. The equilibrium is isolated iff $\det(A) \neq 0$, otherwise every point in the null space of A

$$\text{null}(A) = \{x \in \mathbb{R}^n | Ax = 0\}$$

is an equilibrium of the system.

From the linear system theorem, we know that

- All $\text{Re}\lambda_i < 0$ or $\text{Re}\lambda_i = 0$ with associated Jordan block's rank is 1 \iff origin is "neutrally" stable or marginally stable
- All $\text{Re}\lambda_i < 0 \iff$ origin is stable
- Any $\text{Re}\lambda_i > 0 \iff$ origin is unstable

The LTI system

$$\dot{x} = Ax, \quad x(0) = x_0 \in \mathbb{R}^n$$

with $\det(A) \neq 0$. It has solution

$$x(t) = e^{At}x_0$$

From the Lyapunov stability definitions, we know

- All $\operatorname{Re}\lambda_i < 0$ or $\operatorname{Re}\lambda_i = 0$ with associated Jordan block's rank is 1 \iff origin is stable in the sense of Lyapunov
- All $\operatorname{Re}\lambda_i < 0 \iff$ origin is asymptotically stable (moreover, g.e.s.)
- Any $\operatorname{Re}\lambda_i > 0 \iff$ origin is unstable

The LTI system

$$\dot{x} = Ax, \quad x(0) = x_0 \in \mathbb{R}^n$$

with $\det(A) \neq 0$. It has solution

$$x(t) = e^{At}x_0$$

From the Lyapunov stability definitions, we know

- All $\operatorname{Re}\lambda_i < 0$ or $\operatorname{Re}\lambda_i = 0$ with associated Jordan block's rank is 1 \iff origin is stable in the sense of Lyapunov
- All $\operatorname{Re}\lambda_i < 0 \iff$ origin is asymptotically stable (moreover, g.e.s.)
- Any $\operatorname{Re}\lambda_i > 0 \iff$ origin is unstable

The LTI system

$$\dot{x} = Ax, \quad x(0) = x_0 \in \mathbb{R}^n$$

with $\det(A) \neq 0$. It has solution

$$x(t) = e^{At}x_0$$

From the Lyapunov stability definitions, we know

- All $\operatorname{Re}\lambda_i < 0$ or $\operatorname{Re}\lambda_i = 0$ with associated Jordan block's rank is 1 \iff origin is stable in the sense of Lyapunov
- All $\operatorname{Re}\lambda_i < 0 \iff$ origin is asymptotically stable (moreover, g.e.s.)
- Any $\operatorname{Re}\lambda_i > 0 \iff$ origin is unstable

The LTI system

$$\dot{x} = Ax, \quad x(0) = x_0 \in \mathbb{R}^n$$

with $\det(A) \neq 0$. It has solution

$$x(t) = e^{At}x_0$$

From the Lyapunov stability definitions, we know

- All $\operatorname{Re}\lambda_i < 0$ or $\operatorname{Re}\lambda_i = 0$ with associated Jordan block's rank is 1 \iff origin is stable in the sense of Lyapunov
- All $\operatorname{Re}\lambda_i < 0 \iff$ origin is asymptotically stable (moreover, g.e.s.)
- Any $\operatorname{Re}\lambda_i > 0 \iff$ origin is unstable

Consider a quadratic Lyapunov candidate function

$$V(x) = x^{\top} P x$$

with P is a real symmetric p.d. matrix. The derivative of V along the trajectories of linear system is given by

$$\dot{V} = x^{\top} P A x + x^{\top} A^{\top} P x = -x^{\top} Q x$$

From the Lyapunov stability Theorems, we know

- Q is p.s.d \implies origin is stable in the sense of Lyapunov
- Q is p.d \iff origin is asymptotically stable (moreover, g.e.s.)
- Q is n.d. \implies origin is unstable

Consider a quadratic Lyapunov candidate function

$$V(x) = x^{\top} P x$$

with P is a real symmetric p.d. matrix. The derivative of V along the trajectories of linear system is given by

$$\dot{V} = x^{\top} P A x + x^{\top} A^{\top} P x = -x^{\top} Q x$$

From the Lyapunov stability Theorems, we know

- Q is p.s.d \implies origin is stable in the sense of Lyapunov
- Q is p.d \iff origin is asymptotically stable (moreover, g.e.s.)
- Q is n.d. \implies origin is unstable

Consider a quadratic Lyapunov candidate function

$$V(x) = x^{\top} P x$$

with P is a real symmetric p.d. matrix. The derivative of V along the trajectories of linear system is given by

$$\dot{V} = x^{\top} P A x + x^{\top} A^{\top} P x = -x^{\top} Q x$$

From the Lyapunov stability Theorems, we know

- Q is p.s.d \implies origin is stable in the sense of Lyapunov
- Q is p.d \iff origin is asymptotically stable (moreover, g.e.s.)
- Q is n.d. \implies origin is unstable

Lyapunov Equation

Theorem: A matrix is Hurwitz, that is $\text{Re}(\lambda_i) < 0$ if and only if for any given p.d. symmetric matrix Q , there exists a p.d. symmetric matrix P that satisfies the Lyapunov equation

$$PA + A^T P = -Q$$

Moreover, the solution P is unique.

The proof can be found in Khalil's Nonlinear system Theorem 4.6.

Example:

$$A = \begin{bmatrix} 0 & -1 \\ 1 & -1 \end{bmatrix}, \quad \text{with} \quad Q = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}$$

Remark: For LTI, a.s. implies g.e.s. for unique equilibrium.

Lyapunov Equation

Theorem: A matrix is Hurwitz, that is $\text{Re}(\lambda_i) < 0$ if and only if for any given p.d. symmetric matrix Q , there exists a p.d. symmetric matrix P that satisfies the Lyapunov equation

$$PA + A^T P = -Q$$

Moreover, the solution P is unique.

The proof can be found in Khalil's Nonlinear system Theorem 4.6.

Example:

$$A = \begin{bmatrix} 0 & -1 \\ 1 & -1 \end{bmatrix}, \quad \text{with} \quad Q = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}$$

Remark: For LTI, a.s. implies g.e.s. for unique equilibrium.

Lyapunov Equation

Theorem: A matrix is Hurwitz, that is $\text{Re}(\lambda_i) < 0$ if and only if for any given p.d. symmetric matrix Q , there exists a p.d. symmetric matrix P that satisfies the Lyapunov equation

$$PA + A^T P = -Q$$

Moreover, the solution P is unique.

The proof can be found in Khalil's Nonlinear system Theorem 4.6.

Example:

$$A = \begin{bmatrix} 0 & -1 \\ 1 & -1 \end{bmatrix}, \quad \text{with} \quad Q = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}$$

Remark: For LTI, a.s. implies g.e.s. for unique equilibrium.

Lyapunov Equation

Theorem: A matrix is Hurwitz, that is $\text{Re}(\lambda_i) < 0$ if and only if for any given p.d. symmetric matrix Q , there exists a p.d. symmetric matrix P that satisfies the Lyapunov equation

$$PA + A^T P = -Q$$

Moreover, the solution P is unique.

The proof can be found in Khalil's Nonlinear system Theorem 4.6.

Example:

$$A = \begin{bmatrix} 0 & -1 \\ 1 & -1 \end{bmatrix}, \quad \text{with} \quad Q = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}$$

Remark: For LTI, a.s. implies g.e.s. for unique equilibrium.

Exponentially Stable

Theorem: Let $x = 0$ be an equilibrium point for the autonomous system (1) and $\mathcal{D} \subset \mathbb{R}^n$ be a domain containing $x = 0$. Let $V : \mathcal{D} \rightarrow \mathbb{R}$ be a \mathcal{C}^1 function such that:

$$k_1 \|x\|^a \leq V(x) \leq k_2 \|x\|^a$$

$$\dot{V} \leq -k_3 \|x\|^a$$

for any $t \geq 0$ and any $x \in \mathcal{D}$, where k_1, k_2, k_3 and a are positive constants. Then the origin is *exponentially stable*. If conditions hold globally, then the origin is g.e.s..

Remark: e.s naturally indicates a.s and stable in the sense of Lyapunov.

Exponentially Stable

Theorem: Let $x = 0$ be an equilibrium point for the autonomous system (1) and $\mathcal{D} \subset \mathbb{R}^n$ be a domain containing $x = 0$. Let $V : \mathcal{D} \rightarrow \mathbb{R}$ be a \mathcal{C}^1 function such that:

$$k_1 \|x\|^a \leq V(x) \leq k_2 \|x\|^a$$

$$\dot{V} \leq -k_3 \|x\|^a$$

for any $t \geq 0$ and any $x \in \mathcal{D}$, where k_1, k_2, k_3 and a are positive constants. Then the origin is *exponentially stable*. If conditions hold globally, then the origin is g.e.s..

Remark: e.s naturally indicates a.s and stable in the sense of Lyapunov.

Finite-Time Stable(Supplementary)

Theorem: Let $x = 0$ be an equilibrium point for the autonomous system (1) and $\mathcal{D} \subset \mathbb{R}^n$ be a domain containing $x = 0$. Let $V : \mathcal{D} \rightarrow \mathbb{R}$ be a \mathcal{C}^1 and positive definite function such that:

$$\dot{V} \leq -kV(x)^\alpha$$

for any $t \geq 0$ and any $x \in \mathcal{D}$, where $k > 0$ and $\alpha \in (0, 1)$. Then the origin is *finite-time stable* with the settling time

$$T = \frac{V(0)^{1-\alpha}}{k(1-\alpha)}.$$

If conditions hold globally, then the origin is globally finite time stable.

Lyapunov's Direct Method

Example: Consider the system

$$\dot{x}_1 = x_2 + cx_1 (x_1^2 + x_2^2)$$

$$\dot{x}_2 = -x_1 + cx_2 (x_1^2 + x_2^2)$$

where c is a constant, study the stability property of the system.

Step I: Check the type of system, find the equilibrium $x_e = [0, 0]$.

Step II: Consider a candidate Lyapunov function

$$V(x) = x_1^2 + x_2^2$$

that is positive definite and radially unbounded.

Lyapunov's Direct Method

Example: Consider the system

$$\dot{x}_1 = x_2 + cx_1(x_1^2 + x_2^2)$$

$$\dot{x}_2 = -x_1 + cx_2(x_1^2 + x_2^2)$$

where c is a constant, study the stability property of the system.

Step I: Check the type of system, find the equilibrium $x_e = [0, 0]$.

Step II: Consider a candidate Lyapunov function

$$V(x) = x_1^2 + x_2^2$$

that is positive definite and radially unbounded.

Step III: Calculate the time derivative

$$\dot{V} = 2c(x_1^2 + x_2^2)^2$$

Step IV: draw the conclusion

- $c = 0 \Rightarrow \dot{V} = 0 \Rightarrow$ origin is stable
- $c < 0 \Rightarrow \dot{V}$ is negative definite \Rightarrow origin is g.a.s, but NOT e.s or finite time convergence.
- $c > 0 \Rightarrow \dot{V} > 0$ for any $x \in \mathcal{U} := \{x \in \mathbb{R}^2 | 0 < \|x\| < r, r > 0\}$
 \Rightarrow origin is unstable

Step III: Calculate the time derivative

$$\dot{V} = 2c(x_1^2 + x_2^2)^2$$

Step IV: draw the conclusion

- $c = 0 \Rightarrow \dot{V} = 0 \Rightarrow$ origin is stable
- $c < 0 \Rightarrow \dot{V}$ is negative definite \Rightarrow origin is g.a.s, but NOT e.s or finite time convergence.
- $c > 0 \Rightarrow \dot{V} > 0$ for any $x \in \mathcal{U} := \{x \in \mathbb{R}^2 | 0 < \|x\| < r, r > 0\}$
 \Rightarrow origin is unstable

Step III: Calculate the time derivative

$$\dot{V} = 2c(x_1^2 + x_2^2)^2$$

Step IV: draw the conclusion

- $c = 0 \Rightarrow \dot{V} = 0 \Rightarrow$ origin is stable
- $c < 0 \Rightarrow \dot{V}$ is negative definite \Rightarrow origin is g.a.s, but NOT e.s or finite time convergence.
- $c > 0 \Rightarrow \dot{V} > 0$ for any $x \in \mathcal{U} := \{x \in \mathbb{R}^2 | 0 < \|x\| < r, r > 0\}$
 \Rightarrow origin is unstable