

# Measurement of Central Exclusive Production of $h\bar{h}$ pairs $(h=\pi,K,p)$ with Roman Pot detectors in diffractive proton-proton interactions at $\sqrt{s}=200~{\rm GeV}$

Leszek Adamczyk¹, Łukasz Fulek¹, Włodek Guryn², Mariusz Przybycień¹, and Rafał Sikora¹,†

<sup>1</sup>AGH University of Science and Technology, FPACS, Kraków, Poland <sup>2</sup>Brookhaven National Laboratory, Upton, NY, USA <sup>†</sup>e-mail: rafal.sikora@fis.agh.edu.pl

August 2, 2017

#### Abstract

In this note we present analysis of the Central Exclusive Production process using data from proton-proton collisions collected in 2015. This data was collected using the Roman Pot detectors which ensured efficient triggering and measuring diffractively scattered protons. We describe all intermediate stages of analysis involving extraction of the acceptance and efficiency corrections, comparison of data with Monte Carlo simulations of detector response, and study of systematic uncertainties. Finally, we show the physics outcome of the analysis.

### DRAFT

# Contents

T	Introduction	4
	1.1 Central Exclusive Production	4
	1.2 Double Pomeron Exchange	4
	1.3 Physics motivation for the measurement	
	· · · · · · · · · · · · · · · · · · ·	
	1.3.2 Absorption effects	5
	1.5.5 Size of interaction region	Ü
2	Data set	6
-	2.1 Trigger	6
	2.2 Reconstruction software	7
	2.3 Data format	7
	2.4 Bad runs	7
	2.4.1 RP distance from the beamline	7
	2.1.1 Itt distance from the seaffine	·
3	Event selection	8
	3.1 List of cuts	8
	3.2 Description of cuts	8
	3.2.1 (C1,C2) Primary vertex	8
	3.2.2 (C3) TPC tracks	9
	3.2.3 (C4) RP tracks	9
	3.2.4 (C5) TPC-RP z-vertex matching	9
	3.2.5 (C6) BBC-large signal veto	9
	3.2.6 (C7) TOF clusters limit	9
	3.2.7 (C8) Exclusivity cut (missing $p_T$ cut)	9
	3.2.8 (C9) Particle identification	9
	3.3 Cut flow	9
4		10
	4.1 Trigger efficiency	
	4.1.1 Online veto (BBC-small and ZDC veto)	
	4.1.2 RP triggering efficiency	
	4.1.3 Up and Down RP combination veto	
	4.2 Cuts efficiency	
	4.2.1 TPC z-vertex cut (C2)	
	4.2.2 TPC-RP z-vertex matching (C5)	
	4.2.3 Offline BBC-large veto (C6) and TOF clusters limit (C7)	
	4.2.4 Missing $p_T$ cut (C8)	
	4.2.5 Particle identification (C9)	
	4.3 RP track acceptance and reconstruction efficiency	
	·	10
	· · · · · · · · · · · · · · · · · · ·	10
	4.6 TPC vertex reconstruction efficiency	10
5	Backgrounds	11
J	~	11
		11
	7.2 Energiance background	11
6	Systematic errors	12

List of Figures	13
List of Tables	13
References	14

## 1. Introduction

#### 1.1 Central Exclusive Production

The Central Exclusive Production (CEP) takes place when interacting particles form in the mid-rapidity region a state ("central production") whose all constituents/decay products are measured in the detector ("exclusive"). The initial state particles can either dissociate, excite or stay intact. The latter case of CEP in proton-proton collisions can be written as

$$p + p \rightarrow p + X + p \tag{1.1}$$

and depicted as in Fig. 1.1. Mass and rapidity of state X is given by

$$M_X = \sqrt{s\left(\xi_1 \xi_2 \sin^2{(\alpha/2)} - (1 - \xi_1 - \xi_2) \cos^2{(\alpha/2)}\right)} \stackrel{\alpha = \pi}{=} \sqrt{s\xi_1 \xi_2}, \quad (1.2) \qquad y_X = \frac{1}{2} \ln \frac{\xi_1}{\xi_2}, \quad (1.3)$$

where  $\alpha$  is angle between scattered protons and  $\xi = (p_0 - p)/p_0$  is the fractional momentum loss of proton.

#### 1.2 Double Pomeron Exchange

Reaction from Eq. 1.1 can exhibit purely electromagnetic  $(\gamma-\gamma)$ , mixed  $(\gamma-\mathcal{O})$  or purely strong nature  $(\mathcal{O}-\mathcal{O})$ . The last type is dominant at RHIC energies. It is characterized by the lack of hard scale (if protons are scattered at small angles), therefore perturbative QCD cannot be applied and Regge theory [1] is used instead. An object  $\mathcal{O}$  does not have unequivocal QCD representation - in Regge formalism it is the so-called "trajectory" (Reggeon,  $\mathbb{R}$ ). Reggeon with quantum numbers of vacuum is called "Pomeron" ( $\mathbb{P}$ ) and  $\mathbb{P}$ - $\mathbb{P}$  reaction (Fig. 1.2) is called "Double Pomeron Exchange".

Processes involing IPomeron exchange are referred as diffraction due to cross-section in scattering angle resembling similar shape to instesity pattern of diffracted light. For low values of Mandelstam t (small scattering angles) cross-section takes exponential form

$$\frac{d\sigma}{d|t|} \propto e^{-B|t|},\tag{1.4}$$

where the slope parameter B reflects the size of target at which  $\mathbb{P}$ omerons scatter.

Diffractive events have specific property of the "rapidity gap" which is an angular region free of hadrons. In DIPE two such gaps are present, marked in Fig. 1.1 as  $\Delta \eta_1$  and  $\Delta \eta_2$ .

DIPE is a spin-parity filter - from the fact that scattered particles have all quantum numbers unchanged after the interaction, central states must satisfy

$$I^G J^{PC} = 0^+ \text{even}^{++}.$$
 (1.5)

The lowest order QCD picture of the Pomeron is a pair of oppositely colored gluons (colour singlet). This fact makes the DPE recognized as the gluon-rich environment process which should enhance production of the bound states of gluons ("glueballs") or hybrid mesons.

For detailed introduction to the topic of diffraction see Refs. [2,3].




Figure 1.1: Central Exclusive Production in  $\eta\text{-}\phi$  space.




Figure 1.2: Diagram of DIPE process.

#### 1.3 Physics motivation for the measurement

STAR collected in 2015 large dataset dedicated for the measurement of the Central Diffraction (DIPE in particular). Since that year the experiment was enriched with Roman Pot Phase II\* subsystem and thus gained possibility of detection of forward protons. It enabled studies of properties of the central state with respect to observables related to exchanged IPomerons. No such measurement was performed before at that high c.m.s. energy ( $\sqrt{s} = 200 \text{ GeV}$ , contamination from IReggeon exchanges is small) which makes it particularly attractive. A list of physics issues that can be covered with the study described in this note is briefly introduced below.

#### 1.3.1 DIPE differential cross-sections, mass spectrum

As stated in Sec. 1.2 DIPE is a soft process whose theoretical description is done mainly using phenomenological tools, thus measurement of differential cross-sections is needed to verify various production models.

The main focus is put on the simplest state (and most numerously) produced in DPE, namely a pair of oppositely charged pions,  $\pi^+\pi^-$ . It can be formed either in a non-resonant or resonant mechanism. In the first case the  $\pi^+\pi^-$  continuum is formed by the exchange of the off-shell pion between Pomerons. Currently there are two models of this reaction on the market [4,5], [6]. In the second case the Pomerons directly couple into resonance (e.g.  $f_2(1270)$ ), which then decays to  $\pi^+\pi^-$ . Attempts to calculate cross-section for this production mechanism are presented in Ref. [5] and [7].

Understanding of the mass spectrum in  $\pi^+\pi^-$  channel is important to learn about relative contribution from continuum and resonant production, as well as relative production of resonances. Recognition of resonant states may indicate candidates for low-mass glueballs of  $J^{PC}=0^{++}$ , however presence of underlaying scalar  $q\bar{q}$  states makes this task challenging.

Other channels, like  $K^+K^-$ , are also of great interest. Comparison of the cross-sections for production of  $\pi^+\pi^-$  and  $K^+K^-$  gives information about strength of the Pomeron coupling to different quark flavors. Also, structures in  $d\sigma/dm$  can be easier attributed to resonances by measuring more than one channel and known branching ratios thereof.

Detection of intact protons scattered at very small angle with respect to the beamline enables determination of the reaction plane which makes the Partial Wave Analysis (PWA) possible. It also allows to look at the the cross-sections more differentially, especially with respect to properties of exchanged Pomerons, like carried squared four-momentum t, azimuthal separation of the Pomerons in the transverse plane  $\Delta \varphi$  or relative momentum of Pomerons  $\Delta p_T$ . The last quantity was proposed to distinguish pure  $q\bar{q}$  states from these with gluonic content [8].

#### 1.3.2 Absorption effects

One can imagine in diagram in Fig. 1.2 additional soft lines e.g. between protons in the initial state or one of Pomerons and final state proton. These so-called rescattering effects (or absorption effects) lead to production of hadrons other than these belonging to central state X hence the diffractive signature of an event in form of rapidity gap is no longer present. Measurement of the probability that the state X will remain exclusive and forward protons will remain intact, in other words the rapidity gap survival probability  $S^2$ , would be valuable ingredient for development of absorption models.

#### 1.3.3 Size of interaction region

From the measurement of protons in Roman Pots one is able to reconstruct squared four-momenta transferred in proton-Pomeron vertices and determine the differential cross-section  $d\sigma/d|t|$ . Fit of exponent allows to extract the slope parameter B, which may depend on the Pomeron-Pomeron c.m.s. energy, or in other words on the mass of diffractive system X. Knowledge on the slope parameter gives insight to the volume and distribution of Pomerons inside proton.

## 2. Data set

#### 2.1 Trigger

The main trigger designed for studies of Central Diffraction in run 15 was RP\_CPT2. It was formed of the following conditions combined by logical AND (&&):

- 1. (ET && !IT) || (!ET && IT) = signal in at least one RP on each side of the STAR central detector to ensure presence of two forward-scattered protons; a veto was imposed on simultaneous signal in RPs above and below the beamline, which might have originated either from proton dissociation, or pile-up event, or beam halo proton etc.,
- 2. !BBCE && !BBCW && !ZDCE && !ZDCW = veto on any signal in small BBC tiles or ZDCs on any side of STAR central detector such requirement is in accordance with the double-gap topology of CEP events, it mostly filtered out CEP events with parallel pile-up event(s),
- 3.  $TOF \ge 2$  = at least 2 hits in TOF aim of this condition was to ensure activity in the mid-rapidity; since the lowest multiplicity allowed in CEP is 2, that was the lower threshold of L0 TOF multiplicity.

This trigger was running with an average prescale of 5 and average DAQ rate of 250 Hz, which allowed to collect in total about 560 M events corresponding to 16.5 pb<sup>-1</sup> of integrated luminosity. More information about number of events per run, rates etc. can be found under link provided in Ref. [9], which contains selected data from STAR run log [10]. Luminosity data used in this analysis comes from Ref. [11].

All RP triggers which were intended for usage in diffractive physics analyses or efficiency studies are listed in Tab. 2.1. Components used in definitions of these triggers are outlined in Fig. 2.1. Detailed explanation of all trigger bits can be found in Refs. [12,13]. Explanation of naming convention in Roman Pot system can be found in Ref. [14].




Figure 2.1: Sketch of the trigger components used in definitions of diffractive triggers in run 15.

Trigger name	Definition	Events [M] (available)	Comment
RP_CP	EOR && WOR	73.3 (46.0)	Loose trigger (mostly elastic events) designed for monitoring/trigger efficiency study
RP_CPT	EOR && WOR && !BBCE && !BBCW && !ZDCE && !ZDCW && TOF≥1	38.9 (0.0)	Intended to be main CEP trigger (later switched to RP_CPT2 due to large prescale)
RP_CPT2	(ET && !IT)    (!ET && IT) && !BBCE && !BBCW && !ZDCE && !ZDCW && TOF≥2	556.5 (493.8)	Main CEP trigger Note: On Apr 14 added upper TOF limit (10)
RP_CPX	IT && !BBCE && !BBCW && !ZDCE && !ZDCW && TOF≥2	40.1 (0.0)	The same as RP_CPT2 but only IT configuration
RP_CPEI	ET && IT && !BBCE && !BBCW && !ZDCE && !ZDCW && TOF>2	15.6 (11.8)	Control trigger for CPT2 to estimate effect of !(ET && IT) veto

Table 2.1: Central Diffraction physics triggers and control triggers involving Roman Pot detectors in run 15.

#### 2.2 Reconstruction software

Raw data was processed with STAR libraries in versions SL15k (trigger datasets: production\_pp200trans\_2015, production\_pp200long2\_2015, production\_pp200long3\_2015) and SL16d (production\_pp200long\_2015) [15].

#### 2.3 Data format

The analyzed data was stored in ROOT files in the picoDST format which was in large part a skimmed MuDST (standard STAR format). The picoDST format was introduced in Ref. [16]. PicoDST description files (C++ headers etc.) can be found in the analysis code repository [17].

#### 2.4 Bad runs

Analysis of CEP was performed on the data from runs with completion status "Successful" in the STAR run log. However, based on additional requirements explained below, some runs were omitted from analysis.

#### 2.4.1 RP distance from the beamline




Figure 2.2: Histogram of beam-detector distance |y| (2.2a) and graph showing run-dependence of |y| (2.2b) for all Roman Pots.

## 3. Event selection

Complete list of analysis cuts used for signal extraction is presented in Sec. 3.1. Detailed description of each cut can be found in Sec. 3.2.

#### 3.1 List of $cuts^1$

- C1. Exactly 1 primary vertex with TPC track(s) matched with hits in TOF.
- C2. TPC vertex from C1 is placed within  $|z_{\text{VX}}| < 100 \text{ cm}$ .
- C3. Exactly 2 opposite-sign primary TPC tracks (C3.2) of good quality and within high TPC acceptance region (C3.3) matched with hits in TOF (C3.1).
  - C3.1. Exactly 2 TOF-matched primary tracks,
  - C3.2. Tracks are of opposite signs,
  - C3.3. Both tracks satisfy criteria below:

$$|\eta|<1, \quad p_T>0.2~{\rm GeV}/c, \quad {\rm DCA}<2~{\rm cm}, \quad N_{\rm hits}^{\rm fit}\geq 25, \quad N_{\rm hits}^{\rm dE/dx}\geq 15, \quad N_{\rm hits}^{\rm fit}/N_{\rm hits}^{\rm poss}\geq 0.52,$$

- C4. Exactly 1 RP track on each side of STAR central detector (C4.3) of good quality (C4.1), with local angles consistent with the IP being the track origin (C4.2), lying within fiducial region of high geometrical acceptance (C4.4).
  - C4.1. RP tracks contain only track-points with at least 3 (out of 4) planes used in reconstrucion,
  - C4.2. Local angles [mrad] lie within  $-1.2 < \theta_x^{\text{RP}} < 5.0$  and  $1.5 < |\theta_y^{\text{RP}}| < 4.5$ ,
  - C4.3. Exactly 1 track passing cuts C4.1-C4.2 per side,
  - C4.4. Tracks passing cut C4.3 lie within the  $(p_x, p_y)$  region defined as:  $|p_y| > 0.2 \text{ GeV}/c$ ,  $(p_x/(\text{GeV}/c) + 0.3)^2 + (p_y/(\text{GeV}/c))^2 < 0.46^2$ ,  $(p_x/(\text{GeV}/c) 0.15)^2 + (p_y/(\text{GeV}/c))^2 < 0.42^2$ .
- C5. Vertex z-positions measured in TPC and reconstructed from the difference of proton detection time in west and east RPs are consistent with each other within the resolution  $|z_{\text{VX}}^{\text{TPC}} z_{\text{VX}}^{\text{RP}}| < 36 \text{ cm}$ .
- C6. No signal in any tile of BBC-large (east or west) with ADC > 40 and TDC > 100.
- C7. Maximally 2 reconstructed TOF clusters  $N_{\rm clstrs}^{\rm TOF} \leq 2$ .
- C8. Missing (total) momentum of TPC tracks and RP tracks  $p_T^{
  m miss} < 75~{
  m MeV}/c.$
- C9. Particle (pair) identification:

if 
$$n\sigma_{\mathrm{pion}}^{\mathrm{pair}} > 3$$
 &  $n\sigma_{\mathrm{kaon}}^{\mathrm{pair}} > 3$  &  $n\sigma_{\mathrm{proton}}^{\mathrm{pair}} < 3$  &  $m_{\mathrm{TOF}}^2 > 0.6 \; \mathrm{GeV}/c^2 \rightarrow pp$  elif  $n\sigma_{\mathrm{pion}}^{\mathrm{pair}} > 3$  &  $n\sigma_{\mathrm{kaon}}^{\mathrm{pair}} < 3$  &  $n\sigma_{\mathrm{proton}}^{\mathrm{pair}} > 3$  &  $m_{\mathrm{TOF}}^2 > 0.15 \; \mathrm{GeV}/c^2 \rightarrow KK$  elif  $|n\sigma_{\mathrm{pion}}^{\mathrm{trk1}}| < 3$  &  $|n\sigma_{\mathrm{pion}}^{\mathrm{trk2}}| < 3 \rightarrow \pi\pi$  else event rejected.

#### 3.2 Description of cuts

#### 3.2.1 (C1,C2) Primary vertex

Bla bla

<sup>&</sup>lt;sup>1</sup>Some cuts (e.g. C3) are decomposed to constituent sub-cuts. Cut is formed by the logical AND of all its sub-cuts.

- 3.2.2 (C3) TPC tracks
- 3.2.3 (C4) RP tracks
- 3.2.4 (C5) TPC-RP z-vertex matching
- 3.2.5 (C6) BBC-large signal veto
- 3.2.6 (C7) TOF clusters limit
- 3.2.7 (C8) Exclusivity cut (missing  $p_T$  cut)
- 3.2.8 (C9) Particle identification
- 3.3 Cut flow

# 4. Corrections

- 4.1 Trigger efficiency
- 4.1.1 Online veto (BBC-small and ZDC veto)
- 4.1.2 RP triggering efficiency
- 4.1.3 Up and Down RP combination veto
- 4.2 Cuts efficiency
- 4.2.1 TPC z-vertex cut (C2)
- 4.2.2 TPC-RP z-vertex matching (C5)
- 4.2.3 Offline BBC-large veto (C6) and TOF clusters limit (C7)
- 4.2.4 Missing  $p_T$  cut (C8)
- 4.2.5 Particle identification (C9)
- 4.3 RP track acceptance and reconstruction efficiency
- 4.4 TPC track acceptance and reconstruction efficiency
- 4.5 TOF acceptance, reconstruction and track-matching efficiency
- 4.6 TPC vertex reconstruction efficiency

# 5. Backgrounds

- 5.1 Non-exclusive background
- 5.2 Exclusive background

# 6. Systematic errors

# List of Figures

1.1	Central Exclusive Production in $\eta$ - $\phi$ space	4
1.2	Diagram of DIPE process.	4
2.1	Sketch of the trigger components used in definitions of diffractive triggers in run 15	6
2.2	Beam-detector distance of the Roman Pots in run 15	7

# List of Tables

2.1 Central Diffraction physics triggers and control triggers involving Roman Pot detectors in run 15. 6

## References

- P. D. B. Collins, An Introduction to Regge Theory and High-Energy Physics. Cambridge Monographs on Mathematical Physics. Cambridge Univ. Press, Cambridge, UK, 2009. http://www-spires.fnal.gov/spires/find/books/www?cl=QC793.3.R4C695.
- [2] S. Donnachie, H. G. Dosch, O. Nachtmann, and P. Landshoff, Pomeron physics and QCD, vol. 19. 2002.
- [3] V. Barone and E. Predazzi, *High-Energy Particle Diffraction*, vol. v.565 of *Texts and Monographs in Physics*. Springer-Verlag, Berlin Heidelberg, 2002. http://www-spires.fnal.gov/spires/find/books/www?cl=QC794.6.C6B37::2002.
- [4] P. Lebiedowicz and A. Szczurek, "Exclusive  $pp \to pp\pi^+\pi^-$  reaction: From the threshold to LHC," *Phys. Rev.* **D81** (2010) 036003, arXiv:0912.0190 [hep-ph].
- [5] P. Lebiedowicz, O. Nachtmann, and A. Szczurek, "Central exclusive diffractive production of  $\pi^+\pi^-$  continuum, scalar and tensor resonances in pp and  $p\bar{p}$  scattering within tensor pomeron approach," Phys. Rev. **D93** no. 5, (2016) 054015, arXiv:1601.04537 [hep-ph].
- [6] L. A. Harland-Lang, V. A. Khoze, and M. G. Ryskin, "Modelling exclusive meson pair production at hadron colliders," Eur. Phys. J. C74 (2014) 2848, arXiv:1312.4553 [hep-ph].
- [7] R. Fiore, L. Jenkovszky, and R. Schicker, "Resonance production in Pomeron-Pomeron collisions at the LHC," Eur. Phys. J. C76 no. 1, (2016) 38, arXiv:1512.04977 [hep-ph].
- [8] F. E. Close and A. Kirk, "A Glueball q anti-q filter in central hadron production," *Phys. Lett.* **B397** (1997) 333–338, arXiv:hep-ph/9701222 [hep-ph].
- [9] https://online.star.bnl.gov/rp/pp200/.
- [10] https://online.star.bnl.gov/RunLogRun15/.
- [11] http://www.star.bnl.gov/protected/common/common2015/trigger2015/lumipp200GeV/.
- [12] I. Alekseev and J. H. Lee, "Trigger Definitions for RP phase-II\* for Run15." https://drupal.star.bnl.gov/STAR/system/files/trigger\_updated\_run15\_v2.doc, December, 2014.
- [13] I. Alekseev, "Trigger with Roman Pots Phase II\*." https://drupal.star.bnl.gov/STAR/system/files/trigger4RP2\_v2.pdf, December, 2014.
- [14] R. Sikora, "Naming convention and layout of the Roman Pot Phase II\* subsystem in the STAR experiment." https://drupal.star.bnl.gov/STAR/system/files/RomanPotNomenclature\_0.pdf, October, 2014.
- [15] http://www.star.bnl.gov/public/comp/prod/DataSummary.html.
- [16] L. Adamczyk. https://drupal.star.bnl.gov/STAR/system/files/talk\_42.pdf, March, 2016.
- [17] https://github.com/rafalsikora/CEP-STAR.