

Algebraic Geometry in Complexity Theory

Fulvio Gesmundo

Institut de Mathématiques de Toulouse

Algebraic complexity theory studies complexity of algebraic problems:

- Evaluating polynomials;
- Computing products in algebraic structures (e.g., matrix multiplication);
- Solving polynomial systems.

Algebraic complexity theory studies complexity of algebraic problems:

- Evaluating polynomials;
- Computing products in algebraic structures (e.g., matrix multiplication);
- Solving polynomial systems.

General framework:

- Points in a vector space represent problems.
- The complexity is measured by some *geometric* invariant, for instance membership in an algebraic variety.

Complexity measures for Polynomials

Let $f \in S^d \mathbb{C}^N$ be a homogeneous polynomial of degree d in N variables.

Complexity measures for Polynomials

Let $f \in S^d \mathbb{C}^N$ be a homogeneous polynomial of degree d in N variables.

For instance:

$$h = x_1 x_2 x_3 + x_4 x_5 x_6 \in S^3 \mathbb{C}^6.$$

Complexity measures for Polynomials

Let $f \in S^d \mathbb{C}^N$ be a homogeneous polynomial of degree d in N variables.

For instance:

$$h = x_1 x_2 x_3 + x_4 x_5 x_6 \in S^3 \mathbb{C}^6.$$

We consider two important complexity measures (but there are many others):

- Waring rank.
- Algebraic Branching Program (ABP) Width.

Waring Rank

The **Waring rank** of f , denoted $\text{wr}(f)$, is the smallest r such that

$$f = \ell_1^d + \cdots + \ell_r^d$$

is a sum of powers of linear forms $\ell_i = a_{i1}x_1 + \cdots + a_{iN}x_N$.

Waring Rank

The **Waring rank** of f , denoted $\text{wr}(f)$, is the smallest r such that

$$f = \ell_1^d + \cdots + \ell_r^d$$

is a sum of powers of linear forms $\ell_i = a_{i1}x_1 + \cdots + a_{iN}x_N$.

For instance

$$\begin{aligned} h = x_1x_2x_3 + x_4x_5x_6 = \\ (x_1 + x_2 + x_3)^3 - (x_1 + x_2 - x_3)^3 - (x_1 - x_2 + x_3)^3 + (x_1 - x_2 - x_3)^3 + \\ (x_4 + x_5 + x_6)^3 - (x_4 + x_5 - x_6)^3 - (x_4 - x_5 + x_6)^3 + (x_4 - x_5 - x_6)^3 \end{aligned}$$

Waring Rank

The **Waring rank** of f , denoted $\text{wr}(f)$, is the smallest r such that

$$f = \ell_1^d + \cdots + \ell_r^d$$

is a sum of powers of linear forms $\ell_i = a_{i1}x_1 + \cdots + a_{iN}x_N$.

For instance

$$\begin{aligned} h = x_1x_2x_3 + x_4x_5x_6 = \\ (x_1 + x_2 + x_3)^3 - (x_1 + x_2 - x_3)^3 - (x_1 - x_2 + x_3)^3 + (x_1 - x_2 - x_3)^3 + \\ (x_4 + x_5 + x_6)^3 - (x_4 + x_5 - x_6)^3 - (x_4 - x_5 + x_6)^3 + (x_4 - x_5 - x_6)^3 \end{aligned}$$

shows $\text{wr}(h) \leq 8$ (and equality holds).

ABP Width

The **algebraic branching program width** of f , denoted $\text{abpw}(f)$, is the smallest w such that

$$f = [\ell_{1,1} \ \cdots \ \ell_{1,w}] \begin{bmatrix} \ell_{2,1,1} & \cdots & \\ \vdots & \ddots & \\ & & \ell_{2,w,w} \end{bmatrix} \cdots \begin{bmatrix} \ell_{d-1,1,1} & \cdots & \\ \vdots & \ddots & \\ & & \ell_{d-1,w,w} \end{bmatrix} \begin{bmatrix} \ell_{d,1} \\ \vdots \\ \ell_{d,w} \end{bmatrix}$$

is a product of matrices of size w whose entries are linear forms.

ABP Width

The **algebraic branching program width** of f , denoted $\text{abpw}(f)$, is the smallest w such that

$$f = [\ell_{1,1} \ \cdots \ \ell_{1,w}] \begin{bmatrix} \ell_{2,1,1} & \cdots & \\ \vdots & \ddots & \\ & & \ell_{2,w,w} \end{bmatrix} \cdots \begin{bmatrix} \ell_{d-1,1,1} & \cdots & \\ \vdots & \ddots & \\ & & \ell_{d-1,w,w} \end{bmatrix} \begin{bmatrix} \ell_{d,1} \\ \vdots \\ \ell_{d,w} \end{bmatrix}$$

is a product of matrices of size w whose entries are linear forms.

For instance

$$h = x_1 x_2 x_3 + x_4 x_5 x_6 =$$

$$\begin{bmatrix} x_1 & x_4 \end{bmatrix} \begin{bmatrix} x_2 & 0 \\ 0 & x_5 \end{bmatrix} \begin{bmatrix} x_3 \\ x_6 \end{bmatrix}$$

ABP Width

The **algebraic branching program width** of f , denoted $\text{abpw}(f)$, is the smallest w such that

$$f = [\ell_{1,1} \ \cdots \ \ell_{1,w}] \begin{bmatrix} \ell_{2,1,1} & \cdots & \\ \vdots & \ddots & \\ & & \ell_{2,w,w} \end{bmatrix} \cdots \begin{bmatrix} \ell_{d-1,1,1} & \cdots & \\ \vdots & \ddots & \\ & & \ell_{d-1,w,w} \end{bmatrix} \begin{bmatrix} \ell_{d,1} \\ \vdots \\ \ell_{d,w} \end{bmatrix}$$

is a product of matrices of size w whose entries are linear forms.

For instance

$$h = x_1 x_2 x_3 + x_4 x_5 x_6 =$$

$$\begin{bmatrix} x_1 & x_4 \end{bmatrix} \begin{bmatrix} x_2 & 0 \\ 0 & x_5 \end{bmatrix} \begin{bmatrix} x_3 \\ x_6 \end{bmatrix}$$

shows $\text{abpw}(h) \leq 2$ (and equality holds).

Complexity classes of Polynomials

A complexity class is a collection of (sequences of) polynomials such that a complexity measure has prescribed growth:

Complexity classes of Polynomials

A complexity class is a collection of (sequences of) polynomials such that a complexity measure has prescribed growth:

- $\text{VWaring} = \{(f_n)_{n \in \mathbb{N}} : f_n \in S^{d_n} \mathbb{C}^{N_n} \text{ with } d_n, N_n, \text{wr}(f_n) \text{ are poly-bounded}\}$

Complexity classes of Polynomials

A complexity class is a collection of (sequences of) polynomials such that a complexity measure has prescribed growth:

- $\text{VWaring} = \{(f_n)_{n \in \mathbb{N}} : f_n \in S^{d_n} \mathbb{C}^{N_n} \text{ with } d_n, N_n, \text{wr}(f_n) \text{ are poly-bounded}\}$
For instance

$$f_n = p_n = x_1^n + \cdots + x_n^n \in \text{VWaring}.$$

Complexity classes of Polynomials

A complexity class is a collection of (sequences of) polynomials such that a complexity measure has prescribed growth:

- $\text{VWaring} = \{(f_n)_{n \in \mathbb{N}} : f_n \in S^{d_n} \mathbb{C}^{N_n} \text{ with } d_n, N_n, \text{wr}(f_n) \text{ are poly-bounded}\}$
For instance

$$f_n = p_n = x_1^n + \cdots + x_n^n \in \text{VWaring}.$$

- $\text{VBP} = \{(f_n)_{n \in \mathbb{N}} : f_n \in S^{d_n} \mathbb{C}^{N_n} \text{ with } d_n, N_n, \text{abpw}(f_n) \text{ are poly-bounded}\}$

Complexity classes of Polynomials

A complexity class is a collection of (sequences of) polynomials such that a complexity measure has prescribed growth:

- $\text{VWaring} = \{(f_n)_{n \in \mathbb{N}} : f_n \in S^{d_n} \mathbb{C}^{N_n} \text{ with } d_n, N_n, \text{wr}(f_n) \text{ are poly-bounded}\}$
For instance

$$f_n = p_n = x_1^n + \cdots + x_n^n \in \text{VWaring}.$$

- $\text{VBP} = \{(f_n)_{n \in \mathbb{N}} : f_n \in S^{d_n} \mathbb{C}^{N_n} \text{ with } d_n, N_n, \text{abpw}(f_n) \text{ are poly-bounded}\}$
For instance

$$f_n = m_n = x_1 \cdots x_n \in \text{VBP}$$

$$f_n = p_n \in \text{VBP} \quad \text{using diagonal matrices}$$

$$f_n = \det_n \in \text{VBP}$$

Complexity classes of Polynomials

A complexity class is a collection of (sequences of) polynomials such that a complexity measure has prescribed growth:

- $\text{VWaring} = \{(f_n)_{n \in \mathbb{N}} : f_n \in S^{d_n} \mathbb{C}^{N_n} \text{ with } d_n, N_n, \text{wr}(f_n) \text{ are poly-bounded}\}$
For instance

$$f_n = p_n = x_1^n + \cdots + x_n^n \in \text{VWaring}.$$

- $\text{VBP} = \{(f_n)_{n \in \mathbb{N}} : f_n \in S^{d_n} \mathbb{C}^{N_n} \text{ with } d_n, N_n, \text{abpw}(f_n) \text{ are poly-bounded}\}$
For instance

$$f_n = m_n = x_1 \cdots x_n \in \text{VBP}$$

$$f_n = p_n \in \text{VBP} \quad \text{using diagonal matrices}$$

$$f_n = \det_n \in \text{VBP}$$

Conjecturally hard polynomial. $\text{perm}_n = \sum_{\sigma \in S_n} x_{1\sigma_1} \cdots x_{n\sigma_n}$

Complexity classes of Polynomials

A complexity class is a collection of (sequences of) polynomials such that a complexity measure has prescribed growth:

- $\text{VWaring} = \{(f_n)_{n \in \mathbb{N}} : f_n \in S^{d_n} \mathbb{C}^{N_n} \text{ with } d_n, N_n, \text{wr}(f_n) \text{ are poly-bounded}\}$
For instance

$$f_n = p_n = x_1^n + \cdots + x_n^n \in \text{VWaring}.$$

- $\text{VBP} = \{(f_n)_{n \in \mathbb{N}} : f_n \in S^{d_n} \mathbb{C}^{N_n} \text{ with } d_n, N_n, \text{abpw}(f_n) \text{ are poly-bounded}\}$
For instance

$$f_n = m_n = x_1 \cdots x_n \in \text{VBP}$$

$$f_n = p_n \in \text{VBP} \quad \text{using diagonal matrices}$$

$$f_n = \det_n \in \text{VBP}$$

Conjecturally hard polynomial. $\text{perm}_n = \sum_{\sigma \in S_n} x_{1\sigma_1} \cdots x_{n\sigma_n}$

Valiant's Flagship Conjecture

$\text{perm}_n \notin \text{VBP}$, or equivalently $\text{abpw}(\text{perm}_n)$ is superpolynomial.

[If this conjecture is false, then $P = NP$]

Why geometry?

Let $c(-)$ be one of the complexity measures we defined. Define

$$C^\circ(r) = \left\{ f \in S^d \mathbb{C}^N : c(f) \leq r \right\}.$$

Why geometry?

Let $c(-)$ be one of the complexity measures we defined. Define

$$C^\circ(r) = \left\{ f \in S^d \mathbb{C}^N : c(f) \leq r \right\}.$$

$C^\circ(r)$ is a *constructible* set. In particular, the closure

$$C(r) = \overline{C^\circ(r)}$$

is an algebraic variety, i.e. the solution set of a system of polynomial equations.

Why geometry?

Let $c(-)$ be one of the complexity measures we defined. Define

$$C^\circ(r) = \left\{ f \in S^d \mathbb{C}^N : c(f) \leq r \right\}.$$

$C^\circ(r)$ is a *constructible* set. In particular, the closure

$$C(r) = \overline{C^\circ(r)}$$

is an algebraic variety, i.e. the solution set of a system of polynomial equations.

- Non-membership:
if $h \notin C(r)$ then $h \notin C^\circ(r)$, hence $c(h) > r$.

We can use geometry for lower bounds.

Why geometry?

Let $c(-)$ be one of the complexity measures we defined. Define

$$C^\circ(r) = \left\{ f \in S^d \mathbb{C}^N : c(f) \leq r \right\}.$$

$C^\circ(r)$ is a *constructible* set. In particular, the closure

$$C(r) = \overline{C^\circ(r)}$$

is an algebraic variety, i.e. the solution set of a system of polynomial equations.

- Non-membership:
if $h \notin C(r)$ then $h \notin C^\circ(r)$, hence $c(h) > r$.

We can use geometry for lower bounds.

- Membership:
if $h \in C(r)$ then $h = \lim_{\varepsilon \rightarrow 0} h_\varepsilon$ with $c(h_\varepsilon) \leq r$.

We can use geometry for upper bounds.

Strategy for lower bounds

To prove $c(h) > r$, we look for polynomial equations on $S^d\mathbb{C}^N$ such that:

- they vanishes on $C(r)$ (e.g. on $\{f \in S^4\mathbb{C}^N : \text{abpw}(f) \leq 3\}$);
- does not vanish on h (e.g. on $h = \text{perm}_4$).

Strategy for lower bounds

To prove $c(h) > r$, we look for polynomial equations on $S^d\mathbb{C}^N$ such that:

- they vanishes on $C(r)$ (e.g. on $\{f \in S^4\mathbb{C}^N : \text{abpw}(f) \leq 3\}$);
- does not vanish on h (e.g. on $h = \text{perm}_4$).

If we find one, we obtain the desired lower bound (e.g. $\text{abpw}(\text{perm}_4) \geq 4$).

Strategy for lower bounds

To prove $c(h) > r$, we look for polynomial equations on $S^d\mathbb{C}^N$ such that:

- they vanish on $C(r)$ (e.g. on $\{f \in S^4\mathbb{C}^N : \text{abpw}(f) \leq 3\}$);
- does not vanish on h (e.g. on $h = \text{perm}_4$).

If we find one, we obtain the desired lower bound (e.g. $\text{abpw}(\text{perm}_4) \geq 4$).

Often, instead of explicit polynomial equations we simply study geometric conditions which can be “translated” into equations (even if we rarely do this translation).

Lower bounds for Waring rank

If $c(-) = \text{wr}(-)$ then $C(r)$ is the r -th secant variety of the Veronese variety.
Varieties studied classically by Cayley, Sylvester, Segre, Terracini, etc.

Lower bounds for Waring rank

If $c(-) = \text{wr}(-)$ then $C(r)$ is the r -th secant variety of the Veronese variety. Varieties studied classically by Cayley, Sylvester, Segre, Terracini, etc.

Easy theorem for lower bounds.

If $f \in C(r)$ then the space of k -th order partial derivatives of f has dimension at most r .

Consequence.

If the k -th order partials of f span a space of dimension r , then $\text{wr}(f) \geq r$.

Lower bounds for Waring rank

If $c(-) = \text{wr}(-)$ then $C(r)$ is the r -th secant variety of the Veronese variety. Varieties studied classically by Cayley, Sylvester, Segre, Terracini, etc.

Easy theorem for lower bounds.

If $f \in C(r)$ then the space of k -th order partial derivatives of f has dimension at most r .

Consequence.

If the k -th order partials of f span a space of dimension r , then $\text{wr}(f) \geq r$.

Note.

This is a (matrix) rank condition.

Lower bounds for Waring rank

If $c(-) = \text{wr}(-)$ then $C(r)$ is the r -th secant variety of the Veronese variety. Varieties studied classically by Cayley, Sylvester, Segre, Terracini, etc.

Easy theorem for lower bounds.

If $f \in C(r)$ then the space of k -th order partial derivatives of f has dimension at most r .

Consequence.

If the k -th order partials of f span a space of dimension r , then $\text{wr}(f) \geq r$.

Note.

This is a (matrix) rank condition.

More complicated rank conditions:

- Young flattenings \rightarrow representation theory;
- homological methods \rightarrow commutative algebra;
- apolarity \rightarrow deformation theory.

Lower bounds for Waring rank – cont'd

We use this strategy to prove some theorems:

Theorem.

The sequence $m_n = x_1 \cdots x_n$ is not in VWaring .

Lower bounds for Waring rank – cont'd

We use this strategy to prove some theorems:

Theorem.

The sequence $m_n = x_1 \cdots x_n$ is not in VWaring .

Proof.

For $k < n$, the space of k -th order partials is spanned by the square-free monomials of degree $n - k$ in n variables

Lower bounds for Waring rank – cont'd

We use this strategy to prove some theorems:

Theorem.

The sequence $m_n = x_1 \cdots x_n$ is not in VWaring .

Proof.

For $k < n$, the space of k -th order partials is spanned by the square-free monomials of degree $n - k$ in n variables:

$$\dim \langle k\text{-th order partials of } m_n \rangle = \binom{n}{n-k}.$$

Lower bounds for Waring rank – cont'd

We use this strategy to prove some theorems:

Theorem.

The sequence $m_n = x_1 \cdots x_n$ is not in VWaring .

Proof.

For $k < n$, the space of k -th order partials is spanned by the square-free monomials of degree $n - k$ in n variables:

$$\dim \langle k\text{-th order partials of } m_n \rangle = \binom{n}{n-k}.$$

If $k \approx n/2$ then this is exponential as a function of n .

Therefore $\text{wr}(m_n)$ is not poly-bounded as a function of n .

Lower bounds for Waring rank – cont'd

We use this strategy to prove some theorems:

Theorem.

The sequence $m_n = x_1 \cdots x_n$ is not in VWaring .

Proof.

For $k < n$, the space of k -th order partials is spanned by the square-free monomials of degree $n - k$ in n variables:

$$\dim \langle k\text{-th order partials of } m_n \rangle = \binom{n}{n-k}.$$

If $k \approx n/2$ then this is exponential as a function of n .

Therefore $\text{wr}(m_n)$ is not poly-bounded as a function of n .

Corollary.

$\text{VWaring} \neq \text{VBP}$.

Proof.

m_n belongs to VBP but not to VWaring .

Lower bounds for ABP width

We study geometric properties of the *hypersurface*

$$\mathcal{Z}(f) = \{(x_1, \dots, x_N) \in \mathbb{C}^N : f(x_1, \dots, x_N) = 0\}.$$

Lower bounds for ABP width

We study geometric properties of the *hypersurface*

$$\mathcal{Z}(f) = \{(x_1, \dots, x_N) \in \mathbb{C}^N : f(x_1, \dots, x_N) = 0\}.$$

Theorem. [G.-Ghosal-Ikenmeyer-Lysikov (2022) – restricted version]
If $\text{abpw}(f) \leq w$ then $\{f = 0\} \subseteq \mathbb{C}^N$ contains a linear space of dimension at least $N - w$.

Lower bounds for ABP width

We study geometric properties of the *hypersurface*

$$\mathcal{Z}(f) = \{(x_1, \dots, x_N) \in \mathbb{C}^N : f(x_1, \dots, x_N) = 0\}.$$

Theorem. [G.-Ghosal-Ikenmeyer-Lysikov (2022) – restricted version]
If $\text{abpw}(f) \leq w$ then $\{f = 0\} \subseteq \mathbb{C}^N$ contains a linear space of dimension at least $N - w$.

Idea of the Proof.

From an ABP, get an expression $f = \ell_1 g_1 + \dots + \ell_w g_w$ with $\deg(\ell_j) = 1$. Then

$$\{\ell_1 = \dots = \ell_w = 0\} \subseteq \mathcal{Z}(f).$$

Lower bounds for ABP width

We study geometric properties of the *hypersurface*

$$\mathcal{Z}(f) = \{(x_1, \dots, x_N) \in \mathbb{C}^N : f(x_1, \dots, x_N) = 0\}.$$

Theorem. [G.-Ghosal-Ikenmeyer-Lysikov (2022) – restricted version]
If $\text{abpw}(f) \leq w$ then $\{f = 0\} \subseteq \mathbb{C}^N$ contains a linear space of dimension at least $N - w$.

Idea of the Proof.

From an ABP, get an expression $f = \ell_1 g_1 + \dots + \ell_w g_w$ with $\deg(\ell_j) = 1$. Then

$$\{\ell_1 = \dots = \ell_w = 0\} \subseteq \mathcal{Z}(f).$$

Open problem.

Determine a geometric method which can potentially give super-polynomial lower bounds on abpw .

Matrix multiplication complexity

How costly is it to multiply matrices?

Matrix multiplication complexity

How costly is it to multiply matrices?

$$\begin{aligned}\mu_n : \text{Mat}_{n \times n} \times \text{Mat}_{n \times n} &\rightarrow \text{Mat}_{n \times n} \\ (A, B) &\mapsto AB\end{aligned}$$

Matrix multiplication complexity

How costly is it to multiply matrices?

$$\begin{aligned}\mu_n : \text{Mat}_{n \times n} \times \text{Mat}_{n \times n} &\rightarrow \text{Mat}_{n \times n} \\ (A, B) &\mapsto AB\end{aligned}$$

There is an associated tensor $\mu_n \in \mathbb{C}^{n^2} \otimes \mathbb{C}^{n^2} \otimes \mathbb{C}^{n^2}$.

Matrix multiplication complexity

How costly is it to multiply matrices?

$$\begin{aligned}\mu_n : \text{Mat}_{n \times n} \times \text{Mat}_{n \times n} &\rightarrow \text{Mat}_{n \times n} \\ (A, B) &\mapsto AB\end{aligned}$$

There is an associated tensor $\mu_n \in \mathbb{C}^{n^2} \otimes \mathbb{C}^{n^2} \otimes \mathbb{C}^{n^2}$.

When $n = 2$, this is

$$\begin{aligned}\mu_2 = & (a_{11} \otimes b_{11} + a_{12} \otimes b_{21}) \otimes c_{11} + \\ & (a_{21} \otimes b_{11} + a_{22} \otimes b_{21}) \otimes c_{21} + \\ & (a_{11} \otimes b_{12} + a_{12} \otimes b_{22}) \otimes c_{12} + \\ & (a_{21} \otimes b_{12} + a_{22} \otimes b_{22}) \otimes c_{22}\end{aligned}$$

Matrix multiplication complexity

How costly is it to multiply matrices?

$$\begin{aligned}\mu_n : \text{Mat}_{n \times n} \times \text{Mat}_{n \times n} &\rightarrow \text{Mat}_{n \times n} \\ (A, B) &\mapsto AB\end{aligned}$$

There is an associated tensor $\mu_n \in \mathbb{C}^{n^2} \otimes \mathbb{C}^{n^2} \otimes \mathbb{C}^{n^2}$.

When $n = 2$, this is

$$\begin{aligned}\mu_2 = & (a_{11} \otimes b_{11} + a_{12} \otimes b_{21}) \otimes c_{11} + \\ & (a_{21} \otimes b_{11} + a_{22} \otimes b_{21}) \otimes c_{21} + \\ & (a_{11} \otimes b_{12} + a_{12} \otimes b_{22}) \otimes c_{12} + \\ & (a_{21} \otimes b_{12} + a_{22} \otimes b_{22}) \otimes c_{22}\end{aligned}$$

The **tensor rank** is the relevant measure of complexity: if $T \in U \otimes V \otimes W$ then

$$R(T) = \min\{r : T = u_1 \otimes v_1 \otimes w_1 + \cdots + u_r \otimes v_r \otimes w_r\}.$$

Matrix multiplication complexity

How costly is it to multiply matrices?

$$\begin{aligned}\mu_n : \text{Mat}_{n \times n} \times \text{Mat}_{n \times n} &\rightarrow \text{Mat}_{n \times n} \\ (A, B) &\mapsto AB\end{aligned}$$

There is an associated tensor $\mu_n \in \mathbb{C}^{n^2} \otimes \mathbb{C}^{n^2} \otimes \mathbb{C}^{n^2}$.

When $n = 2$, this is

$$\begin{aligned}\mu_2 = & (a_{11} \otimes b_{11} + a_{12} \otimes b_{21}) \otimes c_{11} + \\ & (a_{21} \otimes b_{11} + a_{22} \otimes b_{21}) \otimes c_{21} + \\ & (a_{11} \otimes b_{12} + a_{12} \otimes b_{22}) \otimes c_{12} + \\ & (a_{21} \otimes b_{12} + a_{22} \otimes b_{22}) \otimes c_{22}\end{aligned}$$

The **tensor rank** is the relevant measure of complexity: if $T \in U \otimes V \otimes W$ then

$$R(T) = \min\{r : T = u_1 \otimes v_1 \otimes w_1 + \cdots + u_r \otimes v_r \otimes w_r\}.$$

Conjecture. The exponent ω of matrix multiplication is 2:

$$R(\mu_n) = O(n^{2+\varepsilon}) \text{ for every } \varepsilon.$$

Overview of upper bounds on ω

- Classical: $R(\mu_n) \leq n^3$ so $\omega \leq 3$.

Overview of upper bounds on ω

- Classical: $R(\mu_n) \leq n^3$ so $\omega \leq 3$.
- [Strassen 1969] $R(\mu_2) \leq 7$;
Consequence: $\omega \leq \log_2(7) \approx 2.81$.

Overview of upper bounds on ω

- Classical: $R(\mu_n) \leq n^3$ so $\omega \leq 3$.

- [Strassen 1969] $R(\mu_2) \leq 7$;

Consequence: $\omega \leq \log_2(7) \approx 2.81$.

- [Bini-Capovani-Lotti-Romani, 1979]

ω is controlled by membership in a variety

$$C(r) = \overline{\{T \in U \otimes V \otimes W : R(T) \leq r\}}.$$

Membership is sufficient for upper bounds! $\omega \leq 2.78$.

Overview of upper bounds on ω

- Classical: $R(\mu_n) \leq n^3$ so $\omega \leq 3$.

- [Strassen 1969] $R(\mu_2) \leq 7$;

Consequence: $\omega \leq \log_2(7) \approx 2.81$.

- [Bini-Capovani-Lotti-Romani, 1979]

ω is controlled by membership in a variety

$$C(r) = \overline{\{T \in U \otimes V \otimes W : R(T) \leq r\}}.$$

Membership is sufficient for upper bounds! $\omega \leq 2.78$.

- [Schönhage, 1981]

Block matrix multiplication tensor to give bounds on a full one: $\omega \leq 2.53$.

Overview of upper bounds on ω

- Classical: $R(\mu_n) \leq n^3$ so $\omega \leq 3$.

- [Strassen 1969] $R(\mu_2) \leq 7$;

Consequence: $\omega \leq \log_2(7) \approx 2.81$.

- [Bini-Capovani-Lotti-Romani, 1979]
 ω is controlled by membership in a variety

$$C(r) = \overline{\{T \in U \otimes V \otimes W : R(T) \leq r\}}.$$

Membership is sufficient for upper bounds! $\omega \leq 2.78$.

- [Schönhage, 1981]
Block matrix multiplication tensor to give bounds on a full one: $\omega \leq 2.53$.
- [Strassen, 1987] Laser method: Find *intermediate tensors* to simulate matrix multiplication: $\omega \leq 2.48$.

Overview of upper bounds on ω

- Classical: $R(\mu_n) \leq n^3$ so $\omega \leq 3$.
- [Strassen 1969] $R(\mu_2) \leq 7$;
Consequence: $\omega \leq \log_2(7) \approx 2.81$.
- [Bini-Capovani-Lotti-Romani, 1979]
 ω is controlled by membership in a variety

$$C(r) = \overline{\{T \in U \otimes V \otimes W : R(T) \leq r\}}.$$

Membership is sufficient for upper bounds! $\omega \leq 2.78$.

- [Schönhage, 1981]
Block matrix multiplication tensor to give bounds on a full one: $\omega \leq 2.53$.
- [Strassen, 1987] Laser method: Find *intermediate tensors* to simulate matrix multiplication: $\omega \leq 2.48$.
- [Coppersmith-Winograd, 1990] $\omega \leq 2.39$.

Overview of upper bounds on ω

- Classical: $R(\mu_n) \leq n^3$ so $\omega \leq 3$.
- [Strassen 1969] $R(\mu_2) \leq 7$;
Consequence: $\omega \leq \log_2(7) \approx 2.81$.
- [Bini-Capovani-Lotti-Romani, 1979]
 ω is controlled by membership in a variety

$$C(r) = \overline{\{T \in U \otimes V \otimes W : R(T) \leq r\}}.$$

Membership is sufficient for upper bounds! $\omega \leq 2.78$.

- [Schönhage, 1981]
Block matrix multiplication tensor to give bounds on a full one: $\omega \leq 2.53$.
- [Strassen, 1987] Laser method: Find *intermediate tensors* to simulate matrix multiplication: $\omega \leq 2.48$.
- [Coppersmith-Winograd, 1990] $\omega \leq 2.39$.
- More improvements by Stothers, Williams, LeGall, Alman-Williams, ...
[Williams-Xu-Xu-Zhou, 2023] $\omega \leq 2.371552$.

Barriers for geometric methods

How far can geometric methods take us?

Barriers for geometric methods

How far can geometric methods take us?

- [Galazka, Efremenko-Garg-Oliveira-Wigderson]
Flattening methods give non-membership for much bigger varieties than the $C(r)$'s.

We cannot hope to prove lower bounds when these varieties fill the ambient space.

[Buczyński, Ballico-Bernardi-G.]

We estimate how far. If $T \in \mathbb{C}^N \otimes \mathbb{C}^N \otimes \mathbb{C}^N$, cannot prove $R(T) \geq 6N$.

Barriers for geometric methods

How far can geometric methods take us?

- [Galazka, Efremenko-Garg-Oliveira-Wigderson]
Flattening methods give non-membership for much bigger varieties than the $C(r)$'s.

We cannot hope to prove lower bounds when these varieties fill the ambient space.

[Buczyński, Ballico-Bernardi-G.]

We estimate how far. If $T \in \mathbb{C}^N \otimes \mathbb{C}^N \otimes \mathbb{C}^N$, cannot prove $R(T) \geq 6N$.

- [Bürgisser-Ikenmeyer-Panova, G.-Ikenmeyer-Panova, Efremenko-Landsberg-Schenck-Weyman, G.-Landsberg]
The *geometric complexity theory program*, in some of its “easy forms”, cannot prove super polynomial lower bounds for abpw

Barriers for geometric methods

How far can geometric methods take us?

- [Galazka, Efremenko-Garg-Oliveira-Wigderson]
Flattening methods give non-membership for much bigger varieties than the $C(r)$'s.

We cannot hope to prove lower bounds when these varieties fill the ambient space.

[Buczyński, Ballico-Bernardi-G.]

We estimate how far. If $T \in \mathbb{C}^N \otimes \mathbb{C}^N \otimes \mathbb{C}^N$, cannot prove $R(T) \geq 6N$.

- [Bürgisser-Ikenmeyer-Panova, G.-Ikenmeyer-Panova, Efremenko-Landsberg-Schenck-Weyman, G.-Landsberg]
The *geometric complexity theory program*, in some of its “easy forms”, cannot prove super polynomial lower bounds for abpw
- [Alman-Williams, Ambainis-Filmus-LeGall, Christandl-Vrana-Zuiddam]
Strassen's laser method, applied to the original Coppersmith-Winograd tensor, cannot prove $\omega < 2.3$.

What can we do?

Lower bounds based on new geometric invariants:

- [Landsberg-Manivel-Ressayre] Degeneracy of dual variety
- [G.-Ghosal-Ikenmeyer-Ghosal] Noether-Lefschetz type conditions
- [Iarrobino, Buczyńska-Buczyński, Jelisiejew-Landsberg-Pal] Classical apolarity, Hilbert scheme of points, deformation theory.

What can we do?

Lower bounds based on new geometric invariants:

- [Landsberg-Manivel-Ressayre] Degeneracy of dual variety
- [G.-Ghosal-Ikenmeyer-Ghosal] Noether-Lefschetz type conditions
- [Iarrobino, Buczyńska-Buczyński, Jelisiejew-Landsberg-Pal] Classical apolarity, Hilbert scheme of points, deformation theory.

Upper bounds via more refined version of laser method:

- [Conner-G.-Landsberg-Ventura]
Use geometry to find better intermediate tensors.
- [Homs-Michałek-Jelisiejew-Seynnaeve]
Refine the laser method using ideas from commutative algebra.

Conclusions and more open directions

- Complexity is (almost) controlled by membership in certain sets $C(r)$
 - Lower bounds via geometric obstructions to membership.
 - Upper bounds via (asymptotic) membership.
- Barriers: Geometric methods have structural limitations.

Conclusions and more open directions

- Complexity is (almost) controlled by membership in certain sets $C(r)$
 - Lower bounds via geometric obstructions to membership.
 - Upper bounds via (asymptotic) membership.
- Barriers: Geometric methods have structural limitations.
- Boundary: To what extent taking closure matters? Debordering in complexity theory

Conclusions and more open directions

- Complexity is (almost) controlled by membership in certain sets $C(r)$
 - Lower bounds via geometric obstructions to membership.
 - Upper bounds via (asymptotic) membership.
- Barriers: Geometric methods have structural limitations.
- Boundary: To what extent taking closure matters? Debordering in complexity theory
- Asymptotic geometry: Strassen developed an asymptotic spectrum of tensors to study matrix multiplication via the laser method. Can we use similar ideas in other settings?