

MA 166: Statistics

Homework 2 (v1.1) ¹

Assigned Monday 31 January 2022

Due Monday 7 February 2022 at 11:59 pm EDT.

Read the footnotes! They give important information about how to present your answers.

1. Suppose that you have a priori knowledge that the continuous random variable X is normally distributed. You make n experimental measurements of X , and you find that fn of the measurements are (miraculously) exactly equal to $+1$, and the other $(1-f)n$ measurements are exactly equal to -1 . The fraction f is greater than $1/2$, so you think that X has a positive mean, but you are really not sure so you decide to do interval estimation to see with what confidence you can make that claim. You may assume that n is large enough that the Central Limit Theorem applies to a good approximation.
 - (a) Use maximum likelihood estimators to find estimates of the mean, μ_e , and the standard deviation, σ_e .
 - (b) We are going to use μ_e and σ_e as the basis for interval estimation in the remainder of this problem ². Suppose that you find you are able to conclude that the mean of X is positive with confidence probability $100(1-\alpha)\%$. Find an expression for n in terms of f and α ³.
 - (c) To make the above concrete, suppose that $f = 0.51$. How large does n need to be to achieve 95% confidence that the mean of X is positive?
2. Suppose that you have a priori knowledge that the continuous random variable Y has the following two-parameter probability density function:

$$f_Y(y) = \begin{cases} \frac{1}{\mu-a} \exp\left(-\frac{y-a}{\mu-a}\right) & \text{for } y \geq a \\ 0 & \text{otherwise,} \end{cases}$$

where it may be assumed that $a \geq 0$ and $\mu > a$.

- (a) Verify that the density function is normalized, and find the theoretical mean and standard deviation of Y in terms of the parameters a and μ .
- (b) Find maximum likelihood estimators, \hat{a}_{mle} and $\hat{\mu}_{\text{mle}}$, for the parameters a and μ . Justify your reasoning carefully, particularly for the calculation of \hat{a}_{mle} .

¹©2022, Bruce M. Boghosian, all rights reserved.

²As we shall see later, σ_e is not the optimum value to use for interval estimation, and the sample standard deviation for the normal distribution is usually defined in a different way. We have not yet covered that material however, so for the purposes of this problem you may proceed as instructed.

³Feel free to use the z_α notation described in the text on page 298

- (c) Now suppose that you take n samples of this data, y_1, \dots, y_n , where n is large enough that the Central Limit Theorem applies to a good approximation. (You may assume that the sample mean that you find is positive.) Find the $100(1-\alpha)\%$ confidence interval for $\mu = E(Y)$ centered at the sample mean. You may leave your answer in terms of \hat{a}_{mle} and $\hat{\mu}_{\text{mle}}$ worked out in part (b).

3. Larsen & Marx, Problem 5.3.6, page 306
4. Larsen & Marx, Problem 5.3.15, page 307