

Properties of Estimators

Unbiasedness

Bruce M. Boghosian



Tufts
UNIVERSITY

School of Arts
and Sciences

Department of Mathematics

Tufts University

©2022, all rights reserved

Bruce M.
Boghosian

Properties of
estimators

Unbiasedness
Example 1

Unbiasedness
Example 2

Unbiasedness
Example 3

Summary

- 1 Properties of estimators
- 2 Unbiasedness Example 1
- 3 Unbiasedness Example 2
- 4 Unbiasedness Example 3
- 5 Summary

Bruce M.
Boghossian

Properties of
estimators

Unbiasedness
Example 1

Unbiasedness
Example 2

Unbiasedness
Example 3

Summary

- For the example of the uniform distribution,

$$f_X(x) = \begin{cases} \frac{1}{b-a} & \text{for } a \leq x \leq b \\ 0 & \text{otherwise} \end{cases}$$

- the MLE estimator was $\hat{a}(\vec{X}) = \min_j X_j$
- the MM estimator was $\hat{a}(\vec{X}) = M_1 - \sqrt{3} \sqrt{M_2 - M_1^2}$, for sample moments

$$M_1 = \frac{1}{n} \sum_j^n X_j \quad \text{and} \quad M_2 = \frac{1}{n} \sum_j^n X_j^2$$

- Which one is “right”?
- There is no single answer to that question. We must instead identify desirable properties of estimators, and see which estimators have which of those properties.

Estimators are themselves random variables!

Bruce M.
Boghossian

Properties of
estimators

Unbiasedness
Example 1

Unbiasedness
Example 2

Unbiasedness
Example 3

Summary

- Estimators such as $\hat{a}(\vec{X})$, are functions of r.v.s.
- Hence estimators are themselves random variables.
- They presumably have probability density functions, though it is not always easy to figure out what they are.
- For estimating the sample mean $\hat{\mu}(\vec{X})$, we were able to use the CLT to study its distribution
- For other estimators, that approach may not be possible
- Because they have density functions, however, we know certain things about them:
 - They have means. We may speak of $E(\hat{a}(\vec{X}))$ & $E(\hat{\mu}(\vec{X}))$.
 - They have variances. We may speak of $\sigma_{\hat{a}}^2$ & $\sigma_{\hat{\mu}}^2$

Bruce M.
Boghosian

Properties of
estimators

Unbiasedness
Example 1

Unbiasedness
Example 2

Unbiasedness
Example 3

Summary

- **Unbiasedness:** Suppose that $\vec{Y} = \{Y_1, Y_2, \dots, Y_n\}$ is a random sample from the continuous PDF $f_Y(y; \theta)$, where θ is an unknown parameter. An estimator $\hat{\theta}(\vec{Y})$ is said to be *unbiased* for θ if $E(\hat{\theta}) = \theta$ for all θ . A similar definition holds for discrete random variables.
- **Efficiency:** Let $\hat{\theta}_1$ and $\hat{\theta}_2$ be two unbiased estimators for parameter θ . If $\text{Var}(\hat{\theta}_1) < \text{Var}(\hat{\theta}_2)$, we say that $\hat{\theta}_1$ is *more efficient* than $\hat{\theta}_2$.
- **Relative efficiency:** The *relative efficiency* of $\hat{\theta}_1$ with respect to $\hat{\theta}_2$ is $\frac{\text{Var}(\hat{\theta}_2)}{\text{Var}(\hat{\theta}_1)}$.

Bruce M.
Boghosian

Properties of
estimators

Unbiasedness
Example 1


Unbiasedness
Example 2

Unbiasedness
Example 3

Summary

- Consider the one-parameter probability density function

$$f_Y(y; \theta) = \begin{cases} \frac{2y}{\theta^2} & \text{if } 0 \leq y \leq \theta \\ 0 & \text{otherwise} \end{cases}$$



- Normalization: $\int_0^\theta dy \frac{2y}{\theta^2} = 1$
- Mean: $\mu = \int_0^\theta dy \frac{2y}{\theta^2} y = \frac{2}{3}\theta$
- Mean square: $E(Y^2) = \int_0^\theta dy \frac{2y}{\theta^2} y^2 = \frac{1}{2}\theta^2$
- Variance: $\text{Var}(Y) = \int_0^\theta dy \frac{2y}{\theta^2} (y - \mu)^2 = \frac{1}{18}\theta^2$
- Standard deviation: $\sigma_Y = \sqrt{\text{Var}(Y)} = \frac{1}{3\sqrt{2}}\theta$

Bruce M.
Boghossian

Properties of
estimators

Unbiasedness
Example 1

Unbiasedness
Example 2

Unbiasedness
Example 3

Summary

- Let $M_1 = \frac{1}{n} \sum_{j=1}^n Y_j$ be the average.
- Set the theoretical mean equal to the average: $\frac{2}{3}\theta_e = M_1$
- Hence $\theta_e = \frac{3}{2}M_1$
- MM estimator is then

$$\hat{\theta}_{\text{mm}}(\vec{Y}) = \frac{3}{2n} \sum_{j=1}^n Y_j$$

- MM estimator is unbiased

$$E(\hat{\theta}_{\text{mm}}(\vec{Y})) = \frac{3}{2n} \sum_{j=1}^n E(Y_j) = \frac{3}{2n} \sum_{j=1}^n \frac{2}{3}\theta = \frac{3}{2n} n \left(\frac{2}{3}\theta \right) = \theta.$$

Bruce M.
Boghossian

Properties of
estimators

Unbiasedness
Example 1

Unbiasedness
Example 2

Unbiasedness
Example 3

Summary

- If $\max_j y_j > \theta$, the likelihood will be zero
- So suppose that $\theta > \max_j y_j$
- Likelihood is $L(\vec{y}; \theta) = \prod_{j=1}^n \left(\frac{2y_j}{\theta^2} \right)$
- This clearly increases as θ decreases, so the MLE estimator is

$$\hat{\theta}_{\text{mle}}(\vec{y}) = \max_j y_j$$

- To calculate $E(\hat{\theta}_{\text{mle}})$, we need $f_{Y_{\max}}(y)$, but we can calculate this using what we know about order statistics.

- Note $F_Y(y) = 0$ for $y \leq 0$, and $F_Y(y) = 1$ for $y \geq \theta$
- For $0 < y < \theta$: $F_Y(y) = \int_0^y dz f_Y(z) = \int_0^y dz \frac{2z}{\theta^2} = \frac{y^2}{\theta^2}$
- Hence the CDF is

$$F_Y(y) = \begin{cases} 0 & \text{if } y \leq 0 \\ \frac{y^2}{\theta^2} & \text{if } 0 \leq y \leq \theta \\ 1 & \text{if } y \geq \theta \end{cases}$$

- From our theorem on order statistics

$$f_{Y_{\max}}(y) = \begin{cases} n \left(\frac{y^2}{\theta^2} \right)^{n-1} \frac{2y}{\theta^2} & \text{if } 0 \leq y \leq \theta \\ 0 & \text{otherwise} \end{cases} = \begin{cases} 2n \frac{y^{2n-1}}{\theta^{2n}} & \text{if } 0 \leq y \leq \theta \\ 0 & \text{otherwise} \end{cases}$$

- Note that this is normalized

$$\int_0^\theta dy f_{Y_{\max}}(y) = \frac{2n}{\theta^{2n}} \frac{\theta^{2n}}{2n} = 1.$$

Bruce M.
Boghossian

Properties of
estimators

Unbiasedness
Example 1

Unbiasedness
Example 2

Unbiasedness
Example 3

Summary

- Expectation value of the MLE estimator is then

$$E(\hat{\theta}_{\text{mle}}) = \int_0^{\theta} dy f_{Y_{\max}}(y)y = \int_0^{\theta} dy 2n \frac{y^{2n-1}}{\theta^{2n}} y = \frac{2n}{\theta^{2n}} \frac{\theta^{2n+1}}{2n+1} = \frac{2n}{2n+1} \theta.$$

- The *MLE estimator is biased* since

$$E(\hat{\theta}_{\text{mle}}(\vec{y})) = \frac{2n}{2n+1} \theta \neq \theta$$

- It is *asymptotically unbiased*, since $\lim_{n \rightarrow \infty} E(\hat{\theta}_{\text{mle}}(\vec{y})) = \theta$.

Bruce M.
Boghossian

Properties of
estimators

Unbiasedness
Example 1

Unbiasedness
Example 2

Unbiasedness
Example 3

Summary

- We can construct an unbiased version of the MLE estimator by defining

$$\hat{\theta}_3(\vec{y}) := \frac{2n+1}{2n} \hat{\theta}_{\text{mle}} := \frac{2n+1}{2n} \max_j y_j.$$

- We can see that $\hat{\theta}_3$ is unbiased since

$$E(\hat{\theta}_3) = \frac{2n+1}{2n} \frac{2n}{2n+1} \theta = \theta,$$

- There is no problem with creating unbiased estimators in this way.
- Note that $\hat{\theta}_3$ is not the MLE estimator, but it is arguably preferable to it.

Example 2: Unbiasedness requirement for a linear estimator

Bruce M.
Boghosian

Properties of
estimators

Unbiasedness
Example 1

Unbiasedness
Example 2

Unbiasedness
Example 3

Summary

- Suppose X_1, X_2, \dots, X_n have PDF $f_X(x, \theta)$, with theoretical mean $E(X) = \theta$.
- Hence $E(X_j) = \theta$ for $j = 1, 2, \dots, n$
- Suppose we construct the estimator $\hat{\theta}(\vec{X}) = \sum_{j=1}^n a_j X_j$.
- $\therefore E(\hat{\theta}(\vec{X})) = \sum_{j=1}^n a_j E(X_j) = \sum_{j=1}^n a_j \mu = \left(\sum_{j=1}^n a_j\right) \mu$
- So $\hat{\theta}$ is unbiased iff $\sum_{j=1}^n a_j = 1$

Example 3: The variance of the normal distribution

Bruce M.
Boghossian

Properties of
estimators

Unbiasedness
Example 1

Unbiasedness
Example 2

Unbiasedness
Example 3

Summary

- Recall the normal distribution with theoretical mean μ and variance $v = \sigma^2$

$$f_X(x) = \frac{1}{\sqrt{2\pi v}} \exp \left[-\frac{(x - \mu)^2}{2v} \right]$$

- Recall that the MLE and MM estimators were

$$\hat{\mu}(\vec{X}) = \bar{X} := \frac{1}{n} \sum_j^n X_j$$

$$\hat{v}(\vec{X}) = \frac{1}{n} \sum_{j=1}^n (X_j - \bar{X})^2$$

- Clearly, the estimator $\hat{\mu}$ is unbiased from the previous example $\sum_{j=1}^n \frac{1}{n} = 1$
- What about the estimator \hat{v} ?

Bruce M.
Boghossian

Properties of
estimators

Unbiasedness
Example 1

Unbiasedness
Example 2

Unbiasedness
Example 3

Summary

■ Lemma 1

$$\begin{aligned} E(X_j \bar{X}) &= E\left(X_j \frac{1}{n} \sum_{k=1}^n X_k\right) = \frac{1}{n} \sum_{k=1}^n E(X_j X_k) \\ &= \frac{1}{n} \sum_{k \neq j}^n E(X_j) E(X_k) + \frac{1}{n} E(X_j^2) \\ &= \frac{n-1}{n} \mu^2 + \frac{1}{n} (\mu^2 + \nu) = \mu^2 + \frac{1}{n} \nu \end{aligned}$$

■ Lemma 2

$$\begin{aligned} E(\bar{X}^2) &= E\left(\frac{1}{n} \sum_{j=1}^n X_j \bar{X}\right) = \frac{1}{n} \sum_{j=1}^n E(X_j \bar{X}) \\ &= \frac{1}{n} \sum_{j=1}^n \left(\mu^2 + \frac{1}{n} \nu\right) = \mu^2 + \frac{1}{n} \nu \end{aligned}$$

Bruce M.
Boghossian

Properties of
estimators

Unbiasedness
Example 1

Unbiasedness
Example 2

Unbiasedness
Example 3

Summary

- Using our lemmas, we have

$$\begin{aligned} E(\hat{v}) &= E \left(\frac{1}{n} \sum_{j=1}^n (X_j - \bar{X})^2 \right) = E \left(\frac{1}{n} \sum_{j=1}^n (X_j^2 - 2X_j\bar{X} + \bar{X}^2) \right) \\ &= \frac{1}{n} \sum_{j=1}^n \left[E(X_j^2) - 2E(X_j\bar{X}) + E(\bar{X}^2) \right] \\ &= \frac{1}{n} \sum_{j=1}^n \left[(\mu^2 + v) - 2 \left(\mu^2 + \frac{1}{n}v \right) + \left(\mu^2 + \frac{1}{n}v \right) \right] \\ &= \frac{n-1}{n} v \end{aligned}$$

- Since $E(\hat{v}) \neq v$, the *variance estimator is biased*, but *asymptotically unbiased*.

Bruce M.
Boghossian

Properties of
estimators

Unbiasedness
Example 1

Unbiasedness
Example 2

Unbiasedness
Example 3

Summary

- We have found $E(\hat{\sigma}^2) = \frac{n-1}{n}\sigma^2$
- We construct the *sample variance* estimator,

$$\hat{S}^2(\vec{X}) := \frac{n}{n-1} \hat{\sigma}^2(\vec{X}) = \frac{1}{n-1} \sum_{j=1}^n (X_j - \bar{X})^2$$

- There is an associated *sample standard deviation* estimator

$$\hat{S}(\vec{X}) := \sqrt{\frac{n}{n-1}} \hat{\sigma}(\vec{X}) = \sqrt{\frac{1}{n-1} \sum_{j=1}^n (X_j - \bar{X})^2}$$

- This is used in interval estimation, instead of the estimated standard deviation.

Bruce M.
Boghossian

Properties of
estimators

Unbiasedness
Example 1

Unbiasedness
Example 2

Unbiasedness
Example 3

Summary

- We have identified some desirable properties of estimators.
- We defined and described the property of *unbiasedness*.
- We presented three examples illustrating this property.