

Bruce M.
Boghosian

Testing
binomial data

Large-sample
testing

Small-sample
testing

Summary

Hypothesis testing and decision rules

Testing binomial data

Bruce M. Boghosian



Tufts
UNIVERSITY

School of Arts
and Sciences

Department of Mathematics

Tufts University

Bruce M.
Boghosian

Testing
binomial data

Large-sample
testing

Small-sample
testing

Summary

1 Testing binomial data

2 Large-sample testing

3 Small-sample testing

4 Summary

Bruce M.
Boghossian

Testing
binomial data

Large-sample
testing

Small-sample
testing

Summary

- We have $Y \in \{0, 1\}$ (0 = tails and 1 = heads).
- Bernoulli distribution for one trial $p_Y(k; p) = p^k(1 - p)^{1-k}$

$$p_Y(0; p) = p^0(1 - p)^{1-0} = 1 - p$$

$$p_Y(1; p) = p^1(1 - p)^{1-1} = p$$

- We have $E(Y) = p$ and $\text{Var}(Y) = p(1 - p)$ for one trial
- Parameter p is unknown.

Bruce M.
Boghossian

Testing
binomial data

Large-sample
testing

Small-sample
testing

Summary

- We conduct n independent Bernoulli trials
- Random vars. $\vec{Y} = \langle Y_1, Y_2, \dots, Y_n \rangle$ where $Y_j \in \{0, 1\}$.
- Outcomes $\vec{k} = \langle k_1, k_2, \dots, k_n \rangle$ where $k_j \in \{0, 1\}$.
- Random var. for total heads is $Y := \sum_{j=1}^n Y_j = n\bar{Y}$
- Outcome for total number of heads is $k := \sum_{j=1}^n k_j = n\bar{k}$
- Binomial distribution for n trials is

$$p_Y(k; p) = \binom{n}{k} p^k (1-p)^{n-k} = \binom{n}{n\bar{k}} p^{n\bar{k}} (1-p)^{n(1-\bar{k})}.$$

- We have $E(Y) = np$ and $\text{Var}(Y) = np(1-p)$ for n trials
- Parameter p is unknown.

Bruce M.
Boghosian

Testing
binomial data

Large-sample
testing

Small-sample
testing

Summary

- Note $0 \leq Y \leq n$ and $E(Y) = np$ and $\sigma_Y = \sqrt{np(1-p)}$
- If n is sufficiently large, $[pn - 3\sigma_Y, pn + 3\sigma_Y] \subset [0, n]$
- Because p is unknown, we use H_0 to make this judgement
- We do a *large-sample test*, relying on CLT, if

$$0 < np_0 - 3\sqrt{np_0(1-p_0)} < np_0 + 3\sqrt{np_0(1-p_0)} < n.$$

$$\therefore 0 < p_0 - 3\sqrt{p_0(1-p_0)/n} < p_0 + 3\sqrt{p_0(1-p_0)/n} < 1.$$

- If above true, normal distribution f_Y obtained from CLT comfortably fits in $[0, n]$ (respectively $f_{\bar{Y}}$ fits in $[0, 1]$)
- If above not true, we must conduct a *small-sample test*.

- Suppose Z is distributed as a standard normal, where

$$Z := \frac{Y - np_0}{\sqrt{np_0(1 - p_0)}} = \frac{\bar{Y} - p_0}{\sqrt{p_0(1 - p_0)/n}}$$

- Take n samples, k_1, \dots, k_n , let $k = \sum_{j=1}^n k_j$, and define

$$z := \frac{k - np_0}{\sqrt{np_0(1 - p_0)}} = \frac{\bar{k} - p_0}{\sqrt{p_0(1 - p_0)/n}}$$

- Use hypothesis testing method for standard normal r.v.
- Find thresholds for not rejecting H_0 if it is true with $100(1 - \alpha)\%$ confidence.

- Take n samples, k_1, \dots, k_n , let $k = \sum_j^n k_j$, and define

$$z := \frac{k - np_0}{\sqrt{np_0(1 - p_0)}} = \frac{\bar{k} - p_0}{\sqrt{p_0(1 - p_0)/n}}$$

- Three tests similar to earlier work on hypothesis testing:
 - To test $H_0 : p = p_0$ versus $H_1 : p > p_0$ at the α level of significance, reject H_0 if $z \geq +z_\alpha$
 - To test $H_0 : p = p_0$ versus $H_1 : p < p_0$ at the α level of significance, reject H_0 if $z \leq -z_\alpha$
 - To test $H_0 : p = p_0$ versus $H_1 : p \neq p_0$ at the α level of significance, reject H_0 if z is either $\leq -z_{\alpha/2}$ or $\geq +z_{\alpha/2}$.

Example: Do people postpone their deaths?

Bruce M.
Boghosian

Testing
binomial data

Large-sample
testing

Small-sample
testing

Summary

- Study of $n = 747$ obituaries
- Only 60, or 8%, died during the three months prior to their birthday.
- If deaths were distributed uniformly, one would expect this figure to be 25%.
- Is the decrease from 25% to 8% statistically significant?

Example: Death postponement (continued)

Bruce M.
Boghossian

Testing
binomial data

Large-sample
testing

Small-sample
testing

Summary

- Define k_j , for $j = 1, \dots, 747$ to be
 - $= 1$ if j th person died in three months prior to birthday
 - $= 0$ if j th person died at any other time of the year
- Then $p_e = \bar{k} = \frac{1}{n} \sum_j^n k_j$ is fraction of deaths three months prior to a birthday
- Take $H_0 : p = 0.25$ since the contrary seems perverse.
- Take $H_1 : p < 0.25$ and demand confidence with $\alpha = 0.05$.

Example: Death postponement (continued)

Bruce M.
Boghossian

Testing
binomial data

Large-sample
testing

Small-sample
testing

Summary

- First note that $np_0 = 747(0.25) = 186.75$ and

$$\sigma = \sqrt{747(0.25)(1 - 0.25)} = 11.83.$$

- np_0 more than 3σ greater than zero and less than $n = 747$.
- Hence large-sample testing is warranted.
- Null hypothesis is that $p = p_0 = 0.25$, and $n = 747$
- Calculate

$$z = \frac{k - np_0}{\sqrt{np_0(1 - p_0)}} = \frac{60 - 747(0.25)}{\sqrt{747(0.25)(1 - 0.25)}} = -10.7.$$

- The above is far, far less than $-z_{0.05} = -1.64$.
- Very strong evidence ($\gg 95\%$ confidence) effect is real.

Bruce M.
Boghossian

Testing
binomial data

Large-sample
testing

Small-sample
testing

Summary

- Example: Experimental drug test with $n = 19$ patients
- Old treatment is known to be 85% effective
- We wish to test $H_0 : p = 0.85$ versus $H_1 : p \neq 0.85$
- For $n = 19$ and $p_0 = 0.85$,
 - $np_0 = 16.15$
 - $\sigma = \sqrt{19(0.85)(0.15)} = 1.556$
- Note that $np_0 + 3\sigma = 16.15 + 3(1.556) = 20.819 > 19$
- Indicates that small-sample testing is necessary

How does small-sample testing work?

Bruce M.
Boghossian

Testing
binomial data

Large-sample
testing

Small-sample
testing

Summary

- List 19 possibilities:

k	$P(Y = k) = \binom{19}{k} (0.85)^k (0.15)^{19-k}$
6	1.99151×10^{-7}
7	2.09582×10^{-6}
8	0.0000178145
9	0.000123382
10	0.000699164
11	0.00324158
12	0.012246
13	0.0373659
14	0.0907457
15	0.171409
16	0.242829
17	0.242829
18	0.152892
19	0.0455994

- Note $P(Y \leq 13) = 0.053 \dots$, and $P(Y = 19) = 0.045 \dots$
- Hence we reject H_0 if $k \leq 13$ or $k = 19$.
- Note that confidence interval is asymmetric.

Bruce M.
Boghosian

Testing
binomial data

Large-sample
testing

Small-sample
testing

Summary

- We have studied the testing of binomial data
- We have described two cases
 - Large-sample testing
 - Small-sample testing
- We have provided an example of each kind of testing.