

Bruce M.
Boghosian

Introduction
to estimation

Confidence
intervals

Properties of
estimators

Bayesian
estimation

Hypothesis
testing

GLR and
GLRT

Small-sample
statistics

Midterm Exam Review

Bruce M. Boghosian



Tufts
UNIVERSITY

School of Arts
and Sciences

Department of Mathematics

Tufts University

Bruce M.
Boghosian

Introduction
to estimation

Confidence
intervals

Properties of
estimators

Bayesian
estimation

Hypothesis
testing

GLR and
GLRT

Small-sample
statistics

- 1 Introduction to estimation
- 2 Confidence intervals
- 3 Properties of estimators
- 4 Bayesian estimation
- 5 Hypothesis testing
- 6 GLR and GLRT
- 7 Small-sample statistics

Bruce M.
Boghossian

Introduction
to estimation

Confidence
intervals

Properties of
estimators

Bayesian
estimation

Hypothesis
testing

GLR and
GLRT

Small-sample
statistics

- Definition of *likelihood* $L(\lambda) = \prod_{j=1}^n f_X(x_j; \lambda)$
- Maximum likelihood estimator $\hat{\lambda}(\vec{x})$ gives most likely value for parameter λ .
- Estimation of parameters if form of pdf is known a priori
- Can be used for discrete or continuous pdfs, discrete or continuous parameters
- The log likelihood $\log L(\lambda)$ is often useful
- Gives only a single result, no confidence interval

Bruce M.
Boghossian

Introduction
to estimation

Confidence
intervals

Properties of
estimators

Bayesian
estimation

Hypothesis
testing

GLR and
GLRT

Small-sample
statistics

- *Method of moments* is another way to creating estimators
- Equate s theoretical moments to s sample moments

$$E(Y) = \int dy f_Y(y; \theta_1, \dots, \theta_s) y = \frac{1}{n} \sum_{j=1}^n y_j$$

$$E(Y^2) = \int dy f_Y(y; \theta_1, \dots, \theta_s) y^2 = \frac{1}{n} \sum_{j=1}^n y_j^2$$

$$\vdots$$

$$E(Y^s) = \int dy f_Y(y; \theta_1, \dots, \theta_s) y^s = \frac{1}{n} \sum_{j=1}^n y_j^s$$

- Yields s simultaneous equations for the s parameters.
- May be different from MLE (e.g., the uniform distribution)
- May be used in combination with MLE.

Bruce M.
Boghossian

Introduction
to estimation

Confidence
intervals

Properties of
estimators

Bayesian
estimation

Hypothesis
testing

GLR and
GLRT

Small-sample
statistics

- Moment-generating functions for proof of *CLT*
- Say $\bar{y} = \frac{1}{n} \sum_{j=1}^n y_j$ is normally distributed with known σ
- Form *standardized random variables* $z_j = \frac{y_j - \bar{y}}{\sigma/\sqrt{n}}$
- Standardized r.v.s distributed like standard normal $f_Z(z)$
- *Z tables* defined so $\int_{z_\alpha}^{\infty} dz f_Z(z) = \alpha$
- Confidence intervals can be symmetric or asymmetric
- $\text{Prob} \left(Y \in \left[\mu - z_{\alpha/2} \frac{\sigma}{\sqrt{n}}, \mu + z_{\alpha/2} \frac{\sigma}{\sqrt{n}} \right] \right) = 1 - \alpha$
- *Margin of error*: Half maximum width of a (usually 95%) confidence interval
- How large does a trial have to be to achieve a certain confidence?

Bruce M.
Boghossian

Introduction
to estimation

Confidence
intervals

Properties of
estimators

Bayesian
estimation

Hypothesis
testing

GLR and
GLRT

Small-sample
statistics

- If MLE and MM estimators differ, which is “correct”?
- Estimators themselves are random variables.
- Estimators as functions of random variables have means and variances.
- For $f_X(x; \theta)$, an *unbiased* estimator has $E(\hat{\theta}(\vec{X})) = \theta$.
- If an estimator is biased, but the bias vanishes as $n \rightarrow \infty$, we say that it is *asymptotically unbiased*.
- Sometimes you can fix biased estimators by applying a correction for finite n .

Bruce M.
Boghossian

Introduction
to estimation

Confidence
intervals

Properties of
estimators

Bayesian
estimation

Hypothesis
testing

GLR and
GLRT

Small-sample
statistics

- Cumulative distribution functions
- Order statistics for distribution of max and min
- *Efficiency* of estimators: Let $\hat{\theta}_1$ and $\hat{\theta}_2$ be two unbiased estimators for parameter θ . If $\text{Var}(\hat{\theta}_1) < \text{Var}(\hat{\theta}_2)$, we say that $\hat{\theta}_1$ is *more efficient* than $\hat{\theta}_2$.
- *Relative efficiency* of estimators: The *relative efficiency* of $\hat{\theta}_1$ with respect to $\hat{\theta}_2$ is $\frac{\text{Var}(\hat{\theta}_2)}{\text{Var}(\hat{\theta}_1)}$.
- Cramér-Rao bound: Absolute efficiency for estimators

Bruce M.
Boghossian

Introduction
to estimation

Confidence
intervals

Properties of
estimators

Bayesian
estimation

Hypothesis
testing

GLR and
GLRT

Small-sample
statistics

- Proof of the Cramér-Rao bound
- *Cauchy-Schwarz inequality*
- *Pearson correlation coefficient*
- Two forms of Cramér-Rao bound
 - First-derivative form
 - Second-derivative form

Properties of estimators: Sufficiency and consistency

Bruce M.
Boghossian

Introduction
to estimation

Confidence
intervals

Properties of
estimators

Bayesian
estimation

Hypothesis
testing

GLR and
GLRT

Small-sample
statistics

- Sufficiency and consistency
- Sufficiency defined by factorization theorem
- Later we learned a second factorization theorem
- Consistency of estimators
- Chebyshev's Theorem for establishing consistency

Bruce M.
Boghosian

Introduction
to estimation

Confidence
intervals

Properties of
estimators

Bayesian
estimation

Hypothesis
testing

GLR and
GLRT

Small-sample
statistics

- Bayes Theorem and examples

$$P(A_j | B) = \frac{P(B | A_j)P(A_j)}{\sum_{k=1}^n P(B | A_k)P(A_k)}.$$

- Updating priors to create new posterior distributions
- Bayesian search strategy
- Bayesian estimation

$$g_{\lambda}(\lambda | W = w_s) = \frac{f_W(w_s | \lambda) f_{\lambda}(\lambda)}{\int_{-\infty}^{+\infty} d\xi f_W(w_s | \xi) f_{\lambda}(\xi)}$$

Bruce M.
Boghossian

Introduction
to estimation

Confidence
intervals

Properties of
estimators

Bayesian
estimation

Hypothesis
testing

GLR and
GLRT

Small-sample
statistics

- State in terms of $z := \frac{\bar{y} - \mu_0}{\sigma / \sqrt{n}}$
- Let y_1, \dots, y_n be a random sample from a normal distribution for which σ is known.
- To test $H_0 : \mu = \mu_0$ versus $H_1 : \mu > \mu_0$ at the α level of significance, reject H_0 if $z \geq z_\alpha$.
- To test $H_0 : \mu = \mu_0$ versus $H_1 : \mu < \mu_0$ at the α level of significance, reject H_0 if $z \leq -z_\alpha$.
- To test $H_0 : \mu = \mu_0$ versus $H_1 : \mu \neq \mu_0$ at the α level of significance, reject H_0 if either $z \leq -z_{\alpha/2}$ or $z \geq +z_{\alpha/2}$.

Bruce M.
Boghosian

Introduction
to estimation

Confidence
intervals

Properties of
estimators

Bayesian
estimation

Hypothesis
testing

GLR and
GLRT

Small-sample
statistics

- Large-sample test if

$$0 < np_0 - 3\sqrt{np_0(1-p_0)} < np_0 + 3\sqrt{np_0(1-p_0)} < n$$

- Otherwise, small-sample test is necessary
- Type I versus Type II errors
- Power curves

Bruce M.
Boghossian

Introduction
to estimation

Confidence
intervals

Properties of
estimators

Bayesian
estimation

Hypothesis
testing

GLR and
GLRT

Small-sample
statistics

- Sets of parameters ω and Ω
- *Generalized Likelihood Ratio (GLR)* is then defined to be

$$\lambda = \frac{\max_{\theta \in \omega} L(\theta)}{\max_{\theta \in \Omega} L(\theta)}$$

- Generalization to many parameters is straightforward.
- Hypothesis testing with the GLR – the GLRT
- Reject H_0 if λ is below a certain threshold.

Bruce M.
Boghossian

Introduction
to estimation

Confidence
intervals

Properties of
estimators

Bayesian
estimation

Hypothesis
testing

GLR and
GLRT

Small-sample
statistics

- Using the sample variance for estimation
- Reviewed gamma and beta functions
- Reviewed gamma and beta distributions
- Sums of gamma distributed r.v.s are gamma distributed
- Sums of squares of normally distributed r.v.s are χ^2 distributed
- Orthogonal matrices – \bar{Y} and S_Y^2 are independent
- Showed that $\frac{(n-1)S^2}{\sigma^2}$ is chi square distributed

Bruce M.
Boghosian

Introduction
to estimation

Confidence
intervals

Properties of
estimators

Bayesian
estimation

Hypothesis
testing

GLR and
GLRT

Small-sample
statistics

- Finding pdf of quotient
- Quotient of two χ^2 r.v.s is F distributed.
- **Def.:** Suppose that U and V are independent chi squared r.v.s with n and m degrees of freedom, respectively. A random variable of the form $\frac{V/m}{U/n}$ is said to have an *F distribution with m and n degrees of freedom*.
- Student T distribution $T_n = \frac{Z}{\sqrt{U/n}}$
- Derived pdf of T_n – fat tails for small samples.
- Learned about T tables in appendices

Normally distributed data – μ and σ^2 both unknown

Bruce M.
Boghossian

Introduction
to estimation

Confidence
intervals

Properties of
estimators

Bayesian
estimation

Hypothesis
testing

GLR and
GLRT

Small-sample
statistics

- Interval estimation of μ using Z ratio
- Interval estimation of μ using T ratio
- Hypothesis testing using Z ratio
- Hypothesis testing using T ratio: One-sample T test
- Let s^2 denote the sample variance from n observations drawn from a normal distribution with unknown mean μ and unknown variance σ^2 . Let $\chi^2 = \frac{(n-1)s^2}{\sigma_0^2}$.
 - To test $H_0 : \sigma^2 = \sigma_0^2$ versus $H_1 : \sigma^2 > \sigma_0^2$ at the α level of significance, reject H_0 if $\chi^2 \geq \chi_{1-\alpha, n-1}^2$.
 - To test $H_0 : \sigma^2 = \sigma_0^2$ versus $H_1 : \sigma^2 < \sigma_0^2$ at the α level of significance, reject H_0 if $\chi^2 \leq \chi_{\alpha, n-1}^2$.
 - To test $H_0 : \sigma^2 = \sigma_0^2$ versus $H_1 : \sigma^2 \neq \sigma_0^2$ at the α level of significance, reject H_0 if χ^2 is either (a) $\leq \chi_{\alpha/2, n-1}^2$ or (b) $\geq \chi_{1-\alpha/2, n-1}^2$.

Bruce M.
Boghosian

Introduction
to estimation

Confidence
intervals

Properties of
estimators

Bayesian
estimation

Hypothesis
testing

GLR and
GLRT

Small-sample
statistics

We reviewed the following topics:

- Introduction to estimation
- Confidence intervals
- Properties of estimators
- Bayesian estimation
- Hypothesis testing
- GLR and GLRT
- Small-sample statistics