

Bruce M.
Boghosian

Introduction
and
motivation

The
multinomial
distribution

Examples

Marginal pdfs
of multinomial
distribution

Summary

Goodness of Fit Tests

Introduction and the Multinomial Distribution

Bruce M. Boghosian



Tufts
UNIVERSITY

School of Arts
and Sciences

Department of Mathematics

Tufts University

Bruce M.
Boghosian

Introduction
and
motivation

The
multinomial
distribution

Examples

Marginal pdfs
of multinomial
distribution

Summary

- 1 Introduction and motivation
- 2 The multinomial distribution
- 3 Examples
- 4 Marginal pdfs of multinomial distribution
- 5 Summary

Bruce M.
Boghosian

Introduction
and
motivation

The
multinomial
distribution

Examples

Marginal pdfs
of multinomial
distribution

Summary

- Most of the methods that we have looked at thus far require a priori knowledge of the probability distribution of the data that we collect.
- We are able to do interval estimation if we have a priori knowledge of the distribution.
- We are able to do hypothesis testing if we have a priori knowledge of the distribution.
- But what if we don't have a priori knowledge of the distribution?

Bruce M.
Boghossian

Introduction
and
motivation

The
multinomial
distribution

Examples

Marginal pdfs
of multinomial
distribution

Summary

- If we are using a test or estimation strategy for normally distributed data, is there at least some method to confirm that this is a reasonable assumption for our data?
- Such a method is called a *goodness-of-fit* test.
- General strategy for such tests
 - Observed data are grouped into k classes.
 - Each class's expected occupancy is calculated on the basis of the presumed model.
 - Occupancies compared with observed occupancies.
 - If comparison indicates substantial difference, conclude that presumed $p_X(k)$ or $f_X(x)$ was incorrect.

Bruce M.
Boghosian

Introduction
and
motivation

The
multinomial
distribution

Examples

Marginal pdfs
of multinomial
distribution

Summary

- GoF tests for distribution with specified parameters
- GoF tests for distribution with unknown parameters
- GoF tests to determine whether or not presumed independent data is indeed independent.

Bruce M.
Boghosian

Introduction
and
motivation

The
multinomial
distribution

Examples

Marginal pdfs
of multinomial
distribution

Summary

- GoF tests are based on statistic that is asymptotically χ^2 .
- That statistic derives from the *multinomial distribution*, so we study that first.
- It is a generalization of the *binomial distribution*.

The binomial distribution (yet again)

Bruce M.
Boghossian

Introduction
and
motivation

The
multinomial
distribution

Examples

Marginal pdfs
of multinomial
distribution

Summary

- Given n independent Bernoulli trials, each with success probability p ,
- Probability that total number of successes is X ,

$$p_X(k) = \text{Prob}(X = k) = \binom{n}{k} p^k (1 - p)^{n-k}.$$

for $k = 0, \dots, n$

- This is because $\binom{n}{k}$ is the number of ways that k successes can occur in n trials.
- A clear generalization is an experiment in which any one of t outcomes can occur for each trial.

Bruce M.
Boghossian

Introduction
and
motivation

The
multinomial
distribution

Examples

Marginal pdfs
of multinomial
distribution

Summary

- Given n independent trials, possible outcomes r_1, \dots, r_t
- Each outcome has respective probability p_1, \dots, p_t .
- These probabilities must sum to unity, $\sum_{j=1}^t p_j = 1$
- Suppose that outcomes are

$X_i = \text{number of times outcome } r_i \text{ appears}$

- For a given set of n trials, we have

$$X_1 = k_1, X_2 = k_2, \dots, X_n = k_n,$$

where $\sum_{j=1}^t k_j = n$.

- Question: Can we find an expression for the quantity

$$p_{X_1, X_2, \dots, X_t}(k_1, k_2, \dots, k_n) = \text{Prob}(X_1 = k_1, X_2 = k_2, \dots, X_t = k_t) ?$$

Bruce M.
Boghosian

Introduction
and
motivation

The
multinomial
distribution

Examples

Marginal pdfs
of multinomial
distribution

Summary

- **Thm.:** The *multinomial distribution* for r.v.s (X_1, X_2, \dots, X_t) with parameters n and (p_1, p_2, \dots, p_t) is

$$p_{X_1, X_2, \dots, X_t}(k_1, k_2, \dots, k_t) = \text{Prob}(X_1 = k_1, X_2 = k_2, \dots, X_t = k_t) \\ = \frac{n!}{k_1! k_2! \dots k_t!} p_1^{k_1} p_2^{k_2} \dots p_t^{k_t}.$$

where $k = 0, \dots, n$, and $\sum_{j=1}^t p_j = 1$, and $\sum_{j=1}^t k_j = n$.

- Note this reduces to the binomial theorem when $t = 2$.
- **Pf.:** A single outcome with k_1, k_2, \dots, k_t has probability $p_1^{k_1} p_2^{k_2} \dots p_t^{k_t}$. So we need only show that the number of possible outcomes with k_1, k_2, \dots, k_t is

$$\frac{n!}{k_1! k_2! \dots k_t!}$$

Bruce M.
Boghossian

Introduction
and
motivation

The
multinomial
distribution

Examples

Marginal pdfs
of multinomial
distribution

Summary

- The number of ways of permuting n objects of t types is

$$\frac{n!}{k_1!k_2!\cdots k_t!},$$

where k_1 is the number of objects of type 1, k_2 is the number of objects of type 2, \dots , k_t is the number of objects of type t . This completes the proof. \square

Bruce M.
Boghossian

Introduction
and
motivation

The
multinomial
distribution

Examples

Marginal pdfs
of multinomial
distribution

Summary

- **Ex.:** Number of ways to arrange n distinguishable balls is $n!$, since there are n ways to place the first ball, $n - 1$ ways to place the second, etc.
- **Ex.:** Number of ways to arrange n balls, k_1 of which are red, k_2 of which are green, and k_3 of which are blue, where $n = k_1 + k_2 + k_3$.
 - Arrange as though they were distinguishable ($n!$ ways).
 - Divide by $k_1!$, number of permutations of the red balls.
 - Divide by $k_2!$, number of permutations of the green balls.
 - Divide by $k_3!$, number of permutations of the blue balls.
- Result is the multinomial coefficient

$$\frac{n!}{k_1! k_2! \cdots k_t!}.$$

Example: A loaded die I

Bruce M.
Boghossian

Introduction
and
motivation

The
multinomial
distribution

Examples

Marginal pdfs
of multinomial
distribution

Summary

- A die has six faces, $i = 1, \dots, 6$.
- Suppose p_i is the probability that a roll results in face i .
- Suppose that $p_i = ci$, where c is to be determined.
- Demand

$$1 = \sum_{i=1}^6 p_i = \sum_{i=1}^6 ci = c \sum_{i=1}^6 i = c \frac{6 \cdot 7}{2} = 21c,$$

so that $c = 1/21$ and $p_i = i/21$.

- **Question:** The die is tossed twelve times. What is the probability that each face will appear exactly twice?

Bruce M.
Boghossian

Introduction
and
motivation

The
multinomial
distribution

Examples

Marginal pdfs
of multinomial
distribution

Summary

- The outcome we want is $X_i = k_i = 2$ for $i = 1, \dots, 6$.
- The result is then

$$\begin{aligned} p_{X_1, X_2, \dots, X_6}(2, 2, \dots, 2) &= \text{Prob}(X_1 = 2, X_2 = 2, \dots, X_6 = 2) \\ &= \frac{12!}{2! 2! \dots 2!} \left(\frac{1}{21}\right)^2 \left(\frac{2}{21}\right)^2 \dots \left(\frac{6}{21}\right)^2 \\ &= 0.00052746 \dots \end{aligned}$$

- Note that if die not loaded, above result would have been

$$\frac{12!}{2! 2! \dots 2!} \left(\frac{1}{6}\right)^{12} = 0.00343829 \dots$$

Bruce M.
Boghossian

Introduction
and
motivation

The
multinomial
distribution

Examples

Marginal pdfs
of multinomial
distribution

Summary

- Five observations are drawn from

$$f_Y(y) = 6y(1 - y)$$

- What is probability that
 - one of the observations lies in $[0, 0.25)$, and
 - none of the observations lies in $[0.25, 0.50)$, and
 - three of the observations lie in $[0.50, 0.75)$, and
 - one of the observations lies in $[0.75, 1.00]$?
- Bin the observations into one of four ranges. Range r_1 is $[0, 0.25)$, range r_2 is $[0.25, 0.50)$, range r_3 is $[0.50, 0.75)$, and range r_4 is $[0.75, 1.00]$.

Bruce M.
Boghossian

Introduction
and
motivation

The
multinomial
distribution

Examples

Marginal pdfs
of multinomial
distribution

Summary

- We have the corresponding probabilities

$$p_1 = \int_0^{1/4} dy \, 6y(1-y) = 5/32$$

$$p_2 = \int_{1/4}^{1/2} dy \, 6y(1-y) = 11/32$$

$$p_3 = \int_{1/2}^{3/4} dy \, 6y(1-y) = 11/32$$

$$p_4 = \int_{3/4}^1 dy \, 6y(1-y) = 5/32$$

- The desired result is

$$\begin{aligned} p_{X_1, X_2, X_3, X_4}(1, 0, 3, 1) &= \text{Prob}(X_1 = 1, X_2 = 0, X_3 = 3, X_4 = 1) \\ &= \frac{5!}{1!0!3!1!} \left(\frac{5}{32}\right)^1 \left(\frac{11}{32}\right)^0 \left(\frac{11}{32}\right)^3 \left(\frac{5}{32}\right)^1 \\ &= 0.0198334 \dots \end{aligned}$$

Bruce M.
Boghossian

Introduction
and
motivation

The
multinomial
distribution

Examples

Marginal pdfs
of multinomial
distribution

Summary

- **Def.:** Given a multivariate distribution of (X_1, X_2, \dots, X_t) , the individual distribution of X_i obtained by summing over all the other values of the random variables, X_j with $j \neq i$, is called the *marginal distribution* of X_i ,

$$p_{X_i}(k_i) = \sum_{j \neq i}^t \sum_{k_j=0}^n p_{X_1, X_2, \dots, X_t}(k_1, k_2, \dots, k_n).$$

- **Thm.:** Let (X_1, X_2, \dots, X_t) be a multinomial random variable with parameters n and (p_1, p_2, \dots, p_t) . Then the marginal pdf for any single one of these random variables, X_i , is binomial with parameters n and (p_i) .

Bruce M.
Boghosian

Introduction
and
motivation

The
multinomial
distribution

Examples

Marginal pdfs
of multinomial
distribution

Summary

- **Pf.:** Dichotomize the outcomes into “result r_i ” or “not result r_i ”. The probability of result r_i is p_i , and the probability of “not result r_i ” is

$$1 - p_i = \sum_{j \neq i}^t p_j.$$

- Then X_i is the number of successes out of n trials, where success is defined as obtaining result r_i , which happens with probability p_i . This is, however, precisely the definition of a binomial random variable with parameters n and p_i . □

Bruce M.
Boghossian

Introduction
and
motivation

The
multinomial
distribution

Examples

Marginal pdfs
of multinomial
distribution

Summary

- In fact, the above proof generalizes.
- The marginal distribution of any subset of multinomial random variables is multinomial.
- Proof can be obtained by the same reasoning and using mathematical induction on t , using the binomial case ($t = 2$) as the anchor case.

- **Ex.:** From past experience, professor expects exam grades to be normally distributed with $\mu = 80$, $\sigma = 5$. Letter grades assigned, with B given for 80 to 89. What is mean and variance of number of students receiving a B?
- Answer for mean

$$\begin{aligned}
 p_B &= \text{Prob}(80 \leq Y \leq 89) \\
 &= \text{Prob}\left(\frac{80 - 80}{5} \leq \frac{Y - 80}{5} \leq \frac{89 - 80}{5}\right) \\
 &= \text{Prob}(0 \leq Z \leq 0.8) \\
 &= 0.4772
 \end{aligned}$$

- Answer for variance

$$\text{Var}(X_B) = np_B(1 - p_B) = 12.47$$

Bruce M.
Boghossian

Introduction
and
motivation

The
multinomial
distribution

Examples

Marginal pdfs
of multinomial
distribution

Summary

- We have motivated and described GoF tests.
- We have related GoF tests to the multinomial distribution.
- We have derived the multinomial distribution and given its probability distribution.
- We have related its algebraic form to the combinatoric problem of permutations of n objects, each of t categories.
- We have shown the marginal distribution of a multinomial r.v. is binomial.
- We have given several examples.