

Bruce M.
Boghosian

Covariance
and
correlation

The bivariate
normal
distribution

Correlation
and causation

Summary

Regression

Covariance and Correlation, the Bivariate Normal Distribution

Bruce M. Boghosian



Tufts
UNIVERSITY

School of Arts
and Sciences

Department of Mathematics

Tufts University

Bruce M.
Boghosian

Covariance
and
correlation

The bivariate
normal
distribution

Correlation
and causation

Summary

- 1 Covariance and correlation
- 2 The bivariate normal distribution
- 3 Correlation and causation
- 4 Summary

Bruce M.
Boghosian

Covariance
and
correlation

The bivariate
normal
distribution

Correlation
and causation

Summary

- Recall the *covariance* of random variables X and Y

$$\text{Cov}(X, Y) = E[(X - \mu_X)(Y - \mu_Y)]$$

- The covariance depends on the units of the variables.
- Make independent of the units by dividing by σ_X and σ_Y , to obtain the *correlation coefficient*,

$$\rho(X, Y) = \frac{\text{Cov}(X, Y)}{\sigma_X \sigma_Y} = E \left[\left(\frac{X - \mu_X}{\sigma_X} \right) \left(\frac{Y - \mu_Y}{\sigma_Y} \right) \right]$$

- This also has the effect of ensuring $\rho(X, Y) \in [-1, +1]$.

Proof that $\rho(X, Y) \in [-1, +1]$

Bruce M.
Boghossian

Covariance
and
correlation

The bivariate
normal
distribution

Correlation
and causation

Summary

- Define the standardized r.v.s, $X^* = \frac{X - \mu_X}{\sigma_X}$ and $Y^* = \frac{Y - \mu_Y}{\sigma_Y}$
- Hence $E(X^*) = E(Y^*) = 0$ and $\text{Var}(X^*) = \text{Var}(Y^*) = 1$.
- Now consider

$$\begin{aligned} 0 \leq \text{Var}(X^* \pm Y^*) &= E[(X^*)^2] + 2E(X^*Y^*) + E[(Y^*)^2] \\ &= \text{Var}(X^*) \pm 2\text{Cov}(X^*, Y^*) + \text{Var}(Y^*) \\ &= 2 \pm 2\rho(X, Y). \end{aligned}$$

- It follows that $-1 \leq \rho(X, Y) \leq +1$. □

- We proved above earlier using Cauchy-Schwarz inequality.

- We have defined the correlation

$$\rho(X, Y) = \frac{E(XY) - E(X)E(Y)}{\sqrt{\text{Var}(X)}\sqrt{\text{Var}(Y)}}$$

- Define the *sample correlation coefficient* by replacing expectation values of moments with sample moments

$$R = \frac{\frac{1}{n} \sum_i^n (X_i - \bar{X}) (Y_i - \bar{Y})}{\sqrt{\frac{1}{n} \sum_i^n (X_i - \bar{X})^2} \sqrt{\frac{1}{n} \sum_i^n (Y_i - \bar{Y})^2}}$$

- In terms of sampled data points (x_j, y_j) , this is written

$$r = \frac{\frac{1}{n} \sum_i^n (x_i - \bar{x}) (y_i - \bar{y})}{\sqrt{\frac{1}{n} \sum_i^n (x_i - \bar{x})^2} \sqrt{\frac{1}{n} \sum_i^n (y_i - \bar{y})^2}}$$

Bruce M.
Boghossian

Covariance
and
correlation

The bivariate
normal
distribution

Correlation
and causation

Summary

- At this point, we have

$$r = \frac{\frac{1}{n} \sum_i^n (x_i - \bar{x})(y_i - \bar{y})}{\sqrt{\frac{1}{n} \sum_i^n (x_i - \bar{x})^2} \sqrt{\frac{1}{n} \sum_i^n (y_i - \bar{y})^2}}$$

$$\hat{\beta}_1 = \frac{\frac{1}{n} \sum_i^n (x_i - \bar{x})(y_i - \bar{y})}{\frac{1}{n} \sum_i^n (x_i - \bar{x})^2}.$$

- Eliminating numerators yields relation between $\hat{\beta}_1$ and r ,

$$\hat{\beta}_1 = r \sqrt{\frac{\sum_i^n (y_i - \bar{y})^2}{\sum_i^n (x_i - \bar{x})^2}}$$

- Mean square error due to lack of linearity

$$\begin{aligned}\sum_i^n (y_i - \hat{y}_i)^2 &= \sum_i^n (y_i - \hat{\beta}_0 - \hat{\beta}_1 x_i)^2 \\ &= \sum_i^n \left[y_i - (\bar{y} - \hat{\beta}_1 \bar{x}) - \hat{\beta}_1 x_i \right]^2 \\ &= \sum_i^n \left[(y_i - \bar{y}) - \hat{\beta}_1 (x_i - \bar{x}) \right]^2,\end{aligned}$$

where we used $\hat{\beta}_0 = \bar{y} - \hat{\beta}_1 \bar{x}$.

- Expand the above and use $\hat{\beta}_1 = \frac{\sum_i (x_i - \bar{x})(y_i - \bar{y})}{\sum_i^n (x_i - \bar{x})^2}$ to find

$$\begin{aligned}\sum_i^n (y_i - \hat{y}_i)^2 &= \sum_i^n (y_i - \bar{y})^2 - 2\hat{\beta}_1 \sum_i^n (x_i - \bar{x})(y_i - \bar{y}) + \hat{\beta}_1^2 \sum_i^n (x_i - \bar{x})^2 \\ &= \sum_i^n (y_i - \bar{y})^2 - \hat{\beta}_1^2 \sum_i^n (x_i - \bar{x})^2\end{aligned}$$

Bruce M.
Boghossian

Covariance
and
correlation

The bivariate
normal
distribution

Correlation
and causation

Summary

- Relationship between $\hat{\beta}_1$ and r

$$\hat{\beta}_1 = r \sqrt{\frac{\sum_i^n (y_i - \bar{y})^2}{\sum_i^n (x_i - \bar{x})^2}}$$

- Mean square error due to lack of linearity

$$\sum_i^n (y_i - \hat{y}_i)^2 = \sum_i^n (y_i - \bar{y})^2 - \hat{\beta}_1^2 \sum_i^n (x_i - \bar{x})^2$$

- Eliminating $\hat{\beta}_1$ and solving for r yields

$$r^2 = \frac{\sum_i^n (y_i - \bar{y})^2 - \sum_i^n (y_i - \hat{\beta}_0 - \hat{\beta}_1 x_i)^2}{\sum_i^n (y_i - \bar{y})^2}.$$

- Define the *coefficient of determination*

$$r^2 = \frac{\sum_i^n (y_i - \bar{y})^2 - \sum_i^n (y_i - \hat{\beta}_0 - \hat{\beta}_1 x_i)^2}{\sum_i^n (y_i - \bar{y})^2}.$$

- This admits a simple interpretation
 - $\sum_i^n (y_i - \bar{y})^2$ is the *total variability* in y .
 - $\sum_i^n (y_i - \hat{\beta}_0 - \hat{\beta}_1 x_i)^2$ is the variability that can not be explained by linear regression.
 - The numerator of r^2 is the variability that *can* be explained by linear regression.
 - The quantity r^2 is the fraction of the variability that can be explained by regression.

Bruce M.
Boghossian

Covariance
and
correlation

The bivariate
normal
distribution

Correlation
and causation

Summary

- Two-dimensional integral of the exponential of a negative-definite quadratic form

$$I(a, b, c, \mu_X, \mu_Y) = \int_{-\infty}^{+\infty} dx \int_{-\infty}^{+\infty} dy \exp [-(ax^2 + 2bxy + cy^2)]$$

- Try to demand that

$$\begin{aligned} ax^2 + 2bxy + cy^2 &= \mu [(x + \kappa y)^2 + (y + \lambda x)^2] \\ &= \mu [(1 + \lambda^2) x^2 + 2(\kappa + \lambda) xy + (1 + \kappa^2) y^2] \end{aligned}$$

- If true for all $x, y \in \mathbb{R}$, we must have

$$\begin{aligned} a &= \mu (1 + \lambda^2) & \lambda &= \frac{a \pm \sqrt{ac - b^2}}{b} \\ b &= \mu (\kappa + \lambda) & \kappa &= \frac{c \pm \sqrt{ac - b^2}}{b} \\ c &= \mu (1 + \kappa^2) & \mu &= \frac{b^2}{a + c \pm 2\sqrt{ac - b^2}} \end{aligned}$$

- We have demonstrated that

$$ax^2 + 2bxy + cy^2 = \mu \left[(x + \kappa y)^2 + (y + \lambda x)^2 \right] = \mu (\xi^2 + \eta^2)$$

where the new variables are $\xi = x + \kappa y$ and $\eta = y + \lambda x$.

- The old and new constants are related as follows

$$\begin{aligned} a &= \mu (1 + \lambda^2) & \lambda &= \frac{a \pm \sqrt{ac - b^2}}{b} \\ b &= \mu (\kappa + \lambda) & \kappa &= \frac{c \pm \sqrt{ac - b^2}}{b} \\ c &= \mu (1 + \kappa^2) & \mu &= \frac{b^2}{a + c \pm 2\sqrt{ac - b^2}} > 0 \end{aligned}$$

- Jacobian is (after a bit of algebra)

$$\frac{\partial(\xi, \eta)}{\partial(x, y)} = \left| \begin{bmatrix} 1 & \kappa \\ \lambda & 1 \end{bmatrix} \right| = |1 - \kappa\lambda| = \frac{\sqrt{ac - b^2}}{\mu}$$

Bruce M.
Boghossian

Covariance
and
correlation

The bivariate
normal
distribution

Correlation
and causation

Summary

- It follows that

$$\begin{aligned} \int_{-\infty}^{+\infty} dx \int_{-\infty}^{+\infty} dy e^{-(ax^2+2bxy+cy^2)} \\ = \frac{\mu}{\sqrt{ac-b^2}} \int_{-\infty}^{+\infty} d\xi \int_{-\infty}^{+\infty} d\eta e^{-\mu(\xi^2+\eta^2)} = \frac{\pi}{\sqrt{ac-b^2}} \end{aligned}$$

- From this we see that

$$f_{X,Y}(x,y) = \frac{\sqrt{ac-b^2}}{\pi} e^{-(ax^2+2bxy+cy^2)}$$

is a normalized bivariate pdf for X and Y .

- We have shown that

$$f_{X,Y}(x, y) = \frac{\sqrt{ac - b^2}}{\pi} e^{-(ax^2 + 2bxy + cy^2)}$$

is a normalized bivariate pdf for X and Y .

- To obtain form given in textbook, rename the constants

$$a = \frac{1}{2(1 - \rho^2)\sigma_X^2}, \quad b = -\frac{\rho}{2(1 - \rho^2)\sigma_X\sigma_Y}, \quad c = \frac{1}{2(1 - \rho^2)\sigma_Y^2}$$

- The above becomes

$$f_{X,Y}(x, y) = \frac{1}{2\pi\sigma_X\sigma_Y\sqrt{1 - \rho^2}} \exp \left\{ -\frac{1}{2} \left(\frac{1}{1 - \rho^2} \right) \left[\frac{(x - \mu_X)^2}{\sigma_X^2} - 2\rho \frac{(x - \mu_X)(y - \mu_Y)}{\sigma_X\sigma_Y} + \frac{(y - \mu_Y)^2}{\sigma_Y^2} \right] \right\}$$

- In the above, we have shifted x and y by means, μ_X and μ_Y , which will not affect normalization.

- Use the bivariate normal distribution in the form

$$f_{X,Y}(x,y) = \frac{1}{2\pi\sigma_X\sigma_Y\sqrt{1-\rho^2}} \exp \left\{ -\frac{1}{2} \left(\frac{1}{1-\rho^2} \right) \left[\frac{(x-\mu_X)^2}{\sigma_X^2} - 2\rho \frac{(x-\mu_X)(y-\mu_Y)}{\sigma_X\sigma_Y} + \frac{(y-\mu_Y)^2}{\sigma_Y^2} \right] \right\}$$

- Either in (x, y) or (ξ, η) coordinates, we can then calculate

$$E(X) = \mu_X$$

$$E(X^2) = \mu_X^2 + \sigma_X^2$$

$$\text{Var}(X) = \sigma_X^2$$

$$E(Y) = \mu_Y$$

$$E(Y^2) = \mu_Y^2 + \sigma_Y^2$$

$$\text{Var}(Y) = \sigma_Y^2$$

$$E(XY) = \mu_X\mu_Y + \rho\sigma_X\sigma_Y$$

$$\text{Cov}(X, Y) = \rho\sigma_X\sigma_Y$$

$$\frac{\text{Cov}(X, Y)}{\sigma_X\sigma_Y} = \rho$$

- We also have

$$E(Y | x) = \mu_Y + \rho \frac{\sigma_Y}{\sigma_X} (x - \mu_X)$$

$$\text{Var}(Y | x) = (1 - \rho^2)\sigma_Y^2$$

Bruce M.
Boghosian

Covariance
and
correlation

The bivariate
normal
distribution

Correlation
and causation

Summary

- The MLEs for μ_X , μ_Y , σ_X^2 , σ_Y^2 and ρ , assuming that all five of them are unknown, are \bar{X} , \bar{Y} , $\frac{1}{n} \sum_i^n (X_i - \bar{X})^2$, $\frac{1}{n} \sum_i^n (Y_i - \bar{Y})^2$, and R , respectively.
- It is also possible to test the null hypothesis $H_0 : \rho = 0$, in order to test for the presence or absence of correlation, using a T test.

Bruce M.
Boghosian

Covariance
and
correlation

The bivariate
normal
distribution

Correlation
and causation

Summary

- Correlation does not indicate causation.
- Two things can be correlated only because they are both correlated with a third thing that is not observed.
- Correlations can be due to the structure of what is observed.

Bruce M.
Boghossian


Covariance
and
correlation

The bivariate
normal
distribution

Correlation
and causation

Summary

- Cristian Calude & Giuseppe Longo, "The Deluge of Spurious Correlations in Big Data," *Foundations of Science* 22/3 (2017) 595-612.
- It turns out that in *any* party of six people, there must either be three who all know each other, or three who are all strangers.



Bruce M.
Boghossian


Covariance
and
correlation

The bivariate
normal
distribution

Correlation
and causation

Summary

- Cristian Calude & Giuseppe Longo, "The Deluge of Spurious Correlations in Big Data," *Foundations of Science* 22/3 (2017) 595-612.
- It turns out that in *any* party of six people, there must either be three who all know each other, or three who are all strangers.



Bruce M.
Boghosian


Covariance
and
correlation

The bivariate
normal
distribution

Correlation
and causation

Summary

- Cristian Calude & Giuseppe Longo, "The Deluge of Spurious Correlations in Big Data," *Foundations of Science* 22/3 (2017) 595-612.
- It turns out that in *any* party of six people, there must either be three who all know each other, or three who are all strangers.



Bruce M.
Boghossian


Covariance
and
correlation

The bivariate
normal
distribution

Correlation
and causation

Summary

- Cristian Calude & Giuseppe Longo, "The Deluge of Spurious Correlations in Big Data," *Foundations of Science* 22/3 (2017) 595-612.
- It turns out that in *any* party of six people, there must either be three who all know each other, or three who are all strangers.



Bruce M.
Boghosian

Covariance
and
correlation

The bivariate
normal
distribution

Correlation
and causation

Summary

- Cristian Calude & Giuseppe Longo, "The Deluge of Spurious Correlations in Big Data," *Foundations of Science* **22/3** (2017) 595-612.
- Ramsey theory is about finding structure and organization in sets of data.
- Ramsey numbers indicate how big a set must be to guarantee the existence of certain minimal structures:
 - $R(3, 3) = 6$ (example on previous slide)
 - $R(4, 5) = 25$
 - $R(3, 3, 3) = 17$
 - $43 \leq R(5, 5) \leq 49$
- Ramsey theory explains why we tend to find structure in seemingly random sets.

Bruce M.
Boghosian

Covariance
and
correlation

The bivariate
normal
distribution

Correlation
and causation

Summary

- Cristian Calude & Giuseppe Longo, "The Deluge of Spurious Correlations in Big Data," *Foundations of Science* **22**/3 (2017) 595-612.
- As data sets increase in size, the ratio of the number of meaningful correlations to the number of spurious correlations will tend to zero.

Bruce M.
Boghosian

Covariance
and
correlation

The bivariate
normal
distribution

Correlation
and causation

Summary

- Correlations can be spurious.
- [Tyler Vigen web page](#)

Bruce M.
Boghosian

Covariance
and
correlation

The bivariate
normal
distribution

Correlation
and causation

Summary

- We defined the *correlation* $\rho(X, Y) \in [-1, +1]$.
- We presented a method of estimating $\rho(X, Y)$ using sample moments.
- We constructed the *Pearson correlation coefficient* R .
- We have made an interpretation of the r^2 as the *coefficient of determination*.
- We have studied bivariate normal distributions.
- We have seen how to parametrize bivariate normal distributions using five parameters – two means, two standard deviations, and the correlation.
- We have discussed some of the problems associated with naive hunting for correlations.