

Bruce M.
Boghosian

Joint and
conditional
probabilities

Sufficiency

Discussion

Summary

Properties of Estimators

Sufficiency

Bruce M. Boghosian



Tufts
UNIVERSITY

School of Arts
and Sciences

Department of Mathematics

Tufts University

Bruce M.
Boghosian

Joint and
conditional
probabilities

Sufficiency

Discussion

Summary

1 Joint and conditional probabilities

2 Sufficiency

3 Discussion

4 Summary

Bruce M.
Boghossian

Joint and
conditional
probabilities


Sufficiency

Discussion

Summary

- *Joint prob., “A and B”*, denoted $P(A \cap B)$ or $P(A, B)$
- *Conditional prob., “A given B”*, denoted $P(A | B)$
- Demand $P(A, B) = P(A | B)P(B) = P(B | A)P(A)$, so

$$P(A | B) = \frac{P(A, B)}{P(B)} \quad \text{and} \quad P(B | A) = \frac{P(A, B)}{P(A)}$$



Bruce M.
Boghossian

Joint and
conditional
probabilities

Sufficiency

Discussion

Summary

- You roll pair of dice, but not allowed to see the outcome.
- You would like to know if the sum is an even number.
- Two people see the outcome, and they each give you information about it.
 - Person A tells you that the sum is ≤ 7 .
 - Person B tells you that the sum is an odd number.
- Whose information was more helpful?

Bruce M.
Boghosian

Joint and
conditional
probabilities

Sufficiency

Discussion

Summary

- Whose information was more helpful?

$$\begin{aligned}
 P(\text{Sum is even} \mid \text{Sum} \leq 7) &= \frac{P(2)+P(4)+P(6)}{P(2)+P(3)+P(4)+P(5)+P(6)+P(7)} \\
 &= \frac{\frac{1}{36} + \frac{3}{36} + \frac{5}{36}}{\frac{1}{36} + \frac{2}{36} + \frac{3}{36} + \frac{4}{36} + \frac{5}{36} + \frac{6}{36}} \\
 &= \frac{9}{21}
 \end{aligned}$$

$$P(\text{Sum is even} \mid \text{Sum is odd}) = 0.$$

- Clearly, Person B's information was more helpful.
- Person B's information was *sufficient*.
- Person A's information was not *sufficient*.

Bruce M.
Boghossian

Joint and
conditional
probabilities

Sufficiency

Discussion

Summary

- Bernoulli PDF: $p_X(k; p) = p^k(1 - p)^{1-k}$ where $k = 0, 1$
- Sample n Bernoulli-distributed random numbers and find

$$X_1 = k_1, \quad X_2 = k_2, \quad X_3 = k_3, \quad \dots, \quad X_n = k_n$$

- Maximum likelihood estimator $\hat{p}(\vec{X}) = \frac{1}{n} \sum_{j=1}^n X_j$.
- Maximum likelihood estimate $p_e = \frac{1}{n} \sum_{j=1}^n k_j$.
- Is \hat{p} a *sufficient estimator* for p ?

Bruce M.
Boghossian

Joint and
conditional
probabilities

Sufficiency

Discussion

Summary

- Is \hat{p} a *sufficient estimator* for p ? What would this mean?
- It means that
 - conditional probability of the observation, given the estimate, is independent of p .
 - $P(X_1 = k_1, \dots, X_n = k_n \mid \hat{p} = p_e)$ is independent of p .
 - while the joint probability $P(X_1 = k_1, \dots, X_n = k_n)$ may depend on \vec{k} and p , when conditioned on the observation made, namely $\hat{p} = p_e$, the dependence on p disappears.
 - everything data can tell us about p contained in est. p_e .
 - prob. of sample can be determined without knowing p .
- Returning to example, is the following independent of p ?

$$P(X_1 = k_1, \dots, X_n = k_n \mid \hat{p} = p_e)$$

Bruce M.
Boghossian

Joint and
conditional
probabilities

Sufficiency

Discussion

Summary

- Is $P(X_1 = k_1, \dots, X_n = k_n \mid \hat{p} = p_e)$ independent of p ?
- We have

$$\begin{aligned} P(X_1 = k_1, \dots, X_n = k_n \mid \hat{p} = p_e) &= \frac{P(X_1 = k_1, \dots, X_n = k_n, \hat{p} = p_e)}{P(\hat{p} = p_e)} \\ &= \frac{P(X_1 = k_1, \dots, X_n = k_n)}{P(\hat{p} = p_e)} \end{aligned}$$

- Let's calculate the numerator and denominator separately.

Bruce M.
Boghossian

Joint and
conditional
probabilities

Sufficiency

Discussion

Summary

■ Numerator:

$$\begin{aligned} P(X_1 = k_1, \dots, X_n = k_n) &= p^{k_1} (1-p)^{1-k_1} \dots p^{k_n} (1-p)^{1-k_n} \\ &= p^{k_1 + \dots + k_n} (1-p)^{(1-k_1) + \dots + (1-k_n)} \\ &= p^{np_e} (1-p)^{n-np_e} \end{aligned}$$

■ Denominator:

$$P(\hat{p} = p_e) = P\left(\sum_{j=1}^n X_j = np_e\right) = \binom{n}{np_e} p^{np_e} (1-p)^{n-np_e}$$

■ Quotient is independent of n so estimator \hat{p} is sufficient:

$$P(X_1 = k_1, \dots, X_n = k_n \mid \hat{p} = p_e) = \frac{p^{np_e} (1-p)^{n-np_e}}{\binom{n}{np_e} p^{np_e} (1-p)^{n-np_e}} = \frac{1}{\binom{n}{np_e}}$$

Bruce M.
Boghosian

Joint and
conditional
probabilities

Sufficiency

Discussion

Summary

- Redo Example 2 with estimator $\hat{p}^*(\vec{X}) = X_1$, so $p_e^* = k_1$.
- Define $\bar{k} := \frac{1}{n} \sum_{j=1}^n k_j$.
- Find the conditional probability of the particular observation, given the estimate

$$\begin{aligned} P(X_1 = k_1, \dots, X_n = k_n \mid \hat{p}^* = k_1) \\ &= \frac{p^{n\bar{k}}(1-p)^{n-n\bar{k}}}{p^{k_1}(1-p)^{1-k_1}} \\ &= p^{n\bar{k}-k_1}(1-p)^{n-1-n\bar{k}+k_1}. \end{aligned}$$

- This depends on p , so the estimator \hat{p} is not sufficient.

Bruce M.
Boghossian

Joint and
conditional
probabilities

Sufficiency

Discussion

Summary

- Recall the PDF:

$$f_Y(y) = \begin{cases} \frac{2y}{\theta^2} & \text{if } 0 \leq y \leq \theta \\ 0 & \text{otherwise} \end{cases}$$

- The method of moments estimator for this is

$$\hat{\theta} = \frac{3}{2}\bar{Y} = \frac{3}{2n} \sum_{j=1}^n Y_j$$

- If $\hat{\theta}$ were sufficient, any two random samples, with same value of θ_e should yield exactly same information about θ .

Bruce M.
Boghossian

Joint and
conditional
probabilities

Sufficiency

Discussion

Summary

- If $\hat{\theta}$ were sufficient, any two random samples, with same value of θ_e should yield exactly same information about θ .
- To demonstrate that is not the case, consider:
 - Case 1: $\vec{y} = \{3, 4, 5\}$ so that $\theta_e = \frac{3}{2} \cdot \frac{1}{3}(3 + 4 + 5) = 6$
 - Case 2: $\vec{y} = \{1, 3, 8\}$ so that $\theta_e = \frac{3}{2} \cdot \frac{1}{3}(1 + 3 + 8) = 6$
- In spite of the fact that $\theta_e = 6$ for both cases, note:
 - Case 1: true θ **could** be = 7, because $y_1, y_2, y_3 < 7$.
 - Case 2: true θ **could not** be = 7, because $y_3 = 8 > 7$.
- Hence, without even calculating the conditional probability, we know $\hat{\theta}$ is not sufficient.

Bruce M.
Boghossian

Joint and
conditional
probabilities

Sufficiency

Discussion

Summary

- Let $X_1 = k_1, \dots, X_n = k_n$ be a random sample of size n from $p_X(k; \theta)$. The statistic $\hat{\theta}(X_1, \dots, X_n)$ is *sufficient* for θ if the likelihood function, $L(\theta)$, factors into the product of the pdf for $\hat{\theta}$ and a constant that does not involve θ ,

$$L(\theta) = \prod_{j=1}^n p_X(k_j; \theta) = p_{\hat{\theta}}(\theta_e; \theta) b(k_1, \dots, k_n).$$

- Example 1 again:

- $L(p) = P(X_1 = k_1, \dots, X_n = k_n) = p^K (1 - p)^{n-K}$
- $f_{\hat{p}}(p) = \binom{n}{np_e} p^K (1 - p)^{n-K}$
- Hence $L(p) = f_{\hat{p}}(p) \left[\binom{n}{np_e} \right]^{-1}$ where $p_e = \frac{1}{n} \sum_{j=1}^n k_j$.

Why is sufficiency desirable

Bruce M.
Boghossian

Joint and
conditional
probabilities

Sufficiency

Discussion

Summary

- For any unknown pdf parameter, there will be an infinity of unbiased estimators.
- Some subset of these will be sufficient estimators, or functions of sufficient estimators.
- The variance of any unbiased estimator based on a sufficient estimator will be lower than that of any unbiased estimator that is not based on a sufficient estimator.
- Hence, sufficient estimators tend to be more efficient.

Bruce M.
Boghossian

Joint and
conditional
probabilities

Sufficiency

Discussion

Summary

- Estimators, by their very nature, discard data, $\hat{\theta}(\vec{X})$
- In doing so, they accomplish a kind of *data reduction*.
- For example, if you can reduce 10^6 normally distributed numbers to a mean and a variance, you have accomplished substantial data reduction.
- You need *all* 10^6 numbers to estimate mean and variance, since you want to estimate those as accurately as possible.
- Once you have $\hat{\mu}(\vec{X})$ and $\hat{\sigma}^2(\vec{X})$, you don't need \vec{X} .
- A sufficient estimator does not *needlessly* discard data.
- If estimator $\hat{\theta}$ is sufficient, everything that can be known about the parameter θ has been extracted from the data, and nothing has been left behind.

Bruce M.
Boghossian

Joint and
conditional
probabilities

Sufficiency

Discussion

Summary

- Given n pieces of data $\vec{X} = \langle X_1, \dots, X_n \rangle$
- An estimator can be unbiased, but not sufficient
- $\hat{\mu}_n(\vec{X}) = X_1$ is unbiased, but not sufficient
 - Unbiased because $E(\hat{\mu}_n) = E(X_1) = \mu$
 - Not sufficient because it wastes $n - 1$ of the numbers in the sample \vec{X}
- An estimator can be sufficient, but not unbiased
 - If $\hat{\theta}$ sufficient for θ , any invertible function of $\hat{\theta}$ is likewise.
 - e.g., $\hat{\theta}_2 = \hat{\theta}^3$ has the same information content as $\hat{\theta}$.
 - One would not expect $E(\hat{\theta}^3) = E[(\hat{\theta})]^3$ so not unbiased

Bruce M.
Boghosian

Joint and
conditional
probabilities

Sufficiency

Discussion

Summary

- We have defined and studied sufficiency of estimators.
- We have looked at examples of estimators that are sufficient and not sufficient.
- We have discussed the relationship between sufficiency and other properties of estimators.