

Bruce M.
Boghosian

Background
and
motivation

Gamma and
beta functions

Gamma and
beta pdfs

The chi
squared
distribution

A linear
algebra
interlude

The
independence
of \bar{Y} and S^2

Summary

Derivation of Student's T Distribution I

Part I: The Chi Square Distribution and related results

Bruce M. Boghosian



Tufts
UNIVERSITY

School of Arts
and Sciences

Department of Mathematics

Tufts University

Bruce M.
Boghossian

Background
and
motivation

Gamma and
beta functions

Gamma and
beta pdfs

The chi
squared
distribution

A linear
algebra
interlude

The
independence
of \bar{Y} and S^2

Summary

- 1 Background and motivation
- 2 Gamma and beta functions
- 3 Gamma and beta pdfs
- 4 The chi squared distribution
- 5 A linear algebra interlude
- 6 The independence of \bar{Y} and S^2
- 7 Summary

Bruce M.
Boghosian

Background
and
motivation

Gamma and
beta functions

Gamma and
beta pdfs

The chi
squared
distribution

A linear
algebra
interlude

The
independence
of \bar{Y} and S^2

Summary

- Recall difference between *variance* and *sample variance*

- $\sigma_Y^2 = \frac{1}{n} \sum_{j=1}^n (Y_j - \bar{Y})^2$

- $S_Y^2 = \frac{1}{n-1} \sum_{j=1}^n (Y_j - \bar{Y})^2$

- where $\bar{Y} := \frac{1}{n} \sum_{k=1}^n Y_k$

- By the CLT, $Z = \frac{\bar{Y} - \mu}{\sigma/\sqrt{n}}$ is normally distributed.

- Question: Is $T = \frac{\bar{Y} - \mu}{S/\sqrt{n}}$ also normally distributed?

- Answer: For very large n , there is little difference in distributions of Z and T .

Large samples and small samples

Bruce M.
Boghosian

Background
and
motivation

Gamma and
beta functions

Gamma and
beta pdfs

The chi
squared
distribution

A linear
algebra
interlude

The
independence
of \bar{Y} and S^2

Summary

- For many years it was believed T was also normally distributed for small n .
- William Sealy Gossett (1876-1937) was the first to realize that it was not.
- Small-sample quality assurance at the Guinness brewery in Dublin
 - Barley and other ingredients came in small batches (small n) from small farms.
 - σ was generally unknown and had to be inferred from the data
 - Gossett noticed distributions of $T = \frac{\bar{Y} - \mu}{S/\sqrt{n}}$ decayed slower than a normal pdf.
 - Distribution was still bell-shaped, but the tails were “thicker”.



Bruce M.
Boghossian

Background
and
motivation

Gamma and
beta functions

Gamma and
beta pdfs

The chi
squared
distribution

A linear
algebra
interlude

The
independence
of \bar{Y} and S^2

Summary

- Review gamma distributions and sums of gamma-distributed random variables
- Understand chi square distribution as a special case of the gamma distribution
- Show sums of squares of iid normal r.v.s, $\sum_{j=1}^n Z_j^2$, are chi square distributed
- Show \bar{Y} and S_Y^2 are independent
- Show $\frac{(n-1)S^2}{\sigma^2}$ is chi square distributed
- Derive pdf of ratio of two iid chi square r.v.s, which is called an F distribution
- Show that $T^2 = \left(\frac{\bar{Y} - \mu}{S/\sqrt{n}} \right)^2$ is F distributed
- Use the above to derive the T distribution pdf f_T

- Definition of the gamma function:

$$\Gamma(r) := \int_0^{\infty} du e^{-u} u^{r-1}$$

- Special cases (first by substitution $u = w^2$, second is elementary):

$$\Gamma(1/2) = \int_0^{\infty} du \frac{e^{-u}}{\sqrt{u}} = \sqrt{\pi} \quad \text{and} \quad \Gamma(1) = \int_0^{\infty} du e^{-u} = 1.$$

- Recurrence formula

$$\Gamma(r+1) := \int_0^{\infty} du e^{-u} u^r = -e^{-u} u^r \Big|_0^{\infty} + r \int_0^{\infty} du e^{-u} u^{r-1}$$

$$\Gamma(r+1) = r\Gamma(r)$$

- Then $\Gamma(2) = 1 \cdot \Gamma(1) = 1!$, $\Gamma(3) = 2 \cdot \Gamma(2) = 2!$, and more generally

$$\Gamma(r+1) = r! \quad \text{if } r \in \mathbb{Z}$$

- Definition

$$B(r, s) := \int_0^1 dt \, t^{r-1} (1-t)^{s-1}$$

- Symmetry

$$B(r, s) = B(s, r)$$

- Relationship to gamma function

$$B(r, s) = \frac{\Gamma(r)\Gamma(s)}{\Gamma(r+s)}$$

Bruce M.
Boghossian

Background
and
motivation

Gamma and
beta functions

Gamma and
beta pdfs

The chi
squared
distribution

A linear
algebra
interlude

The
independence
of \bar{Y} and S^2

Summary

Proof of relationship between beta and gamma functions

■ First note

$$\Gamma(r)\Gamma(s) = \int_0^\infty du e^{-u} u^{r-1} \int_0^\infty dv e^{-v} v^{s-1} = \int_0^\infty du \int_0^\infty dv e^{-u-v} u^{r-1} v^{s-1}$$

■ Change variables $u = zt$ and $v = z(1 - t)$, so $z = u + v$ and $t = u/(u + v)$

■ Jacobian is $J = \begin{vmatrix} t & z \\ 1-t & -z \end{vmatrix} = z$

$$\begin{aligned} \Gamma(r)\Gamma(s) &= \int_0^\infty dz \int_0^1 dt z e^{-z} (zt)^{r-1} [z(1-t)]^{s-1} \\ &= \int_0^\infty dz e^{-z} z^{r+s-1} \int_0^1 dt t^{r-1} (1-t)^{s-1} \\ &= \Gamma(r+s) B(r, s), \end{aligned}$$

from which the desired result immediately follows. □

- Definition of the two-parameter gamma pdf for $y \geq 0$

$$f_Y(y) = \frac{\lambda^r}{\Gamma(r)} y^{r-1} e^{-\lambda y}$$

- Normalization follows immediately from definition of gamma function
- Expectation:

$$E(Y) = \int_0^\infty dy \frac{\lambda^r}{\Gamma(r)} y^{r-1} e^{-\lambda y} y = \frac{r}{\lambda} \int_0^\infty dy \frac{\lambda^{r+1}}{\Gamma(r+1)} y^{(r+1)-1} e^{-\lambda y}$$

$$E(Y) = \frac{r}{\lambda}$$

■ Mean sq.:

$$E(Y^2) = \int_0^{\infty} dy \frac{\lambda^r}{\Gamma(r)} y^{r-1} e^{-\lambda y} y^2 = \frac{r(r+1)}{\lambda^2} \int_0^{\infty} dy \frac{\lambda^{r+2}}{\Gamma(r+2)} y^{r+1} e^{-\lambda y}$$

$$E(Y^2) = \frac{r(r+1)}{\lambda^2}$$

■ Variance:

$$\text{Var}(Y) = E(Y^2) - [E(Y)]^2 = \frac{r(r+1)}{\lambda^2} - \left(\frac{r}{\lambda}\right)^2$$

$$\text{Var}(Y) = \frac{r}{\lambda^2}$$

- Definition of the two-parameter beta pdf for $y \geq 0$

$$f_{\Theta}(\theta) = \frac{1}{B(r, s)} \theta^{r-1} (1 - \theta)^{s-1}$$

- Normalization follows immediately from definition of beta function
- Expectation:

$$E(\Theta) = \frac{B(r+1, s)}{B(r, s)} = \frac{\Gamma(r+1)\Gamma(s)}{\Gamma(r+s+1)} \cdot \frac{\Gamma(r+s)}{\Gamma(r)\Gamma(s)} = \frac{r\Gamma(r)\Gamma(r+s)}{(r+s)\Gamma(r+s)\Gamma(r)}$$

$$E(\Theta) = \frac{r}{r+s}$$

■ Mean square:

$$E(\Theta^2) = \frac{B(r+2, s)}{B(r, s)} = \frac{\Gamma(r+2)\Gamma(s)}{\Gamma(r+s+2)} \cdot \frac{\Gamma(r+s)}{\Gamma(r)\Gamma(s)} = \frac{r(r+1)\Gamma(r)\Gamma(r+s)}{(r+s)(r+s+1)\Gamma(r+s)\Gamma(r)}$$

$$E(\Theta^2) = \frac{r(r+1)}{(r+s)(r+s+1)}$$

■ Variance:

$$\text{Var}(\Theta) = E(\Theta^2) - [E(\Theta)]^2 = \frac{r(r+1)}{(r+s)(r+s+1)} - \left(\frac{r}{r+s}\right)^2$$

$$\text{Var}(\Theta) = \frac{rs}{(r+s)^2(r+s+1)}$$

Bruce M.
Boghossian

Background
and
motivation

Gamma and
beta functions

Gamma and
beta pdfs

The chi
squared
distribution

A linear
algebra
interlude

The
independence
of \bar{Y} and S^2

Summary

- Suppose $U, V \in \mathbb{R}$ are independent r.v.s with pdfs $f_U(u)$ and $f_V(v)$
- We wish to find the pdf of the sum $U + V$. Begin with the cdf

$$\begin{aligned} F_{U+V}(t) &= P(U + V < t) \\ &= \int_{-\infty}^{+\infty} du \int_{-\infty}^{t-u} dv f_U(u) f_V(v). \end{aligned}$$

- Differentiating both sides with respect to t yields

$$f_{U+V}(t) = \int_{-\infty}^{+\infty} du f_U(u) f_V(t - u).$$

Sums of gamma-distributed r.v.s

- Suppose U and V are independent gamma-distributed r.v.s with parameters (r, λ) and (s, λ) , respectively.
- Then $f_{U+V}(t)$ is given by a convolution

$$\begin{aligned} f_{U+V}(t) &= \int_{-\infty}^{+\infty} du f_U(u) f_V(t-u) = \int_0^t du \left[\frac{\lambda^r}{\Gamma(r)} u^{r-1} e^{-\lambda u} \right] \left[\frac{\lambda^s}{\Gamma(s)} (t-u)^{s-1} e^{-\lambda(t-u)} \right] \\ &= e^{-\lambda t} \frac{\lambda^{r+s}}{\Gamma(r)\Gamma(s)} \int_0^t du u^{r-1} (t-u)^{s-1} = e^{-\lambda t} \frac{\lambda^{r+s}}{\Gamma(r)\Gamma(s)} t^r t^{s-1} \int_0^1 dz z^{r-1} (1-z)^{s-1} \\ &= \frac{\lambda^{r+s}}{\Gamma(r)\Gamma(s)} t^{r+s-1} B(r, s) e^{-\lambda t} = \frac{\lambda^{r+s}}{\Gamma(r)\Gamma(s)} t^{r+s-1} \frac{\Gamma(r)\Gamma(s)}{\Gamma(r+s)} e^{-\lambda t} \end{aligned}$$

- So $U + V$ is also gamma-distributed, with parameters $(r + s, \lambda)$,

$$f_{U+V}(t) = \frac{\lambda^{r+s}}{\Gamma(r+s)} t^{r+s-1} e^{-\lambda t}$$

- **Thm.:** Let $U = \sum_{j=1}^n Z_j^2$ where the Z_j are iid standard normal.
- Then U is gamma-distributed with parameters $r = n/2$ and $\lambda = 1/2$,

$$f_U(u) = \frac{\left(\frac{1}{2}\right)^{n/2}}{\Gamma\left(\frac{n}{2}\right)} u^{(n/2)-1} e^{-u/2} \quad \text{where } u > 0$$

- **Pf.:** First take $n = 1$. For all $u \geq 0$,

$$F_{Z^2}(u) = P(Z^2 \leq u) = P(-\sqrt{u} \leq Z \leq +\sqrt{u}) = 2P(0 \leq Z \leq \sqrt{u}).$$

or

$$F_{Z^2}(u) = \frac{2}{\sqrt{2\pi}} \int_0^{\sqrt{u}} dz e^{-z^2/2}$$

Bruce M.
Boghossian

Background
and
motivation

Gamma and
beta functions

Gamma and
beta pdfs

The chi
squared
distribution

A linear
algebra
interlude

The
independence
of \bar{Y} and S^2

Summary

- **Pf. (continued):** We have

$$F_{Z^2}(u) = \frac{2}{\sqrt{2\pi}} \int_0^{\sqrt{u}} dz e^{-z^2/2}$$

- Differentiate to find that Z is gamma-distributed with parameters $r = 1/2$ and $\lambda = 1/2$,

$$f_{Z^2}(u) = \frac{2}{\sqrt{2\pi}\sqrt{u}} e^{-u/2} = \frac{\left(\frac{1}{2}\right)^{1/2}}{\Gamma\left(\frac{1}{2}\right)} u^{(1/2)-1} e^{-u/2}$$

- Hence, if $U = \sum_{j=1}^n Z_j^2$, it must be that U is gamma-distributed with parameters $r = n/2$ and $\lambda = 1/2$.

Bruce M.
Boghossian

Background
and
motivation

Gamma and
beta functions

Gamma and
beta pdfs

The chi
squared
distribution

A linear
algebra
interlude

The
independence
of \bar{Y} and S^2

Summary

- **Def.:** The pdf of $U = \sum_{j=1}^n Z_j^2$, where Z_j are iid standard normal, is called the *chi squared distribution with n degrees of freedom*.

$$f_{Z^2}(u) = \frac{\left(\frac{n}{2}\right)^{1/2}}{\Gamma\left(\frac{1}{2}\right)} u^{(n/2)-1} e^{-u/2}$$

- The chi squared distribution is a special case of the gamma distribution with parameters $n/2$ and $1/2$.

Bruce M.
Boghossian

Background
and
motivation

Gamma and
beta functions

Gamma and
beta pdfs

The chi
squared
distribution

A linear
algebra
interlude

The
independence
of \bar{Y} and S^2

Summary

- Column vector $v = \begin{bmatrix} v_1 \\ \vdots \\ v_n \end{bmatrix}$
- Transpose is row vector $v^T = [v_1 \ \cdots \ v_n]$
- Square of length of vector is given by

$$\|v\|^2 = v_1^2 + \cdots + v_n^2 = [v_1 \ \cdots \ v_n] \begin{bmatrix} v_1 \\ \vdots \\ v_n \end{bmatrix} = v^T v$$

- Hence length of vector is given by

$$\|v\| = \sqrt{v^T v}$$

- Consider a linear transformation of the vector, $u = Av$, where A is an $n \times n$ matrix, and demand that it preserve length

$$0 = u^T u - v^T v = (Av)^T (Av) - v^T v = v^T A^T A v - v^T v = v^T (A^T A - I) v.$$

- If we require the above to be true for all vectors v , it must be that

$$A^T A = I$$

- A matrix with this property is called an *orthogonal matrix*.
- If the square matrix A is nonsingular, postmultiplying both sides of the the above by A^{-1} yields

$$A^T = A^{-1}.$$

and premultiplying both sides of this by A yields

$$A A^T = I$$

- Let a_j denote the j th column of A
- The equation $A^T A = I$ indicated

$$a_j^T a_k = \begin{cases} 1 & \text{if } j = k \\ 0 & \text{otherwise} \end{cases}$$

- Hence the rows and columns of an orthogonal matrix are unit vectors.

Bruce M.
Boghossian

Background
and
motivation

Gamma and
beta functions

Gamma and
beta pdfs

The chi
squared
distribution

A linear
algebra
interlude

The
independence
of \bar{Y} and S^2

Summary

- Given an orthogonal matrix A , we have $A^T A = I$
- Take the determinant of both sides and use the theorems on determinants
 - $\det(AB) = \det(A) \det(B)$
 - $\det(A^T) = \det(A)$
- The result is

$$\det(A^T A) = \det(A^T) \det(A) = [\det(A)]^2 = \det(I) = 1$$

and hence

$$\det(A) = \pm 1.$$

- A transformation with $\det(A) = +1$ is a *proper orthogonal transformation*
- A transformation with $\det(A) = -1$ is an *improper orthogonal transformation*

Bruce M.
Boghossian

Background
and
motivation

Gamma and
beta functions

Gamma and
beta pdfs

The chi
squared
distribution

A linear
algebra
interlude

The
independence
of \bar{Y} and S^2

Summary

- The transformation $u = Av$ can be written $u_i = \sum_{j=1}^n A_{ij}v_j$
- The (i, j) th element of the Jacobian matrix is A_{ij} so the matrix is A ,

$$\frac{\partial u_i}{\partial v_j} = A_{ij}$$

- Jacobian factor for transforming n -dimensional integral over the v is

$$J = |\det(A)| = |\pm 1| = 1.$$

- Hence if we write $du = du_1 \cdots du_n$ and $dv = dv_1 \cdots dv_n$,

$$\int du f(u) = \int dv f(Av).$$

The transformation from X to Z I

- Let $X_j = \frac{Y_j - \mu}{\sigma}$ for $j = 1, \dots, n$
- We know the X_j are $N(0, 1)$ (standard normal)
- Let A be an orthogonal matrix whose last row is $\left[\frac{1}{\sqrt{n}} \quad \frac{1}{\sqrt{n}} \quad \cdots \quad \frac{1}{\sqrt{n}} \right]$
- Then

$$Z_n = \frac{X_1}{\sqrt{n}} + \cdots + \frac{X_n}{\sqrt{n}} = \sqrt{n} \bar{X}$$

Bruce M.
Boghosian

Background
and
motivation

Gamma and
beta functions

Gamma and
beta pdfs

The chi
squared
distribution

A linear
algebra
interlude

The
independence
of \bar{Y} and S^2

Summary

- Also, for variables z and x with $z = Ax$, we have

$$\|z\|^2 = z_1^2 + \cdots + z_n^2 = x_1^2 + \cdots + x_n^2 = \|x\|^2$$

- Since $\|x\|^2 = \|z\|^2$ and the Jacobian is one, the multivariate pdfs transform as

$$\begin{aligned} f_{X_1, \dots, X_n}(x_1, \dots, x_n) &= (2\pi)^{-n/2} \exp \left[-\frac{1}{2}(x_1^2 + \cdots + x_n^2) \right] \\ &= (2\pi)^{-n/2} \exp \left[-\frac{1}{2}(z_1^2 + \cdots + z_n^2) \right] = f_{Z_1, \dots, Z_n}(z_1, \dots, z_n) \end{aligned}$$

- Hence the Z_j are also iid $N(0, 1)$ (standard normal) r.v.s

The transformation from X to Z III

Bruce M.
Boghossian

Background
and
motivation

Gamma and
beta functions

Gamma and
beta pdfs

The chi
squared
distribution

A linear
algebra
interlude

The
independence
of \bar{Y} and S^2

Summary

- Finally we note that

$$\sum_{j=1}^n Z_j^2 = \sum_{j=1}^{n-1} Z_j^2 + n\bar{X}^2 = \sum_{j=1}^n X_j^2 = \sum_{j=1}^n (X_j - \bar{X})^2 + n\bar{X}^2$$

- Hence we have

$$\frac{1}{n-1} \sum_{j=1}^{n-1} Z_j^2 = \frac{1}{n-1} \sum_{j=1}^n (X_j - \bar{X})^2 = S^2$$

- Hence S^2 is independent of \bar{X}^2 , and hence of \bar{X} .
- Since $Y_j = \mu + \sigma X_j$, S^2 is independent of \bar{Y} .



Bruce M.
Boghossian

Background
and
motivation

Gamma and
beta functions

Gamma and
beta pdfs

The chi
squared
distribution

A linear
algebra
interlude

The
independence
of \bar{Y} and S^2

Summary

- We have provided some background and motivation for small-sample statistics
- We have reviewed the gamma and beta functions
- We have reviewed the gamma and beta pdfs
- We have learned about the chi squared distribution
- We have reviewed the linear algebra of orthogonal matrices
- We have proven the independence of \bar{Y} and S^2