

Bruce M.
Boghosian

Uniform
distribution

Definition
Likelihood and
maximum
likelihood
Estimators

The normal
distribution

Definition
Likelihood and
maximum
likelihood
Estimators

Summary

Maximum Likelihood Estimation:

The Uniform Distribution and the Normal Distribution

Bruce M. Boghosian



Tufts
UNIVERSITY

School of Arts
and Sciences

Department of Mathematics

Tufts University

©2022, all rights reserved

Bruce M.
Boghosian

Uniform
distribution

Definition
Likelihood and
maximum
likelihood
Estimators

The normal
distribution

Definition
Likelihood and
maximum
likelihood
Estimators

Summary

- 1 Uniform distribution
 - Definition
 - Likelihood and maximum likelihood
 - Estimators
- 2 The normal distribution
 - Definition
 - Likelihood and maximum likelihood
 - Estimators
- 3 Summary

Bruce M.
Boghossian

Uniform
distribution

Definition
Likelihood and
maximum
likelihood
Estimators


The normal
distribution

Definition
Likelihood and
maximum
likelihood
Estimators

Summary

- $X \in \mathbb{R}$ is a *continuous random variable*
- $X \in \mathbb{R}$ has the *uniform probability density function*,

$$f_X(x) = \begin{cases} \frac{1}{b-a} & \text{if } x \in [a, b] \\ 0 & \text{otherwise} \end{cases}$$



- Given experimental results $\vec{x} = \{x_1, x_2, \dots, x_n\}$, can we estimate a and b ?

Bruce M.
Boghossian

Uniform
distribution

Definition
Likelihood and
maximum
likelihood
Estimators

The normal
distribution

Definition
Likelihood and
maximum
likelihood
Estimators

Summary

■ Normalization:

$$\int_{\mathbb{R}} dx f_X(x) = \int_a^b dx \frac{1}{b-a} = \frac{b-a}{b-a} = 1$$

■ Mean:

$$\int_{\mathbb{R}} dx f_X(x)x = \int_a^b dx \frac{x}{b-a} = \frac{b+a}{2}$$

■ Variance:

$$\int_{\mathbb{R}} dx f_X(x)x^2 - \left(\frac{b+a}{2}\right)^2 = \int_a^b dx \frac{x^2}{b-a} - \left(\frac{b+a}{2}\right)^2 = \frac{(b-a)^2}{12}$$

■ Distribution has two parameters, a and b .

Bruce M.
Boghossian

Uniform
distribution
Definition
Likelihood and
maximum
likelihood
Estimators

The normal
distribution
Definition
Likelihood and
maximum
likelihood
Estimators

Summary

- We can be certain that $a \leq x_j \leq b$ for all $j = 1, \dots, n$.
- We seek two estimators, $\hat{a}(\vec{x})$ and $\hat{b}(\vec{x})$.
- Neither a nor b is the mean in this case, so these estimators should be unrelated to the average of the data.

Bruce M.
Boghossian

Uniform
distribution

Definition
Likelihood and
maximum
likelihood
Estimators


The normal
distribution

Definition
Likelihood and
maximum
likelihood
Estimators

Summary

- Given n iid samples of X from uniform distribution,

$$f_X(x) = \begin{cases} \frac{1}{b-a} & \text{if } x \in [a, b] \\ 0 & \text{otherwise} \end{cases}$$



- The *likelihood* is given by

$$L(a, b; \vec{x}) = \prod_{j=1}^n f_X(x_j) = \begin{cases} \left(\frac{1}{b-a}\right)^n & \text{if } x_j \in [a, b] \text{ for all } j = 1, \dots, n \\ 0 & \text{if } x_j \notin [a, b] \text{ for any } j = 1, \dots, n \end{cases}$$

- Choose $a \leq \min_j x_j$ and $b \geq \max_j x_j$ so result is $\left(\frac{1}{b-a}\right)^n$.
- Maximize result by choosing *estimates*

$$a_e = \min_j x_j \quad \text{and} \quad b_e = \max_j x_j.$$

Bruce M.
Boghosian

Uniform
distribution

Definition
Likelihood and
maximum
likelihood
Estimators

The normal
distribution

Definition
Likelihood and
maximum
likelihood
Estimators

Summary

- Maximize result by choosing *estimates*

$$a_e = \min_j x_j \quad \text{and} \quad b_e = \max_j x_j.$$

- The maximum likelihood *estimators* for a and b are then

$$\hat{a}(\vec{x}) = \min_j x_j$$

$$\hat{b}(\vec{x}) = \max_j x_j$$

- Note that $\hat{a}(\vec{x})$ always overestimates the true a .
- Note that $\hat{b}(\vec{x})$ always underestimates the true b .

Bruce M.
Boghossian

Uniform
distribution

Definition
Likelihood and
maximum
likelihood
Estimators

The normal
distribution

Definition
Likelihood and
maximum
likelihood
Estimators

Summary

- $X \in \mathbb{R}$ is a *continuous random variable*
- $X \in \mathbb{R}$ has the *normal probability density function*,

$$f_X(x) = \frac{1}{\sqrt{2\pi v}} \exp \left[-\frac{(x - \mu)^2}{2v} \right],$$

which is normalized, with mean μ and variance $v = \sigma^2$.

- Distribution has two parameters μ and v .
- Given experimental results $\vec{x} = \{x_1, x_2, \dots, x_n\}$, can we estimate μ and v ?

Bruce M.
Boghossian

Uniform
distribution

Definition
Likelihood and
maximum
likelihood
Estimators

The normal
distribution

Definition
Likelihood and
maximum
likelihood
Estimators

Summary

- The *likelihood* is given by

$$L(\mu, \nu; \vec{x}) = \prod_{j=1}^n f_X(x_j) = \prod_{j=1}^n \frac{1}{\sqrt{2\pi\nu}} \exp \left[-\frac{(x_j - \mu)^2}{2\nu} \right]$$

- The *log likelihood* is

$$\begin{aligned} \ln L(\mu, \nu; \vec{x}) &= \sum_{j=1}^n \left[-\frac{1}{2} \ln(2\pi\nu) - \frac{(x_j - \mu)^2}{2\nu} \right] \\ &= -\frac{n}{2} \ln(2\pi\nu) - \frac{1}{2\nu} \sum_{j=1}^n (x_j - \mu)^2 \end{aligned}$$

- We must find the maximum with respect to *both* μ and ν .

Bruce M.
Boghossian

Uniform
distribution

Definition
Likelihood and
maximum
likelihood
Estimators

The normal
distribution

Definition
Likelihood and
maximum
likelihood
Estimators

Summary

- *Log likelihood* is

$$\ln L(\mu, v; \vec{x}) = -\frac{n}{2} \ln(2\pi v) - \frac{1}{2v} \sum_{j=1}^n (x_j - \mu)^2$$

- Set partial derivatives to zero

$$0 = \frac{\partial}{\partial \mu} \ln L(\mu, v; \vec{x}) = \frac{1}{v} \sum_{j=1}^n (x_j - \mu) = \frac{1}{v} \left(\sum_{j=1}^n x_j - n\mu \right)$$

$$0 = \frac{\partial}{\partial v} \ln L(\mu, v; \vec{x}) = -\frac{n}{2v} + \frac{1}{2v^2} \sum_{j=1}^n (x_j - \mu)^2$$

- Solving for location of maximum (μ_e, v_e) yields

$$\mu_e = \frac{1}{n} \sum_{j=1}^n x_j \quad \text{and} \quad v_e = \frac{1}{n} \sum_{j=1}^n \left[x_j - \left(\frac{1}{n} \sum_{k=1}^n x_k \right) \right]^2$$

Bruce M.
Boghossian

Uniform
distribution

Definition
Likelihood and
maximum
likelihood
Estimators

The normal
distribution

Definition
Likelihood and
maximum
likelihood
Estimators

Summary

- Maximum likelihood estimators for a and b are

$$\hat{\mu}(\vec{x}) = \bar{x} = \frac{1}{n} \sum_{j=1}^n x_j$$

$$\hat{\nu}(\vec{x}) = \frac{1}{n} \sum_{j=1}^n \left[x_j - \left(\frac{1}{n} \sum_{k=1}^n x_k \right) \right]^2$$

Bruce M.
Boghossian

Uniform
distribution

Definition
Likelihood and
maximum
likelihood
Estimators

The normal
distribution

Definition
Likelihood and
maximum
likelihood
Estimators

Summary

- We have extended our study of likelihood and maximum likelihood estimation to continuous random variables.
- We have applied the method to both the uniform distribution and the normal distribution.