

Bruce M.
Boghosian

The
Generalized
Likelihood
Ratio (GLR)

Example: GLR
for uniform
distribution

Hypothesis
testing with
GLR

Example: GLR
for Binomial
distribution

Summary

Hypothesis testing and decision rules

The Generalized Likelihood Ratio

Bruce M. Boghosian



Tufts
UNIVERSITY

School of Arts
and Sciences

Department of Mathematics

Tufts University

Bruce M.
Boghosian

The
Generalized
Likelihood
Ratio (GLR)

Example: GLR
for uniform
distribution

Hypothesis
testing with
GLR

Example: GLR
for Binomial
distribution

Summary

- 1 The Generalized Likelihood Ratio (GLR)
- 2 Example: GLR for uniform distribution
- 3 Hypothesis testing with GLR
- 4 Example: GLR for Binomial distribution
- 5 Summary

Bruce M.
Boghosian

The
Generalized
Likelihood
Ratio (GLR)

Example: GLR
for uniform
distribution

Hypothesis
testing with
GLR

Example: GLR
for Binomial
distribution

Summary

- Suppose y_1, y_2, \dots, y_n is random sample from the uniform pdf on $[0, \theta]$
- The parameter θ is unknown
- We wish to conduct a hypothesis test at level of significance α between the alternatives
 - $H_0 : \theta = \theta_0$
 - $H_1 : \theta < \theta_0$

Bruce M.
Boghossian

The
Generalized
Likelihood
Ratio (GLR)

Example: GLR
for uniform
distribution

Hypothesis
testing with
GLR

Example: GLR
for Binomial
distribution

Summary

- Define ω to be the set of parameter values possible under the constraints of H_0
- Define Ω to be the set of all unknown parameters
- In the example of the uniform distribution on $[0, \theta]$,
 - $\omega = \{\theta \mid \theta = \theta_0\}$
 - $\Omega = \{\theta \mid 0 < \theta \leq \theta_0\}$
- Note that $\omega \subset \Omega$

Bruce M.
Boghossian

The
Generalized
Likelihood
Ratio (GLR)

Example: GLR
for uniform
distribution

Hypothesis
testing with
GLR

Example: GLR
for Binomial
distribution

Summary

- Let L be the likelihood function which, in our example, is

$$L(\theta) = \prod_j^n f_Y(y_j; \theta) = \begin{cases} \left(\frac{1}{\theta}\right)^n & \text{if } y_{\max} \leq \theta \\ 0 & \text{otherwise} \end{cases}$$

- Maximize likelihood over all $\theta \in \omega$, in other words $\theta = \theta_0$
 - This will be L evaluated at θ_0
 - $\max_{\theta \in \omega} L(\theta) = \left(\frac{1}{\theta_0}\right)^n$
- Maximize likelihood over all $\theta \in \Omega$.
 - This will be L evaluated at the maximum likelihood estimate
 - $\max_{\theta \in \Omega} L(\theta) = \left(\frac{1}{y_{\max}}\right)^n$

Bruce M.
Boghosian

The
Generalized
Likelihood
Ratio (GLR)

Example: GLR
for uniform
distribution

Hypothesis
testing with
GLR

Example: GLR
for Binomial
distribution

Summary

- *Generalized Likelihood Ratio (GLR)* is then defined to be

$$\lambda = \frac{\max_{\theta \in \omega} L(\theta)}{\max_{\theta \in \Omega} L(\theta)}$$

- For our example, we have

$$\lambda = \frac{\max_{\theta \in \omega} L(\theta)}{\max_{\theta \in \Omega} L(\theta)} = \frac{\left(\frac{1}{\theta_0}\right)^n}{\left(\frac{1}{y_{\max}}\right)^n} = \left(\frac{y_{\max}}{\theta_0}\right)^n$$

- Note that λ will be positive, but always strictly ≤ 1 .
- Values of λ near one are compatible with H_0
- Values of λ near zero suggest that we reject H_0

Bruce M.
Boghossian

- Let y_1, y_2, \dots, y_n be random sample from $f_Y(y; \theta_1, \dots, \theta_k)$
- The GLR is defined to be

$$\lambda = \frac{\max_{\theta \in \omega} L(\theta_1, \dots, \theta_k)}{\max_{\theta \in \Omega} L(\theta_1, \dots, \theta_k)}$$

- Note that the generalization to k parameters is, in principle, straightforward.

The
Generalized
Likelihood
Ratio (GLR)

Example: GLR
for uniform
distribution

Hypothesis
testing with
GLR

Example: GLR
for Binomial
distribution

Summary

Bruce M.
Boghossian

The
Generalized
Likelihood
Ratio (GLR)

Example: GLR
for uniform
distribution

Hypothesis
testing with
GLR

Example: GLR
for Binomial
distribution

Summary

- Recall that we have determined
 - Note that λ will be positive, but always strictly ≤ 1 .
 - Values of λ near one are compatible with H_0
 - Values of λ near zero suggest that we reject H_0
- GLR Test (GLRT) rejects H_0 whenever $0 < \lambda \leq \lambda^*$
- Here λ^* is chosen to satisfy

$$P(0 < \Lambda \leq \lambda^* \mid H_0 \text{ is true}) = \alpha,$$

where Λ is the random variable associated with λ .

- Note that if we knew $f_\Lambda(\lambda \mid H_0)$, then λ^* could be determined from

$$\alpha = \int_0^{\lambda^*} d\lambda f_\Lambda(\lambda \mid H_0)$$

Bruce M.
Boghosian

The
Generalized
Likelihood
Ratio (GLR)

Example: GLR
for uniform
distribution

Hypothesis
testing with
GLR

Example: GLR
for Binomial
distribution

Summary

- We would like to choose our cutoff so that

$$\alpha = \int_0^{\lambda^*} d\lambda f_{\Lambda}(\lambda | H_0)$$

- Unfortunately, the pdf $f_{\Lambda}(\lambda | H_0)$ may not be so easy to determine
- For the case of $\lambda = \left(\frac{y_{\max}}{\theta_0}\right)^n$, we have

$$\begin{aligned} \alpha &= P(\Lambda \leq \lambda^* | H_0 \text{ is true}) \\ &= P\left[\left(\frac{Y_{\max}}{\theta_0}\right)^n \leq \lambda^* | H_0 \text{ is true}\right] \\ &= P\left(Y_{\max} \leq \theta_0 \sqrt[n]{\lambda^*} | H_0 \text{ is true}\right) \end{aligned}$$

Bruce M.
Boghossian

The
Generalized
Likelihood
Ratio (GLR)

Example: GLR
for uniform
distribution

Hypothesis
testing with
GLR

Example: GLR
for Binomial
distribution

Summary

- From last page

$$\begin{aligned}
 \alpha &= P\left(Y_{\max} \leq \theta_0 \sqrt[n]{\lambda^*} \mid H_0 \text{ is true}\right) \\
 &= F_{Y_{\max}}\left(\theta_0 \sqrt[n]{\lambda^*}\right) \\
 &= \left[F_Y\left(\theta_0 \sqrt[n]{\lambda^*}\right)\right]^n \\
 &= \left(\frac{\theta_0 \sqrt[n]{\lambda^*}}{\theta_0}\right)^n \\
 &= \lambda^*
 \end{aligned}$$

- So reject H_0 if $\lambda \leq \lambda^* = \alpha$, or equivalently $y_{\max} \leq \theta_0 \sqrt[n]{\alpha}$

Bruce M.
Boghossian

The
Generalized
Likelihood
Ratio (GLR)

Example: GLR
for uniform
distribution

Hypothesis
testing with
GLR

Example: GLR
for Binomial
distribution

Summary

- Bernoulli trial $p_X(k; p) = p^k(1 - p)^{1-k}$
- Likelihood is

$$L(p) = \prod_j^n p^{k_j}(1 - p)^{1-k_j} = p^k(1 - p)^{n-k}$$

where $k = \sum_j^n k_j$.

- Log likelihood $\ln L(p) = k \ln p + (n - k) \ln(1 - p)$
- Max likelihood $0 = \frac{d}{dp} \ln L(p) = \frac{k}{p} - \frac{n-k}{1-p}$ so $p_e = \frac{k}{n}$
- Max likelihood estimator $\hat{p}(\vec{X}) = \frac{1}{n} \sum_j^n X_j = \bar{X}$

Bruce M.
Boghossian

The
Generalized
Likelihood
Ratio (GLR)

Example: GLR
for uniform
distribution

Hypothesis
testing with
GLR


Example: GLR
for Binomial
distribution

Summary

- Test $H_0 : p = p_0$ versus $H_1 : p \neq p_0$
- $\max_{p \in \omega} L(p) = L(p_0) = p_0^k (1 - p_0)^{n-k}$
- $\max_{p \in \Omega} L(p) = L(\hat{p}) = \left(\frac{k}{n}\right)^k \left(1 - \frac{k}{n}\right)^{n-k}$
- GLR is then

$$\lambda = \frac{\max_{p \in \omega} L(p)}{\max_{p \in \Omega} L(p)} = \frac{p_0^k (1 - p_0)^{n-k}}{\left(\frac{k}{n}\right)^k \left(1 - \frac{k}{n}\right)^{n-k}} = \left(\frac{p_0}{k/n}\right)^k \left(\frac{1 - p_0}{1 - k/n}\right)^{n-k}$$

- Example for $n = 200$ and $p_0 = 0.6$ (so $np_0 = 120$):



Bruce M.
Boghossian

The
Generalized
Likelihood
Ratio (GLR)

Example: GLR
for uniform
distribution

Hypothesis
testing with
GLR

Example: GLR
for Binomial
distribution

Summary

- Demand that

$$\alpha = \text{Prob}(\lambda < \lambda^*) = \text{Prob}\left(\left(\frac{p_0}{k/n}\right)^k \left(\frac{1-p_0}{1-k/n}\right)^{n-k} < \lambda^*\right)$$

- Given p_0 , work out correspondence between λ^* and α .
- **Ex.:** For $n = 200$ and $p_0 = 0.60$, and taking $\lambda^* = 0.25$
 - Reject H_0 if $\lambda < \lambda^*$
 - Reject H_0 if $k \leq 108$ or $k \geq 132$

$$\begin{aligned} \alpha &= \sum_{k=0}^{108} \binom{200}{k} (0.6)^k (1-0.6)^{200-k} \\ &\quad + \sum_{k=132}^{200} \binom{200}{k} (0.6)^k (1-0.6)^{200-k} = 0.0972 \end{aligned}$$

- About 90% confidence we will not reject H_0 if it is true.

Bruce M.
Boghosian

The
Generalized
Likelihood
Ratio (GLR)

Example: GLR
for uniform
distribution

Hypothesis
testing with
GLR

Example: GLR
for Binomial
distribution

Summary

- **Ex.:** For $n = 200$ and $p_0 = 0.60$, and taking $\lambda^* = 0.15$
 - Reject H_0 if $\lambda < \lambda^*$
 - Reject H_0 if $k \leq 106$ or $k \geq 134$

$$\alpha = \sum_{k=0}^{106} \binom{200}{k} (0.6)^k (1 - 0.6)^{200-k} + \sum_{k=134}^{200} \binom{200}{k} (0.6)^k (1 - 0.6)^{200-k} = 0.0513$$

- About 95% confidence we will not reject H_0 if it is true.

Bruce M.
Boghosian

The
Generalized
Likelihood
Ratio (GLR)

Example: GLR
for uniform
distribution

Hypothesis
testing with
GLR

Example: GLR
for Binomial
distribution

Summary

- **Ex.:** For $n = 200$ and $p_0 = 0.60$, and taking $\lambda^* = 0.0369$
 - Reject H_0 if $\lambda < \lambda^*$
 - Reject H_0 if $k \leq 102$ or $k \geq 139$

$$\alpha = \sum_{k=0}^{102} \binom{200}{k} (0.6)^k (1 - 0.6)^{200-k} + \sum_{k=139}^{200} \binom{200}{k} (0.6)^k (1 - 0.6)^{200-k} = 0.00913$$

- About 99% confidence we will not reject H_0 if it is true.

Bruce M.
Boghosian

The
Generalized
Likelihood
Ratio (GLR)


Example: GLR
for uniform
distribution

Hypothesis
testing with
GLR

Example: GLR
for Binomial
distribution

Summary

- **Ex.:** For $n = 200$ and $p_0 = 0.60$, plot α versus λ^*
- Smaller α results from smaller thresholds λ^* .
- Discrete “steps” due to discreteness of r.v. $k \in \mathbb{Z}$.



Bruce M.
Boghosian

The
Generalized
Likelihood
Ratio (GLR)

Example: GLR
for uniform
distribution

Hypothesis
testing with
GLR

Example: GLR
for Binomial
distribution

Summary

- We introduced and defined Generalized Likelihood Ratio
- We demonstrated how to use GLR for hypothesis testing
 - For uniform distribution
 - For binomial distribution
- Reject H_0 when $\lambda < \lambda^*$.
- Hence smaller λ^* mean smaller α , and greater confidence against rejecting H_0 if it is true.