

Bruce M.
Boghosian

A loose end
from last time

Bayes'
Theorem

Example

Bayesian
search
strategy

Application to
estimation

Summary

Bayesian estimation

Bruce M. Boghosian



Tufts
UNIVERSITY

School of Arts
and Sciences

Department of Mathematics

Tufts University

Bruce M.
Boghosian

A loose end
from last time

Bayes'
Theorem

Example

Bayesian
search
strategy

Application to
estimation

Summary

1 A loose end from last time

2 Bayes' Theorem

3 Example

4 Bayesian search strategy

5 Application to estimation

6 Summary

Bruce M.
Boghossian

A loose end
from last time

Bayes'
Theorem

Example

Bayesian
search
strategy

Application to
estimation

Summary

- Recall MLE estimator $\hat{\sigma}^2$ for the variance of the normal is not unbiased,

$$E \left[\hat{\sigma}^2(\vec{Y}) \right] = \frac{n-1}{n} \sigma^2$$

- but it is *asymptotically unbiased* since

$$E \left[\lim_{n \rightarrow \infty} \left(\hat{\sigma}^2(\vec{Y}) - \sigma^2 \right) \right] = \lim_{n \rightarrow \infty} \left(\frac{n-1}{n} \sigma^2 - \sigma^2 \right) = 0.$$

Bruce M.
Boghossian

A loose end
from last time

Bayes'
Theorem

Example

Bayesian
search
strategy

Application to
estimation

Summary

- Given n possible events $\{A_j\}_{j=1}^n$, each with some corresponding probability
- These events *partition* the set of all such possibilities A :
 $A = \bigcup_{j=1}^n A_j = A_1 \cup A_2 \cup \dots \cup A_n$ (Every possible outcome is accounted for)
 $A_j \cap A_k = \emptyset$ if $j \neq k$ (Outcomes are mutually exclusive)
- We may have an initial guesses for the *prior distribution*, $\{P(A_j)\}_{j=1}^n$.
- Now event let B be some new observation related to the above events.
- We wish to update the $\{P(A_j)\}_{j=1}^n$, given the observation B .

Bayesian estimation: The problem

Bruce M.
Boghossian

A loose end
from last time

Bayes'
Theorem

Example

Bayesian
search
strategy

Application to
estimation

Summary

- We wish to update the $\{P(A_j)\}_{j=1}^n$, given the observation B .
- That is, we wish to find the *posterior distribution*, $\{P(A_j | B)\}_{j=1}^n$.
- If there is a subsequent guess, the process could be repeated.
- In this way, the algorithm “learns” from new information.
- According to the “Bayesians”, you should not suppose that a new piece of information will give you all the answers you are seeking. Rather, the new information should update prior beliefs.

- Recall from the definition of conditional probability

$$P(A_j, B) = P(A_j | B)P(B) = P(B | A_j)P(A_j).$$

$$\therefore P(A_j | B) = \frac{P(B | A_j)P(A_j)}{P(B)}$$

- Then recall that, since the $\{A_j\}_{j=1}^n$ partition A ,

$$P(B) = \sum_{k=1}^n P(B | A_k)P(A_k)$$

- This gives us *Bayes' Theorem*,

$$P(A_j | B) = \frac{P(B | A_j)P(A_j)}{\sum_{k=1}^n P(B | A_k)P(A_k)}.$$

Bruce M.
Boghossian

A loose end
from last time

Bayes'
Theorem

Example

Bayesian
search
strategy

Application to
estimation

Summary

■ *Bayes' Theorem*

$$P(A_j | B) = \frac{P(B | A_j)P(A_j)}{\sum_{k=1}^n P(B | A_k)P(A_k)}.$$

- Note numerator is one term of denominator, and all terms are positive.
- Result will always be in $[0, 1]$, as expected.
- Gives the desired $P(A_j | B)$ in terms of things that we might actually know, or be able to measure, or at least be able to estimate.
- The results of thinking in this way can defy intuition.
- An enormous body of scientific conclusions are gleaned in exactly this way.
- Key question: How do we use this theorem in the context of a real problem?

Example: Librarians and Farmers

Bruce M.
Boghossian

A loose end
from last time

Bayes'
Theorem

Example

Bayesian
search
strategy

Application to
estimation

Summary

- Example due to D. Kahneman and A. Tversky, as presented in “Thinking Fast and Slow” by D. Kahneman, FSG Publishers, New York (2011) pp. 6-7.
- **Given information:** “An individual has been described by a neighbor as follows: ‘Steve is very shy and withdrawn, invariably helpful but with very little interest in people or the world of reality. A meek and tidy soul, he has a need for order and structure, and a passion for detail.’ ”
- **Question:** “Is Steve more likely to be a librarian or a farmer?”
- According to Kahneman, the vast majority of people who are asked this question say that Steve is more likely to be a librarian, almost surely due to the stereotypical view of librarians as having these personality traits.
- Does this make sense? Are we missing any information?

Example: Librarians and Farmers (continued)

Bruce M.
Boghossian

A loose end
from last time

Bayes'
Theorem

Example

Bayesian
search
strategy

Application to
estimation

Summary

- Does this make sense? Are we missing information?
- Kahneman: There are $20\times$ as many farmers as librarians in the US.
- Consider a sample of 105 people consisting of 100 farmers and 5 librarians.
- Say 40% of librarians have this personality trait and only 10% of farmers.
- That means 2 librarians and 10 farmers in our sample have the trait.
- Given that a randomly selected person has this trait, there is only a $2/12 = 1/6$ chance that they are a librarian, and a $10/12 = 5/6$ chance that they are a farmer.
- Kahneman: "Because there are so many farmers, it is almost certain that more 'meek and tidy' will be found on tractors than at library information desks."
- Clearly we *were* missing information. The ratio of farmers to librarians *matters* to the question as posed! Most people don't even consider this possibility!

Bruce M.
Boghossian

A loose end
from last time

Bayes'
Theorem

Example

Bayesian
search
strategy

Application to
estimation

Summary

- Decide on your events and give them variable names:

- B = Steve is a “meek and tidy soul”
- A_1 = Steve is a librarian
- A_2 = Steve is a farmer

- We know that $P(A_1) = \frac{1}{21}$ and $P(A_2) = \frac{20}{21}$.

- We estimate $P(B | A_1) = \frac{4}{10}$, and $P(B | A_2) = \frac{1}{10}$.

- Then the probability that Steve is a librarian given that he is a “meek and tidy soul” is

$$P(A_1 | B) = \frac{P(B | A_1)P(A_1)}{P(B | A_1)P(A_1) + P(B | A_2)P(A_2)} = \frac{\frac{4}{10} \cdot \frac{1}{21}}{\frac{4}{10} \cdot \frac{1}{21} + \frac{1}{10} \cdot \frac{20}{21}} = \frac{4}{24} = \frac{1}{6},$$

in agreement with our earlier calculation.

Bruce M.
Boghossian

A loose end
from last time

Bayes'
Theorem

Example

Bayesian
search
strategy

Application to
estimation

Summary

- $A_j :=$ Event that missing object is in region r_j
- $B_j :=$ Event that object would be found if it was in r_j , and r_j was searched
- $P(A_j | B_j^C)$ is probability that the item is in r_j , given
- Start with *prior* estimates for the probabilities $\{P(A_j)\}_{j=1}^n$. This could be as simple as the uniform distribution $\forall j \in \{1, \dots, n\} : P(A_j) = \frac{1}{n}$.
- Suppose a search is undertaken in r_j , and nothing is found there.

- Apply Bayes' Theorem, and note that $P(B_j^C | A_j^C) = 1$

$$\begin{aligned} P(A_j | B_j^C) &= \frac{P(B_j^C | A_j) P(A_j)}{P(B_j^C | A_j) P(A_j) + P(B_j^C | A_j^C) P(A_j^C)} \\ &= \frac{[1 - P(B_j | A_j)] P(A_j)}{[1 - P(B_j | A_j)] P(A_j) + [1 - P(A_j)]} = \left[\frac{1 - P(B_j | A_j)}{1 - P(B_j | A_j) P(A_j)} \right] P(A_j) \end{aligned}$$

- Hence, after unsuccessful search in r_j , prior for $P(A_j)$ is updated to obtain

$$P^*(A_j) = P(A_j | B_j^C) = \left[\frac{1 - P(B_j | A_j)}{1 - P(B_j | A_j) P(A_j)} \right] P(A_j) < P(A_j).$$

Updating the priors for the other regions

- Rescale all the other priors, $P(A_k)$ where $k \neq j$,

$$P^*(A_k) = P(A_k | B_j^C) = \alpha_j P(A_k)$$

- Demand normalization $1 = \sum_{m=1}^n P^*(A_m)$ to find

$$\alpha_j = \frac{1}{1 - P(B_j | A_j)P(A_j)}$$

- So the other priors update as follows

$$P^*(A_k) = P(A_k | B_j^C) = \left[\frac{1}{1 - P(B_j | A_j)P(A_j)} \right] P(A_k) > P(A_k).$$

Bayesian search strategy algorithm

Bruce M.
Boghosian

A loose end
from last time

Bayes'
Theorem

Example

Bayesian
search
strategy

Application to
estimation

Summary

- For all $m = 1, \dots, n$:
 - Determine the probabilities that a search in r_m will find the object if it is in r_m .
 - Guess the priors $P(A_m)$.
- Now repeat until the object is found:
 - Suppose that the value of m for which $P(A_m)$ is highest is $m = j$.
 - Conduct a search in region r_j
 - If the search is unsuccessful, update the priors and repeat.

- Let W be a statistic dependent on a parameter θ . Call its pdf $f_W(w | \theta)$.
- Suppose that θ is the value of a random variable Θ .
- Denote the prior distribution of Θ by
 - $p_\Theta(\theta)$ if Θ is discrete
 - $f_\Theta(\theta)$ if Θ is continuous
- Posterior distribution of Θ , given that $W = w$ is the quotient

$$g_\Theta(\theta | W = w) = \begin{cases} \frac{p_W(w | \theta) f_\Theta(\theta)}{\int d\xi p_W(w | \xi) f_\Theta(\xi)} & \text{if } W \text{ is discrete} \\ \frac{p_W(w | \theta) f_\Theta(\theta)}{\int d\xi f_W(w | \xi) f_\Theta(\xi)} & \text{if } W \text{ is continuous} \end{cases}$$

- If Θ is discrete
 - Replace the $f_\Theta(\theta)$ by $p_\Theta(\theta)$.
 - Replace the integrals over θ by sums.

Bruce M.
Boghossian

A loose end
from last time

Bayes'
Theorem

Example

Bayesian
search
strategy

Application to
estimation

Summary

- We have reviewed Bayes' Theorem
- We have given an example of Bayesian reasoning
- We have studied Bayesian search strategy
- We have studied Bayesian estimation