

Bruce M.
Boghosian

Introduction
and
motivation

Pearson's
Goodness of
Fit Test

Benford's Law

Summary

Goodness of Fit Tests

All Parameters Known

Bruce M. Boghosian



Tufts
UNIVERSITY

School of Arts
and Sciences

Department of Mathematics

Tufts University

Bruce M.
Boghosian

Introduction
and
motivation

Pearson's
Goodness of
Fit Test

Benford's Law

Summary

- 1 Introduction and motivation
- 2 Pearson's Goodness of Fit Test
- 3 Benford's Law
- 4 Summary

Bruce M.
Boghosian

Introduction
and
motivation

Pearson's
Goodness of
Fit Test

Benford's Law

Summary

- Suppose that you can specify both of
 - the model distribution for your data, and
 - all parameters of that distribution.
- You would like to check GoF for your data to
 - the known model distribution,
 - with the known parameters.

Bruce M.
Boghosian

Introduction
and
motivation

Pearson's
Goodness of
Fit Test

Benford's Law

Summary

- Let $f_Y(y)$ be the true pdf.
- Let $f_0(y)$ be the presumed pdf.
- Null and alternative hypotheses:

$$H_0 : f_Y(y) = f_0(y)$$

$$H_1 : f_Y(y) \neq f_0(y)$$

Bruce M.
Boghossian

Introduction
and
motivation

Pearson's
Goodness of
Fit Test

Benford's Law

Summary

- Let $p_X(k)$ be the true probability distribution.
- Let $p_0(k)$ be the presumed probability distribution.
- Null and alternative hypotheses:

$$H_0 : p_X(k) = p_0(k)$$

$$H_1 : p_X(k) \neq p_0(k)$$

- Another way to describe H_0 and H_1 for discrete r.v.s:

$$H_0 : p_1 = p_{1_0}, p_2 = p_{2_0}, \dots, p_t = p_{t_0}$$

$$H_1 : p_j \neq p_{j_0} \text{ for at least one } j \in \{1, \dots, t\}$$

Bruce M.
Boghosian

Introduction
and
motivation

Pearson's
Goodness of
Fit Test

Benford's Law

Summary

- **Thm.:** Let r_1, \dots, r_t be the set of possible outcomes associated with each of n independent trials, where $P(r_i) = p_i$ for $i = 1, \dots, t$. Let the r.v. X_i be the number of times r_i occurs for $i = 1, \dots, t$.

- The r.v.

$$D = \sum_{i=1}^t \frac{(X_i - np_i)^2}{np_i}$$

has approximately a χ^2 distribution with $t - 1$ degrees of freedom. (For the approximation to be adequate, the t classes should be defined so that $np_i \geq 5$, for all i .)

Bruce M.
Boghosian

Introduction
and
motivation

Pearson's
Goodness of
Fit Test

Benford's Law

Summary

- **Thm. (continued):** Let k_1, \dots, k_t be the observed frequencies for outcomes r_1, \dots, r_t , respectively, and let $np_{1_0}, \dots, np_{t_0}$ be the corresponding expected frequencies, based on the null hypothesis.
- At the α level of significance, $H_0 : f_Y(y) = f_0(y)$ (or similar discrete H_0) is rejected if

$$d = \sum_{i=1}^t \frac{(k_i - np_{i_0})^2}{np_{i_0}} \geq \chi_{1-\alpha, t-1}^2,$$

where, again, $np_{i_0} \geq 5$ for all $i = 1, \dots, t$.

Bruce M.
Boghossian

Introduction
and
motivation

Pearson's
Goodness of
Fit Test

Benford's Law

Summary

- **Pf.:** A full proof is beyond the scope of this course.
- We can, however, motivate the case where $t = 2$.

$$\begin{aligned} D &= \frac{(X_1 - np_1)}{np_1} + \frac{(X_2 - np_2)}{np_2} \\ &= \frac{(X_1 - np_1)}{np_1} + \frac{[n - X_1 - n(1 - p_1)]}{n(1 - p_1)} \\ &= \frac{(X_1 - np_1)^2}{np_1(1 - p_1)} \\ &= \left[\frac{X_1 - E(X_1)}{\sqrt{\text{Var}(X_1)}} \right]^2 \end{aligned}$$

- Note D is the square of a variable that is asymptotically a standard normal, so it is χ^2 distributed with $2 - 1 = 1$ degrees of freedom.

- **Comment (given without proof):** A decision rule based on D is asymptotically equivalent to the GLRT of

$$H_0 : p_1 = p_{1_0}, \dots, p_t = p_{t_0}.$$

- Likelihood function

$$L(p_1, \dots, p_t) = \prod_{j=1}^n \frac{n!}{k_{1j}! \dots k_{tj}!} p_1^{k_{1j}} \dots p_t^{k_{tj}}$$

$$\begin{aligned} \therefore \ln L(p_1, \dots, p_t) &= \sum_{j=1}^n \left[\ln \left(\frac{n!}{k_{1j}! \dots k_{tj}!} \right) + k_{1j} \ln p_1 + \dots + k_{tj} \ln p_t \right] \\ &= K_1 \ln p_1 + \dots + K_t \ln p_t + \sum_{j=1}^n \ln \left(\frac{n!}{k_{1j}! \dots k_{tj}!} \right) \end{aligned}$$

where $K_i := \sum_{j=1}^n k_{ij}$.

- We have

- $\Omega = \{(p_1, \dots, p_t) \mid p_1 + \dots + p_t = 1\}$

- $\omega = \{(p_1, \dots, p_t) \mid p_1 = p_{1_0}, \dots, p_t = p_{t_0}\}$

- To find $\max_{\Omega} L(p_1, \dots, p_t)$, use Lagrange multiplier μ

$$0 = \frac{\partial}{\partial p_i} [\ln L(p_1, \dots, p_t) - \mu (p_1 + \dots + p_t)] = \frac{K_i}{p_i} - \mu,$$

so $p_i = \frac{K_i}{\mu}$, whence the normalization condition gives

$$p_i = \kappa_i := \frac{K_i}{K_1 + \dots + K_t}.$$

- It follows that

$$\max_{\Omega} L(p_1, \dots, p_t) = L(\kappa_1, \dots, \kappa_t).$$

Bruce M.
Boghosian

Introduction
and
motivation

Pearson's
Goodness of
Fit Test

Benford's Law

Summary

- We also have

$$\max_{\omega} L(p_1, \dots, p_t) = L(p_{1_0}, \dots, p_{t_0})$$

- So the GLR is

$$\lambda = \frac{\max_{\omega} L(p_1, \dots, p_t)}{\max_{\Omega} L(p_1, \dots, p_t)} = \frac{L(p_{1_0}, \dots, p_{t_0})}{L(\kappa_1, \dots, \kappa_t)}.$$

where

$$\kappa_i := \frac{K_i}{K_1 + \dots + K_t}.$$

- Relating this to D in the asymptotic limit of large n is not straightforward.

Bruce M.
Boghosian

Introduction
and
motivation

Pearson's
Goodness of
Fit Test

Benford's Law

Summary

- Many years ago, the astronomer Simon Newcomb noticed that the first pages of tables of logarithms are more smudged from use than later pages.
- Leading digit of number in scientific notation is 1 to 9.
- In tables of numbers, including seemingly random data and statistics of various sorts, you might think that each leading digit would appear with probability $1/9 \approx 11.1\%$.
- Instead numbers are observed to lead off with the digit 1 about 30% of the time, with digit 2 about 17.6% of the time, etc. They lead off with 9 only 4.6% of the time.
- This claim was checked by Frank Benford in the 1930s. It is called *Benford's Law*, or the *Newcomb-Benford Law*.

Bruce M.
Boghossian

Introduction
and
motivation

Pearson's
Goodness of
Fit Test

Benford's Law

Summary

- T.P. Hill (1998)
- Benford's Law applies to dimensioned data – with units.
- The probability distribution of real numbers in scientific notation surely can not depend on these units, so it must be invariant under scaling, whence

$$P(cx) = f(c)P(x)$$

- If $\int dx P(x) = 1$, then $\int dx P(cx) = 1/c$, so $f(c) = 1/c$,

$$P(cx) = \frac{1}{c}P(x).$$

- Differentiate with respect to c : $xP'(cx) = -P(x)/c^2$.
- Set $c = 1$ to obtain $xP'(x) = -P(x)$.
- This has solution $P(x) = 1/x$.

Bruce M.
Boghosian

Introduction
and
motivation

Pearson's
Goodness of
Fit Test

Benford's Law

Summary

- $P(x) = 1/x$ is not normalizable for all x , but we need to use it only in finite intervals.
- The probability that the leading digit is k is then

$$p_D(k) = \frac{\int_k^{k+1} dx P(x)}{\int_1^{10} dx P(x)} = \frac{\log(k+1) - \log k}{\log 10 - \log 1} = \log_{10} \left(1 + \frac{1}{k} \right)$$

- Results are consistent with what is often observed

d	$p_D(k)$
1	0.301030
2	0.176091
3	0.124939
4	0.0969100
5	0.0791812
6	0.0669468
7	0.0579919
8	0.0511525
9	0.0457575

Bruce M.
Boghossian

Introduction
and
motivation

Pearson's
Goodness of
Fit Test

Benford's Law

Summary

- Benford's Law is sometimes used in financial audits
- In first 355 digits of university's operating budget, observed digit frequency is

d	k_d
1	111
2	60
3	46
4	29
5	26
6	22
7	21
8	20
9	20

- Form the statistic

$$d = \frac{[111 - 355(0.301)]^2}{355(0.301)} + \dots + \frac{[20 - 355(0.046)]^2}{355(0.046)} = 2.49$$

- Note $d < \chi^2_{0.95,8} = 15.507$ so we fail to reject H_0 .

Bruce M.
Boghosian

Introduction
and
motivation

Pearson's
Goodness of
Fit Test

Benford's Law

Summary

- We have motivated GoF tests with known parameters.
- We have stated and justified Pearson's GoF test.
- We have derived Benford's Law.
- We have studied GoF of data to Benford's Law.