

Bruce M.
Boghosian

Motivation

Bernoulli trials

The Poisson
distribution

Uniform
distribution

Standard
normal
distribution

Summary

Maximum Likelihood Estimation

Bruce M. Boghosian



Department of Mathematics

Tufts University

Bruce M.
Boghosian

Motivation

Bernoulli trials

The Poisson
distribution

Uniform
distribution

Standard
normal
distribution

Summary

- 1 Motivation
- 2 Bernoulli trials
- 3 The Poisson distribution
- 4 Uniform distribution
- 5 Standard normal distribution
- 6 Summary

Bruce M.
Boghosian

Motivation

Bernoulli trials

The Poisson
distribution

Uniform
distribution

Standard
normal
distribution

Summary

- Suppose we know the results of a repeated random experiment.
- We have a priori knowledge of the *form* of the probability function.
- We do not know the values of the *parameter(s)* of the distribution.
- Can we use the experimental results to estimate the parameter(s)?
- In this lecture, we learn to do so by maximizing a quantity called *likelihood*.
- This method is called *maximum likelihood estimation*.
- It is best to learn this method by example, so we present examples using a variety of probability functions.

A random experiment

Bruce M.
Boghosian

Motivation

Bernoulli trials

The Poisson
distribution

Uniform
distribution

Standard
normal
distribution

Summary

- A coin lands on heads with probability p , tails with probability $1 - p$.
- You are not given the value of p .
- You flip the coin five times and find the sequence *HTHHT*.
- Suppose this outcome is *all you know* about the coin.
- What is your best guess for p ?

A posteriori calculation of probability of the outcome

Bruce M.
Boghossian

Motivation

Bernoulli trials


The Poisson
distribution

Uniform
distribution

Standard
normal
distribution

Summary

- We have done the experiment and we know the result *HTHHT*.
- The coin flips may be assumed to have been independent.
- The *likelihood* of result *HTHHT* is $L(p) = p(1-p)pp(1-p) = p^3(1-p)^2$.
- Note that $L(p)$ is not a probability density function!
- For which value of p is $L(p)$ maximized?



Bruce M.
Boghosian

Motivation

Bernoulli trials

The Poisson
distribution

Uniform
distribution

Standard
normal
distribution

Summary

- We have $L(p) = p^3(1 - p)^2$
- Derivative $L'(p) = p^2(1 - p)(3 - 5p)$ has roots $p = 0$, $p = 3/5$ and $p = 1$.
- Second derivative $L''(p) = 2p(10p^2 - 12p + 3)$, so $L''(3/5) = -18/25 < 0$
- Second derivative is negative at $p = 3/5$, indicating a maximum at that point.
- *Maximum likelihood* occurs for $p = \boxed{p_e = 3/5}$.

- Define random variable for each coin toss,

$$X := \begin{cases} 1 & \text{if toss results in heads (with probability } p) \\ 0 & \text{if toss results in tails (with probability } 1 - p). \end{cases}$$

- Discrete probability function for one toss, where $k \in \{0, 1\}$,

$$p_X(k) = \text{Prob}(X = k) = p^k(1 - p)^{1-k}$$

- Normalization: $\sum_{k=0}^1 p_X(k) = (1 - p) + p = 1$
- Mean: $\sum_{k=0}^1 p_X(k)k = (1 - p)0 + p1 = p$
- Variance: $\sum_{k=0}^1 p_X(k)k^2 - p^2 = (1 - p)0^2 + p1^2 - p^2 = p - p^2 = p(1 - p)$

Defining the likelihood for n tosses

Bruce M.
Boghosian

Motivation

Bernoulli trials

The Poisson
distribution

Uniform
distribution

Standard
normal
distribution

Summary

- *Likelihood* of n tosses with $\vec{X} = \langle X_1, X_2, \dots, X_n \rangle$ equal to $\vec{k} = \langle k_1, k_2, \dots, k_n \rangle$
- Product of discrete probability functions for observed data using parameter p ,

$$\begin{aligned} L(p; \vec{k}) &:= \text{Prob}(\vec{X} = \vec{k}) = \prod_{j=1}^n p_{X_j}(k_j) \\ &= p^{k_1}(1-p)^{1-k_1} p^{k_2}(1-p)^{1-k_2} \dots p^{k_n}(1-p)^{1-k_n} \\ &= \prod_{j=1}^n p^{k_j}(1-p)^{1-k_j} = p^K(1-p)^{n-K} \end{aligned}$$

where $K := \sum_{j=1}^n k_j$.

- We now wish to find the value of p that maximizes $L(p; \vec{k})$.

Maximizing the likelihood for n tosses

Bruce M.
Boghossian

Motivation

Bernoulli trials

The Poisson
distribution

Uniform
distribution

Standard
normal
distribution

Summary

- Defining $K := \sum_{j=1}^n k_j$, the likelihood function is

$$L(p; \vec{k}) := p^K (1-p)^{n-K}.$$

- Note that \vec{k} and hence K is known from the experimental outcome.
- We maximize the *log likelihood* with respect to p ,

$$\ln L(p; \vec{k}) = K \ln p + (n-K) \ln(1-p)$$

- Setting derivative to zero yields

$$0 = \frac{d}{dp} \ln L(p; \vec{k}) = \frac{K}{p} - \frac{n-K}{1-p}.$$

- *Maximum likelihood* is $p = p_e := \frac{K}{n}$, so

$$p_e = \frac{1}{n} \sum_{j=1}^n k_j$$

Bruce M.
Boghossian

Motivation

Bernoulli trials

The Poisson
distribution

Uniform
distribution

Standard
normal
distribution

Summary

- Maximum likelihood occurs when p is $p_e = \frac{1}{n} \sum_{j=1}^n k_j$
- Note that this is a function of the outcomes \vec{k} that estimates the parameter p .
- Considered as a function of \vec{k} yielding p , this is called an *estimator*,

$$\hat{p}(\vec{k}) = \frac{1}{n} \sum_{j=1}^n k_j.$$

- In this case, $\hat{p}(\vec{k})$ is just the *average* of the experimental outcomes \vec{k} .
- Here and henceforth, we use the “hat” to denote estimator functions.
- More generally, $L(p; \vec{k})$ is maximized for $p = \hat{p}(\vec{k})$.
- This approach is called *maximum likelihood estimation*.
- It estimates one or more parameters of known probability functions.
- Note that there must be a priori knowledge of the form of $p_X(k)$.

Bruce M.
Boghossian

Motivation

Bernoulli trials

The Poisson
distribution

Uniform
distribution

Standard
normal
distribution

Summary

- Sample space S is nonnegative integers.
- Poisson random variable $X \in S = \{0, 1, 2, \dots\}$
- We have

$$p_X(k) = \text{Prob}(X = k) = e^{-\lambda} \frac{\lambda^k}{k!}$$

- Normalization

$$\sum_{k=0}^{\infty} p_X(k) = \sum_{k=0}^{\infty} e^{-\lambda} \frac{\lambda^k}{k!} = e^{-\lambda} \sum_{k=0}^{\infty} \frac{\lambda^k}{k!} = e^{-\lambda} e^{+\lambda} = 1.$$

- Mean

$$E(X) = \sum_{k=0}^{\infty} p_X(k)k = e^{-\lambda} \sum_{k=0}^{\infty} \frac{\lambda^k}{k!} k = \lambda$$

The Poisson distribution (continued)

Bruce M.
Boghosian

Motivation
Bernoulli trials

The Poisson
distribution

Uniform
distribution

Standard
normal
distribution

Summary

■ Variance

$$E(X^2) - E(X)^2 = \sum_{k=0}^{\infty} p_X(k)k^2 - \lambda^2 = e^{-\lambda} \sum_{k=0}^{\infty} \frac{\lambda^k}{k!} k^2 - \lambda^2 = \lambda$$

■ Again, note that this is a one-parameter probability function.

■ The mean and variance are both equal to the parameter λ .

■ Now, suppose we are given $n = 50$ samples from this distribution

3, 4, 3, 5, 3, 4, 5, 0, 5, 5, 5, 4, 2, 2, 3, 2, 2, 5, 5, 5, 5, 6, 5, 2, 5,
2, 2, 0, 6, 3, 6, 7, 2, 4, 5, 1, 3, 3, 5, 1, 7, 2, 0, 8, 4, 5, 8, 2, 3, 5

■ Can we estimate λ using maximum likelihood estimation?

Defining the likelihood for n samples

Bruce M.
Boghosian

Motivation

Bernoulli trials

The Poisson
distribution

Uniform
distribution

Standard
normal
distribution

Summary

- Given the results of an experiment $\vec{k} = \langle k_1, k_2, \dots \rangle$ where $k_j \in \{0, 1, 2, \dots\}$
- Define the likelihood

$$L(\lambda; \vec{k}) := \text{Prob}(\vec{X} = \vec{k}) = \prod_{j=1}^n p_{X_j}(k_j) = \prod_{j=1}^n e^{-\lambda} \frac{\lambda^{k_j}}{k_j!}$$

- Easier to maximize the log likelihood

$$\ln L(\lambda; \vec{k}) = \sum_{j=1}^n [-\lambda + k_j \ln \lambda - \ln(k_j!)] = -n\lambda + K \ln \lambda - \sum_{j=1}^n \ln(k_j!)$$

where, once again, $K := \sum_{j=1}^n k_j$.

Bruce M.
Boghossian

Motivation
Bernoulli trials

The Poisson
distribution

Uniform
distribution

Standard
normal
distribution

Summary

■ Log likelihood

$$\ln L(\lambda; \vec{k}) = -n\lambda + K \ln \lambda - \sum_{j=1}^n \ln(k_j!)$$

■ Log likelihood maximized for

$$0 = \frac{d}{d\lambda} \ln L(\lambda; \vec{k}) = -n + \frac{K}{\lambda}$$

■ Result is $\lambda = \lambda_e := \frac{K}{n}$, or

$$\lambda_e = \frac{1}{n} \sum_{j=1}^n k_j$$

Bruce M.
Boghossian

Motivation

Bernoulli trials

The Poisson
distribution

Uniform
distribution

Standard
normal
distribution

Summary

- Maximum likelihood estimator for the Poisson distribution is

$$\hat{\lambda}(\vec{k}) = \frac{1}{n} \sum_{j=1}^n k_j$$

- For the 50 points shown earlier, $\hat{\lambda}(\vec{k}) = 3.78$.
- Actual value of λ used to sample the points was 4.

3, 4, 3, 5, 3, 4, 5, 0, 5, 5, 5, 4, 2, 2, 3, 2, 2, 5, 5, 5, 5, 6, 5, 2, 5,
2, 2, 0, 6, 3, 6, 7, 2, 4, 5, 1, 3, 3, 5, 1, 7, 2, 0, 8, 4, 5, 8, 2, 3, 5

Bruce M.
Boghosian

Motivation

Bernoulli trials

The Poisson
distribution

Uniform
distribution

Standard
normal
distribution

Summary

- So far, these results are not terribly surprising.
- The mean of the Bernoulli trials is p .
- The mean of the Poisson distribution is λ .
- Both MLE analyses estimate the parameter to be the sample mean.

Bruce M.
Boghossian

Motivation
Bernoulli trials

The Poisson
distribution


Uniform
distribution

Standard
normal
distribution

Summary

- Suppose $X \in \mathbb{R}$ has the continuous *probability density function*,

$$f_X(x) = \begin{cases} \frac{1}{b-a} & \text{if } X \in [a, b] \\ 0 & \text{otherwise} \end{cases}$$



- Normalization: $\int_{\mathbb{R}} dx f_X(x) = \int_a^b dx \frac{1}{b-a} = \frac{b-a}{b-a} = 1$
- Mean: $\int_{\mathbb{R}} dx f_X(x)x = \int_a^b dx \frac{x}{b-a} = \frac{b+a}{2}$
- Variance: $\int_{\mathbb{R}} dx f_X(x)x^2 - \left(\frac{b+a}{2}\right)^2 = \int_a^b dx \frac{x^2}{b-a} - \left(\frac{b+a}{2}\right)^2 = \frac{(b-a)^2}{12}$
- This is a two-parameter distribution, with parameters a and b .
- Given experimental results $\vec{x} = \{x_1, x_2, \dots, x_n\}$, can we estimate a and b ?

Defining the likelihood for n samples

Bruce M.
Boghossian

Motivation
Bernoulli trials

The Poisson
distribution


Uniform
distribution

Standard
normal
distribution

Summary

- Suppose $X \in \mathbb{R}$ has the continuous *probability density function*,

$$f_X(x) = \begin{cases} \frac{1}{b-a} & \text{if } X \in [a, b] \\ 0 & \text{otherwise} \end{cases}$$



- Likelihood is

$$L(a, b; \vec{x}) = \prod_{j=1}^n f_X(x_j) = \begin{cases} \left(\frac{1}{b-a}\right)^n & \text{if } x_j \in [a, b] \text{ for all } j = 1, \dots, n \\ 0 & \text{if } x_j \notin [a, b] \text{ for any } j = 1, \dots, n \end{cases}$$

- Choose $a \leq \min_j x_j$ and $b \geq \max_j x_j$ so result is $\left(\frac{1}{b-a}\right)^n$.
- Maximize result by choosing $a_e = \min_j x_j$ and $b_e = \max_j x_j$.

Bruce M.
Boghosian

Motivation

Bernoulli trials

The Poisson
distribution

Uniform
distribution

Standard
normal
distribution

Summary

- Maximize $L(a, b; \vec{x})$ by choosing $a_e = \min_j x_j$ and $b_e = \max_j x_j$.
- Maximum likelihood estimators for a and b are

$$\hat{a}(\vec{x}) = \min_j x_j$$

$$\hat{b}(\vec{x}) = \max_j x_j$$

Bruce M.
Boghosian

Motivation

Bernoulli trials

The Poisson
distribution

Uniform
distribution

Standard
normal
distribution

Summary

- Suppose $X \in \mathbb{R}$ has the continuous *probability density function*,

$$f_X(x) = \frac{1}{\sqrt{2\pi v}} \exp \left[-\frac{(x - \mu)^2}{2v} \right],$$

which we recognize to be normalized, with mean μ and variance v .

- This is a two-parameter distribution, with parameters μ and v .
- Given experimental results $\vec{x} = \{x_1, x_2, \dots, x_n\}$, can we estimate μ and v ?

■ Likelihood is

$$L(\mu, \nu; \vec{x}) = \prod_{j=1}^n f_X(x_j) = \prod_{j=1}^n \frac{1}{\sqrt{2\pi\nu}} \exp \left[-\frac{(x_j - \mu)^2}{2\nu} \right]$$

■ Log likelihood is

$$\begin{aligned} \ln L(\mu, \nu; \vec{x}) &= \sum_{j=1}^n \left[-\frac{1}{2} \ln(2\pi\nu) - \frac{(x_j - \mu)^2}{2\nu} \right] \\ &= -\frac{n}{2} \ln(2\pi\nu) - \frac{1}{2\nu} \sum_{j=1}^n (x_j - \mu)^2 \end{aligned}$$

■ We must find maximum with respect to both μ and ν .

Maximum likelihood estimation of μ and ν

Bruce M.
Boghossian

Motivation

Bernoulli trials

The Poisson
distribution

Uniform
distribution

Standard
normal
distribution

Summary

- Log likelihood is $\ln L(\mu, \nu; \vec{x}) = -\frac{n}{2} \ln(2\pi\nu) - \frac{1}{2\nu} \sum_{j=1}^n (x_j - \mu)^2$
- Set partial derivatives to zero

$$0 = \frac{\partial}{\partial \mu} \ln L(\mu, \nu; \vec{x}) = \frac{1}{\nu} \sum_{j=1}^n (x_j - \mu) = \frac{1}{\nu} \left(\sum_{j=1}^n x_j - n\mu \right)$$

$$0 = \frac{\partial}{\partial \nu} \ln L(\mu, \nu; \vec{x}) = -\frac{n}{2\nu} + \frac{1}{2\nu^2} \sum_{j=1}^n (x_j - \mu)^2$$

- Solving for location of maximum (μ_e, ν_e) yields

$$\mu_e = \frac{1}{n} \sum_{j=1}^n x_j \quad \text{and} \quad \nu_e = \frac{1}{n} \sum_{j=1}^n \left[x_j - \left(\frac{1}{n} \sum_{k=1}^n x_k \right) \right]^2$$

Bruce M.
Boghosian

- Maximum likelihood estimators for a and b are

$$\hat{\mu}(\vec{x}) = \frac{1}{n} \sum_{j=1}^n x_j$$

$$\hat{\nu}(\vec{x}) = \frac{1}{n} \sum_{j=1}^n \left[x_j - \left(\frac{1}{n} \sum_{k=1}^n x_k \right) \right]^2$$

Motivation

Bernoulli trials

The Poisson
distribution

Uniform
distribution

Standard
normal
distribution

Summary

Bruce M.
Boghosian

Motivation

Bernoulli trials

The Poisson
distribution

Uniform
distribution

Standard
normal
distribution

Summary

- We have learned the method of *maximum likelihood estimation*.
- Allows estimation of parameters if form of the distribution is known a priori.
- We have seen examples with discrete and continuous probability functions.
- We have seen examples with one and two parameters.
- We have seen examples where using the log likelihood is useful and not useful.
- In all cases, we have calculated *estimator functions*.