

Bruce M.
Boghosian

Least squares
with linear
algebra

Motivation for
the linear
model

Regression
curves

The Linear
Model

Properties of
estimators

Estimating $\hat{\sigma}^2$

Summary

Regression

The Linear Model

Bruce M. Boghosian



Tufts
UNIVERSITY

School of Arts
and Sciences

Department of Mathematics

Tufts University

Bruce M.
Boghossian

Least squares
with linear
algebra

Motivation for
the linear
model

Regression
curves

The Linear
Model

Properties of
estimators

Estimating $\hat{\sigma}^2$

Summary

- 1 Least squares with linear algebra
- 2 Motivation for the linear model
- 3 Regression curves
- 4 The Linear Model
- 5 Properties of estimators
- 6 Estimating $\hat{\sigma}^2$
- 7 Summary

Bruce M.
Boghossian

Least squares
with linear
algebra

Motivation for
the linear
model

Regression
curves

The Linear
Model

Properties of
estimators

Estimating $\hat{\sigma}^2$

Summary

- Some of you have learned an alternative linear algebraic approach to the method of least squares.
- Write the equations as follows

$$a + bx_1 = y_1$$

$$a + bx_2 = y_2$$

$$\vdots$$

$$a + bx_n = y_n$$

- Cast in rectangular matrix form $Ax = y$, where

$$\underbrace{\begin{pmatrix} 1 & x_1 \\ 1 & x_2 \\ \vdots & \vdots \\ 1 & x_n \end{pmatrix}}_A \underbrace{\begin{pmatrix} a \\ b \end{pmatrix}}_x = \underbrace{\begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{pmatrix}}_y$$

- n equations for two unknowns – generally no solution

Bruce M.
Boghossian

Least squares
with linear
algebra

Motivation for
the linear
model

Regression
curves

The Linear
Model

Properties of
estimators

Estimating $\hat{\sigma}^2$

Summary

- We can not generally solve $Ax = y$.
- So instead we minimize

$$\begin{aligned}\|Ax - y\|^2 &= (Ax - y)^T (Ax - y) \\ &= (x^T A^T - y^T)(Ax - y) \\ &= x^T A^T Ax - 2x^T A^T y + y^T y\end{aligned}$$

- Minimize with respect to components of x to obtain

$$2A^T Ax - 2A^T y = 0.$$

- Result is

$$(A^T A)x = A^T y$$

- Note that $A^T A$ is a 2×2 square matrix

Bruce M.
Boghossian

Least squares
with linear
algebra

Motivation for
the linear
model

Regression
curves

The Linear
Model

Properties of
estimators

Estimating $\hat{\sigma}^2$

Summary

- Write out the result $(A^T A) x = A^T y$

$$\begin{pmatrix} 1 & 1 & \cdots & 1 \\ x_1 & x_2 & \cdots & x_n \end{pmatrix} \begin{pmatrix} 1 & x_1 \\ 1 & x_2 \\ \vdots & \vdots \\ 1 & x_n \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 1 & 1 & \cdots & 1 \\ x_1 & x_2 & \cdots & x_n \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{pmatrix}$$

- But the above is equivalent to

$$\begin{pmatrix} n & \sum_i^n x_i \\ \sum_i^n x_i & \sum_i^n x_i^2 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} \sum_i^n y_i \\ \sum_i^n x_i y_i \end{pmatrix}$$

which is exactly what we derived last time.

Bruce M.
Boghossian

Least squares
with linear
algebra

Motivation for
the linear
model

Regression
curves

The Linear
Model

Properties of
estimators

Estimating $\hat{\sigma}^2$

Summary

- We studied least-squares fits from a geometric perspective.
- Our points (x_i, y_i) were given.
- The coordinates x_i and y_i were numbers.
- We found polynomial that minimizes least-squares error L .

Bruce M.
Boghossian

Least squares
with linear
algebra

Motivation for
the linear
model

Regression
curves

The Linear
Model

Properties of
estimators

Estimating $\hat{\sigma}^2$

Summary

- More likely, horizontal and/or vertical coordinates are r.v.s.
- Example:
 - We might consider (x_i, Y_i) , so Y_i is a r.v.
 - For each given x_i there is a distribution for Y_i .
- Example:
 - We might consider (X_i, Y_i) , so both are r.v.s.
 - For each i , there is a bivariate distribution of X_i and Y_i .

Bruce M.
Boghossian

Least squares
with linear
algebra

Motivation for
the linear
model

Regression
curves

The Linear
Model

Properties of
estimators

Estimating $\hat{\sigma}^2$

Summary

- **Def.:** Let $f_{Y|x}(y)$ be the pdf of r.v. Y for a given value x , and let $E(Y|x)$ denote the expected value associated with $f_{Y|x}(y)$. The function

$$y = E(Y|x)$$

is called the *regression curve* of Y on x .

Bruce M.
Boghossian

Least squares
with linear
algebra

Motivation for
the linear
model

Regression
curves

The Linear
Model

Properties of
estimators

Estimating $\hat{\sigma}^2$

Summary

- Suppose that, for $0 \leq x \leq 1$ and $0 \leq y \leq 1$,

$$f_{Y|x}(y) = 2 \left(\frac{1 - x - y + 3xy}{1 + x} \right).$$

- Note that this is normalized,

$$\int_0^1 dy f_{Y|x}(y) = 2 \int_0^1 dy \left(\frac{1 - x - y + 3xy}{1 + x} \right) = 1.$$

- Also note that

$$E(Y|x) = \int_0^1 dy f_{Y|x}(y)y = 2 \int_0^1 dy \left(\frac{1 - x - y + 3xy}{1 + x} \right) y = \frac{x + 1/3}{x + 1}.$$

Bruce M.
Boghossian

Least squares
with linear
algebra

Motivation for
the linear
model

Regression
curves

The Linear
Model

Properties of
estimators

Estimating $\hat{\sigma}^2$

Summary

- For $0 \leq x \leq 1$ and $0 \leq y \leq 1$, we have

$$f_{Y|x}(y) = 2 \left(\frac{1 - x - y + 3xy}{1 + x} \right) \quad \text{and} \quad E(Y|x) = \frac{x + 1/3}{x + 1}.$$

- This means, for example, that when $x = 1/2$, we have

$$f_{Y|1/2}(y) = \frac{2}{3}(1 + y) \quad \text{and} \quad E(Y|1/2) = \frac{5}{9}.$$

- One can check that, as expected,

$$E(Y|1/2) = \int_0^1 dy f_{Y|1/2}(y)y = \int_0^1 dy (2y)y = \frac{2}{3}.$$

Bruce M.
Boghossian

Least squares
with linear
algebra

Motivation for
the linear
model

Regression
curves

The Linear
Model

Properties of
estimators

Estimating $\hat{\sigma}^2$

Summary

■ Special case:

- $f_{Y|x}(y)$ is a normal pdf for all x .
- The standard deviation of $f_{Y|x}(y)$, σ , is the same for all x .
- The means of the conditional Y distributions are collinear,

$$y = E(Y|x) = \beta_0 + \beta_1 x.$$

- All conditional distributions represent independent r.v.s.
- β_0 , β_1 , and σ^2 are unknown and should be estimated.

Bruce M.
Boghossian

Least squares
with linear
algebra

Motivation for
the linear
model

Regression
curves

The Linear
Model

Properties of
estimators

Estimating $\hat{\sigma}^2$

Summary

- We have $f_{Y|x}(y) = \frac{1}{\sqrt{2\pi} \sigma} \exp \left[-\frac{1}{2} \left(\frac{y - \beta_0 - \beta_1 x}{\sigma} \right)^2 \right]$
- Likelihood is then

$$\begin{aligned} L &= \prod_{i=1}^n f_{Y_i|x_i}(y_i) = \prod_{i=1}^n \frac{1}{\sqrt{2\pi} \sigma} \exp \left[-\frac{1}{2} \left(\frac{y_i - \beta_0 - \beta_1 x_i}{\sigma} \right)^2 \right] \\ &= \left(\frac{1}{\sqrt{2\pi} \sigma} \right)^n \exp \left[-\frac{1}{2} \sum_{i=1}^n \left(\frac{y_i - \beta_0 - \beta_1 x_i}{\sigma} \right)^2 \right] \end{aligned}$$

- Log likelihood is then

$$-2 \ln L = n \ln(2\pi) + n \ln(\sigma^2) + \frac{1}{\sigma^2} \sum_{i=1}^n (y_i - \beta_0 - \beta_1 x_i)^2$$

Bruce M.
Boghossian

Least squares
with linear
algebra

Motivation for
the linear
model

Regression
curves

The Linear
Model

Properties of
estimators

Estimating $\hat{\sigma}^2$

Summary

- Find maximum with respect to β_0 , β_1 and σ^2

$$0 = \frac{\partial(-2 \ln L)}{\partial \beta_0} = \frac{2}{\sigma^2} \sum_{i=1}^n (y_i - \beta_0 - \beta_1 x_i) (-1)$$

$$0 = \frac{\partial(-2 \ln L)}{\partial \beta_1} = \frac{2}{\sigma^2} \sum_{i=1}^n (y_i - \beta_0 - \beta_1 x_i) (-x_i)$$

$$0 = \frac{\partial(-2 \ln L)}{\partial \sigma^2} = \frac{n}{\sigma^2} - \frac{2}{\sigma^4} \sum_{i=1}^n (y_i - \beta_0 - \beta_1 x_i)^2$$

- Result for estimators is (where $\hat{Y}_i := \hat{\beta}_0 + \hat{\beta}_1 x_i$)

$$\hat{\beta}_1 = \frac{n \sum_i^n x_i Y_i - (\sum_i^n x_i) (\sum_i^n Y_i)}{n (\sum_i^n x_i^2) - (\sum_i^n x_i)^2}$$

$$\hat{\beta}_0 = \bar{Y} - \hat{\beta}_1 \bar{x}$$

$$\hat{\sigma}^2 = \frac{1}{n} \sum_i^n (Y_i - \hat{Y}_i)^2$$

Bruce M.
Boghossian

Least squares
with linear
algebra

Motivation for
the linear
model

Regression
curves

The Linear
Model

Properties of
estimators

Estimating $\hat{\sigma}^2$

Summary

- Let Y_1, \dots, Y_n be any set of independent random variables with means μ_1, \dots, μ_n and variances $\sigma_1^2, \dots, \sigma_n^2$, respectively. Let a_1, \dots, a_n be any set of constants. Then $Y = a_1 Y_1 + \dots + a_n Y_n$ is normally distributed with mean $\mu = \sum_i^n a_i \mu_i$ and variance $\sigma^2 = \sum_i^n a_i^2 \sigma_i^2$.
- Proof is straightforward using moment-generating functions.

Bruce M.
Boghossian

Least squares
with linear
algebra

Motivation for
the linear
model

Regression
curves

The Linear
Model

Properties of
estimators

Estimating $\hat{\sigma}^2$

Summary

- **Thm.:** $\hat{\beta}_0$ and $\hat{\beta}_1$ are both normally distributed.
- **Thm.:** $\hat{\beta}_0$ and $\hat{\beta}_1$ are both unbiased,

$$E(\hat{\beta}_0) = \beta_0$$

$$E(\hat{\beta}_1) = \beta_1$$

- **Thm.:** The variances of the estimators are

$$\text{Var}(\hat{\beta}_0) = \frac{\sigma^2 \sum_i^n x_i^2}{n \sum_i^n (x_i - \bar{x})^2}$$

$$\text{Var}(\hat{\beta}_1) = \frac{\sigma^2}{n \sum_i^n (x_i - \bar{x})^2}$$

Bruce M.
Boghossian

Least squares
with linear
algebra

Motivation for
the linear
model

Regression
curves

The Linear
Model

Properties of
estimators

Estimating $\hat{\sigma}^2$

Summary

- **Pf. (for $\hat{\beta}_1$):** Note that

$$\begin{aligned}\hat{\beta}_1 &= \frac{n \sum_i^n x_i Y_i - (\sum_i^n x_i) (\sum_i^n Y_i)}{n (\sum_i^n x_i^2) - (\sum_i^n x_i)^2} = \frac{\sum_i^n x_i Y_i - \left(\frac{1}{n} \sum_i^n x_i\right) (\sum_i^n Y_i)}{\left(\sum_i^n x_i^2\right) - n \left(\frac{1}{n} \sum_i^n x_i\right) \left(\frac{1}{n} \sum_i^n Y_i\right)} \\ &= \frac{\sum_i^n x_i Y_i - \bar{x} (\sum_i^n Y_i)}{(\sum_i^n x_i^2) - n \bar{x}^2} = \frac{\sum_i^n (x_i - \bar{x}) Y_i}{(\sum_i^n x_i^2) - n \bar{x}^2}\end{aligned}$$

- This is a linear combination of normally distributed r.v.s, and thus normally distributed.

Bruce M.
Boghossian

Least squares
with linear
algebra

Motivation for
the linear
model

Regression
curves

The Linear
Model

Properties of
estimators

Estimating $\hat{\sigma}^2$

Summary

- **Pf. (for $\hat{\beta}_1$):** Using the same form for $\hat{\beta}_1$ used above,

$$\begin{aligned} E(\hat{\beta}_1) &= E\left(\frac{\sum_i^n (x_i - \bar{x}) Y_i}{(\sum_i^n x_i^2) - n\bar{x}^2}\right) \\ &= \frac{\sum_i^n (x_i - \bar{x}) E(Y_i)}{(\sum_i^n x_i^2) - n\bar{x}^2} \\ &= \frac{\sum_i^n (x_i - \bar{x}) (\beta_0 + \beta_1 x_i)}{(\sum_i^n x_i^2) - n\bar{x}^2} \\ &= \beta_1. \end{aligned}$$

Bruce M.
Boghossian

Least squares
with linear
algebra

Motivation for
the linear
model

Regression
curves

The Linear
Model

Properties of
estimators

Estimating $\hat{\sigma}^2$

Summary

- **Pf. (for $\hat{\beta}_1$):** Using the same form for $\hat{\beta}_1$ used above,

$$\begin{aligned}\text{Var}(\hat{\beta}_1) &= \text{Var}\left(\frac{\sum_i^n (x_i - \bar{x}) Y_i}{\left(\sum_i^n x_i^2\right) - n\bar{x}^2}\right) \\ &= \sum_i^n \left(\frac{(x_i - \bar{x})}{\left(\sum_j^n x_j^2\right) - n\bar{x}^2}\right)^2 \text{Var}(Y_i) \\ &= \frac{\sigma^2}{n \sum_i^n (x_i - \bar{x})^2}\end{aligned}$$

Bruce M.
Boghossian

Least squares
with linear
algebra

Motivation for
the linear
model

Regression
curves

The Linear
Model

Properties of
estimators

Estimating $\hat{\sigma}^2$

Summary

- **Thm.:** Let $(x_1, Y_1), \dots, (x_n, Y_n)$ satisfy the assumptions of the Linear Model. Then
 - $\hat{\beta}_1$, \bar{Y} , and $\hat{\sigma}^2$ are mutually independent.
 - $\frac{n\hat{\sigma}^2}{\sigma^2}$ has a χ^2 distribution with $n - 2$ degrees of freedom.
- **Corr.:** Let $\hat{\sigma}^2$ be the MLE for σ^2 in a simple linear model. Then
 - $\left(\frac{n}{n-2}\right) \hat{\sigma}^2$ is an unbiased estimator for σ^2 .
 - The random variables \hat{Y}^2 and $\hat{\sigma}^2$ are independent.

Bruce M.
Boghossian

Least squares
with linear
algebra

Motivation for
the linear
model

Regression
curves

The Linear
Model

Properties of
estimators

Estimating $\hat{\sigma}^2$

Summary

■ Calculating $\sum_i^n (y_i - \hat{y}_i)^2 = \sum_i^n (y_i - \hat{\beta}_0 - \hat{\beta}_1 x_i)^2$

$$\begin{aligned} &= \sum_i^n (y_i - \bar{y}_i)^2 - \hat{\beta}_1^2 \sum_i^n (x_i - \bar{x}_i)^2 \\ &= \sum_i^n y_i^2 - \frac{1}{n} \sum_i^n y_i^2 - \frac{[\sum_i^n x_i y_i - \frac{1}{n} (\sum_i^n x_i) (\sum_i^n y_i)]^2}{(\sum_i^n x_i^2) - \frac{1}{n} (\sum_i^n x_i)^2} \\ &= \sum_i^n y_i^2 - \hat{\beta}_0 \sum_i^n y_i - \hat{\beta}_1 \sum_i^n x_i y_i. \end{aligned}$$

Bruce M.
Boghossian

Least squares
with linear
algebra

Motivation for
the linear
model

Regression
curves

The Linear
Model

Properties of
estimators

Estimating $\hat{\sigma}^2$

Summary

- Let $(x_1, Y_1), \dots, (x_n, Y_n)$ satisfy the assumptions of the Linear Model, and let

$$S^2 = \frac{1}{n-2} \sum_i^n \left(y_i - \hat{\beta}_0 - \hat{\beta}_1 x_i \right)^2$$

- Then the following has a Student T distribution with $n-2$ df.

$$T_{n-2} = \frac{\hat{\beta}_1 - \beta_1}{S / \sqrt{\sum_i^n (x_i - \bar{x})^2}}$$

Bruce M.
Boghossian

Least squares
with linear
algebra

Motivation for
the linear
model

Regression
curves

The Linear
Model

Properties of
estimators

Estimating $\hat{\sigma}^2$

Summary

- We reexamined the application of linear algebra to least-squares fitting.
- We have examined the least-squares method with random variables for the dependent variable.
- We have defined and motivated the Linear Model.
- We have defined regression curves.
- We worked out ML estimators for the three parameters of the linear model.
- We showed that $\hat{\beta}_1$ is Student T distributed with $n - 2$ degrees of freedom.