

Bruce M.
Boghosian

Motivation

Bayes'
Theorem

Example

Summary

Bayesian estimation

Motivation and Bayes' Theorem

Bruce M. Boghosian



Tufts
UNIVERSITY

School of Arts
and Sciences

Department of Mathematics

Tufts University

Bruce M.
Boghosian

Motivation

Bayes'
Theorem

Example

Summary

1 Motivation

2 Bayes' Theorem

3 Example

4 Summary

Bruce M.
Boghosian

Motivation

Bayes'
Theorem

Example

Summary

- Given n possible events $\{A_j\}_{j=1}^n$, each with some corresponding probability
- These events *partition* the set of all such possibilities A :

$$A = \bigcup_{j=1}^n A_j = A_1 \cup A_2 \cup \dots \cup A_n \quad (\text{Every possible outcome is accounted for})$$

$$A_j \cap A_k = \emptyset \text{ if } j \neq k \quad (\text{Outcomes are mutually exclusive})$$
- We have initial guess for *prior distribution*, $\{P(A_j)\}_{j=1}^n$.
- Let event B be a new observation related to above events.
- We wish to update $\{P(A_j)\}_{j=1}^n$, given observation B .

Bruce M.
Boghosian

Motivation

Bayes'
Theorem

Example

Summary

- We wish to update $\{P(A_j)\}_{j=1}^n$, given observation B .
- We wish to find *posterior distribution*, $\{P(A_j | B)\}_{j=1}^n$.
- If there is a subsequent guess, process could be repeated.
- In this way, the algorithm “learns” from new information.
- According to “Bayesians”, you should not suppose that a new piece of information will give you all the answers you are seeking.
- Rather, new information should update prior beliefs.

- Recall from the definition of conditional probability

$$P(A_j, B) = P(A_j | B)P(B) = P(B | A_j)P(A_j).$$

$$\therefore P(A_j | B) = \frac{P(B | A_j)P(A_j)}{P(B)}$$

- Then recall that, since the $\{A_j\}_{j=1}^n$ partition A ,

$$P(B) = \sum_{k=1}^n P(B | A_k)P(A_k)$$

- This gives us *Bayes' Theorem*,

$$P(A_j | B) = \frac{P(B | A_j)P(A_j)}{\sum_{k=1}^n P(B | A_k)P(A_k)}.$$

■ *Bayes' Theorem*

$$P(A_j | B) = \frac{P(B | A_j)P(A_j)}{\sum_{k=1}^n P(B | A_k)P(A_k)}.$$

- Note numerator is one term of denominator, and all terms are positive.
- Result will always be in $[0, 1]$, as expected.
- Gives the desired $P(A_j | B)$ in terms of things that we might actually know, or be able to measure, or at least be able to estimate.
- The results of thinking in this way can defy intuition.
- An enormous body of scientific conclusions are gleaned in exactly this way.
- Key question: How do we use this theorem in the context of a real problem?

Bruce M.
Boghossian

Motivation

Bayes'
Theorem

Example

Summary

- Example due to D. Kahneman and A. Tversky, as presented in “Thinking Fast and Slow” by D. Kahneman, FSG Publishers, New York (2011) pp. 6-7.
- **Given information:** “An individual has been described by a neighbor as follows: ‘Steve is very shy and withdrawn, invariably helpful but with very little interest in people or the world of reality. A meek and tidy soul, he has a need for order and structure, and a passion for detail.’ ”
- **Question:** “Is Steve more likely to be librarian or farmer?”
- According to Kahneman, the vast majority of people who are asked this question say that Steve is more likely to be a librarian, almost surely due to the stereotypical view of librarians as having these personality traits.
- Does this make sense? Are we missing any information?

Example: Librarians and Farmers (continued)

Bruce M.
Boghosian

Motivation

Bayes'
Theorem

Example

Summary

- Does this make sense? Are we missing information?
- There are $20\times$ as many farmers as librarians in the US.
- Sample of 105 people – 100 farmers and 5 librarians.
- Say 40% librarians have this trait & only 10% farmers.
- Hence 2 librarians & 10 farmers in our sample have trait.
- Given that a randomly selected person has this trait, there is only a $\frac{2}{12} = \frac{1}{6}$ chance that they are a librarian, and a $\frac{10}{12} = \frac{5}{6}$ chance that they are a farmer.
- Kahneman: “Because there are so many farmers, it is almost certain that more ‘meek and tidy’ will be found on tractors than at library information desks.”
- Clearly we *were* missing information. The ratio of farmers to librarians *matters* to the question as posed! Most people don’t even consider this possibility!

Bruce M.
Boghosian

Motivation

Bayes'
Theorem

Example

Summary

- Decide on your events and give them variable names:
 - B = Steve is a “meek and tidy soul”
 - A_1 = Steve is a librarian
 - A_2 = Steve is a farmer
- We know that $P(A_1) = \frac{1}{21}$ and $P(A_2) = \frac{20}{21}$.
- We estimate $P(B | A_1) = \frac{4}{10}$, and $P(B | A_2) = \frac{1}{10}$.
- Then the probability that Steve is a librarian given that he is a “meek and tidy soul” is

$$P(A_1 | B) = \frac{P(B | A_1)P(A_1)}{P(B | A_1)P(A_1) + P(B | A_2)P(A_2)} = \frac{\frac{4}{10} \cdot \frac{1}{21}}{\frac{4}{10} \cdot \frac{1}{21} + \frac{1}{10} \cdot \frac{20}{21}} = \frac{4}{24} = \frac{1}{6},$$

in agreement with our earlier calculation.

Bruce M.
Boghosian

Motivation

Bayes'
Theorem

Example

Summary

- We have reviewed Bayes' Theorem
- We have given an example of Bayesian reasoning
- We have motivated Bayesian parameter estimation