

Ion Channel Aperture Arrays as Categorical Filters: The Five Names Theorem, Catalytic Power, and Saturation Dynamics

Kundai Farai Sachikonye

Chair of Experimental Bioinformatics, Technical University of Munich

AIme Registry for Artificial Intelligence

Maximus-von-Imhof-Forum 3, 85354 Freising, Germany

kundai.sachikonye@wzw.tum.de

May 2026

Abstract. We derive ion channel aperture arrays as mathematical necessities of bounded partition-based computation in aqueous lipid-bilayer substrates. The argument proceeds from the Bounded Phase Space axiom through five theorems to a unique aperture implementation. A partition boundary separating two regions of phase space with finite Liouville measure presents any crossing particle with a Born desolvation energy barrier of approximately $40 k_B T$ in a hydrophobic slab of dielectric constant $\epsilon_m \approx 2$; spontaneous crossing is therefore exponentially suppressed ($\sim e^{-40} \approx 10^{-17}$) and computation requires specialised permeation structures. We prove the Five Names Theorem: any selective permeation structure at the partition boundary of a bounded aqueous dynamical system is simultaneously a geometric aperture with transmission function $\kappa \in [0, 1]$, a mediator of virtual sub-states in the backward trajectory completion algorithm, an information catalyst for the crossing transformation, a Maxwell demon for S-entropy coordinate measurement, and a physically instantiated subtask node in the cascade-fabric routing graph. These five characterisations are shown to be pairwise equivalent. In an aqueous lipid bilayer at physiological temperature, the unique physical realisation satisfying all five is an ion-channel protein: a water-filled protein pore that collapses the Born barrier, provides a selectivity filter (S-entropy matching gate), and is structurally invariant across individual ion transit events. The catalytic power $\kappa(\mathcal{X}; \mathbf{t})$ of a channel with target S-entropy coordinates $\mathbf{t}$ for an input system with coordinates $\mathbf{S}(\mathcal{X})$ equals the spectral overlap integral between their Koopman spectra. For an array of n channels in parallel, the array catalytic power satisfies $\kappa_{\text{arr}} = 1 - \prod_{i=1}^n (1 - \kappa_i)$ (the Saturation Theorem), which converges to unity if and only if $\sum_{i=1}^n \kappa_i = \infty$. The action potential is identified as a virtual sub-state: a finite-lifetime Koopman excitation that enables the membrane to transition between conductance eigenstates without permanently occupying either. Validation against patch-clamp conductance data confirms predicted catalytic powers to within experimental uncertainty.

Keywords: ion channel, Born energy barrier, Maxwell demon, catalytic power, Five Names Theorem, virtual sub-state, action potential, cascade saturation, bounded phase space, aperture array

1. Introduction

The lipid bilayer membrane is, as established in the first companion paper [Sachikonye, 2026a], the necessary and sufficient partition boundary for computation in bounded aqueous systems. A partition boundary that is completely impermeable separates two phase space regions permanently: no information can cross, and no computation is possible. A boundary that is fully permeable can-

not sustain a partition: the two regions merge into one. Useful computation requires a boundary that is *selectively* permeable — one that passes some trajectories and blocks others according to a computable criterion.

The question of how selective permeability arises in a lipid bilayer has been answered in biology by five decades of electrophysiology: ion channels are specialised protein structures that create water-filled pores spanning the membrane, equipped with

selectivity filters and voltage- or ligand-controlled gates [Hille, 2001, Neher and Sakmann, 1976, Doyle et al., 1998]. The standard account is evolutionary: channels evolved over billions of years to satisfy physiological requirements. This account is correct but incomplete: it does not explain *why* the resulting structures have the specific mathematical properties they do, nor why those properties are precisely what is needed for the partition computation described in this paper series.

We give a first-principles derivation. Starting from the Born energy barrier for ion crossing of a hydrophobic slab [Parsegian, 1969], we show that the bilayer is exponentially impermeable to free ions (§3). We then derive, from the operational requirements of partition-based computation, the set of properties that any permeation structure must satisfy (§4). The Five Names Theorem shows that these properties are not five independent requirements but five descriptions of a single mathematical object. In an aqueous bilayer, this object is uniquely realised by an ion channel (§5). The catalytic power κ — the probability that an input system with given S-entropy coordinates passes through a given aperture — is derived from the Koopman spectral overlap (§6). The action potential is identified as a virtual sub-state mediating transitions between conductance eigenstates (§7). The Saturation Theorem for parallel arrays is proved in §8.

2. Bounded Phase Space Framework

We state the foundational axiom and collect the results from companion papers [Sachikonye, 2026a,b] needed below.

Axiom 1 (Bounded Phase Space). Every persistent physical system $\mathcal{X}$ occupies a region $\Omega \subset \mathbb{R}^{2d}$ with finite Liouville measure $\mu(\Omega) = \int_{\Omega} d^d q d^d p / h^d < \infty$, admitting a hierarchy of refining measurable partitions $P_0 \supset P_1 \supset \dots$.

From [Sachikonye, 2026a]: The partition boundary $\partial\Omega$ separating interior Ω_{in} from exterior Ω_{out} is a lipid bilayer membrane of thickness $d = 4.0$ nm and bending modulus $\kappa_b = 20 k_B T$.

From [Sachikonye, 2026b]: Every bounded oscillatory system has a discrete Koopman spectrum $\Sigma(\mathcal{X}) = \{(\omega_k, A_k, \phi_k)\}$, and its S-entropy coordinates $\mathbf{S}(\mathcal{X}) = (S_k, S_t, S_e) \in [0, 1]^3$ are the unique minimal sufficient statistics for categorical identification.

3. Partition Boundary Opacity

3.1. The Born Energy Barrier

A monovalent ion of radius r and charge Ze crossing from water ($\varepsilon_w \approx 80$) into a hydrophobic slab ($\varepsilon_m \approx 2$) of thickness d experiences a desolvation energy barrier. The Born model [Born, 1920] gives, for an infinite slab, the midpoint energy:

$$\Delta E_{\text{Born}} = \frac{(Ze)^2}{8\pi\varepsilon_0 r} \left(\frac{1}{\varepsilon_m} - \frac{1}{\varepsilon_w} \right). \quad (1)$$

For a slab of finite thickness d , the image-charge series gives a correction; the full result for $d = 4$ nm, $r = 0.095$ nm (Na^+) was computed by Parsegian [1969]:

$$\Delta E_{\text{Born}}^{(\text{slab})} \approx 40 k_B T \quad (T = 310 \text{ K}). \quad (2)$$

Theorem 3.1 (Partition Boundary Opacity). *The spontaneous crossing rate of monovalent ions across a lipid bilayer membrane in the absence of specialised permeation structures is:*

$$k_{\text{cross}} = \nu_0 \exp(-\Delta E_{\text{Born}}/k_B T) \approx 10^9 \text{ s}^{-1} \times e^{-40} \approx 4 \times 10^{-9} \text{ s}^{-1} \quad (3)$$

where $\nu_0 \approx k_B T/h \approx 6 \times 10^{12} \text{ s}^{-1}$ is the attempt frequency (thermal rate at the transition state). The corresponding permeability coefficient is $P_{\text{ion}} \approx 10^{-14} \text{ cm s}^{-1}$, seventeen orders of magnitude below the water permeability coefficient $P_{\text{water}} \approx 10^3 \text{ cm s}^{-1}$.

Proof. Equation (3) follows directly from equation (2) and Kramers' transition-state rate theory [Kramers, 1940] with $\nu_0 = k_B T/h$ (the thermal frequency). The permeability coefficient $P = k_{\text{cross}} \times d = 4 \times 10^{-9} \text{ s}^{-1} \times 4 \times 10^{-9} \text{ m} \approx 10^{-17} \text{ m s}^{-1} = 10^{-15} \text{ cm s}^{-1}$, consistent with the experimental lower bound of $P_{\text{Na}^+} \leq 10^{-14} \text{ cm s}^{-1}$ measured for pure bilayers [Hille, 2001, Walter and Gutknecht, 1983]. $\square$

Corollary 3.2 (Computational Stasis). *A partition boundary with no specialised permeation structures supports zero information flux between Ω_{in} and Ω_{out} on any timescale relevant to partition-based computation ($\tau \lesssim 1$ s).*

Proof. At the spontaneous crossing rate (3), the expected number of spontaneous crossings per square micrometre of membrane per second is $k_{\text{cross}} \times \rho_{\text{ion}} \times d \approx 10^{-17} \text{ m s}^{-1} \times 0.15 \times 10^3 \text{ mol L}^{-1} \times N_A \approx 10^{-6} \text{ ion } \mu\text{m}^{-2} \text{ s}^{-1}$. Since each crossing carries at

most $k_B T \ln 2$ bits of information [Landauer, 1961], the information flux is $\lesssim 10^{-6} k_B T \ln 2 \mu\text{m}^{-2} \text{s}^{-1}$, negligible compared to the $\sim 10^{20} \text{ ops s}^{-1} \text{ cm}^{-2}$ capacity of the fluid-phase membrane [Sachikonye, 2026a]. $\square$

Corollary 3.2 establishes the necessity of specialised permeation structures. We now derive the properties such structures must have.

4. The Five Names Theorem

4.1. Five Definitions of an Aperture

We give five definitions, each capturing a different aspect of a selective permeation structure at the partition boundary.

Definition 4.1 (Geometric Aperture). A *geometric aperture* at the partition boundary $\partial\Omega$ is a pair (A, κ) where $A \subseteq \partial\Omega$ is a measurable region (the *pore domain*) and $\kappa : [0, 1]^3 \rightarrow [0, 1]$ is a *transmission function* mapping an input system's S-entropy coordinates $\mathbf{S}(\mathcal{X})$ to the probability that it crosses $\partial\Omega$ through A .

Definition 4.2 (Virtual Sub-State Mediator). A *virtual sub-state mediator* is a structure V at $\partial\Omega$ that enables passage of a trajectory from Ω_{in} to Ω_{out} (or vice versa) via a transient state $|v\rangle$ satisfying:

- (i) $|v\rangle$ has non-zero Koopman amplitude: $A_v > 0$;
- (ii) $|v\rangle$ is not an eigenstate of the partition operator $\hat{P}$ of either Ω_{in} or Ω_{out} : $\hat{P}|v\rangle \neq \lambda|v\rangle$;
- (iii) $|v\rangle$ has finite lifetime: $\tau_v = A_v^{-1}(\text{d}A_v/\text{d}t)^{-1} < \infty$.

Definition 4.3 (Information Catalyst). An *information catalyst* for the crossing transformation $T : \Omega_{\text{in}} \rightarrow \Omega_{\text{out}}$ is an agent C satisfying:

- (i) *Necessity*: T has zero rate without C (Theorem 3.1);
- (ii) *Catalytic invariance*: the structure of C is unchanged by each transit event: $|C\rangle_{\text{before}} = |C\rangle_{\text{after}}$;
- (iii) *Information conservation*: $H(C_{\text{before}}) = H(C_{\text{after}})$, where H is the Shannon entropy of the channel's conformational state distribution.

Definition 4.4 (Maxwell Demon). A *Maxwell demon* at the partition boundary is an agent D that:

- (i) Measures the S-entropy coordinates $\mathbf{S}(\mathcal{X})$ of each approaching particle;

- (ii) Conditionally permits crossing with probability $\kappa(\mathbf{S}(\mathcal{X}))$ based on the measurement outcome;
- (iii) Operates at energy cost $\geq k_B T \ln 2$ per binary decision [Szilárd, 1929, Landauer, 1961].

Definition 4.5 (Cascade Subtask Node). A *cascade subtask node* in the backward trajectory completion algorithm is a node u in the routing graph $G = (V, E)$ of the cascade fabric such that:

- (i) u lies on the boundary between two ternary routing sub-trees;
- (ii) Passage through u is conditional: a trajectory routed to u is forwarded to the sub-tree on the other side with probability κ_u and reflected with probability $1 - \kappa_u$;
- (iii) κ_u is computed from the S-entropy distance between the trajectory's coordinates and the sub-tree's target category.

4.2. Equivalence

Theorem 4.6 (Five Names). *For any structure $\mathcal{A}$ at the partition boundary $\partial\Omega$ of a bounded aqueous dynamical system, the following are equivalent:*

- (G) $\mathcal{A}$ is a geometric aperture (Definition 4.1);
- (V) $\mathcal{A}$ is a virtual sub-state mediator (Definition 4.2);
- (I) $\mathcal{A}$ is an information catalyst (Definition 4.3);
- (M) $\mathcal{A}$ is a Maxwell demon (Definition 4.4);
- (S) $\mathcal{A}$ is a cascade subtask node (Definition 4.5).

Proof. We prove the cycle $\mathbf{G} \Rightarrow \mathbf{V} \Rightarrow \mathbf{I} \Rightarrow \mathbf{M} \Rightarrow \mathbf{S} \Rightarrow \mathbf{G}$.

G $\Rightarrow$ V: A geometric aperture with $0 < \kappa < 1$ has a pore domain $A \subseteq \partial\Omega$. A particle traversing A occupies A transiently during transit, with duration $\tau_v \approx L_{\text{pore}}/v_{\text{th}}$ where L_{pore} is the pore length and $v_{\text{th}} = \sqrt{2k_B T/m}$ is the thermal velocity. During this transit:

- (i) The particle's trajectory has non-zero Koopman amplitude at frequencies corresponding to the transit: $A_v = |F(\omega_{\text{transit}})|^2 > 0$ since $\omega_{\text{transit}} = v_{\text{th}}/L_{\text{pore}} > 0$.
- (ii) The transit state is not an eigenstate of $\hat{P}_{\text{in}}$ or $\hat{P}_{\text{out}}$ since the particle is in neither Ω_{in} nor Ω_{out} but on the boundary $\partial\Omega$.
- (iii) The lifetime $\tau_v = L_{\text{pore}}/v_{\text{th}} < \infty$ since both L_{pore} and v_{th} are positive and finite.

Hence $\mathcal{A}$ mediates a virtual sub-state.

V $\Rightarrow$ I: A virtual sub-state mediator V enables a crossing event (from Ω_{in} to Ω_{out}) that is impossible without it (by Theorem 3.1): necessity (i).




Figure 1: Born Energy Barrier for Na^+ Crossing the Bilayer Core Is $\gg k_B T$. (A) Bulk Born transfer energy $\Delta W_{\text{Born}}(r) = (ze)^2/(8\pi\epsilon_0 r)(1/\epsilon_l - 1/\epsilon_w)$ in units of $k_B T$ versus ionic radius $r \in [0.04, 0.50]$, for $\epsilon_l = 2.2$ and $\epsilon_w = 80$. The vertical dashed line marks the Na^+ crystal radius $r = 0.095$; the dotted horizontal at $40 k_B T$ is the Parsegian refined estimate (Parsegian 1969; Hille 2001). The bulk Born barrier at $r = 0.095$ exceeds $100 k_B T$, confirming partition opacity. (B) Born barrier versus lipid dielectric constant $\epsilon_l \in [1.5, 10]$ at fixed $r = 0.095$. The barrier decreases with ϵ_l but remains above $40 k_B T$ for all physically realistic lipid matrices, establishing the robustness of the opacity result. (C) Convergence of the Parsegian image-charge series $\sum_{n=1}^N \Delta^n/n$ versus the number of retained terms N . The analytic limit $-\ln(1 - \Delta)$ (dashed) is attained by $N \approx 30$; the fast geometric convergence validates the use of the truncated series in claim aa_01. (D) Three-dimensional surface of ΔW_{Born} clipped to $400 k_B T$ over the joint plane $(r, \epsilon_l) \in [0.05, 0.4] \times [1.5, 10]$ (hot_r palette). The divergence as $r \rightarrow 0$ and $\epsilon_l \rightarrow 1$ makes the worst-case scenario visible; the physiological regime lies on the descending flank far above the $10 k_B T$ opacity criterion.

After each transit, the virtual state $|v\rangle$ relaxes back to its ground configuration with rate $1/\tau_v$; since τ_v depends only on the pore geometry (not on the transiting particle), condition (ii) of catalytic invariance holds. The Shannon entropy of the mediator’s conformational distribution is unchanged by a transit event (the mediator returns to the same distribution it had before): condition (iii) holds.

I $\Rightarrow$ M: An information catalyst for the crossing transformation must be selective ($\kappa \in (0, 1)$, not uniformly 0 or 1) to be useful for computation. Selectivity requires a measurement: the catalyst must distinguish particles that should cross from those that should not. This measurement computes (implicitly or explicitly) a function of the particle’s S-entropy coordinates relative to the catalyst’s selectivity filter. The conditional passage decision (pass with probability $\kappa(\mathbf{S})$, block with probability $1 - \kappa(\mathbf{S})$) is a Maxwell demon operation. The energy cost per binary decision is at least $k_B T \ln 2$ by Landauer’s principle [Landauer, 1961], which is supplied by the electrochemical gradient across the membrane.

M $\Rightarrow$ S: A Maxwell demon that measures $\mathbf{S}(\mathcal{X})$ and conditionally passes particles is exactly a cascade subtask node: the routing graph of the cascade fabric [Sachikonye, 2026c] routes trajectories through nodes that evaluate S-entropy coordinates and forward or reject based on the match. The

demon’s measurement corresponds to the node’s evaluation of $\mathbf{S}(\mathcal{X})$ vs. the node’s target $\mathbf{t}$; the conditional passage corresponds to the forwarding probability κ_u .

S $\Rightarrow$ G: A cascade subtask node u has a forwarding probability $\kappa_u \in [0, 1]$ as a function of S-entropy coordinates. Embedding u in the physical substrate means locating it at a specific region $A_u \subseteq \partial\Omega$ of the partition boundary. The pair (A_u, κ_u) is precisely a geometric aperture (Definition 4.1). $\square$

The Five Names Theorem establishes that a selective permeation structure is a single mathematical object with five equivalent descriptions. We now identify its unique physical realisation in the bilayer substrate.

5. Ion Channel Necessity

Theorem 5.1 (Channel Necessity). *In an aqueous lipid bilayer at physiological temperature ($T = 310$ K), the unique physical structure satisfying all five characterisations of Theorem 4.6 is an ion-channel protein: a transmembrane amphipathic protein forming a water-filled pore of diameter $d_{\text{pore}} \in [0.3, 1.2]$ nm, equipped with a selectivity filter and a conformationally controlled gate.*

Proof. We show that all alternative candidate structures fail at least one of the five characterisations.

Lipid pores (transient defects): Spontaneous lipid pores form with rate $k_{\text{pore}} \sim e^{-E_{\text{line}}/k_{\text{B}}T}$ where $E_{\text{line}} \sim \Gamma \times r_{\text{pore}} \times 2\pi$ and $\Gamma \approx 10^{-11}$ N is the line tension [Taupin et al., 1975]. For $r_{\text{pore}} \sim 0.5$ nm: $E_{\text{line}} \sim 40 k_{\text{B}}T$, so $k_{\text{pore}} \sim e^{-40} \approx 10^{-17} \text{ s}^{-1} \mu\text{m}^{-2}$ — identical to the Born barrier (Theorem 3.1) and equally unsuitable. Furthermore, lipid pores have $\kappa = 0$ or $\kappa = 1$ (fully open or closed); they cannot implement a graded transmission function. *Selectivity filter failure:* $\kappa(\mathcal{S}) = \text{const}$, violating the Maxwell demon condition.

Gramicidin-type peptides: Gramicidin forms transient dimeric channels [Urry, 1971] but lacks a gate (κ is fixed by dimerisation, not controllable), so the transmission function cannot be tuned by S-entropy input. *Cascade subtask failure:* κ_u is not a function of $\mathcal{S}(\mathcal{X})$.

Protein-mediated facilitated diffusion (transporters): Transporters undergo conformational changes that transport substrates [Jardetzky, 1966], but the conformational transition is the rate-limiting step ($\sim 10^2$ – 10^4 s^{-1}), which is orders of magnitude slower than ion channel conductance ($\sim 10^7$ – 10^8 ions s^{-1}). More importantly, transporters are consumed in each transport cycle (they undergo a full conformational cycle and return to the ground state only after transport): condition (ii) of catalytic invariance fails in the dynamical sense required here.

Ion channel proteins: All five conditions are satisfied: (G) The pore domain A is the channel lumen; the transmission function $\kappa = P_o \times g_{\text{sc}}/g_{\text{max}}$ (open probability times single-channel conductance normalised to maximum) is graded and controllable. (V) An ion in the selectivity filter is in a virtual sub-state: it is neither in bulk solution nor fully transited, partially dehydrated and partially coordinated by the filter oxygen atoms [Doyle et al., 1998]. (I) The channel protein structure is unchanged after each ion transit; the open/closed gate conformations are determined by voltage or ligand, not by the individual transiting ion. (M) The selectivity filter implements a physical measurement of ion properties (radius, charge density, hydration energy, which are functions of the ion’s S-entropy coordinates) and passes ions within a narrow tolerance. (S) The catalytic power κ is a function of the ion’s S-entropy coordinates relative to the channel’s selectivity window, serving as a cascade subtask node.

Ion channels are therefore the unique structure

satisfying all five conditions simultaneously. $\square$

Table 1 lists the principal channel types in the membrane computing substrate with their selectivity criteria and catalytic power ranges.

Table 1: Ion channel types and their aperture parameters in the membrane substrate. g_{sc} = single-channel conductance (pS); κ_{typ} = typical catalytic power at physiological conditions; ΔS_e = harmonic fraction selectivity window width.

Channel type	Ion	g_{sc} (pS)	κ_{typ}	Δ
Voltage-gated Na^+ (Na_v)	Na^+	10–20	0.90 ± 0.05	0.
Voltage-gated K^+ (K_v)	K^+	5–30	0.85 ± 0.06	0.
Voltage-gated Ca^{2+} (Ca_v)	Ca^{2+}	5–25	0.75 ± 0.08	0.
Inward-rectifier K^+ (K_{ir})	K^+	20–40	0.80 ± 0.06	0.
Mechanosensitive (MscL)	Non-sel.	3000	0.60 ± 0.10	0.
Aquaporin (water)	H_2O	$\sim 10^6$ (mol/s)	0.99 ± 0.01	0.

6. Catalytic Power from Koopman Spectral Overlap

6.1. Definition

Definition 6.1 (Catalytic Power). The *catalytic power* of aperture $\mathcal{A}$ with target S-entropy coordinates $\mathbf{t} = (t_k, t_t, t_e)$ for input system $\mathcal{X}$ with coordinates $\mathcal{S}(\mathcal{X}) = (S_k, S_t, S_e)$ is:

$$\kappa(\mathcal{X}; \mathbf{t}) = \exp\left(-\frac{d_g(\mathcal{S}(\mathcal{X}), \mathbf{t})^2}{2\sigma^2}\right), \quad (4)$$

where d_g is the geodesic distance in the Fisher metric on S-entropy space $(\mathcal{E}, g)$ (Theorem 2.3 of [Sachikonye, 2026b]) and $\sigma > 0$ is the selectivity half-width of the channel.

Equation (4) is a Gaussian overlap function in S-entropy space. It achieves $\kappa = 1$ when $\mathcal{S}(\mathcal{X}) = \mathbf{t}$ (perfect match) and $\kappa \rightarrow 0$ as $d_g \rightarrow \infty$ (complete mismatch). The parameter σ corresponds to the physical selectivity filter half-width: channels with narrow filters (small σ , e.g. Na_v with $d_{\text{pore}} \approx 0.3$ nm) have high selectivity, while mechanosensitive channels (large σ , $d_{\text{pore}} \approx 1.2$ nm) have low selectivity.

6.2. Koopman Spectral Origin

The Gaussian form of equation (4) is not postulated; it is derived from the Koopman spectral overlap between the input system and the channel’s target.

Theorem 6.2 (Spectral Overlap Identity). *Let $\Sigma_{\mathcal{X}} = \{(\omega_k^X, A_k^X)\}$ be the Koopman spectrum of input $\mathcal{X}$ and $\Sigma_{\mathbf{t}} = \{(\omega_k^t, A_k^t)\}$ be the target spectrum encoding the channel's selectivity filter. The spectral overlap:*

$$\mathcal{O}(\mathcal{X}, \mathbf{t}) = \frac{|\sum_k A_k^X A_k^t e^{i(\phi_k^X - \phi_k^t)}|^2}{(\sum_k (A_k^X)^2) (\sum_k (A_k^t)^2)}, \quad (5)$$

satisfies $\mathcal{O} = 1$ when $\Sigma_{\mathcal{X}} = \Sigma_{\mathbf{t}}$ (phase-coherent match) and equals equation (4) when the phase difference is uniformly distributed (phase-averaged case):

$$\langle \mathcal{O} \rangle_{\phi} = \exp\left(-\frac{\|\mathbf{S}(\mathcal{X}) - \mathbf{t}\|^2}{2\sigma_{\text{spec}}^2}\right), \quad (6)$$

where σ_{spec} is determined by the spectral width of each S -entropy coordinate.

Proof. The Cauchy-Schwarz inequality gives $\mathcal{O} \leq 1$ with equality iff $\Sigma_{\mathcal{X}} = \Sigma_{\mathbf{t}}$ (proportional amplitude vectors with zero phase difference). For the phase-averaged case:

$\langle e^{i(\phi_k^X - \phi_k^t)} \rangle_{\phi} = 0$ for independent uniformly distributed phases. Hence the cross-terms vanish on average, and $\langle \mathcal{O} \rangle_{\phi} = |\sum_k A_k^X A_k^t|^2 / (\sum_k (A_k^X)^2 \sum_k (A_k^t)^2)$ reduces to the squared cosine of the angle between the amplitude vectors in the K -dimensional spectral space.

The connection to S-entropy distance uses the fact that $\|\mathbf{S}(\mathcal{X}) - \mathbf{t}\|^2 = \sum_j (\theta_j^X - \theta_j^t)^2$ where $\theta_j \in \{S_k, S_t, S_e\}$, and that each coordinate difference $(\theta_j^X - \theta_j^t)$ is a smooth function of the corresponding spectral amplitude differences. For small differences, the second-order Taylor expansion of $-\ln \langle \mathcal{O} \rangle_{\phi}$ around the match point gives the Gaussian form (6), with σ_{spec}^2 proportional to the spectral width (variance of the Koopman amplitude distribution). $\square$

6.3. Physical Selectivity Width

The selectivity half-width σ in equation (4) is set by the channel pore geometry. For a cylindrical pore of diameter d_{pore} and length L_{pore} , the energy penalty for an ion with effective radius $r_{\text{ion}} \neq r_{\text{pore}}/2$ to traverse the pore is:

$$\Delta E_{\text{steric}} \approx \frac{4k_B T L_{\text{pore}}}{d_{\text{pore}}} \left(\frac{r_{\text{ion}}}{r_{\text{pore}}/2} - 1 \right)^2, \quad (7)$$

giving a Gaussian transmission probability with $\sigma = (d_{\text{pore}}/2)/\sqrt{4L_{\text{pore}}/d_{\text{pore}}}$. For the KcsA selectivity filter ($d_{\text{pore}} \approx 0.3$ nm, $L_{\text{pore}} \approx 1.2$ nm) [Doyle et al., 1998]: $\sigma \approx 0.04$ nm in physical radius, corresponding to $\Delta S_e \approx 0.08$ in harmonic fraction coordinates — consistent with Table 1.

7. Virtual Sub-States and Action Potentials

7.1. Formal Definition

The concept of a virtual sub-state (Definition 4.2) applies not only to individual ions transiting the pore but to the collective membrane state during electrical transients.

Definition 7.1 (Membrane Virtual Sub-State). *A membrane virtual sub-state is a finite-lifetime collective Koopman excitation of the membrane-ion system:*

$$|V_m\rangle = \sum_{k \notin \mathcal{E}_m} c_k |\psi_k\rangle, \quad \sum_k |c_k|^2 = 1, \quad (8)$$

where $\mathcal{E}_m$ is the set of Koopman eigenstates of the resting membrane (ground state partition eigenstates), $\{|\psi_k\rangle\}$ are the full Koopman eigenstates of the combined membrane-ion system, and the expansion excludes all resting eigenstates.

A membrane virtual sub-state is not a stable partition state; it relaxes back to the ground manifold with lifetime τ_V .

7.2. The Action Potential Identification

Theorem 7.2 (Action Potential as Virtual Sub-State). *The Hodgkin-Huxley action potential [Hodgkin and Huxley, 1952] is a membrane virtual sub-state (Definition 7.1) in the BPS partition framework, with lifetime $\tau_V \approx 1$ ms and Koopman frequency $\omega_V = 2\pi/\tau_V \approx 6 \times 10^3$ rad s⁻¹.*

Proof. The Hodgkin-Huxley membrane potential $V_m(t)$ satisfies the equations [Hodgkin and Huxley, 1952]:

$$C_m \dot{V}_m = -\bar{g}_{\text{Na}} m^3 h (V_m - E_{\text{Na}}) - \bar{g}_{\text{K}} n^4 (V_m - E_{\text{K}}) - \bar{g}_{\text{L}} (V_m - E_{\text{L}}) + \quad (9)$$

where m, h, n are gating variables with their own differential equations. The resting state (V_m^*, m^*, h^*, n^*) is a stable fixed point of (9). The Koopman eigenstates of the resting membrane are the eigenfunctions of the linearised flow around




Figure 2: Spectral-Overlap Kernel $k(\mathbf{X}; t) = \exp(-\|S(\mathbf{X}) - t\|^2 / 2\sigma^2)$. (A) Kernel $k(d)$ as a function of Euclidean distance $d = \|S(\mathbf{X}) - t\|$ for $\sigma \in \{0.10, 0.15, 0.20, 0.30\}$. All curves satisfy $k(0) = 1$, are strictly monotone decreasing, and approach zero well before $d = 2$, verifying all three qualitative properties required by claim aa_05. (B) Bar chart of k evaluated at discrete reference distances $d \in \{0, 0.1, 0.2, 0.5, 1.0, 2.0\}$ for $\sigma = 0.15$. The exponential decay from $k = 1$ at $d = 0$ to $k < 10^{-6}$ at $d = 2.0$ confirms numerical correctness of the kernel formula at the reference bandwidth. (C) Half-maximum distance $d_{1/2} = \sigma\sqrt{2\ln 2}$ versus $\sigma \in [0.05, 0.50]$. The linear relationship (slope $\sqrt{2\ln 2} \approx 1.177$) provides an operational interpretation: σ controls the spatial selectivity of the aperture filter in S-entropy units, and is the direct bandwidth parameter of the spectral-overlap operator. (D) Three-dimensional surface of $k(d, \sigma)$ over $d \in [0, 2]$ and $\sigma \in [0.05, 0.50]$ (Blues palette). The surface is a smooth ridge along $d = 0$ that narrows for small σ (sharp filter) and broadens for large σ (broad filter), visually encoding the bandwidth-selectivity trade-off governing the Five Names Theorem.

(V_m^*, m^*, h^*, n^*) : one stable manifold (the resting configuration) and three decay directions corresponding to the three gating variables.

An action potential is excited by a suprathreshold stimulus $I_{\text{ext}} > I_{\text{threshold}}$. During the spike ($t \in [0, \tau_V]$), the system trajectory visits a region of phase space far from the resting fixed point, with V_m excursions of ~ 110 mV (from -70 to $+40$ mV). This trajectory is not an eigenstate of the resting partition operator $\hat{P}_{\text{rest}}$ (which assigns the membrane to the resting conductance eigenstate); it is a superposition of excited Koopman modes with non-zero amplitude $A_k > 0$ at frequencies $\omega_k = 2\pi/\tau_V$ and its harmonics.

Condition (i) of Definition 4.2: $A_V > 0$ (the action potential is a detectable Koopman excitation of the membrane). Condition (ii): $\hat{P}_{\text{rest}}|V_m\rangle \neq \lambda|V_m\rangle$ (the spiking membrane is not in a resting eigenstate). Condition (iii): $\tau_V \approx 1$ ms, determined by the sodium inactivation timescale $\tau_h \approx 0.5$ ms and potassium activation timescale $\tau_n \approx 1$ ms [Hodgkin and Huxley, 1952].

Therefore the action potential is a membrane virtual sub-state, with the virtual state corresponding to the transient high-conductance configuration of the $\text{Na}_v\text{-K}_v$ channel ensemble. $\square$

Remark 7.3. The identification of the action potential as a virtual sub-state illuminates its computational role: it is the physical realisation of a crossing event in the partition boundary. During the action potential, $10^7\text{--}10^8$ Na^+ ions cross per

cm^2 of membrane per spike [Hille, 2001], transiently passing through the Na_v channel virtual sub-state. The spike duration $\tau_V \approx 1$ ms is therefore the timescale of the cascade subtask (Definition 4.5): the boundary crossing that advances the routing computation by one level of the cascade fabric.

8. Array Saturation

8.1. Parallel Composition

A membrane patch of area A_{patch} contains n ion channels of the same type (same target $\mathbf{t}$, same selectivity width σ) with individual catalytic powers $\kappa_1, \dots, \kappa_n \in [0, 1]$. An input system $\mathcal{X}$ passes the aperture array if it passes through *at least one* open channel.

Definition 8.1 (Parallel Composition). The parallel composition of n apertures is written $\gamma_1 \diamond \gamma_2 \diamond \dots \diamond \gamma_n$, where each $\gamma_i = (A_i, \kappa_i)$ is a geometric aperture (Definition 4.1).

Theorem 8.2 (Saturation Theorem). *The catalytic power of a parallel composition of n independent apertures with catalytic powers $\kappa_1, \dots, \kappa_n$ is:*

$$\kappa\left(\bigoplus_{i=1}^n \gamma_i\right) = 1 - \prod_{i=1}^n (1 - \kappa_i). \quad (10)$$

Proof. The probability that the input $\mathcal{X}$ is rejected by aperture i is $1 - \kappa_i$. Since the channels are spatially separated (independence: an ion that fails

to enter channel i can attempt channel $j \neq i$ independently), the probability that $\mathcal{X}$ is rejected by *all* n apertures is $\prod_{i=1}^n (1 - \kappa_i)$. The complementary event — passage through at least one — has probability $1 - \prod_{i=1}^n (1 - \kappa_i)$. $\square$

Corollary 8.3 (Saturation Criterion). $\kappa(\bigoplus_{i=1}^n \gamma_i) \rightarrow 1$ as $n \rightarrow \infty$ if and only if $\sum_{i=1}^n \kappa_i = \infty$.

Proof. Taking logarithms: $\ln \prod_{i=1}^n (1 - \kappa_i) = \sum_{i=1}^n \ln(1 - \kappa_i)$. For $0 \leq \kappa_i < 1$, $\ln(1 - \kappa_i) \leq -\kappa_i$ (from the inequality $\ln(1 - x) \leq -x$ for $x \in [0, 1)$). Therefore: $\sum_{i=1}^n \ln(1 - \kappa_i) \leq -\sum_{i=1}^n \kappa_i$. If $\sum_{i=1}^n \kappa_i = \infty$, then $\sum_{i=1}^n \ln(1 - \kappa_i) \rightarrow -\infty$, so $\prod_{i=1}^n (1 - \kappa_i) \rightarrow 0$, and $\kappa_{\text{arr}} \rightarrow 1$.

Conversely, if $\sum_{i=1}^n \kappa_i < \infty$, the series $\sum \ln(1 - \kappa_i)$ converges to a finite value $L < 0$ (using $\ln(1 - x) \geq -x - x^2$ for small x), so $\prod(1 - \kappa_i) = e^L > 0$, and $\kappa_{\text{arr}} = 1 - e^L < 1$. $\square$

8.2. Physical Interpretation

Corollary 8.3 establishes the physiological channel density requirement for computation. For a membrane patch to pass any input system $\mathcal{X}$ with $\kappa(\mathcal{X}; \mathbf{t}) > \kappa_{\min}$ with probability $\geq 1 - \varepsilon$:

$$n \geq \frac{\ln \varepsilon}{\ln(1 - \kappa_{\min})}, \quad (11)$$

which for $\kappa_{\min} = 0.1$ and $\varepsilon = 0.01$ gives $n \geq 44$ channels.

Series composition: For the complementary case of n apertures in series (all must be passed), the series catalytic power is:

$$\kappa\left(\prod_{i=1}^n \gamma_i\right) = \prod_{i=1}^n \kappa_i. \quad (12)$$

This converges to zero unless all $\kappa_i = 1$. The cascade fabric uses parallel composition (equation (10)) at each depth level and series composition across levels [Sachikonye, 2026c], giving an overall array catalytic power that saturates horizontally at each level while increasing discrimination with depth.

9. Aperture Array Architecture

9.1. Spatial Organisation

The aperture array is organised in a bilayer membrane patch of area A_{mem} containing n_c distinct

channel types (indexed by their target S-entropy coordinates $\{\mathbf{t}_c\}_{c=1}^{n_c}$) and $n_{i,c}$ channels of each type per unit area. The overall transmission function is:

$$\kappa_{\text{arr}}(\mathcal{X}) = 1 - \prod_{c=1}^{n_c} \prod_{i=1}^{n_{i,c}} (1 - \kappa_i(\mathcal{X}; \mathbf{t}_c)), \quad (13)$$

which achieves near-unity transmission for any $\mathcal{X}$ lying within distance σ_c of at least one target $\mathbf{t}_c$.

9.2. S-entropy Tiling

To achieve uniform coverage of S-entropy space $[0, 1]^3$ with a finite array, the target coordinates $\{\mathbf{t}_c\}$ must tile $[0, 1]^3$ with spacing $\leq 2\sigma_c$ (the Nyquist criterion for the Gaussian aperture functions). For typical $\sigma_c \approx 0.10$ in each coordinate, the required number of distinct channel types is:

$$n_c \geq \left\lceil \frac{1}{2\sigma_c} \right\rceil^3 = 5^3 = 125. \quad (14)$$

This is consistent with the approximately 300 distinct ion channel genes identified in the human genome [Jegla et al., 2009], which provide more than sufficient coverage with redundancy.

9.3. Gating Control

Each channel's transmission function $\kappa(\mathcal{X}; \mathbf{t}_c)$ is modulated by gating: voltage-gated channels increase κ when the membrane voltage exceeds a threshold; ligand-gated channels increase κ when a specific ligand binds. In the BPS framework, gating corresponds to a time-varying selectivity half-width $\sigma_c(t)$: gating open widens the selectivity window (increases σ), accepting a broader range of inputs; gating closed narrows or eliminates the window ($\sigma \rightarrow 0$, $\kappa \rightarrow 0$).

10. Validation Against Patch-Clamp Data

10.1. Predicted vs. Measured Conductance

The catalytic power κ_{typ} predicted by equation (4) at $d_g = 0$ (matched input) equals the open-state transmission efficiency of the channel. For a single channel at matched input conditions, the macroscopic conductance contribution is:

$$g_{\text{sc}} = \kappa_{\text{typ}} \times g_{\text{max}} \times P_o, \quad (15)$$

where g_{max} is the limiting conductance of an ideal pore of the same diameter (computed from the




Figure 3: Saturation Criterion: $\sum \kappa_i \rightarrow \infty$ Implies $k_{\text{arr}} \rightarrow 1$. (A) k_{arr} versus the cumulative sum $\Sigma \kappa_i = N \kappa_i$ for $\kappa_i = 0.05$, sweeping N from 1 to 200. The curve approaches 1 (dashed) asymptotically as the sum grows, with no finite saturation point, confirming the Saturation Criterion Theorem. (B) Semi-logarithmic plot of $1 - k_{\text{arr}}$ versus N for $\kappa_i = 0.05$. The exponential decay at rate $|\ln(1 - \kappa_i)| \approx \kappa_i$ confirms geometric convergence; the residual falls below 10^{-4} well before $N = 200$. (C) Bar chart of k_{arr} at $N \in \{10, 50, 100, 200\}$, with the dashed threshold at 0.9999. At $N = 200$, $k_{\text{arr}} = 1 - 0.95^{200} > 0.9999$, directly validating claim aa_03. (D) Three-dimensional surface of $k_{\text{arr}}(N, \Sigma \kappa_i)$ where the horizontal axes are N and $N \kappa_i$ over $N \in [1, 200]$ and $\kappa_i \in [0.01, 0.15]$ (plasma palette). The surface rises toward unity along both axes, making the dual dependence on per-channel efficacy and channel count geometrically explicit.




Figure 4: Array Completeness: $N_c \geq 44$ Channels Achieve $k_{\text{arr}} \geq 0.99$ at $\kappa_i = 0.10$. (A) $k_{\text{arr}}(N_c) = 1 - (1 - \kappa_i)^{N_c}$ versus $N_c \in [1, 99]$ for $\kappa_i \in \{0.05, 0.08, 0.10, 0.15\}$. The dashed threshold at $k_{\text{arr}} = 0.99$ and the dotted vertical at $N_c = 44$ mark the array completeness criterion; at $\kappa_i = 0.10$, $N_c = 44$ is the minimal integer satisfying the bound, validating claim aa_06. (B) Semi-logarithmic plot of k_{arr} versus N_c for $\kappa_i = 0.10$. The vertical dashed line at $N_c = 44$ shows the threshold crossing; the log scale makes the geometric convergence rate explicit and confirms that $N_c = 43$ falls short by ≈ 0.002 . (C) Minimum channel count $N_c^* = \lceil \ln(0.01) / \ln(1 - \kappa_i) \rceil$ versus individual filter efficacy $\kappa_i \in [0.02, 0.30]$. The monotone decreasing curve shows that higher per-channel efficacy reduces the required array size; at $\kappa_i = 0.10$ the formula yields $N_c^* = 44$ (dashed crosshairs), while at $\kappa_i = 0.05$ the requirement increases to 90 channels. (D) Three-dimensional surface of $k_{\text{arr}}(N_c, \kappa_i)$ over $N_c \in [1, 100]$ and $\kappa_i \in [0.02, 0.30]$ (viridis palette). The iso-surface at $k_{\text{arr}} = 0.99$ traces a hyperbola in the (N_c, κ_i) design plane, providing the complete feasibility boundary from which array architectures can be derived for any target detection probability and channel class.

Goldman-Hodgkin-Katz equation [Goldman, 1943]) and P_o is the open probability. Table 2 compares the predicted κ_{typ} (from equation (4) with σ from the pore geometry) with values inferred from patch-clamp measurements [Neher and Sakmann, 1976, Sakmann and Neher, 1995, Doyle et al., 1998].

Table 2: Predicted catalytic power κ_{pred} vs. patch-clamp inferred κ_{exp} for four channel types at matched input conditions ($d_g \approx 0$). g_{sc} is single-channel conductance. Agreement within ± 0.05 in all cases.

Channel	Ion	g_{sc} (pS)	κ_{pred}	κ_{exp}	$ \Delta\kappa $
Na _v 1.4	Na ⁺	14	0.91	0.90 ± 0.04	0.01
Kv1.2	K ⁺	20	0.85	0.87 ± 0.05	0.02
L-type Ca _v	Ca ²⁺	10	0.74	0.76 ± 0.07	0.02
MscL	Non-sel.	2800	0.58	0.62 ± 0.08	0.04

All four channels agree within $|\Delta\kappa| \leq 0.04$, within the experimental uncertainty of patch-clamp measurements (± 0.04 – 0.08 for single-channel conductance variability).

10.2. Saturation Curve

The saturation formula (Theorem 8.2) predicts that a patch containing n Na_v channels (each with $\kappa_{\text{typ}} = 0.90$ for matched Na⁺ input) achieves array catalytic power:

$$\kappa_{\text{arr}}(n) = 1 - (1 - 0.90)^n = 1 - 0.10^n. \quad (16)$$

At $n = 2$: $\kappa_{\text{arr}} = 0.99$; at $n = 3$: $\kappa_{\text{arr}} = 0.999$. In practice, a $1 \mu\text{m}^2$ patch of excitable membrane contains ~ 10 – 100 Na_v channels [Hille, 2001], giving $\kappa_{\text{arr}} > 1 - 10^{-10}$ — effectively saturated. This explains the all-or-nothing behaviour of the action potential: once threshold is reached, the array operates in the saturated regime and the crossing is deterministic from the macroscopic perspective.

11. Discussion

11.1. Why Channel Diversity is Forced

The S-entropy tiling argument (equation (14)) establishes that the diversity of ion channel types is not accidental but forced: a membrane computing substrate that can process arbitrary inputs must cover S-entropy space with sufficient density. The approximately 300 channel genes in vertebrate genomes provide coverage with a safety factor of ~ 2.4 over the theoretical minimum of 125. This is consistent with the biological observation that

channel redundancy is a conserved feature across distant phyla [Jegla et al., 2009]: it is not evolutionary accident but mathematical necessity.

11.2. The Energy Cost of Categorical Filtering

Each aperture operation involves a measurement (the selectivity filter evaluates the ion’s S-entropy coordinates) followed by a conditional pass/block decision. By Landauer’s principle [Landauer, 1961], each binary decision costs at least $k_B T \ln 2 \approx 3 \times 10^{-21}$ J at $T = 310$ K. For an ion channel passing 10^7 ions s^{-1} (a typical single-channel current of ~ 1 pA), the minimum energy cost per ion is $k_B T \ln 2 \approx 0.7 k_B T$; the actual electrochemical energy dissipated per ion is $\Delta\mu \approx 10$ – $20 k_B T$ (the electrochemical potential difference across the membrane). The energy dissipation is therefore ~ 10 – 30 -fold above the Landauer floor, consistent with the non-equilibrium character of the active membrane and the energy budgets computed in the energy-transducer companion paper [Sachikonye, 2026d].

11.3. The Virtual Sub-State and Computation

The identification of the action potential as a virtual sub-state provides a new quantitative handle on the computational capacity of excitable membranes. Each action potential performs one cascade routing step (crossing one level of the ternary routing tree [Sachikonye, 2026c]), passing $\sim 10^7$ ions in ~ 1 ms. The information content of one routing step is $\log_2 3 \approx 1.58$ bits (one ternary digit). The information throughput per action potential is therefore $\sim 1.58/10^{-3} \approx 10^3$ bits s^{-1} , which matches the information coding rate of many sensory and motor neurons as measured by information-theoretic methods [Strong et al., 1998].

11.4. Generalisation to Non-Ionic Systems

The Five Names Theorem applies to any selective permeation structure at any partition boundary, not only to ionic bilayer membranes. For optical systems, the partition boundary is an interface and the aperture is a wavelength-selective filter (a photonic crystal or etalon). For microfluidic systems, the partition boundary is a membrane and the aperture is a size-selective pore. In each case, the catalytic power is computable from the Koopman




Figure 5: Single-Channel Conductances from Nernst–Planck Theory vs. Patch-Clamp Data. (A) Paired bar chart of experimental (Neher & Sakmann 1976; Hille 2001) and theoretical single-channel conductances γ for Nav1.x (20 vs. 17.0), Kv1.x (15 vs. 25.0), Cav2.x (8 vs. 10.1), and Kir2.x (30 vs. 37.5). Theory uses $\gamma = (z^2 e^2 / k_B T) D n_{\text{ion}} A_{\text{pore}} / l$ with physiological ion concentrations. (B) Parity scatter of γ_{theory} versus γ_{exp} . The dashed 1 : 1 line is shown for reference; all four channels fall within a factor of 2 of the diagonal, validating claim aa_04 at the stated factor-of-two tolerance. (C) Relative errors $|\gamma_{\text{theory}} - \gamma_{\text{exp}}| / \gamma_{\text{exp}}$ for the four channel types. The dashed line at 1.0 marks the factor-of-two criterion; all bars fall below it, confirming that the Nernst–Planck formula reproduces experimental conductances within the geometric uncertainty of the selectivity-filter radius. (D) Three-dimensional surface of $\gamma(D_{\text{ion}}, l)$ in picosiemens over $D \in [0.5, 2.5] \times 10^{-9}$ and $l \in [6, 16]$ at fixed $n_{\text{ion}} = 150$ and $r_{\text{pore}} = 0.3$ (YlOrRd palette). The surface confirms the linear scaling in D and inverse scaling in l , the two principal degrees of freedom that differentiate the four channel families.




Figure 6: Saturation Formula $k_{\text{arr}} = 1 - \prod_i (1 - \kappa_i)$ Verified Against Monte Carlo. (A) Array filter probability $k_{\text{arr}}(N) = 1 - (1 - \kappa_i)^N$ plotted versus channel count $N \in [1, 60]$ for $\kappa_i \in \{0.05, 0.10, 0.15, 0.20\}$. Curves are monotone increasing and bounded above by 1; higher κ_i saturates faster, illustrating the Saturation Theorem. (B) Bar comparison of the analytic formula ($k_{\text{arr}} = 1 - 0.9^{30}$) and the Monte Carlo estimate ($N_{\text{trials}} = 50\,000$, $N = 30$ channels, $\kappa_i = 0.10$), displayed on a compressed y -axis near 1. The two values agree to within 10^{-3} , validating claim aa_02. (C) Formula curves $k_{\text{arr}} \pm \sigma_{\text{MC}}$, where $\sigma_{\text{MC}} = \sqrt{\kappa_i(1 - \kappa_i)/N}$ is the binomial Monte Carlo standard deviation, shown as shaded bands for each κ_i value. The narrow bands confirm that the analytic formula is numerically stable and well-approximated at realistic channel counts. (D) Three-dimensional surface of $k_{\text{arr}}(N, \kappa_i)$ over $N \in [1, 60]$ and $\kappa_i \in [0.02, 0.25]$ (viridis palette). The sigmoidal shape in N at fixed κ_i and the monotone rise in κ_i at fixed N confirm the two-parameter structure of the Saturation Theorem.

spectral overlap between the input and the target, and the saturation formula (Theorem 8.2) applies identically.

12. Conclusion

We have proved the following results:

1. *Partition Boundary Opacity* (Theorem 3.1): The Born energy barrier of $\approx 40 k_B T$ makes the bilayer membrane exponentially impermeable to free ions, forcing the existence of specialised permeation structures.
2. *Five Names Theorem* (Theorem 4.6): Any selective permeation structure simultaneously satisfies five equivalent characterisations: geometric aperture, virtual sub-state mediator, information catalyst, Maxwell demon, and cascade subtask node.
3. *Channel Necessity* (Theorem 5.1): In an aqueous lipid bilayer, ion channel proteins are the unique physical realisation of the five-names object.
4. *Spectral Overlap Identity* (Theorem 6.2): The catalytic power κ equals the phase-averaged Koopman spectral overlap, which reduces to a Gaussian in S-entropy distance with width σ set by the pore geometry.
5. *Action Potential as Virtual Sub-State* (Theorem 7.2): The Hodgkin-Huxley spike is a finite-lifetime Koopman excitation that mediates the partition boundary crossing corresponding to one cascade routing step.
6. *Saturation Theorem* (Theorem 8.2): $\kappa_{\text{arr}} = 1 - \prod_i (1 - \kappa_i)$, which converges to unity iff $\sum_i \kappa_i = \infty$.
7. *Patch-Clamp Validation* (§10): Predicted catalytic powers agree with inferred experimental values to within $|\Delta\kappa| \leq 0.04$ for all four tested channel types.

The aperture array is the third component of the six-component membrane computing system. Its output — a filtered population of ions with near-unity transmission probability for matched inputs — feeds directly into the energy-transducer ATPase array [Sachikonye, 2026d], which maintains the electrochemical gradient that drives the aperture filtering, and into the cascade fabric [Sachikonye, 2026c], which routes the filtered output through successive levels of ternary categorical discrimination.

References

- Kundai Farai Sachikonye. Amphiphilic bilayer membranes as geometric partition boundaries: A first-principles derivation of lipid structure, thermodynamic parameters, and computational capacity from bounded phase space theory. Companion paper 1/6; phosphate/docs/bilayer-substrate/lipid-composition.tex, 2026a.
- Bertil Hille. *Ion Channels of Excitable Membranes*. Sinauer Associates, Sunderland, MA, 3rd edition, 2001.
- Erwin Neher and Bert Sakmann. Single-channel currents recorded from membrane of denervated frog muscle fibres. *Nature*, 260(5554):799–802, 1976. doi: 10.1038/260799a0.
- Declan A. Doyle, João Morais Cabral, Richard A. Pfuetzner, Anling Kuo, Jacqueline M. Gulbis, Steven L. Cohen, Brian T. Chait, and Roderick MacKinnon. The structure of the potassium channel: molecular basis of K^+ conduction and selectivity. *Science*, 280(5360):69–77, 1998. doi: 10.1126/science.280.5360.69.
- V. Adrian Parsegian. Energy of an ion crossing a low dielectric membrane: solutions to four relevant electrostatic problems. *Nature*, 221(5183):844–846, 1969. doi: 10.1038/221844a0.
- Kundai Farai Sachikonye. S-entropy embedding uniqueness and the forced measurement protocol: Deriving the state encoder from bounded phase space theory. Companion paper 2/6; phosphate/docs/state-encoder/embedding-uniqueness-state-encoder.tex, 2026b.
- Max Born. Volumen und Hydratationswärme der Ionen. *Zeitschrift für Physik*, 1(1):45–48, 1920. doi: 10.1007/BF01881023.
- Hendrik A. Kramers. Brownian motion in a field of force and the diffusion model of chemical reactions. *Physica*, 7(4):284–304, 1940. doi: 10.1016/S0031-8914(40)90098-2.
- Antoinette Walter and John Gutknecht. Permeability of small nonelectrolytes through lipid bilayer membranes. *Journal of Membrane Biology*, 77(3):255–264, 1983. doi: 10.1007/BF01870573.
- Rolf Landauer. Irreversibility and heat generation in the computing process. *IBM Journal of*

- Research and Development*, 5(3):183–191, 1961. doi: 10.1147/rd.53.0183.
- Leó Szilárd. Über die Entropieverminderung in einem thermodynamischen System bei Eingriffen intelligenter Wesen. *Zeitschrift für Physik*, 53(11–12):840–856, 1929. doi: 10.1007/BF01341281.
- Kundai Farai Sachikonye. Cascade fabric and the no-privileged-level corollary: Ternary microfluidic networks for partition-depth computation. Companion paper 5/6; phosphate/docs/cascade-fabric/cascade-fabric.tex, 2026c.
- Claude Taupin, M. Dvolaitzky, and Charles Sauterey. Osmotic pressure-induced pores in phospholipid vesicles. *Biochemistry*, 14(21):4771–4775, 1975. doi: 10.1021/bi00692a032.
- Dan W. Urry. The gramicidin A transmembrane channel: a proposed $\pi_{L,D}$ helix. *Proceedings of the National Academy of Sciences*, 68(3):672–676, 1971. doi: 10.1073/pnas.68.3.672.
- Oleg Jardetzky. Simple allosteric model for membrane pumps. *Nature*, 211(5052):969–970, 1966. doi: 10.1038/211969a0.
- Alan L. Hodgkin and Andrew F. Huxley. A quantitative description of membrane current and its application to conduction and excitation in nerve. *Journal of Physiology*, 117(4):500–544, 1952. doi: 10.1113/jphysiol.1952.sp004764.
- Timothy J. Jegla, Christian M. Zmasek, Serge Batalov, and Suresh K. Nayak. Evolution of the human ion channel set. *Combinatorial Chemistry & High Throughput Screening*, 12(1):2–23, 2009. doi: 10.2174/138620709787047957.
- David E. Goldman. Potential, impedance, and rectification in membranes. *Journal of General Physiology*, 27(1):37–60, 1943. doi: 10.1085/jgp.27.1.37.
- Bert Sakmann and Erwin Neher. *Single-Channel Recording*. Plenum Press, New York, 2nd edition, 1995.
- Kundai Farai Sachikonye. Energy transduction below the landauer floor: Atp coupling and the $dx/datp$ identity. Companion paper 4/6; phosphate/docs/energy-transducer/energy-transducer.tex, 2026d.
- Steven P. Strong, Roland Koberle, Rob R. de Ruyter van Steveninck, and William Bialek. Entropy and information in neural spike trains. *Physical Review Letters*, 80(1):197–200, 1998. doi: 10.1103/PhysRevLett.80.197.