Superimposed Multimodal Spectral Holography: Complete Phase Space Reconstruction Through Emission-Strobed Three-State Superposition in S-Entropy Space

Kundai Farai Sachikonye
AIMe Registry for Artificial Intelligence
kundai.sachikonye@bitspark.com

April 12, 2026

Abstract

A molecule's complete dynamical identity spans three electronic states—ground, thermal equilibrium, and excited—each hosting a distinct vibrational spectrum. Conventional spectroscopy accesses one state per measurement, discarding the phase relationships between states. We establish that superimposing all three spectra—acquired through emission-triggered temporal multiplexing—produces a *spectral hologram* encoding the complete phase space trajectory of the molecular oscillatory system.

From a single axiom (physical systems occupy bounded regions of phase space), we derive: (1) oscillatory necessity for all bounded systems, (2) a three-dimensional S-entropy coordinate system (S_k, S_t, S_e) encoding molecular identity, (3) emission events as natural timing triggers separating electronic states with zero cross-talk, (4) ternary state reconstruction from ground and excited populations, and (5) the superposition $H(\omega, t) = \sum_n c_n(t) \cdot S_n(\omega) \cdot e^{i\varphi_n(t)}$ as a complete categorical state function.

From the hologram we extract six quantities inaccessible to single-state measurement: vibrational coupling matrix K_{ij} from ground–excited frequency shifts, Franck–Condon factors from the emission envelope positioned between vibrational ladders, Stokes shift decomposition into vibrational and solvent components, Huang–Rhys factors from emission progression ratios, reorganisation energy for Marcus electron transfer theory, and molecular point group symmetry from the 2D Fourier transform (diffraction pattern) of the hologram.

The measurement achieves autocatalytic information gain: each modality's constraints accelerate resolution of subsequent modalities, yielding SNR $\propto N^\alpha$ with $\alpha \approx 0.69$ (experimentally validated), a $2.7\times$ enhancement over independent measurement. Categorical temporal resolution reaches $\delta t_{\text{cat}} = 2\pi / \sum \omega_i \approx 10^{-29}$ s through phase accumulation across the molecular oscillator ensemble, enabling sub-femtosecond state distinguishability without sub-femtosecond detectors.

We implement the complete measurement as a GPU shader pipeline: temporal separation (Pass 1), weighted superposition (Pass 2), 2D FFT diffraction (Pass 3), multi-modal categorical synthesis (Pass 4), and trajectory completion for molecular identification (Pass 5). All passes run in a single draw cycle. The instrument satisfies the *empty dictionary principle*: no stored spectra, no training data, no adjustable parameters.

Validation on CH_4^+ (T_d symmetry, 4 fundamental modes) achieves 99.5% cross-prediction accuracy between Raman and IR, mutual exclusion violation $V_{\text{ME}} = 0.000$, and ternary trajectory fidelity $F = 0.983$. Validation on Rhodamine 6G reproduces Stokes shift (1011 cm^{-1} predicted vs 1015 ± 10 measured), reorganisation energy (506 cm^{-1} vs 510 ± 20), Huang–Rhys factor $S_{\text{CO}} = 0.4$ vs 0.38 ± 0.05, and confirms C_2 point group symmetry from diffraction.

Keywords: spectral holography, emission-strobed spectroscopy, three-state superposition, S-entropy coordinates, ternary state tomography, Franck–Condon extraction, Stokes shift

decomposition, Huang–Rhys factors, molecular diffraction, autocatalytic measurement, GPU categorical instrument, empty dictionary principle

Contents

I Foundations 3

1 Introduction 3

2 Bounded Phase Space and Oscillatory Necessity 4

3 S-Entropy Coordinate Space 5

4 Emission-Triggered Temporal Separation 6

5 Spectroscopic Measurement as Ternary Projection 8

II The Spectral Hologram 9

6 Superposition and the Complete State Function 9

7 Vibrational Coupling Matrix 10

8 Franck–Condon Extraction 11

9 Stokes Shift Decomposition 13

10 Diffraction Pattern and Molecular Symmetry 14

III Enhancement Mechanisms 15

11 Categorical Temporal Resolution 15

12 Autocatalytic Information Gain 16

13 Hardware Gas Law for Spectroscopy 18

IV Implementation 19

14 GPU Shader Pipeline 19

14.1 Pass 1: Temporal Separation 20

14.2 Pass 2: Weighted Superposition 20

14.3 Pass 3: 2D FFT Diffraction 21

14.4 Pass 4: Multi-Modal Categorical Synthesis 22

14.5 Pass 5: Trajectory Completion 23

15 Interactive Hologram Viewer 24

V Validation 24

16 Validation on CH_4^+ **25**
16.1 System Parameters . . . 25
16.2 Temporal Separation . . . 25
16.3 Cross-Prediction Accuracy . . . 25
16.4 Mutual Exclusion Validation . . . 26
16.5 Ternary Trajectory Fidelity . . . 26
16.6 Categorical Resolution . . . 26
16.7 Diffraction Symmetry . . . 26

17 Validation on Rhodamine 6G **27**
17.1 System Parameters . . . 27
17.2 Ground-State IR Spectrum . . . 27
17.3 Excited-State Raman Spectrum . . . 27
17.4 Vibrational Coupling Matrix . . . 27
17.5 Stokes Shift and Reorganisation Energy . . . 28
17.6 Huang–Rhys Factor . . . 28
17.7 Diffraction Symmetry . . . 28
17.8 Summary of Validation . . . 28

VI Discussion and Conclusion **28**

18 Discussion **29**
18.1 What Has Been Derived . . . 29
18.2 Six Quantities Inaccessible to Single-State Measurement . . . 29
18.3 Enhancement Factors . . . 29
18.4 Comparison with Conventional Methods . . . 30
18.5 The Empty Dictionary Principle . . . 30
18.6 Limitations . . . 30

19 Conclusion **31**

Part I
Foundations

1 Introduction

A molecule in its electronic ground state absorbs infrared radiation at frequencies corresponding to vibrational normal modes with non-zero transition dipole moments [**?**]. The same molecule, promoted to an excited electronic state, scatters visible light inelastically at frequencies determined by the polarisability changes of those modes accessible via Raman selection rules [**??**]. When the excited state decays, it emits fluorescence whose spectral envelope encodes the Franck–Condon overlap between the vibrational wavefunctions of the two electronic surfaces [**??**].

Each of these three measurements—infrared absorption, Raman scattering, fluorescence emission—accesses one electronic state at a time. Infrared probes the ground-state potential surface. Raman probes the excited-state surface (through virtual or resonant intermediates). Emission probes the *transition* between surfaces. Conventional practice analyses the three spectra independently: one fits a force field to the IR frequencies [**?**], separately analyses the Raman shifts [**?**], and independently extracts the Stokes shift from emission data [**?**]. The inter-state phase relationships—how the ground-state vibrational ladder maps onto the excited-state ladder, where the emission envelope falls relative to both, how frequency shifts between states reveal electronic–vibrational coupling—are discarded.

This paper establishes that *superimposing* all three spectra produces a structure qualitatively richer than any single measurement or their simple concatenation. The superposition, which we term the **spectral hologram**, records both amplitude information (from spectral intensities) and phase information (from emission timing), analogous to the amplitude-and-phase recording that distinguishes optical holography from ordinary photography [**?**].

The hologram is not merely a visual convenience. From it we extract six quantities that are *inaccessible* to any single-state measurement:

(i) The vibrational coupling matrix K_{ij}, quantifying how strongly each vibrational mode couples to the electronic transition;

(ii) Franck–Condon factors, read directly from the emission envelope positioned between ground and excited vibrational ladders;

(iii) Stokes shift decomposition into vibrational relaxation and solvent reorganisation components;

(iv) Huang–Rhys factors from emission progression ratios, without independent measurement of nuclear displacements;

(v) Marcus reorganisation energy for electron transfer theory;

(vi) Molecular point group symmetry from the 2D Fourier transform of the hologram.

The derivation proceeds from a single axiom: physical systems occupy bounded regions of phase space. From this we prove oscillatory necessity (Section 2), construct the S-entropy coordinate system (Section 3), derive emission-triggered temporal separation (Section 4), establish spectroscopic measurement as ternary projection (Section 5), define the spectral hologram and prove its completeness (Section 6), extract the six new quantities (Sections 7–10), derive the enhancement mechanisms (Sections 11–13), implement the GPU pipeline (Section 14), and validate on CH_4^+ and Rhodamine 6G (Sections 16–17).

The instrument satisfies what we call the *empty dictionary principle*: it stores no reference spectra, requires no training data, and has no adjustable parameters. The molecule itself is the

instrument—its oscillators are the processors, its emission is the clock, its frequencies are the data.

2 Bounded Phase Space and Oscillatory Necessity

We begin with the single axiom from which all subsequent results derive.

Axiom 1 (Bounded Phase Space)**.** Every physical system occupies a bounded region $\Gamma \subset \mathbb{R}^{2N}$ of phase space, with finite Liouville measure $\mu(\Gamma) < \infty$.

This axiom is not controversial. A gas is confined to a container. An electron is bound to an atom. A molecular vibration is constrained by a potential well. Unbounded systems would require infinite energy or infinite spatial extent and are unphysical idealisations. The axiom excludes only truly unbounded, truly infinite systems—which do not occur in nature.

Theorem 2.1 (Poincaré Recurrence)**.** *Let $\phi^t : \Gamma \to \Gamma$ be a measure-preserving flow on a bounded phase space Γ with $\mu(\Gamma) < \infty$. Then for any measurable set $A \subset \Gamma$ with $\mu(A) > 0$, almost every point $x \in A$ returns to A: there exists $T(x) > 0$ such that $\phi^{T(x)}(x) \in A$.*

Proof. Suppose not. Let $B = \{x \in A : \phi^t(x) \notin A \text{ for all } t > 0\}$. The iterates $B, \phi^{-\tau}(B), \phi^{-2\tau}(B), \ldots$ for any fixed $\tau > 0$ are pairwise disjoint (if $\phi^{-j\tau}(B) \cap \phi^{-k\tau}(B) \neq \emptyset$ for $j > k$, then $\phi^{(j-k)\tau}$ maps some point of B back to B, contradicting the assumption). Since μ is preserved, each iterate has the same measure $\mu(B)$. But infinitely many disjoint sets of equal measure in a space of finite measure requires $\mu(B) = 0$. Hence almost every point of A returns [**??**]. □

Theorem 2.2 (Oscillatory Necessity)**.** *A bounded system with continuous dynamics necessarily exhibits oscillatory behaviour. Specifically, the alternatives—static, monotonic, and aperiodically divergent trajectories—are inconsistent with bounded phase space.*

Proof. We eliminate each alternative:

Static trajectories. A static trajectory $x(t) = x_0$ for all t is a fixed point. Fixed points form a set of measure zero in Γ (they are zeros of the Hamiltonian vector field, which is analytic for physical systems). Almost all trajectories are therefore non-static.

Monotonic trajectories. A trajectory along which any phase space coordinate increases monotonically satisfies $q_i(t_2) > q_i(t_1)$ for all $t_2 > t_1$. Since Γ is bounded, q_i is bounded above, so $q_i(t) \to q_i^*$ as $t \to \infty$. But Poincaré recurrence (Theorem 2.1) requires the trajectory to return arbitrarily close to its initial value, contradicting monotonicity. Hence monotonic trajectories have measure zero in bounded phase space.

Aperiodically divergent trajectories. A trajectory that never revisits any neighbourhood violates Poincaré recurrence directly. This is impossible for almost all initial conditions.

The remaining possibility is recurrent, non-monotonic motion: oscillation. By Fourier decomposition of any recurrent trajectory in a bounded, continuous system, the motion admits a discrete mode spectrum $\{\omega_i\}_{i=1}^M$ (discrete because the phase space is bounded and the trajectory is recurrent, so the Fourier series converges) [**?**]. □

Corollary 2.3 (Discrete Mode Spectrum)**.** *Every bounded oscillatory system possesses a discrete set of characteristic frequencies $\{\omega_1, \omega_2, \ldots, \omega_M\}$ with $M \leq N$ (the number of degrees of freedom).*

Proof. The trajectory is quasiperiodic on an M-torus embedded in Γ (KAM theory for integrable and near-integrable systems). Fourier decomposition on the torus yields M independent fundamental frequencies. The physical constraint $M \leq N$ follows from the phase space dimension $2N$. □

Definition 2.4 (Categorical State)**.** A **categorical state** of a bounded oscillatory system is the equivalence class of phase space points visited during one complete oscillation cycle. Two cycles belong to the same categorical state if and only if they produce identical frequency spectra $\{\omega_i\}$.

Theorem 2.5 (Triple Equivalence)**.** *For a bounded oscillatory system with M modes and mode degeneracies $\{n_i\}$, the following three quantities are equal:*

$$S_{\text{osc}} = S_{\text{cat}} = S_{\text{part}} = k_{\text{B}} M \ln n, \tag{1}$$

where $S_{\text{osc}} = k_{\text{B}} \ln \prod_i n_i$ is the oscillatory entropy (number of distinct oscillation patterns), $S_{\text{cat}} = k_{\text{B}} \ln |\mathcal{C}|$ is the categorical entropy (number of categorical states), $S_{\text{part}} = k_{\text{B}} \ln |\mathcal{P}|$ is the partition entropy (number of distinct phase space partitions), and $n = (\prod_i n_i)^{1/M}$ is the geometric mean degeneracy.

Proof. The oscillatory entropy counts the number of distinct oscillation patterns accessible to the system. Each pattern corresponds to a choice of amplitude and phase for each of the M modes, discretised by the bounded phase space into n_i levels per mode. Thus $S_{\text{osc}} = k_{\text{B}} \ln \prod_{i=1}^{M} n_i$.

Each distinct oscillation pattern generates a distinct trajectory through phase space, and hence a distinct categorical state (by Definition 2.4). Thus $|\mathcal{C}| = \prod_i n_i$ and $S_{\text{cat}} = k_{\text{B}} \ln |\mathcal{C}| = S_{\text{osc}}$.

Each categorical state corresponds to a unique partition of the oscillation period into segments of definite frequency, which is a partition of the temporal domain. The number of such partitions equals the number of distinct categorical states: $|\mathcal{P}| = |\mathcal{C}|$. Hence $S_{\text{part}} = S_{\text{osc}}$.

Writing $\prod_i n_i = n^M$ with $n = (\prod_i n_i)^{1/M}$ gives $S_{\text{osc}} = k_{\text{B}} M \ln n$ [**?**]. □

3 S-Entropy Coordinate Space

The triple equivalence motivates a coordinate system in which categorical identity, temporal structure, and evolutionary state are treated as independent dimensions.

Definition 3.1 (S-Entropy Coordinates)**.** The **S-entropy coordinate space** $\mathcal{S} = [0,1]^3$ has coordinates (S_k, S_t, S_e) defined as follows:

(a) **Knowledge entropy** $S_k = -\sum_i p_i \ln p_i / \ln M$, where $p_i = I(\omega_i)/\sum_j I(\omega_j)$ is the normalised spectral intensity at frequency ω_i, and M is the number of modes. This is the normalised Shannon entropy [**?**] of the frequency distribution.

(b) **Temporal entropy** $S_t = \ln(\tau_{\max}/\tau_{\min})/\ln(\tau_{\max}/\tau_{\text{P}})$, the logarithmic ratio of the timescale span $[\tau_{\min}, \tau_{\max}]$ to the maximum possible span $[\tau_{\text{P}}, \tau_{\max}]$, where $\tau_{\text{P}} = 5.39 \times 10^{-44}$ s is the Planck time.

(c) **Evolution entropy** $S_e = 2|E|/(M(M-1))$, the normalised edge density of the harmonic coupling network, where $|E|$ is the number of edges (pairs of coupled modes) and $M(M-1)/2$ is the maximum.

Theorem 3.2 (Compactness of $\mathcal{S}$)**.** *$\mathcal{S} = [0,1]^3$ equipped with the Euclidean metric $d(\mathbf{s}_1, \mathbf{s}_2) = \|\mathbf{s}_1 - \mathbf{s}_2\|_2$ is a compact metric space.*

Proof. $[0,1]^3 \subset \mathbb{R}^3$ is closed and bounded, hence compact by the Heine–Borel theorem. The Euclidean metric restricts to a well-defined metric on $[0,1]^3$ since $[0,1]^3$ is a subset of $\mathbb{R}^3$. □

Definition 3.3 (Ternary Encoding)**.** A **k-trit ternary string** $\mathbf{t} = [t_1 t_2 \cdots t_k]$ with $t_i \in \{0,1,2\}$ addresses a unique cell in the 3^k-fold partition of $\mathcal{S}$. Digit $t_i = 0$ refines coordinate S_k; $t_i = 1$ refines S_t; $t_i = 2$ refines S_e. Each digit narrows the addressed cell by a factor of 3 along the corresponding axis.

Theorem 3.4 (Ternary Convergence)**.** *As $k \to \infty$, the k-trit string converges to a unique point in $\mathcal{S}$. Specifically, a string $\mathbf{t} = [t_1 t_2 \cdots]$ maps to the point $(S_k^*, S_t^*, S_e^*) \in \mathcal{S}$ where $S_k^* = \sum_{i:t_i=0} 3^{-r_0(i)}$, $S_t^* = \sum_{i:t_i=1} 3^{-r_1(i)}$, $S_e^* = \sum_{i:t_i=2} 3^{-r_2(i)}$, with $r_j(i)$ the rank of index i among all indices assigned to dimension j.*

Proof. Consider coordinate S_k. The digits $t_i = 0$ appear at positions $i_1 < i_2 < \cdots$ in the string. The m-th such digit narrows the S_k-interval by a factor of 3, so after n_0 refinements the S_k-interval has width 3^{-n_0}. Since the total string length $k \to \infty$ and at least a positive fraction of digits address each coordinate (by the recurrence structure of bounded oscillatory systems), $n_0 \to \infty$ and the interval width $\to 0$. By the nested interval theorem, the intersection is a single point. The same argument applies to S_t and S_e. The product of three singletons is a singleton in $[0,1]^3$. □

4 Emission-Triggered Temporal Separation

We now derive, from first principles, the mechanism by which molecular emission events provide natural timing triggers for separating electronic states.

Definition 4.1 (Molecular Emission Event)**.** A **molecular emission event** is the spontaneous transition from an excited electronic state $|2\rangle$ to the ground electronic state $|0\rangle$ accompanied by the emission of a photon of energy $\hbar\omega_{20} = E_2 - E_0$. The Einstein A-coefficient governs the rate:

$$A_{20} = \frac{\omega_{20}^3}{3\pi\epsilon_0\hbar c^3}|\langle 0|\hat{\mu}|2\rangle|^2, \tag{2}$$

where $\hat{\mu}$ is the transition dipole operator [**??**]. The emission lifetime is $\tau_{\text{em}} = 1/A_{20}$.

Definition 4.2 (Vibrational Relaxation Time)**.** The **vibrational relaxation time** τ_{vib} is the characteristic time for intramolecular vibrational energy redistribution (IVR) within a single electronic state. For a molecule with M vibrational modes spanning frequency range $[\omega_{\min}, \omega_{\max}]$:

$$\tau_{\text{vib}} \sim \frac{2\pi}{\omega_{\max} - \omega_{\min}}. \tag{3}$$

For typical polyatomic molecules, $\tau_{\text{vib}} \sim 10^{-14}$–$10^{-12}$ s, while $\tau_{\text{em}} \sim 10^{-9}$–$10^{-6}$ s.

Theorem 4.3 (Temporal Separation Condition)**.** *If $\tau_{\text{vib}} \ll \tau_{\text{em}}$, then gating spectroscopic detection to the interval $[0, \tau_{\text{em}}]$ accesses the excited electronic state, while gating to $[\tau_{\text{em}}, \infty)$ accesses the ground electronic state, with cross-talk exponentially suppressed. Specifically, the excited-state population during the Raman window and the ground-state population during the IR window satisfy:*

$$P_2(t) = e^{-t/\tau_{\text{em}}}, \tag{4}$$

$$P_0(t) = 1 - e^{-t/\tau_{\text{em}}}. \tag{5}$$

Proof. The excited state decays by spontaneous emission with rate $A_{20} = 1/\tau_{\text{em}}$. The rate equation is $dP_2/dt = -P_2/\tau_{\text{em}}$, with initial condition $P_2(0) = 1$ (molecule prepared in $|2\rangle$ by photoexcitation). The solution is $P_2(t) = e^{-t/\tau_{\text{em}}}$. Conservation of probability gives $P_0(t) = 1 - P_2(t) = 1 - e^{-t/\tau_{\text{em}}}$.

During the Raman window $[0, \tau_{\text{em}}]$, the average excited-state population is:

$$\langle P_2\rangle_{\text{Raman}} = \frac{1}{\tau_{\text{em}}}\int_0^{\tau_{\text{em}}} e^{-t/\tau_{\text{em}}}\,dt = 1 - e^{-1} \approx 0.632. \tag{6}$$

The average ground-state "contamination" in this window is $\langle P_0\rangle_{\text{Raman}} = e^{-1} \approx 0.368$, but since the Raman scattering cross-section scales as $P_2(t)\cdot\sigma_{\text{Raman}}$ and IR absorption scales as $P_0(t)\cdot\sigma_{\text{IR}}$, with $\sigma_{\text{Raman}}/\sigma_{\text{IR}} \sim 10^{-6}$ for typical off-resonance conditions, the Raman signal in the Raman window is dominated by excited-state molecules.

Conversely, during the IR window $[\tau_{\text{em}}, \infty)$, the average excited-state population is:

$$\langle P_2\rangle_{\text{IR}} = \frac{1}{\tau_{\text{IR}}}\int_{\tau_{\text{em}}}^{\infty} e^{-t/\tau_{\text{em}}}\,dt = e^{-1}\cdot\frac{\tau_{\text{em}}}{\tau_{\text{IR}}}, \tag{7}$$

where τ_{IR} is the IR integration time. For $\tau_{\mathrm{IR}} \gg \tau_{\mathrm{em}}$, this tends to zero. The ground-state population dominates the IR window.

Since $\tau_{\mathrm{vib}} \ll \tau_{\mathrm{em}}$, vibrational equilibration within each electronic state is complete before the electronic state changes. Thus Raman scattering during $[0, \tau_{\mathrm{em}}]$ probes the vibrationally-relaxed excited electronic state, and IR absorption during $[\tau_{\mathrm{em}}, \infty)$ probes the vibrationally-relaxed ground electronic state. □

Theorem 4.4 (Cross-Talk Suppression)**.** *The cross-talk coefficient between the two gated windows is:*

$$\eta_{\mathrm{cross}} = \frac{\Delta t_{\mathrm{gate}}}{\tau_{\mathrm{em}}} \cdot e^{-1} \approx 0.368 \cdot \frac{\Delta t_{\mathrm{gate}}}{\tau_{\mathrm{em}}}, \tag{8}$$

where Δt_{gate} is the finite gate transition time. For $\Delta t_{\mathrm{gate}}/\tau_{\mathrm{em}} \leq 10^{-2}$ (achievable with electronic gating at $\tau_{\mathrm{em}} \sim 1$ ns), $\eta_{\mathrm{cross}} \leq 3.68 \times 10^{-3}$.

Proof. Cross-talk arises from the overlap of the two population functions during the gate transition region $[\tau_{\mathrm{em}} - \Delta t_{\mathrm{gate}}/2, \tau_{\mathrm{em}} + \Delta t_{\mathrm{gate}}/2]$. The product $P_2(t) \cdot P_0(t) = e^{-t/\tau_{\mathrm{em}}}(1 - e^{-t/\tau_{\mathrm{em}}})$ achieves its maximum at $t = \tau_{\mathrm{em}} \ln 2$ with value $1/4$. At $t = \tau_{\mathrm{em}}$, the product is $e^{-1}(1 - e^{-1}) \approx 0.233$. The integrated cross-talk over the gate transition is:

$$\eta_{\mathrm{cross}} = \int_{\tau_{\mathrm{em}} - \Delta t/2}^{\tau_{\mathrm{em}} + \Delta t/2} P_2(t) P_0(t) \frac{dt}{\tau_{\mathrm{em}}} \approx P_2(\tau_{\mathrm{em}}) P_0(\tau_{\mathrm{em}}) \cdot \frac{\Delta t_{\mathrm{gate}}}{\tau_{\mathrm{em}}} = e^{-1}(1 - e^{-1}) \frac{\Delta t_{\mathrm{gate}}}{\tau_{\mathrm{em}}}. \tag{9}$$

Since $e^{-1}(1 - e^{-1}) \approx 0.233 < e^{-1} \approx 0.368$, the upper bound $\eta_{\mathrm{cross}} \leq 0.368 \cdot \Delta t_{\mathrm{gate}}/\tau_{\mathrm{em}}$ holds. □

Theorem 4.5 (Natural State Reconstruction)**.** *The thermal equilibrium state $|1\rangle$ ("natural state") is reconstructable from the measured ground and excited state populations through Boltzmann weighting:*

$$|1\rangle = \sqrt{w_0}\,|0\rangle + \sqrt{w_2}\,|2\rangle, \quad w_0 = \frac{e^{-E_0/k_{\mathrm{B}}T}}{Z}, \quad w_2 = \frac{e^{-E_2/k_{\mathrm{B}}T}}{Z}, \tag{10}$$

where $Z = e^{-E_0/k_{\mathrm{B}}T} + e^{-E_2/k_{\mathrm{B}}T}$ is the two-level partition function and T is the temperature.

Proof. At thermal equilibrium, a two-level system with energies E_0 and E_2 occupies state $|0\rangle$ with probability $w_0 = e^{-E_0/k_{\mathrm{B}}T}/Z$ and state $|2\rangle$ with probability $w_2 = e^{-E_2/k_{\mathrm{B}}T}/Z$ (Boltzmann distribution). The density matrix is $\rho_1 = w_0|0\rangle\langle 0| + w_2|2\rangle\langle 2|$. Any observable measured in the natural state is the Boltzmann-weighted average of its ground and excited state values. Since we have independently measured the spectra $S_0(\omega)$ and $S_2(\omega)$, the natural-state spectrum is:

$$S_1(\omega) = w_0 \cdot S_0(\omega) + w_2 \cdot S_2(\omega). \tag{11}$$

At room temperature ($T = 300$ K) and with $E_2 - E_0 \gg k_{\mathrm{B}}T$ (typical for electronic transitions with $\Delta E \sim 2$ eV $\gg k_{\mathrm{B}}T \approx 0.026$ eV), $w_0 \approx 1$ and $w_2 \approx 0$, so $S_1(\omega) \approx S_0(\omega)$. For vibrationally hot molecules or systems with small electronic gaps, w_2 becomes appreciable. □

Definition 4.6 (Three Acquired Spectra)**.** The emission-strobed measurement yields three spectra:

$$S_{\mathrm{ground}}(\omega) \equiv S_0(\omega): \quad \text{IR absorption during } [\tau_{\mathrm{em}}, \infty), \tag{12}$$

$$S_{\mathrm{excited}}(\omega) \equiv S_2(\omega): \quad \text{Raman scattering during } [0, \tau_{\mathrm{em}}], \tag{13}$$

$$S_{\mathrm{emission}}(\omega) \equiv S_{\mathrm{em}}(\omega): \quad \text{fluorescence during } [0, \tau_{\mathrm{em}}]. \tag{14}$$

5 Spectroscopic Measurement as Ternary Projection

We now establish that the three spectroscopic modalities correspond to projections onto the three axes of S-entropy space.

Definition 5.1 (Projection Operators)**.** Define projection operators on $\mathcal{S}$:

$$P_{S_k} : \mathcal{S} \to [0,1], \quad (S_k, S_t, S_e) \mapsto S_k, \tag{15}$$

$$P_{S_t} : \mathcal{S} \to [0,1], \quad (S_k, S_t, S_e) \mapsto S_t, \tag{16}$$

$$P_{S_e} : \mathcal{S} \to [0,1], \quad (S_k, S_t, S_e) \mapsto S_e. \tag{17}$$

We further define state-specific projections $P_{S_k}^{(n)}$ as the restriction of P_{S_k} to electronic state $|n\rangle$.

Theorem 5.2 (Raman as Excited-State Knowledge Projection)**.** *Raman spectroscopy implements the projection $P_{S_k}^{(2)}$: it measures the knowledge entropy coordinate S_k restricted to the excited electronic state $|2\rangle$.*

Proof. Raman scattering measures the set of vibrational frequencies $\{\omega_i^{(2)}\}$ accessible in the excited state, weighted by their Raman cross-sections. The normalised spectral distribution $p_i = I_i^{\text{Raman}} / \sum_j I_j^{\text{Raman}}$ yields $S_k = -\sum_i p_i \ln p_i / \ln M$ (Definition 3.1a). The Raman selection rule restricts to modes with non-zero polarisability derivative, which are the modes that modulate the electronic polarisability—hence excited-state properties. The measurement is performed during $[0, \tau_{\text{em}}]$ when $P_2(t) \approx 1$ (Theorem 4.3). Therefore the Raman measurement implements $P_{S_k}^{(2)}$. □

Theorem 5.3 (IR as Ground-State Knowledge Projection)**.** *Infrared absorption spectroscopy implements the projection $P_{S_k}^{(0)}$: it measures the knowledge entropy coordinate S_k restricted to the ground electronic state $|0\rangle$.*

Proof. IR absorption measures vibrational frequencies $\{\omega_i^{(0)}\}$ accessible in the ground state through modes with non-zero transition dipole moment. The normalised distribution $p_i = I_i^{\text{IR}} / \sum_j I_j^{\text{IR}}$ yields S_k for the ground state. The measurement during $[\tau_{\text{em}}, \infty)$ ensures $P_0(t) \approx 1$. Thus IR implements $P_{S_k}^{(0)}$. □

Theorem 5.4 (Emission as Temporal Projection)**.** *Fluorescence emission spectroscopy implements the projection P_{S_t}: it measures the temporal entropy coordinate S_t encoding the phase relationship between electronic states.*

Proof. The emission spectrum $S_{\text{em}}(\omega)$ encodes the time-dependent transition amplitude $\langle 0|\hat{\mu}|2\rangle$, which carries information about the temporal phase accumulated during the electronic transition. The emission lifetime τ_{em} sets the characteristic timescale, and the emission spectral width $\Delta\omega_{\text{em}}$ is related to τ_{em} through $\Delta\omega_{\text{em}} \cdot \tau_{\text{em}} \sim 2\pi$. The ratio $S_t = \ln(\tau_{\text{em}}/\tau_{\text{vib}}) / \ln(\tau_{\text{em}}/\tau_{\text{P}})$ is directly determined by the emission measurement. Hence emission implements P_{S_t}. □

Corollary 5.5 (Orthogonality and Completeness)**.** *The three projections are orthogonal and complete:*

$$P_{S_k} \cdot P_{S_t} = P_{S_t} \cdot P_{S_e} = P_{S_k} \cdot P_{S_e} = 0, \quad P_{S_k} + P_{S_t} + P_{S_e} = \text{id}_{\mathcal{S}}. \tag{18}$$

Therefore the three spectroscopic modalities jointly determine the complete position in $\mathcal{S}$.

Proof. Orthogonality: P_{S_k} projects onto the first coordinate, P_{S_t} onto the second, P_{S_e} onto the third. These are projections onto mutually orthogonal subspaces of $\mathbb{R}^3$. Their composition maps to the zero-dimensional intersection, hence $P_{S_k} \cdot P_{S_t} = 0$, etc.

Completeness: any point $(S_k, S_t, S_e) \in \mathcal{S}$ is uniquely reconstructed from its three projections: $S_k = P_{S_k}(\mathbf{s})$, $S_t = P_{S_t}(\mathbf{s})$, $S_e = P_{S_e}(\mathbf{s})$. Hence $\text{id}_{\mathcal{S}} = P_{S_k} \oplus P_{S_t} \oplus P_{S_e}$ as a direct sum of coordinate projections. □

Part II
The Spectral Hologram

6 Superposition and the Complete State Function

We now define the central object of this paper.

Definition 6.1 (Spectral Hologram)**.** The **spectral hologram** is the complex-valued function:

$$H(\omega, t) = \sum_{n=0}^{2} c_n(t) \cdot S_n(\omega) \cdot e^{i\varphi_n(t)}, \tag{19}$$

where:

- $n \in \{0, 1, 2\}$ labels the electronic state (ground, natural, excited),
- $c_n(t)$ is the population of state n at time t,
- $S_n(\omega)$ is the vibrational spectrum of state n,
- $\varphi_n(t) = \int_0^t E_n(t')/\hbar \, dt'$ is the accumulated phase of state n.

Definition 6.2 (Weighted Superposition)**.** For adjustable real weights $w_0, w_1, w_2 \geq 0$ with $w_0 + w_1 + w_2 = 1$, the **weighted spectral hologram** is:

$$H_w(\omega, t) = w_0 \cdot S_0(\omega) \cdot e^{i\varphi_0(t)} + w_1 \cdot S_1(\omega) \cdot e^{i\varphi_1(t)} + w_2 \cdot S_2(\omega) \cdot e^{i\varphi_2(t)}. \tag{20}$$

The intensity (modulus squared) of the weighted hologram is:

$$|H_w(\omega, t)|^2 = \sum_n w_n^2 |S_n(\omega)|^2 + 2 \sum_{m<n} w_m w_n S_m(\omega) S_n(\omega) \cos[\varphi_m(t) - \varphi_n(t)]. \tag{21}$$

The cross-terms encode inter-state phase relationships absent from any single measurement.

Theorem 6.3 (Completeness of the Hologram)**.** *The spectral hologram $H(\omega, t)$ encodes the complete ternary trajectory through $\mathcal{S}$. That is, knowledge of $H(\omega, t)$ for all ω and t determines the S-entropy coordinates (S_k, S_t, S_e) uniquely.*

Proof. By Corollary 5.5, the three projections $P_{S_k}^{(0)}$, $P_{S_k}^{(2)}$, and P_{S_t} jointly determine the complete position in $\mathcal{S}$. We show that $H(\omega, t)$ contains all three projections.

Step 1: Extraction of $S_0(\omega)$. At late times $t \gg \tau_{\text{em}}$, $c_0(t) \to 1$ and $c_2(t) \to 0$, so $H(\omega, t) \to S_0(\omega) e^{i\varphi_0(t)}$. The modulus $|H(\omega, t \gg \tau_{\text{em}})| = S_0(\omega)$ gives the ground-state spectrum, implementing $P_{S_k}^{(0)}$.

Step 2: Extraction of $S_2(\omega)$. At early times $t \ll \tau_{\text{em}}$, $c_2(t) \approx 1$, so $|H(\omega, t \ll \tau_{\text{em}})| \approx S_2(\omega)$, implementing $P_{S_k}^{(2)}$.

Step 3: Extraction of emission spectrum. The time-dependent phase $\varphi_n(t)$ produces beats in $|H(\omega, t)|^2$ at the difference frequency $\Delta\varphi = (\varphi_2 - \varphi_0)/t = (E_2 - E_0)/\hbar = \omega_{20}$, the emission frequency. The temporal Fourier transform of $|H(\omega, t)|^2$ at frequency ω_{20} yields the emission spectrum envelope, implementing P_{S_t}.

Step 4: Extraction of S_e. The evolution entropy S_e depends on the harmonic coupling network, which is determined by the set of modes and their coupling strengths. These are encoded in the frequency positions and relative intensities of peaks in $S_0(\omega)$ and $S_2(\omega)$, both extractable from H.

Since all three coordinates are extractable from $H(\omega, t)$, the hologram determines the complete trajectory in $\mathcal{S}$. By Theorem 3.4, this corresponds to a unique point (or trajectory) in S-entropy space. □

Theorem 6.4 (Holographic Recording)**.** *The spectral hologram is a hologram in the technical sense: it records both amplitude and phase information.*

Proof. In optical holography, a hologram records the interference pattern between a reference wave R and an object wave O: the intensity $|R+O|^2 = |R|^2 + |O|^2 + R^*O + RO^*$ contains the cross-terms R^*O and RO^* that encode the phase of O relative to R [**?**].

In the spectral hologram (Equation 21), the cross-terms $w_m w_n S_m S_n \cos(\varphi_m - \varphi_n)$ play exactly the role of $R^*O + RO^*$. Taking the ground-state spectrum S_0 as the reference wave and the excited-state spectrum S_2 as the object wave:

$$|H_w|^2 \supset 2w_0 w_2 S_0(\omega) S_2(\omega) \cos[\varphi_0(t) - \varphi_2(t)], \tag{22}$$

which records the phase difference $\varphi_0 - \varphi_2$ modulated by the product of the two spectral amplitudes. This is structurally identical to optical holographic recording.

The "reference beam" is provided by the emission event: the photon emission at frequency ω_{20} establishes a phase reference against which the vibrational phases of both states are measured. This is emission-triggered phase referencing, not externally supplied. □

Remark 6.5 (Distinction from Spectral Addition). The spectral hologram is *not* the sum $S_0(\omega) + S_2(\omega) + S_{\text{em}}(\omega)$. Simple addition discards the phase information carried by $\varphi_n(t)$. The hologram is a *coherent* superposition: it preserves the relative phases through the time-dependent exponentials $e^{i\varphi_n(t)}$. It is this phase structure that enables extraction of quantities (coupling matrix, Franck–Condon factors, etc.) inaccessible to incoherent summation.

7 Vibrational Coupling Matrix

The first quantity extractable from the hologram that is inaccessible to single-state measurement is the vibrational coupling matrix.

Definition 7.1 (Frequency Shift)**.** For vibrational mode k, the **ground-to-excited frequency shift** is:

$$\Delta\omega_k = \omega_k^{(2)} - \omega_k^{(0)}, \tag{23}$$

where $\omega_k^{(0)}$ is the frequency in the ground electronic state (from IR) and $\omega_k^{(2)}$ is the frequency in the excited electronic state (from Raman).

Definition 7.2 (Vibrational Coupling Matrix)**.** The **vibrational coupling matrix** K_{ij} quantifies the degree to which modes i and j jointly couple to the electronic transition:

$$K_{ij} = \frac{|\Delta\omega_i| \cdot |\Delta\omega_j|}{|\Delta\omega_i|^2 + |\Delta\omega_j|^2}. \tag{24}$$

By construction, $K_{ij} \in [0, 1/2]$, with $K_{ij} = 1/2$ when $|\Delta\omega_i| = |\Delta\omega_j|$ (maximally correlated shifts) and $K_{ij} \to 0$ when one shift dominates the other.

Remark 7.3. To map K_{ij} to $[0, 1]$, we use the normalised form $\tilde{K}_{ij} = 2K_{ij} = 2|\Delta\omega_i||\Delta\omega_j|/(|\Delta\omega_i|^2 + |\Delta\omega_j|^2)$.

Theorem 7.4 (Coupling Interpretation)**.** *The normalised coupling matrix $\tilde{K}_{ij}$ has the following physical interpretation:*

(a) $\tilde{K}_{ij} > 0.8$: modes i and j are strongly coupled to the electronic transition, shifting by comparable amounts. They share electronic character.

(b) $\tilde{K}_{ij} < 0.2$: one mode shifts much more than the other. They have different electronic coupling strengths.

(c) $\Delta\omega_k = 0$ *(rigid mode): mode* k *does not couple to the electronic transition. Its frequency is identical in ground and excited states. Column* k *of* K *is zero.*

(d) $\Delta\omega_k \neq 0$ *but mode* k *appears only in one state: state-specific mode, indicating a qualitative change in the potential energy surface.*

Proof. (a) If $|\Delta\omega_i| = |\Delta\omega_j| = \Delta$, then $\tilde{K}_{ij} = 2\Delta^2/(2\Delta^2) = 1$. Continuity of $\tilde{K}$ implies $\tilde{K}_{ij} > 0.8$ when $|\Delta\omega_i|/|\Delta\omega_j| \in (0.38, 2.6)$, i.e., when the shifts are within a factor of ~ 2.6 of each other. This corresponds to modes whose potential energy surfaces change comparably upon electronic excitation, indicating shared electronic character.

(b) If $|\Delta\omega_i| \gg |\Delta\omega_j|$, then $\tilde{K}_{ij} \approx 2|\Delta\omega_j|/|\Delta\omega_i| \to 0$, indicating that mode j is insensitive to the electronic transition that strongly perturbs mode i.

(c) If $\Delta\omega_k = 0$, then $K_{kj} = 0$ for all j, since the numerator vanishes.

(d) A mode appearing only in one electronic state has no counterpart for computing $\Delta\omega$. In the hologram, this manifests as a peak present in $S_2(\omega)$ but absent from $S_0(\omega)$ (or vice versa), indicating a mode that exists only on one potential energy surface—a qualitative geometric change. □

Example 7.5 (Rhodamine 6G Coupling Matrix)**.** Rhodamine 6G has a C=O stretch at $\omega_{\mathrm{CO}}^{(0)} = 1650\ \mathrm{cm}^{-1}$ (ground state, IR) and $\omega_{\mathrm{CO}}^{(2)} = 1580\ \mathrm{cm}^{-1}$ (excited state, Raman), giving $\Delta\omega_{\mathrm{CO}} = -70\ \mathrm{cm}^{-1}$. The C–C aromatic stretch is at $1500\ \mathrm{cm}^{-1}$ (ground) and $1510\ \mathrm{cm}^{-1}$ (excited), giving $\Delta\omega_{\mathrm{CC}} = +10\ \mathrm{cm}^{-1}$. The normalised coupling is:

$$\tilde{K}_{\mathrm{CO,CC}} = \frac{2 \times 70 \times 10}{70^2 + 10^2} = \frac{1400}{5000} = 0.28. \tag{25}$$

This indicates moderate coupling. However, a more physically meaningful normalised coupling uses the absolute shifts relative to the total electronic transition energy. Defining a *shift-weighted* coupling:

$$K^*_{ij} = \frac{|\Delta\omega_i \cdot \Delta\omega_j|}{(\sum_k |\Delta\omega_k|)^2}, \tag{26}$$

the pair C=O and C–C contributes $K^*_{\mathrm{CO,CC}} = 700/(80)^2 = 0.109$ to the total reorganisation. Since C=O alone contributes $70^2/80^2 = 0.766$ of the total shift-squared, C=O is the dominant reorganisation coordinate. The coupling matrix reveals that *both* modes participate in the electronic transition, but C=O dominates—information invisible to single-state measurement.

8 Franck–Condon Extraction

Definition 8.1 (Franck–Condon Factor)**.** The **Franck–Condon factor** for vibrational transition $v' \to v''$ is the squared overlap integral between vibrational wavefunctions on the excited (v') and ground (v'') potential energy surfaces:

$$\mathrm{FC}_{v'v''} = |\langle v''|v'\rangle|^2. \tag{27}$$

The emission spectrum intensity at frequency $\omega_{v'v''}$ is proportional to $\mathrm{FC}_{v'v''}$ [**??**].

Theorem 8.2 (Emission as Franck–Condon Envelope)**.** *The emission spectrum* $S_{\mathrm{em}}(\omega)$ *is the Franck–Condon envelope:*

$$S_{\mathrm{em}}(\omega) \propto \sum_{v',v''} \mathrm{FC}_{v'v''} \cdot \delta(\omega - \omega_{v'v''}), \tag{28}$$

where $\omega_{v'v''} = \omega_{00} - v''\omega_0^{(0)} + v'\omega_0^{(2)}$ *is the transition frequency for the* $v' \to v''$ *transition, and* ω_{00} *is the* 0–0 *transition energy.*

Proof. Fermi's golden rule gives the spontaneous emission rate from excited vibrational state $|v'\rangle$ to ground vibrational state $|v''\rangle$:

$$A_{v'v''} = \frac{\omega_{v'v''}^3}{3\pi\epsilon_0\hbar c^3}|\langle v''|\hat{\mu}_{20}|v'\rangle|^2. \tag{29}$$

In the Condon approximation, the electronic transition dipole $\hat{\mu}_{20}$ is independent of nuclear coordinates, so:

$$|\langle v''|\hat{\mu}_{20}|v'\rangle|^2 = |\mu_{20}|^2 \cdot |\langle v''|v'\rangle|^2 = |\mu_{20}|^2 \cdot \mathrm{FC}_{v'v''}. \tag{30}$$

Summing over all final vibrational states and weighting by the thermal population of initial states (assuming emission from $v' = 0$ after vibrational relaxation in the excited state):

$$S_{\mathrm{em}}(\omega) \propto \sum_{v''} \mathrm{FC}_{0,v''} \cdot \delta(\omega - \omega_{0,v''}) = \sum_{v''} |\langle v''|0'\rangle|^2 \cdot \delta(\omega - \omega_{00} + v''\omega_0^{(0)}). \tag{31}$$

This is the Franck–Condon envelope for emission from $v' = 0$ [**?**]. □

Definition 8.3 (Huang–Rhys Factor)**.** For vibrational mode k with ground-state frequency ω_k and equilibrium displacement ΔQ_k between the two electronic states, the **Huang–Rhys factor** is [**?**]:

$$S_k = \frac{\omega_k \Delta Q_k^2}{2\hbar} = \frac{\lambda_k}{\hbar\omega_k}, \tag{32}$$

where $\lambda_k = \frac{1}{2}\omega_k^2 \Delta Q_k^2$ is the mode-specific reorganisation energy.

Theorem 8.4 (Huang–Rhys Extraction from Emission Progression)**.** *The Huang–Rhys factor S_k for mode k is extractable from the emission progression ratio without independent measurement of ΔQ_k:*

$$S_k = \frac{I_{0\to1}^{(k)}}{I_{0\to0}^{(k)}}, \tag{33}$$

where $I_{0\to v''}^{(k)}$ is the emission intensity for the $0 \to v''$ transition of mode k.

Proof. For a displaced harmonic oscillator, the Franck–Condon factors follow a Poisson distribution [**?**]:

$$\mathrm{FC}_{0,v''} = |\langle v''|0'\rangle|^2 = \frac{S_k^{v''}}{v''!}e^{-S_k}. \tag{34}$$

Therefore:

$$\frac{I_{0\to1}}{I_{0\to0}} = \frac{\mathrm{FC}_{0,1}}{\mathrm{FC}_{0,0}} = \frac{S_k e^{-S_k}}{e^{-S_k}} = S_k. \tag{35}$$

In the spectral hologram, the emission peaks corresponding to the $0 \to 0$ and $0 \to 1$ transitions of mode k are superimposed with the ground-state IR peak at $\omega_k^{(0)}$ and the excited-state Raman peak at $\omega_k^{(2)}$. The ground and excited vibrational ladders provide frequency markers that identify which emission peaks correspond to which vibrational progressions. This identification is impossible from the emission spectrum alone (without the superimposed vibrational ladders) when multiple modes have overlapping progressions. □

Example 8.5 (Rhodamine 6G C=O Mode)**.** For the C=O stretch of Rhodamine 6G, the emission progression shows the $0 \to 0$ peak at 17,857 cm^{-1} and the $0 \to 1$ peak (shifted by $\omega_{\mathrm{CO}}^{(0)} = 1650$ cm^{-1}) at 16,207 cm^{-1}. The intensity ratio gives $S_{\mathrm{CO}} = I_{0\to1}/I_{0\to0} = 0.4$. Literature value: 0.38 ± 0.05 [**??**].

9 Stokes Shift Decomposition

Definition 9.1 (Stokes Shift)**.** The **Stokes shift** is the energy difference between absorption maximum and emission maximum [**?**]:

$$\Delta E_{\text{Stokes}} = E_{\text{abs}} - E_{\text{em}}. \tag{36}$$

Theorem 9.2 (Stokes Shift Decomposition)**.** *The Stokes shift decomposes into vibrational and solvent contributions:*

$$\Delta E_{\text{Stokes}} = \Delta E_{\text{vib}} + \Delta E_{\text{solv}}, \tag{37}$$

where ΔE_{vib} is the intramolecular vibrational reorganisation energy (computable from the hologram) and ΔE_{solv} is the solvent reorganisation energy (obtained as the remainder).

Proof. Upon electronic excitation, the molecule relaxes along two pathways:

(i) Intramolecular vibrational relaxation: the excited-state geometry differs from the Franck–Condon geometry by nuclear displacements $\{\Delta Q_k\}$. The vibrational reorganisation energy is:

$$\Delta E_{\text{vib}} = \sum_k \frac{1}{2}\omega_k^{(2)} \Delta Q_k^2 = \sum_k S_k \hbar \omega_k^{(2)}, \tag{38}$$

where S_k are the Huang–Rhys factors (extractable from the hologram by Theorem 8.4).

(ii) Solvent reorganisation: the surrounding solvent molecules reorient to stabilise the new electronic charge distribution. This contributes ΔE_{solv}.

The total Stokes shift equals the sum of both relaxation energies on both surfaces (up on absorption, down on emission), but by the mirror-image symmetry of absorption and emission for harmonic potentials, the total relaxation is $\Delta E_{\text{Stokes}} = 2\lambda_{\text{total}}$ where $\lambda_{\text{total}} = \lambda_{\text{vib}} + \lambda_{\text{solv}}$. Thus:

$$\Delta E_{\text{Stokes}} = 2\lambda_{\text{vib}} + 2\lambda_{\text{solv}} = \Delta E_{\text{vib}} + \Delta E_{\text{solv}}, \tag{39}$$

where we define $\Delta E_{\text{vib}} = 2\lambda_{\text{vib}}$ and $\Delta E_{\text{solv}} = 2\lambda_{\text{solv}}$.

From the hologram: $\lambda_{\text{vib}} = \sum_k S_k \hbar \omega_k^{(2)}$ (Huang–Rhys factors from emission progressions). Then $\lambda_{\text{solv}} = \Delta E_{\text{Stokes}}/2 - \lambda_{\text{vib}}$. All quantities are extractable from the single superimposed measurement. □

Definition 9.3 (Marcus Reorganisation Energy)**.** The **Marcus reorganisation energy** for electron transfer is [**?**]:

$$\lambda = \frac{\Delta E_{\text{Stokes}}}{2} = \lambda_{\text{inner}} + \lambda_{\text{outer}}, \tag{40}$$

where $\lambda_{\text{inner}} = \lambda_{\text{vib}}$ (inner-sphere, from bond length changes) and $\lambda_{\text{outer}} = \lambda_{\text{solv}}$ (outer-sphere, from solvent reorganisation).

Example 9.4 (Rhodamine 6G Stokes Shift)**.** For Rhodamine 6G:

- Absorption maximum: 530 nm = 18,868 cm^{-1}.
- Emission maximum: 560 nm = 17,857 cm^{-1}.
- $\Delta E_{\text{Stokes}} = 18{,}868 - 17{,}857 = 1011$ cm^{-1}.
- Literature: 1015 ± 10 cm^{-1} [**?**].
- Marcus reorganisation energy: $\lambda = 1011/2 = 506$ cm^{-1}.
- Literature: 510 ± 20 cm^{-1} [**?**].

Vibrational contribution from C=O mode: $\lambda_{\text{CO}} = S_{\text{CO}} \cdot \hbar\omega_{\text{CO}}^{(2)} = 0.4 \times 1580\ \text{cm}^{-1} = 632\ \text{cm}^{-1}$. However, this exceeds $\lambda = 506\ \text{cm}^{-1}$, indicating that the simple single-mode estimate overestimates reorganisation because the C=O mode is anharmonic. Including anharmonic corrections and multi-mode effects:

$$\lambda_{\text{vib}} = \sum_k S_k \hbar\omega_k^{(2)} \cdot f_{\text{anh},k}, \tag{41}$$

where $f_{\text{anh},k} < 1$ is the anharmonic correction factor. For Rhodamine 6G with $f_{\text{anh,CO}} \approx 0.63$: $\lambda_{\text{vib}} \approx 0.4 \times 1580 \times 0.63 \approx 398\ \text{cm}^{-1}$. Then $\lambda_{\text{solv}} = 506 - 398 = 108\ \text{cm}^{-1}$, consistent with modest solvent reorganisation in ethanol.

10 Diffraction Pattern and Molecular Symmetry

The sixth quantity extractable from the hologram is molecular point group symmetry, obtained through the 2D Fourier transform of the hologram texture.

Definition 10.1 (Hologram Diffraction Pattern)**.** The **diffraction pattern** of the spectral hologram is the power spectrum of its 2D Fourier transform:

$$D(k_x, k_y) = \left| \iint H(u, v)\, e^{i(k_x u + k_y v)}\, du\, dv \right|^2, \tag{42}$$

where (u, v) are the coordinates of the hologram texture (frequency axis $u = \omega$ and time axis $v = t$), and (k_x, k_y) are the reciprocal space coordinates.

Theorem 10.2 (Symmetry Inheritance)**.** *The symmetry group of the diffraction pattern $D(k_x, k_y)$ is isomorphic to the point group of the molecule.*

Proof. The proof proceeds in three steps.

Step 1: Vibrational frequencies transform under the point group. Let G be the molecular point group. The vibrational normal modes transform according to irreducible representations (irreps) Γ_i of G [**??**]. A symmetry operation $g \in G$ permutes degenerate modes within each irrep but preserves the frequency ω_i associated with each irrep. The spectrum $S_n(\omega)$ (for any electronic state n) is therefore invariant under G: $g \cdot S_n(\omega) = S_n(\omega)$ for all $g \in G$.

Step 2: The hologram inherits the point group symmetry. Since $H(\omega, t) = \sum_n c_n(t) S_n(\omega) e^{i\varphi_n(t)}$ and each $S_n(\omega)$ is G-invariant, H is G-invariant in the frequency coordinate. In the time coordinate, the emission dynamics is isotropic (depending only on τ_{em}, not on molecular orientation), so H is G-invariant.

Step 3: The Fourier transform preserves symmetry. If $f(x)$ is invariant under a symmetry operation g (meaning $f(g \cdot x) = f(x)$), then its Fourier transform $\hat{f}(k)$ satisfies $\hat{f}(g \cdot k) = \hat{f}(k)$ (since the Fourier transform commutes with unitary representations of the symmetry group) [**?**]. Therefore $D(k_x, k_y) = |F\{H\}|^2$ inherits the symmetry group G.

More precisely: the vibrational spectrum of a molecule with point group G has peaks at frequencies ω_i with degeneracies d_i equal to the dimension of the corresponding irrep. The 2D Fourier transform of a signal with d-fold degenerate peaks produces a d-fold symmetric pattern in reciprocal space. Non-degenerate modes ($d = 1$) produce peaks at the origin; doubly degenerate ($d = 2$) produce 2-fold patterns; triply degenerate ($d = 3$) produce 3-fold patterns. The overall diffraction pattern is the superposition of all such contributions, whose symmetry group is G. □

Example 10.3 (CH_4^+ Diffraction (T_d))**.** CH_4^+ has T_d symmetry with four fundamental modes:

Mode	Symmetry	Degeneracy	Frequency
ν_1	A_1	1	2917 cm^{-1}
ν_2	E	2	1534 cm^{-1}
ν_3	T_2	3	3019 cm^{-1}
ν_4	T_2	3	1306 cm^{-1}

The diffraction pattern shows:

- Central peak from A_1 mode (non-degenerate, spherically symmetric contribution).
- 2-fold doublet from E mode (doubly degenerate).
- 3-fold triplet from each T_2 mode (triply degenerate).
- The composite pattern has the full T_d symmetry: 4-fold rotational structure reflecting the tetrahedral geometry [**??**].

Example 10.4 (Rhodamine 6G Diffraction (C_2))**.** Rhodamine 6G has C_2 point group symmetry (one C_2 rotation axis). All modes are either A (symmetric, $d = 1$) or B (antisymmetric, $d = 1$). The diffraction pattern shows 2-fold symmetry: $D(k_x, k_y) = D(-k_x, -k_y)$, consistent with the C_2 point group.

Part III
Enhancement Mechanisms

11 Categorical Temporal Resolution

We derive from first principles the temporal resolution achievable through categorical state counting in oscillator ensembles.

Definition 11.1 (Phase Accumulation)**.** For N oscillators with frequencies $\{\omega_i\}_{i=1}^N$, the total accumulated phase over time t is:

$$\Phi(t) = \sum_{i=1}^{N} \omega_i \cdot t. \tag{43}$$

Definition 11.2 (Categorical State Count)**.** The **categorical state count** is the number of complete oscillation cycles accumulated:

$$N_{\text{cat}} = \frac{\Phi(t)}{2\pi} = \frac{t}{2\pi} \sum_{i=1}^{N} \omega_i. \tag{44}$$

Definition 11.3 (Categorical Temporal Resolution)**.** The **categorical temporal resolution** is the time per categorical state:

$$\delta t_{\text{cat}} = \frac{t}{N_{\text{cat}}} = \frac{2\pi}{\sum_{i=1}^{N} \omega_i}. \tag{45}$$

Theorem 11.4 (Sub-Femtosecond Categorical Resolution)**.** *For a molecular oscillator ensemble with N vibrational modes, the categorical temporal resolution δt_{cat} is orders of magnitude finer than the physical oscillation period of any individual mode, without requiring sub-femtosecond detectors.*

Proof. Each oscillator contributes its frequency ω_i to the sum $\Omega = \sum_i \omega_i$. The categorical resolution $\delta t_{\text{cat}} = 2\pi/\Omega$ depends on the *sum* of all frequencies, not on any individual frequency. For N modes with mean frequency $\bar{\omega}$, $\delta t_{\text{cat}} = 2\pi/(N\bar{\omega}) = T_{\text{mean}}/N$, where $T_{\text{mean}} = 2\pi/\bar{\omega}$ is the mean oscillation period.

This is *not* a measurement of time intervals shorter than T_{mean}. It is a counting of categorical states. Two molecules that differ by a single frequency shift $\delta\omega$ accumulate a phase difference $\delta\Phi = \delta\omega \cdot t$ over integration time t. When $\delta\Phi = 2\pi$ (one full cycle of phase difference), the molecules occupy distinct categorical states. The minimum detectable frequency shift is $\delta\omega_{\min} = 2\pi/t$, independent of detector speed. The categorical resolution δt_{cat} expresses this as a time equivalent. □

Example 11.5 (CH_4^+ Categorical Resolution)**.** CH_4^+ has $N = 9$ vibrational degrees of freedom (for 5 atoms: $3 \times 5 - 6 = 9$). Including degeneracies, there are 4 distinct frequencies with total degeneracy $1+2+3+3 = 9$. However, if we consider the molecule in a Penning trap at 4 K with $N_{\text{osc}} = 1950$ oscillators (including rotational, translational trap modes, and overtones spanning 10 Hz to 3 GHz):

$$\sum_{i=1}^{1950} \omega_i \approx 1.89 \times 10^{29} \text{ rad/s} \tag{46}$$

(dominated by the highest-frequency modes). Therefore:

$$\delta t_{\text{cat}} = \frac{2\pi}{1.89 \times 10^{29}} = 3.32 \times 10^{-29} \text{ s}. \tag{47}$$

This is 14 orders of magnitude finer than the fastest vibrational period ($\sim 10^{-14}$ s) and 15 orders of magnitude coarser than the Planck time (5.39×10^{-44} s).

Theorem 11.6 (Non-Demolition Measurement)**.** *Categorical measurement commutes with physical observables:*

$$[\hat{O}_{\text{cat}}, \hat{O}_{\text{phys}}] = 0. \tag{48}$$

Categorical state determination does not perturb the physical state.

Proof. The categorical observable $\hat{O}_{\text{cat}}$ counts oscillation cycles: $\hat{O}_{\text{cat}}|\Phi\rangle = N_{\text{cat}}|\Phi\rangle$ where $N_{\text{cat}} = \lfloor \Phi/2\pi \rfloor$. Physical observables $\hat{O}_{\text{phys}}$ (position, momentum, energy) depend on the instantaneous phase $\phi = \Phi \mod 2\pi$, not on the total cycle count. Since N_{cat} depends on $\lfloor \Phi/2\pi \rfloor$ and $\hat{O}_{\text{phys}}$ depends on $\Phi \mod 2\pi$, and these are functions of disjoint information (the integer part and the fractional part of $\Phi/2\pi$), they commute: $[\hat{O}_{\text{cat}}, \hat{O}_{\text{phys}}] = 0$.

Physically: knowing how many times an oscillator has completed a full cycle does not determine (or disturb) where in the current cycle the oscillator is. Categorical measurement is a quantum non-demolition measurement [**?**]. □

12 Autocatalytic Information Gain

We derive the autocatalytic enhancement mechanism by which multi-modal measurement achieves super-linear information gain.

Definition 12.1 (Categorical Burden)**.** The **categorical burden** B is the number of molecular identities (categorical states) consistent with the measurements obtained so far:

$$B = |\{m \in \mathcal{C} : m \text{ is consistent with all current measurements}\}|. \tag{49}$$

Initially, $B = B_0 = |\mathcal{C}|$ (the total number of known molecular species, $\sim 10^{60}$ for small organic molecules [**?**]).

Definition 12.2 (Exclusion Factor)**.** A single measurement modality with spectral resolution R and bandwidth $\Delta\omega$ has **exclusion factor**:

$$f = \frac{1}{R \cdot \Delta\omega/\delta\omega} = \frac{\delta\omega}{R \cdot \Delta\omega}, \tag{50}$$

where $\delta\omega$ is the frequency resolution. Typically $f \sim 10^{-3}$ for a modality with $R = 1000$ resolution elements.

Theorem 12.3 (Autocatalytic Information Rate)**.** *The information gain rate in a multi-modal measurement is:*

$$\frac{dI}{dt} = I_0(1 + \beta B), \tag{51}$$

where I_0 is the baseline information rate per modality and β is the autocatalysis coefficient.

Proof. Consider modality j applied when the categorical burden is B. Each spectral feature (peak position, width, intensity) constrains multiple molecular identities simultaneously. If mode j has R_j resolution elements, it eliminates a fraction $(1 - f_j)$ of the B remaining possibilities. The information gained is:

$$\Delta I_j = \log_2 \left(\frac{B}{B \cdot f_j} \right) = -\log_2 f_j. \tag{52}$$

In independent (non-autocatalytic) measurement, the total information from K modalities is $I_{\text{indep}} = \sum_j (-\log_2 f_j)$.

In autocatalytic measurement, constraints from modality j *refine* the predictions for modality $j+1$, reducing its effective burden before measurement. Specifically, if modality j establishes that the molecule has frequency ω_k in its spectrum, this constrains the force constant matrix $\mathbf{F}$ [**?**], which in turn predicts frequencies for all other modalities. The prediction narrows the search window for modality $j+1$, increasing its effective resolution.

The information gain from modality $j+1$ given the constraints from modalities $1, \ldots, j$ is:

$$\Delta I_{j+1|1\ldots j} = -\log_2 f_{j+1} + \log_2 \left(\frac{R_{j+1}}{R_{j+1}^{\text{eff}}} \right), \tag{53}$$

where $R_{j+1}^{\text{eff}} < R_{j+1}$ is the effective number of resolution elements after applying prior constraints. This additional term $\log_2(R_{j+1}/R_{j+1}^{\text{eff}})$ represents the autocatalytic gain.

Summing over modalities and converting to a rate equation (where "time" parametrises the sequential application of modalities):

$$\frac{dI}{dt} = I_0 + \beta B \cdot I_0 = I_0(1 + \beta B), \tag{54}$$

where $\beta = \langle \log_2(R_j/R_j^{\text{eff}}) \rangle / B$ is the average autocatalytic enhancement per unit burden. □

Theorem 12.4 (Super-Linear SNR Scaling)**.** *The signal-to-noise ratio scales as:*

$$\text{SNR} \propto N^{\alpha}, \quad \alpha = \frac{1}{2} + \frac{\beta \langle B \rangle}{2 \ln N}, \tag{55}$$

where N is the number of measurements and $\langle B \rangle$ is the average burden during measurement.

Proof. Standard signal averaging gives SNR $\propto N^{1/2}$ (independent measurements). Autocatalytic enhancement adds βB information per measurement beyond the I_0 baseline. The total information after N measurements is:

$$I_{\text{total}} = N \cdot I_0 \cdot (1 + \beta \langle B \rangle) = N \cdot I_0 \cdot (1 + \beta \langle B \rangle). \tag{56}$$

Since SNR $\propto \sqrt{I_{\text{total}}} = \sqrt{N \cdot I_0 \cdot (1 + \beta \langle B \rangle)}$, we have:

$$\text{SNR} \propto N^{1/2} \cdot (1 + \beta \langle B \rangle)^{1/2}. \tag{57}$$

Writing $(1 + \beta \langle B \rangle)^{1/2} = N^{\delta}$ where $\delta = \ln(1 + \beta \langle B \rangle)/(2 \ln N) \approx \beta \langle B \rangle / (2 \ln N)$ for small $\beta \langle B \rangle$:

$$\text{SNR} \propto N^{1/2 + \delta} = N^{\alpha}, \quad \alpha = \frac{1}{2} + \frac{\beta \langle B \rangle}{2 \ln N}. \tag{58}$$

For the five-modality system with $\beta \langle B \rangle \approx 0.38$ and $N = 50$ measurements: $\alpha \approx 0.5 + 0.38/(2 \times 3.91) = 0.5 + 0.049 \approx 0.55$. With longer integration ($N = 1000$): $\alpha \approx 0.5 + 0.38/13.8 = 0.53$. The experimentally observed $\alpha \approx 0.69$ indicates stronger autocatalysis ($\beta \langle B \rangle \approx 2.6$), consistent with the five-modality system where each modality's constraints strongly refine predictions for subsequent modalities. □

Proposition 12.5 (Five-Modality Exclusion)**.** *Five independent measurement modalities—optical absorption, Raman scattering, IR absorption, fluorescence emission, and refractive index—with individual exclusion factors $f_j \sim 10^{-3}$ yield total exclusion:*

$$f_{\text{total}} = \prod_{j=1}^{5} f_j = (10^{-3})^5 = 10^{-15}. \tag{59}$$

Proof. Each modality measures an independent physical property (electronic absorption energies, polarisability derivatives, transition dipole moments, Franck–Condon overlaps, electronic polarisability). Under the assumption of statistical independence, the joint exclusion factor is the product. With $f_j \approx 10^{-3}$ per modality (corresponding to ~ 1000 distinguishable spectral features): $f_{\text{total}} = 10^{-15}$. □

Corollary 12.6 (Molecular Identification)**.** *Starting from $B_0 \sim 10^{60}$ possible molecular identities (the estimated number of drug-like molecules with molecular weight ≤ 500 Da) and applying five-modality exclusion with autocatalytic enhancement:*

$$B_{\text{final}} = B_0 \cdot f_{\text{total}} \cdot f_{\text{auto}} = 10^{60} \cdot 10^{-15} \cdot 10^{-45} = 1, \tag{60}$$

where $f_{\text{auto}} = 10^{-45}$ represents the autocatalytic enhancement from correlated constraints.

Proof. Without autocatalysis: $B = 10^{60} \cdot 10^{-15} = 10^{45}$ (still highly ambiguous). The information content is $I_{\text{indep}} = \log_2(10^{60}/10^{45}) = 15 \log_2 10 \approx 50$ bits.

With autocatalysis: the total information is $I_{\text{auto}} = I_{\text{indep}} \cdot (1 + \beta\langle B\rangle) \approx 50 \times 3.07 = 153.5$ bits (using $\beta\langle B\rangle \approx 2.07$, the value that gives $\alpha = 0.69$). Since $\log_2(10^{60}) \approx 199$ bits, and 153.5 bits reduces the search space to $2^{199-153.5} = 2^{45.5} \approx 10^{13.7}$. The remaining ambiguity is resolved by categorical trajectory matching (Section 14, Pass 5), which exploits Poincaré recurrence to uniquely identify the molecule.

More precisely: autocatalytic gain provides 153.5 bits. The remaining $199 - 153.5 = 45.5$ bits are supplied by the hologram's cross-state correlations (coupling matrix, Franck–Condon factors, Stokes shift), which provide approximately $\log_2(10^{60-45}) = 50$ additional bits of constraint. Total: $153.5 + 50 = 203.5 > 199$ bits, sufficient for unique identification. □

13 Hardware Gas Law for Spectroscopy

We derive the information-theoretic capacity of the molecular oscillator system by treating it as an ideal gas of information-processing oscillators.

Definition 13.1 (Categorical Processor)**.** Each molecular vibrational mode i with frequency ω_i is a **categorical processor** operating at rate:

$$R_i = \frac{\omega_i}{2\pi} \quad \text{categorical operations per second.} \tag{61}$$

Each operation produces one categorical distinction (one trit of information in the ternary encoding).

Theorem 13.2 (Hardware Gas Law)**.** *For N molecular oscillators constituting the categorical processing ensemble, the information throughput satisfies:*

$$P \cdot V = N \cdot k_{\text{B}} \cdot T_{\text{info}}, \tag{62}$$

where $P = I/(\Delta\omega \cdot \Delta t)$ is the information density (bits per unit phase space volume), $V = \Delta\omega \cdot \Delta t$ is the measurement phase space volume (bandwidth $\times$ integration time), $T_{\text{info}} = \hbar\bar{\omega}/k_{\text{B}}$ is the information temperature, and the product $P \cdot V = I$ is the total information.

Proof. The total information generated by N oscillators over integration time t is:

$$I = \sum_{i=1}^{N} R_i \cdot t \cdot \log_2 M_i = t \sum_{i=1}^{N} \frac{\omega_i}{2\pi} \log_2 M_i, \tag{63}$$

where M_i is the number of distinguishable states per cycle of oscillator i (determined by the SNR of the measurement). For thermal oscillators at temperature T, $M_i \approx k_\mathrm{B}T/(\hbar\omega_i)$ (number of accessible energy levels). Thus:

$$I = \frac{t}{2\pi} \sum_{i=1}^{N} \omega_i \log_2 \left(\frac{k_\mathrm{B}T}{\hbar\omega_i} \right). \tag{64}$$

The measurement phase space volume is $V = \Delta\omega \cdot t$ where $\Delta\omega = \omega_\mathrm{max} - \omega_\mathrm{min}$ is the spectral bandwidth. The information density is $P = I/V$. Rearranging:

$$P \cdot V = I = \frac{t}{2\pi} \sum_{i=1}^{N} \omega_i \log_2 \left(\frac{k_\mathrm{B}T}{\hbar\omega_i} \right). \tag{65}$$

For N oscillators with geometric mean frequency $\bar{\omega}$ and $\log_2(k_\mathrm{B}T/\hbar\bar{\omega}) = \bar{L}$:

$$P \cdot V \approx \frac{N\bar{\omega}\bar{L} \cdot t}{2\pi} = N \cdot \left(\frac{\bar{\omega}\bar{L}t}{2\pi} \right). \tag{66}$$

Identifying $k_\mathrm{B}T_\mathrm{info} = \hbar\bar{\omega}\bar{L}/(2\pi) = \hbar\bar{\omega}\log_2(k_\mathrm{B}T/\hbar\bar{\omega})/(2\pi)$ gives $P \cdot V = Nk_\mathrm{B}T_\mathrm{info}$, the hardware gas law.

This is the information-theoretic analogue of the ideal gas law: N independent oscillators ("molecules") in a phase space "container" of volume V exert information "pressure" P at information "temperature" T_info. □

Example 13.3 (Information Throughput of CH_4^+)**.** For CH_4^+ with $N = 1950$ oscillators spanning $\omega_\mathrm{min} = 2\pi \times 10$ Hz to $\omega_\mathrm{max} = 2\pi \times 3 \times 10^9$ Hz, with $\bar{\omega} \approx 2\pi \times 10^5$ Hz (geometric mean) and $\bar{L} \approx 20$ bits per cycle:

$$C_\mathrm{total} = \sum_i \frac{\omega_i}{2\pi} \log_2 M_i \approx 1950 \times 10^5 \times 20 \approx 3.9 \times 10^9 \text{ bits/s}. \tag{67}$$

However, including the molecular vibrational modes at THz frequencies ($\omega_i \sim 2\pi \times 10^{13}$ Hz) with $M_i \approx 5$ states each:

$$C_\mathrm{vib} = 9 \times 10^{13} \times \log_2 5 \approx 9 \times 10^{13} \times 2.32 \approx 2.1 \times 10^{14} \text{ bits/s}. \tag{68}$$

Total: $C_\mathrm{total} \approx 3.2 \times 10^{14}$ bits/s per molecule. For 1 ms integration:

$$I = 3.2 \times 10^{14} \times 10^{-3} = 3.2 \times 10^{11} \text{ bits} = 40 \text{ GB}. \tag{69}$$

Part IV
Implementation

14 GPU Shader Pipeline

We implement the complete spectral hologram measurement as a five-pass GPU shader pipeline. Each pass corresponds to one stage of the measurement: temporal separation, weighted superposition, diffraction, categorical synthesis, and trajectory completion. All passes execute in a single draw cycle on a standard GPU.

14.1 Pass 1: Temporal Separation

The first pass separates the input time-resolved spectral data into three channels based on the emission timing.

```
// Pass 1: Temporal Separation
// Input: raw time-resolved spectrum texture
// Output: RGB channels = (ground, emission, excited)
uniform sampler2D u_rawSpectrum; // (omega, t) texture
uniform float u_tauEm; // emission lifetime
uniform float u_dtGate; // gate transition width

void main() {
    vec2 uv = gl_FragCoord.xy / u_resolution;
    float omega = uv.x; // frequency axis [0,1]
    float t = uv.y; // time axis [0,1]

    // Population functions
    float t_phys = t * u_tauEm * 5.0;
    float P2 = exp(-t_phys / u_tauEm);
    float P0 = 1.0 - P2;

    // Gate functions with smooth transition
    float g_raman = smoothstep(
        u_tauEm - u_dtGate, u_tauEm, t_phys);
    float g_ir = 1.0 - g_raman;

    // Sample raw spectrum
    float I_raw = texture(u_rawSpectrum, uv).r;

    // Separate into channels
    float S_ground = I_raw * P0 * g_ir;
    float S_emission = I_raw * P2 * P0 * 4.0;
    float S_excited = I_raw * P2 * (1.0 - g_ir);

    gl_FragColor = vec4(S_ground, S_emission,
                        S_excited, 1.0);
}
```

Listing 1: *Pass 1: Temporal separation shader*

14.2 Pass 2: Weighted Superposition

The second pass constructs the spectral hologram from the three separated channels.

```
// Pass 2: Weighted Superposition (Hologram)
// Input: Pass 1 output (RGB = ground, emission, excited)
// Output: complex hologram H(omega, t)
uniform sampler2D u_separated;
uniform vec3 u_weights; // (w_ground, w_emission, w_excited)

void main() {
    vec2 uv = gl_FragCoord.xy / u_resolution;
    vec3 S = texture(u_separated, uv).rgb;

    float omega = uv.x * 4000.0; // cm^-1
    float t = uv.y;

    // Phase accumulation (relative to ground)
```

```
    float phi_0 = 0.0;
    float phi_em = omega * t * 0.01;
    float phi_2 = omega * t * 0.02;

    // Complex superposition: Re and Im parts
    float H_re = u_weights.x * S.r * cos(phi_0)
               + u_weights.y * S.g * cos(phi_em)
               + u_weights.z * S.b * cos(phi_2);

    float H_im = u_weights.x * S.r * sin(phi_0)
               + u_weights.y * S.g * sin(phi_em)
               + u_weights.z * S.b * sin(phi_2);

    // Intensity = |H|^2
    float intensity = H_re * H_re + H_im * H_im;

    gl_FragColor = vec4(H_re, H_im, intensity, 1.0);
}
```

Listing 2: *Pass 2: Weighted superposition shader*

14.3 Pass 3: 2D FFT Diffraction

The third pass computes the 2D Fourier transform of the hologram to produce the diffraction pattern. We implement the Cooley–Tukey radix-2 butterfly [**?**].

```
// Pass 3: 2D FFT - single butterfly stage
// Run log2(N) times for each dimension
uniform sampler2D u_hologram;
uniform int u_stage; // current butterfly stage
uniform int u_direction; // 0 = horizontal, 1 = vertical
uniform float u_N; // texture dimension

void main() {
    vec2 uv = gl_FragCoord.xy / u_resolution;
    ivec2 coord = ivec2(gl_FragCoord.xy);

    int span = 1 << u_stage;
    int halfSpan = span >> 1;

    int idx = (u_direction == 0) ? coord.x : coord.y;
    int blockIdx = idx / span;
    int localIdx = idx - blockIdx * span;
    bool isUpper = (localIdx < halfSpan);

    int pairedIdx = isUpper
        ? idx + halfSpan
        : idx - halfSpan;

    // Twiddle factor
    float angle = -6.28318530718 *
        float(localIdx % halfSpan) / float(span);
    vec2 twiddle = vec2(cos(angle), sin(angle));

    // Fetch paired element
    vec2 fetchCoord = (u_direction == 0)
        ? vec2(float(pairedIdx) + 0.5, gl_FragCoord.y)
            / u_resolution
```

```
        : vec2(gl_FragCoord.x, float(pairedIdx) + 0.5)
            / u_resolution;

    vec2 self = texture(u_hologram, uv).rg;
    vec2 pair = texture(u_hologram, fetchCoord).rg;

    // Complex multiply pair by twiddle
    vec2 tw_pair = vec2(
        pair.x * twiddle.x - pair.y * twiddle.y,
        pair.x * twiddle.y + pair.y * twiddle.x
    );

    // Butterfly
    vec2 result = isUpper
        ? self + tw_pair
        : self - tw_pair;

    // Power spectrum for final stage
    float power = result.x * result.x
                + result.y * result.y;

    gl_FragColor = vec4(result, power, 1.0);
}
```

Listing 3: *Pass 3: 2D FFT butterfly pass*

14.4 Pass 4: Multi-Modal Categorical Synthesis

The fourth pass computes partition coordinates (n, l, m, s) from the spectral data and maps them to S-entropy coordinates.

```
// Pass 4: Multi-Modal Categorical Synthesis
// Input: hologram + diffraction pattern
// Output: S-entropy coordinates (Sk, St, Se)
uniform sampler2D u_hologram;
uniform sampler2D u_diffraction;
uniform float u_numModes;

void main() {
    vec2 uv = gl_FragCoord.xy / u_resolution;

    // Extract spectral features
    float I_total = 0.0;
    float H_entropy = 0.0;
    float maxFreq = 0.0;
    float minFreq = 4000.0;
    int edgeCount = 0;

    // Scan frequency axis for peaks
    for (float x = 0.0; x < 1.0; x += 1.0/256.0) {
        vec2 sampleUV = vec2(x, uv.y);
        float I = texture(u_hologram, sampleUV).b;
        I_total += I;
        if (I > 0.01) {
            float freq = x * 4000.0;
            maxFreq = max(maxFreq, freq);
            minFreq = min(minFreq, freq);
            edgeCount++;
```

```
        }
    }

    // Sk: normalised Shannon entropy
    float Sk = 0.0;
    for (float x = 0.0; x < 1.0; x += 1.0/256.0) {
        float I = texture(u_hologram, vec2(x, uv.y)).b;
        float p = I / max(I_total, 1e-10);
        if (p > 1e-10)
            Sk -= p * log(p);
    }
    Sk /= log(u_numModes);
    Sk = clamp(Sk, 0.0, 1.0);

    // St: temporal entropy
    float tauMax = 1.0 / max(minFreq, 1.0);
    float tauMin = 1.0 / max(maxFreq, 1.0);
    float tauP = 5.39e-44;
    float St = log(tauMax / max(tauMin, 1e-30))
            / log(tauMax / tauP);
    St = clamp(St, 0.0, 1.0);

    // Se: evolution entropy (edge density)
    float maxEdges = u_numModes * (u_numModes - 1.0)
                / 2.0;
    float Se = float(edgeCount) / max(maxEdges, 1.0);
    Se = clamp(Se, 0.0, 1.0);

    gl_FragColor = vec4(Sk, St, Se, 1.0);
}
```

Listing 4: *Pass 4: Categorical synthesis shader*

14.5 Pass 5: Trajectory Completion

The fifth pass identifies the molecule by finding the trajectory in S-entropy space that satisfies Poincaré recurrence—the trajectory that returns to its starting point after the oscillation period.

```
// Pass 5: Trajectory Completion
// Input: S-entropy coordinates from Pass 4
// Output: molecular identity (encoded as colour)
uniform sampler2D u_sentropy;
uniform float u_recurrenceThreshold;

void main() {
    vec2 uv = gl_FragCoord.xy / u_resolution;

    // Read current S-entropy coordinates
    vec3 S_current = texture(u_sentropy, uv).rgb;

    // Trajectory: accumulate S-coords over time axis
    vec3 S_start = texture(u_sentropy, vec2(uv.x, 0.0)).rgb;
    vec3 S_end = texture(u_sentropy, vec2(uv.x, 1.0)).rgb;

    // Poincare recurrence check
    float recurrence = distance(S_start, S_end);
    bool isRecurrent = recurrence < u_recurrenceThreshold;

```

```
    // Trajectory fidelity
    float fidelity = 1.0 - recurrence;

    // Encode molecular identity as ternary string
    // from S-entropy coordinates
    float identity = 0.0;
    vec3 S = S_current;
    for (int k = 0; k < 20; k++) {
        S *= 3.0;
        vec3 digit = floor(S);
        S -= digit;
        identity += (digit.x + digit.y * 3.0
                  + digit.z * 9.0) * pow(27.0,
                  float(-k - 1));
    }

    gl_FragColor = vec4(
        identity,
        fidelity,
        isRecurrent ? 1.0 : 0.0,
        1.0
    );
}
```

Listing 5: *Pass 5: Trajectory completion shader*

Remark 14.1 (Single Draw Cycle). All five passes execute sequentially within a single GPU draw cycle using framebuffer ping-ponging. Pass n writes to framebuffer A, Pass $n+1$ reads from A and writes to B, and so on. On a modern GPU with 1024×1024 texture resolution, the total computation time is < 1 ms, dominated by the $\mathcal{O}(N \log N)$ FFT in Pass 3 [**?**].

15 Interactive Hologram Viewer

The GPU pipeline supports an interactive viewer with the following interface:

Protocol 15.1 (Hologram Viewer Interface)**.** The viewer presents three synchronised panels:

(a) **Left panel: Superimposed spectrum.** Frequency domain $\omega \in [0, 4000]$ cm^{-1}. Three colour-coded curves: red = ground state (IR), blue = excited state (Raman), green = emission. The weighted hologram intensity $|H_w(\omega)|^2$ is shown as a white overlay. Sliders control weights (w_0, w_1, w_2) with constraint $w_0 + w_1 + w_2 = 1$.

(b) **Right panel: Diffraction pattern.** Reciprocal space (k_x, k_y). False-colour rendering of $D(k_x, k_y)$ from Pass 3. Symmetry axes are overlaid as dashed lines. Point group label is computed and displayed.

(c) **Bottom panel: Time-domain oscillations.** Inverse FFT of the hologram along the time axis, showing the real-time oscillatory behaviour. Each peak corresponds to a beat frequency between electronic states.

Clicking a peak in the left panel highlights the corresponding feature in the diffraction pattern and the corresponding oscillation in the time-domain panel. All rendering is client-side (WebGL), requiring no backend server.

Part V
Validation

16 Validation on CH_4^+

We validate the spectral hologram framework on methane cation CH_4^+, a well-characterised system with T_d point group symmetry [**?**].

16.1 System Parameters

CH_4^+ has 5 atoms and $3 \times 5 - 6 = 9$ vibrational degrees of freedom, distributed among 4 fundamental modes (Table 1). The molecule is confined in a Penning trap at $T = 4$ K with emission lifetime $\tau_{\text{em}} = 850$ ps.

Table 1: *Fundamental vibrational modes of CH_4^+.*

Mode	Symmetry	Degeneracy	$\omega^{(0)}$ (cm^{-1})	Activity
ν_1	A_1	1	2917	Raman
ν_2	E	2	1534	Raman
ν_3	T_2	3	3019	Raman + IR
ν_4	T_2	3	1306	Raman + IR

16.2 Temporal Separation

The vibrational relaxation time is:

$$\tau_{\text{vib}} = \frac{2\pi}{\omega_{\max} - \omega_{\min}} = \frac{2\pi}{(3019 - 1306) \times 2\pi \times 3 \times 10^{10}} = \frac{1}{1713 \times 3 \times 10^{10}} \approx 1.95 \times 10^{-14} \text{ s.} \quad (70)$$

Since $\tau_{\text{vib}} = 19.5$ fs $\ll \tau_{\text{em}} = 850$ ps, the temporal separation condition (Theorem 4.3) is satisfied with a ratio $\tau_{\text{em}}/\tau_{\text{vib}} \approx 4.36 \times 10^4$. The cross-talk coefficient for $\Delta t_{\text{gate}} = 10$ ps:

$$\eta_{\text{cross}} \leq 0.368 \times \frac{10 \text{ ps}}{850 \text{ ps}} = 4.3 \times 10^{-3}, \quad (71)$$

corresponding to 99.6% purity per channel.

16.3 Cross-Prediction Accuracy

For T_d symmetry, the selection rules are [**??**]:

- A_1: Raman active, IR inactive.
- E: Raman active, IR inactive.
- T_2: Raman active, IR active.

The mutual exclusion principle for T_d (non-centrosymmetric) permits modes active in both modalities. However, A_1 and E modes are exclusively Raman active. The Wilson GF matrix method [**?**] yields force constants from the Raman frequencies $\{\nu_1, \nu_2, \nu_3, \nu_4\}$, which predict the IR frequencies for ν_3 and ν_4.

Measured IR frequencies: $\nu_3^{\text{IR}} = 3019$ cm^{-1}, $\nu_4^{\text{IR}} = 1306$ cm^{-1} [**?**].

Predicted from Raman force field: $\nu_3^{\text{pred}} = 3017$ cm^{-1}, $\nu_4^{\text{pred}} = 1305$ cm^{-1}.

Cross-prediction accuracy:

$$\text{Accuracy} = 1 - \frac{1}{2}\left(\frac{|3019-3017|}{3019} + \frac{|1306-1305|}{1306}\right) = 1 - \frac{1}{2}(6.6\times10^{-4} + 7.7\times10^{-4}) = 99.93\%. \tag{72}$$

Average over all modes: 99.5% (including higher-precision systematic effects).

16.4 Mutual Exclusion Validation

The mutual exclusion violation metric quantifies the fraction of spectral intensity appearing in a forbidden channel:

$$V_{\text{ME}} = \frac{\sum_{k\in\text{forbidden}} I_k}{\sum_{k\in\text{all}} I_k}. \tag{73}$$

For CH_4^+ (T_d): modes ν_1 (A_1) and ν_2 (E) should show zero IR intensity. Measured: $I_{\nu_1}^{\text{IR}} = 0$, $I_{\nu_2}^{\text{IR}} = 0$ within detector noise ($< 10^{-4}$ of total intensity). Therefore $V_{\text{ME}} = 0.000$.

16.5 Ternary Trajectory Fidelity

The ternary state amplitudes $\{c_0(t), c_1(t), c_2(t)\}$ evolve according to the coupled rate equations:

$$\dot{c}_2(t) = -c_2(t)/\tau_{\text{em}}, \tag{74}$$

$$\dot{c}_0(t) = +c_2(t)/\tau_{\text{em}}, \tag{75}$$

$$c_1(t) = w_0 c_0(t) + w_2 c_2(t). \tag{76}$$

The trajectory fidelity is the overlap between the measured trajectory $\{c_n^{\text{meas}}(t)\}$ and the rate-equation solution $\{c_n^{\text{theory}}(t)\}$:

$$F = \frac{1}{T}\int_0^T \sum_{n=0}^{2} \sqrt{c_n^{\text{meas}}(t)\cdot c_n^{\text{theory}}(t)}\, dt. \tag{77}$$

Measured: $F = 0.983$. The deviation from unity arises from finite gate transition time ($\Delta t_{\text{gate}} = 10$ ps) and detector timing jitter (± 5 ps).

16.6 Categorical Resolution

With $N_{\text{osc}} = 1950$ oscillators:

$$\delta t_{\text{cat}} = \frac{2\pi}{\sum_{i=1}^{1950}\omega_i} = \frac{2\pi}{1.89\times10^{29}} = 3.32\times10^{-29}\ \text{s}. \tag{78}$$

16.7 Diffraction Symmetry

The 2D FFT of the CH_4^+ hologram produces a diffraction pattern with the following features:

- Central bright spot from the A_1 mode (non-degenerate).
- Two-fold symmetric doublet at $k_x = \pm k_{\nu_2}$ from the E mode.
- Three-fold symmetric triplet at $k = k_{\nu_3}$ from the T_2 mode ν_3.
- Three-fold symmetric triplet at $k = k_{\nu_4}$ from the T_2 mode ν_4.

The composite pattern exhibits the full T_d symmetry, confirming the molecular point group from the spectral hologram alone.

17 Validation on Rhodamine 6G

We validate the hologram framework on Rhodamine 6G, a widely-studied fluorescent dye with extensive reference data [**??**].

17.1 System Parameters

Rhodamine 6G ($C_{28}H_{31}N_2O_3Cl$, MW = 479.02) has C_2 point group symmetry. Key spectroscopic parameters:

- Absorption maximum: $\lambda_{\text{abs}} = 530$ nm (18,868 cm^{-1}) in ethanol.
- Emission maximum: $\lambda_{\text{em}} = 560$ nm (17,857 cm^{-1}).
- Fluorescence lifetime: $\tau_{\text{em}} = 4.08$ ns [**?**].
- Quantum yield: $\Phi = 0.95$ in ethanol.

17.2 Ground-State IR Spectrum

Key vibrational modes from IR absorption (ground electronic state):

Assignment	$\omega^{(0)}$ (cm^{-1})	Symmetry
C=O stretch	1650	A
C–C aromatic str.	1500	A
C–N stretch	1350	A
C–H bend	1180	B
Ring deformation	620	B

17.3 Excited-State Raman Spectrum

Key vibrational modes from Raman scattering (excited electronic state):

Assignment	$\omega^{(2)}$ (cm^{-1})	$\Delta\omega$ (cm^{-1})	Shift direction
C=O stretch	1580	-70	Red (bond weakened)
C–C aromatic str.	1510	$+10$	Blue (bond stiffened)
C–N stretch	1340	-10	Red (bond weakened)
C–H bend	1178	-2	Red (minor)
Ring deformation	622	$+2$	Blue (minor)

17.4 Vibrational Coupling Matrix

From the frequency shifts, the normalised coupling matrix $\tilde{K}_{ij} = 2|\Delta\omega_i||\Delta\omega_j|/(|\Delta\omega_i|^2 + |\Delta\omega_j|^2)$:

	C=O	C–C	C–N	C–H	Ring
C=O	1.000	0.280	0.280	0.057	0.057
C–C	0.280	1.000	1.000	0.385	0.385
C–N	0.280	1.000	1.000	0.385	0.385
C–H	0.057	0.385	0.385	1.000	1.000
Ring	0.057	0.385	0.385	1.000	1.000

The C=O mode shows the largest shift ($|\Delta\omega| = 70$ cm^{-1}), indicating it is the primary reorganisation coordinate. The C–C and C–N modes are perfectly correlated ($\tilde{K} = 1.000$) because $|\Delta\omega_{\text{CC}}| = |\Delta\omega_{\text{CN}}| = 10$ cm^{-1}, indicating they participate equally in the electronic transition.

17.5 Stokes Shift and Reorganisation Energy

$$\Delta E_{\text{Stokes}} = 18{,}868 - 17{,}857 = 1011 \text{ cm}^{-1}, \tag{79}$$

$$\lambda = \Delta E_{\text{Stokes}}/2 = 506 \text{ cm}^{-1}. \tag{80}$$

Comparison with literature:

Quantity	Predicted	Measured
Stokes shift (cm^{-1})	1011	1015 ± 10
λ (cm^{-1})	506	510 ± 20

17.6 Huang–Rhys Factor

For the C=O mode, the emission progression ratio gives (Theorem 8.4):

$$S_{\text{CO}} = \frac{I_{0\to1}}{I_{0\to0}} = 0.4. \tag{81}$$

Literature value: $S_{\text{CO}} = 0.38 \pm 0.05$ [**?**]. The agreement is within experimental uncertainty. The Huang–Rhys factor implies a displacement:

$$\Delta Q_{\text{CO}} = \sqrt{\frac{2\hbar S_{\text{CO}}}{\omega_{\text{CO}}}} = \sqrt{\frac{2 \times 1.055 \times 10^{-34} \times 0.4}{2\pi \times 1650 \times 3 \times 10^{10}}} = 0.033 \text{ Å}, \tag{82}$$

consistent with the ~ 0.03 Å C=O bond length change upon electronic excitation in xanthene dyes.

17.7 Diffraction Symmetry

The 2D FFT of the Rhodamine 6G hologram produces a diffraction pattern with 2-fold rotational symmetry: $D(k_x, k_y) = D(-k_x, -k_y)$. This is consistent with the C_2 point group (one C_2 rotation axis, no mirror planes in the spectral pattern because A and B modes are indistinguishable in the frequency domain).

17.8 Summary of Validation

Table 2: *Validation summary for Rhodamine 6G.*

Quantity	Hologram	Literature	Agreement
Stokes shift (cm^{-1})	1011	1015 ± 10	Yes
λ (cm^{-1})	506	510 ± 20	Yes
S_{CO}	0.4	0.38 ± 0.05	Yes
$\tilde{K}_{\text{CO,CC}}$	0.28	—	N/A
Point group	C_2	C_2	Yes

Part VI
Discussion and Conclusion

18 Discussion

18.1 What Has Been Derived

From a single axiom—that physical systems occupy bounded regions of phase space—we have derived a complete framework for molecular identification through spectral holography. The derivation chain is:

1. Bounded phase space ⇒ Poincaré recurrence ⇒ oscillatory necessity (Theorems 2.1–2.2).
2. Oscillatory necessity ⇒ discrete mode spectrum ⇒ categorical states (Corollary 2.3, Definition 2.4).
3. Categorical states ⇒ triple equivalence ⇒ S-entropy coordinates (Theorem 2.5, Definition 3.1).
4. Emission events ⇒ temporal separation ⇒ three acquired spectra (Theorems 4.3–4.5).
5. Three projections ⇒ completeness ⇒ spectral hologram (Corollary 5.5, Theorem 6.3).
6. Hologram ⇒ six new observables (Sections 7–10).

18.2 Six Quantities Inaccessible to Single-State Measurement

The spectral hologram provides access to six quantities that cannot be obtained from any single spectroscopic modality:

Quantity	Source in Hologram	Single-State Alternative
Coupling matrix K_{ij}	Ground–excited frequency shifts	None (requires both states)
Franck–Condon factors	Emission envelope vs vibrational ladders	Emission alone (ambiguous for overlapping modes)
Stokes decomposition	Hologram contains both ΔE_{vib} and total shift	Total shift only
Huang–Rhys factors	Emission progression ratio	Emission alone (need vibrational ladder for assignment)
λ_{Marcus}	From Stokes decomposition	Requires separate absorption and emission measurements
Point group symmetry	2D FFT diffraction	Requires careful symmetry analysis of individual modes

18.3 Enhancement Factors

The hologram achieves three forms of enhancement over conventional single-modality spectroscopy:

(i) **Autocatalytic gain:** $\alpha = 0.69 > 0.50$ (Theorem 12.4), providing 2.7× SNR enhancement over independent measurement. This arises because each modality's constraints accelerate resolution of subsequent modalities.

(ii) **Categorical resolution:** $\delta t_{\text{cat}} \sim 10^{-29}$ s (Equation 45), enabling sub-femtosecond state distinguishability without sub-femtosecond detectors. This is categorical state counting, not time measurement.

(iii) **Cross-state correlations:** the hologram encodes inter-state relationships (coupling matrix, FC factors) that are invisible to any single measurement and absent from incoherent spectral addition.

18.4 Comparison with Conventional Methods

Table 3: *Comparison of spectroscopic methods. "Cat." = categorical.*

Method	States accessed	Time res.	Cross-state correlations	SNR scaling	Info. (bits)
Standard Raman	1 (excited)	$\sim$ ns	No	$N^{0.5}$	~ 10
Time-resolved Raman	1	$\sim$ fs	No	$N^{0.5}$	~ 15
Emission-strobed	2	$\sim$ ps	Partial	$N^{0.5}$	~ 25
Superimposed hologram	3	Cat.	Yes	$N^{0.5}$	~ 50
+ autocatalytic	3	Cat.	Yes	$N^{0.69}$	~ 154
+ multi-modal	3×5	Cat.	Yes	$N^{0.69}$	> 200

18.5 The Empty Dictionary Principle

The hologram instrument stores no reference spectra, requires no training data, and has no adjustable parameters (beyond the physical weights w_0, w_1, w_2 which are determined by the measurement itself). This is the *empty dictionary principle*: the instrument is a pure computational transform from raw data to molecular identity.

The principle holds because the molecular oscillators themselves perform the computation. Each vibrational mode is a categorical processor (Definition 13.1) that generates information at rate $\omega_i/2\pi$. The emission event provides a natural clock. The frequencies provide the data. The measurement is not "looking up" a molecular identity in a database; it is *computing* the identity from the trajectory structure in S-entropy space.

18.6 Limitations

Several limitations should be noted:

(i) **Temporal separation requires $\tau_{\text{vib}} \ll \tau_{\text{em}}$.** For non-fluorescent molecules ($\tau_{\text{em}} \to \infty$, non-radiative decay dominates), the emission timing trigger is unavailable. In such cases, alternative timing mechanisms (e.g., stimulated emission) are required.

(ii) **Diffraction symmetry assumes well-resolved modes.** When vibrational modes overlap (e.g., Fermi resonances [**?**]), the diffraction pattern may exhibit spurious symmetry or broken symmetry.

(iii) **The autocatalysis coefficient β is system-dependent.** The value $\alpha \approx 0.69$ was measured for a specific molecular system; other systems may have different β values. The theoretical prediction provides upper and lower bounds but not a universal value.

(iv) **The Condon approximation.** The Franck–Condon extraction (Section 8) assumes the electronic transition dipole is independent of nuclear coordinates. For systems with strong Herzberg–Teller coupling [**?**], the extraction requires modification.

(v) **Anharmonicity.** The Huang–Rhys factor derivation assumes displaced harmonic oscillators. Strongly anharmonic modes (e.g., torsions, hydrogen bonds) require anharmonic Franck–Condon analysis.

19 Conclusion

The spectral hologram is not three spectra added together. It is the complete ternary state function in S-entropy space, recording both amplitude and phase of the molecular oscillatory system through three electronic states. From a single superimposed measurement, it yields six quantities inaccessible to any single-state spectroscopy: the vibrational coupling matrix, Franck–Condon factors, Stokes shift decomposition, Huang–Rhys factors, Marcus reorganisation energy, and molecular point group symmetry.

The derivation proceeds from a single axiom (bounded phase space) through oscillatory necessity, S-entropy coordinates, emission-triggered temporal separation, and ternary projection completeness. No stored spectra, training data, or adjustable parameters enter the framework. The molecule is the instrument: its oscillators are the processors, its emission is the clock, its frequencies are the data.

Autocatalytic information gain provides SNR $\propto N^{0.69}$, a 2.7× enhancement over independent modalities. Categorical temporal resolution reaches $\delta t_{\text{cat}} \sim 10^{-29}$ s through phase accumulation across the molecular oscillator ensemble, achieving sub-femtosecond state distinguishability without sub-femtosecond detectors.

Validation on CH_4^+ achieves 99.5% cross-prediction accuracy, zero mutual exclusion violation, and 0.983 trajectory fidelity. Validation on Rhodamine 6G reproduces Stokes shift, reorganisation energy, and Huang–Rhys factors within experimental uncertainty, and correctly identifies the C_2 point group from diffraction.

The hologram is empty. It stores nothing. It derives everything.

References

Andreas C. Albrecht. On the theory of Raman intensities. *The Journal of Chemical Physics*, 34 (5):1476–1484, 1961.

Vladimir I. Arnold. *Mathematical Methods of Classical Mechanics*. Springer-Verlag, New York, 2nd edition, 1989.

Ludwig Boltzmann. Über die Beziehung zwischen dem zweiten Hauptsatze der mechanischen Wärmetheorie und der Wahrscheinlichkeitsrechnung respektive den Sätzen über das Wärmegleichgewicht. *Wiener Berichte*, 76:373–435, 1877.

Max Born and Emil Wolf. *Principles of Optics*. Cambridge University Press, Cambridge, 7th edition, 1999.

Edward U. Condon. Nuclear motions associated with electron transitions in diatomic molecules. *Physical Review*, 32(6):858–872, 1928.

James W. Cooley and John W. Tukey. An algorithm for the machine calculation of complex Fourier series. *Mathematics of Computation*, 19(90):297–301, 1965.

F. Albert Cotton. *Chemical Applications of Group Theory*. Wiley-Interscience, New York, 3rd edition, 1990.

Franz Duschinsky. On the interpretation of electronic spectra of polyatomic molecules. *Acta Physicochimica URSS*, 7:551–566, 1937.

Albert Einstein. Zur Quantentheorie der Strahlung. *Physikalische Zeitschrift*, 18:121–128, 1917.

Enrico Fermi. Über den Ramaneffekt des Kohlendioxyds. *Zeitschrift für Physik*, 71(3–4):250–259, 1931.

James Franck. Elementary processes of photochemical reactions. *Transactions of the Faraday Society*, 21:536–542, 1926.

Daniel C. Harris and Michael D. Bertolucci. *Symmetry and Spectroscopy: An Introduction to Vibrational and Electronic Spectroscopy*. Dover, New York, 1989.

Gerhard Herzberg. *Molecular Spectra and Molecular Structure. II. Infrared and Raman Spectra of Polyatomic Molecules*. Van Nostrand, New York, 1945.

Robert C. Hilborn. Einstein coefficients, cross sections, f values, dipole moments, and all that. *American Journal of Physics*, 50(11):982–986, 1982.

Kun Huang and Avril Rhys. Theory of light absorption and non-radiative transitions in F-centres. *Proceedings of the Royal Society of London. Series A*, 204(1078):406–423, 1950.

III Johnson, Russell D. NIST computational chemistry comparison and benchmark database, NIST standard reference database number 101. https://cccbdb.nist.gov/, 2022. Release 22, May 2022.

Sunghwan Kim et al. PubChem 2023 update. *Nucleic Acids Research*, 51(D1):D1373–D1380, 2023.

Robert F. Kubin and Aaron N. Fletcher. Fluorescence quantum yields of some rhodamine dyes. *Journal of Luminescence*, 27(4):455–462, 1982.

Rolf Landauer. Irreversibility and heat generation in the computing process. *IBM Journal of Research and Development*, 5(3):183–191, 1961.

Derek A. Long. *The Raman Effect: A Unified Treatment of the Theory of Raman Scattering by Molecules*. Wiley, Chichester, 2002.

Douglas Magde, Roger Wong, and Paul G. Seybold. Fluorescence quantum yields and their relation to lifetimes of rhodamine 6G and fluorescein in nine solvents: Improved absolute standards for quantum yields. *Photochemistry and Photobiology*, 75(4):327–334, 2002.

Rudolph A. Marcus. On the theory of oxidation-reduction reactions involving electron transfer. I. *The Journal of Chemical Physics*, 24(5):966–978, 1956.

Takeshi Oka. Observation of the infrared spectrum of H_3^+. *Physical Review Letters*, 45(7):531–534, 1980.

Henri Poincaré. Sur le problème des trois corps et les équations de la dynamique. *Acta Mathematica*, 13:1–270, 1890.

C. V. Raman and K. S. Krishnan. A new type of secondary radiation. *Nature*, 121(3048):501–502, 1928.

Claude E. Shannon. A mathematical theory of communication. *Bell System Technical Journal*, 27(3):379–423, 1948.

George Gabriel Stokes. On the change of refrangibility of light. *Philosophical Transactions of the Royal Society of London*, 142:463–562, 1852.

Edgar Bright Wilson, Jr., John Courtney Decius, and Paul C. Cross. *Molecular Vibrations: The Theory of Infrared and Raman Vibrational Spectra*. McGraw-Hill, New York, 1955.