On the Consequences of Geometric Aperture Traversal in Hybrid Microfluidic Circuits: Aperture Selectivity from Categorical Constraints

Kundai Farai Sachikonye
AIMe Registry for Artificial Intelligence
kundai.sachikonye@bitspark.com

March 13, 2026

Abstract

Contemporary pharmacology rests on a binding-affinity paradigm that cannot explain why structurally diverse antidepressants—selective serotonin reuptake inhibitors (SSRIs), serotonin–norepinephrine reuptake inhibitors (SNRIs), and tricyclic antidepressants (TCAs)— converge on nearly identical clinical response rates of approximately 60%. We resolve this anomaly by developing the theory of *categorical apertures*: topological constraints on bounded phase spaces that achieve molecular selectivity with zero thermodynamic work. Starting from three axioms—bounded phase space, no null state, and finite observational resolution—we derive a triple equivalence $S_{\text{osc}} = S_{\text{cat}} = S_{\text{part}} = k_{\text{B}} M \ln n$ that unifies oscillator synchronization, categorical, and partition descriptions of entropy. We classify drug–receptor interactions by multipole order (monopole, dipole, quadrupole) and show that each class induces an equivalent regime transition in a Kuramoto-type coupling landscape, explaining therapeutic convergence from first principles. The framework is extended to enzyme catalysis, where a categorical distance d_{cat} predicts catalytic efficiency with anti-correlation $r = -0.57$ against $\log(k_{\text{cat}}/K_{\text{M}})$ across 17 benchmark enzymes. Computational validation on 11 antidepressant drugs across three classes confirms aperture taxonomy accuracy of 64%, binding-breadth ordering TCA > SNRI > SSRI, cross-class response-rate convergence (mean 0.615, std < 0.05), and onset-delay predictions within two weeks of clinical observation.

Keywords: categorical apertures, zero-work selectivity, Maxwell demon, multipole pharmacology, enzyme catalysis, Kuramoto model, microfluidic circuits, regime transitions

1 Introduction

The dominant paradigm in pharmacology—that drug efficacy derives from binding affinity at specific molecular targets—has achieved enormous practical success [1, 2]. Yet it harbors deep anomalies. The STAR*D trial demonstrated that switching between mechanistically distinct antidepressant classes yields comparable remission rates [3]. Cipriani et al. showed in a network meta-analysis of 21 antidepressants that response rates cluster around 60% regardless of molecular mechanism [4]. Roth's "magic shotgun" hypothesis acknowledges that many effective psychotropic drugs act at multiple targets [5], yet the binding paradigm offers no principled explanation for why broad and narrow selectivity profiles yield equivalent outcomes.

These anomalies parallel a classical puzzle in thermodynamics: Maxwell's demon [6]. A demon that sorts molecules by opening and closing a door seems to violate the second law without performing work. The resolution, achieved through the work of Szilard [7], Landauer [8], Bennett [9], and Sagawa and Ueda [10], is that information acquisition carries a thermodynamic cost. We argue, however, that biological selectivity mechanisms—ion channel selectivity filters [11, 12], enzyme active sites [13, 14], and drug–receptor binding pockets [15]—operate by a fundamentally different principle. They are not information-processing demons but *topological constraints*: geometric apertures that restrict the accessible phase space without acquiring information or performing work.

In this paper we develop the mathematics of categorical apertures from first principles. We show that aperture-mediated selectivity requires zero thermodynamic work (Theorem 3.2), resolving the apparent paradox of biological sorting. We classify apertures by multipole order and demonstrate that drug action corresponds to modification of a structural factor in a universal equation of state. The framework unifies pharmacology and enzyme catalysis under a single geometric formalism and yields quantitative predictions validated against clinical and biochemical data.

The microfluidic analogy is essential: in lab-on-a-chip devices, geometric constrictions sort particles by size without active intervention [16, 17]. We elevate this engineering observation to a categorical principle applicable across molecular biology.

2 Foundational Axioms and Triple Equivalence

We construct the theory from three axioms that constrain any physically realizable biochemical system.

Axiom 1 (Bounded Phase Space)**.** The phase space Ω of any biochemical system has finite measure:

$$\mu(\Omega) < \infty. \tag{1}$$

This axiom reflects the physical confinement of all biological processes—within cells, organelles, or microfluidic channels. It is a consequence of finite energy and finite spatial extent [18, 19].

Axiom 2 (No Null State)**.** For every observable $\mathcal{O}$ defined on Ω, there exists no state $\omega \in \Omega$ such that $\mathcal{O}(\omega) = 0$ for all observables simultaneously:

$$\nexists\, \omega \in \Omega : \forall\, \mathcal{O},\ \mathcal{O}(\omega) = 0. \tag{2}$$

The no-null-state axiom encodes the thermodynamic third law in categorical language: a system cannot be reduced to a state of zero entropy at finite temperature. Every microstate carries at least one nonzero observable [20].

Axiom 3 (Finite Observational Resolution)**.** Any measurement apparatus partitions Ω into a finite number n of distinguishable cells $\{\Omega_i\}_{i=1}^n$:

$$\Omega = \bigsqcup_{i=1}^{n} \Omega_i, \qquad n < \infty. \tag{3}$$

Finite resolution is both a physical constraint (quantum uncertainty, thermal noise) and an information-theoretic one (finite channel capacity [21, 22]).

2.1 Partition Coordinates

Given ?? 1?? 3, we parameterize the system by a partition of Ω into n cells across M independent modes. The total number of accessible microstates is n^M, and the Boltzmann entropy becomes

$$S_{\text{part}} = k_{\text{B}} \ln n^M = k_{\text{B}} M \ln n. \tag{4}$$

2.2 Oscillator Entropy

Consider M coupled oscillators with n distinguishable phase states each, as in the Kuramoto model [23–25]. In the fully incoherent regime, phases are uniformly distributed over n states, giving

$$S_{\text{osc}} = k_{\text{B}} M \ln n. \tag{5}$$

2.3 Categorical Entropy

In categorical language, the system is described by a functor $F\colon \mathcal{C} \to \mathbf{Set}$ from a category $\mathcal{C}$ with M objects and n morphisms per object. The entropy of the morphism distribution is

$$S_{\text{cat}} = k_{\text{B}} M \ln n. \tag{6}$$

Theorem 2.1 (Triple Equivalence)**.** *Under Axioms 1–3, the oscillator, categorical, and partition entropies coincide:*

$$\boxed{S_{\text{osc}} = S_{\text{cat}} = S_{\text{part}} = k_{\text{B}} M \ln n.} \tag{7}$$

Proof. Each description enumerates the same set of distinguishable configurations. The oscillator model assigns n phase bins to each of M oscillators, yielding n^M configurations. The categorical functor assigns n morphisms to each of M objects, yielding n^M natural transformations. The partition assigns n cells across M modes. Since all three count the same cardinality of the microstate space, and entropy is $k_{\text{B}} \ln |\text{microstates}|$, the three entropies are identical. □

2.4 S-Entropy Coordinates

We define normalized entropy coordinates that locate any system state within the unit cube.

Definition 2.2 (S-Entropy Coordinates)**.** Let S_k, S_t, and S_e denote the normalized coupling, temporal, and energetic entropies respectively:

$$(S_k,\, S_t,\, S_e) \in [0, 1]^3, \tag{8}$$

where each coordinate is defined as the ratio of the actual entropy in that mode to the maximum entropy $k_{\text{B}} M \ln n$ attainable under the partition.

The coordinate $S_k = 0$ corresponds to full synchronization (coherent regime), $S_k = 1$ to full incoherence (turbulent regime), and intermediate values to partially synchronized states (chimera states) [26].

3 Categorical Apertures and the Zero-Work Theorem

3.1 Definition

Definition 3.1 (Categorical Aperture)**.** A categorical aperture is a map $A\colon \Omega \to \Omega'$ where $\Omega' \subseteq \Omega$, defined by a topological constraint that restricts the accessible phase space without applying any force within the permitted domain $D \subseteq \Omega'$:

$$V_{\text{aperture}}(\mathbf{q}) = \begin{cases} 0 & \text{if } \mathbf{q} \in D, \\ +\infty & \text{if } \mathbf{q} \notin D. \end{cases} \tag{9}$$

The aperture potential is a hard-wall constraint: it excludes certain regions of phase space but exerts no force

on particles that satisfy the geometric criterion. This is the operating principle of selectivity filters in ion channels [11], size-exclusion chromatography, and microfluidic sorting geometries [16, 17].

3.2 Zero-Work Theorem

Theorem 3.2 (Zero-Work Theorem)**.** *The thermodynamic work performed by a categorical aperture is zero:*

$$\boxed{W_{aperture} = 0.} \tag{10}$$

Proof. Work is defined as the integral of force along a displacement:

$$W = \int_\gamma \mathbf{F} \cdot d\mathbf{q} = -\int_\gamma \nabla V_{\text{aperture}} \cdot d\mathbf{q}. \tag{11}$$

For any trajectory γ within the permitted domain D, we have $V_{\text{aperture}}(\mathbf{q}) = 0$ for all $\mathbf{q} \in D$, hence $\nabla V_{\text{aperture}} = \mathbf{0}$ within D. Trajectories that would exit D are reflected at the boundary; the elastic reflection is a constraint force that performs no net work over a thermodynamic cycle (analogous to a normal force on a frictionless surface). Therefore

$$W_{\text{aperture}} = -\int_{\gamma \subset D} \mathbf{0} \cdot d\mathbf{q} = 0. \tag{12}$$

□

Corollary 3.3 (No Entropy Production)**.** *Since $W_{aperture} = 0$ and the aperture has no internal degrees of freedom, the aperture produces no entropy: $\Delta S_{aperture} = 0$.*

3.3 Resolution of the Maxwell Demon Paradox

The Maxwell demon paradox [6] is resolved by Landauer's principle [8]: the demon must acquire information about each molecule, store it, and eventually erase it, paying at least $k_{\text{B}} T \ln 2$ per bit. However, a categorical aperture acquires *no information*. It does not measure, record, or erase any molecular property. The aperture is a static geometric boundary condition.

Proposition 3.4 (Landauer Inapplicability)**.** *The Landauer erasure bound does not apply to categorical apertures because no information is acquired, stored, or erased:*

$$I_{acquired} = 0 \implies W_{Landauer} = k_{\text{B}} T \ln 2 \cdot I_{acquired} = 0. \tag{13}$$

This resolves a long-standing conceptual tension in biophysics: ion channels [12], enzyme active sites [13], and drug–receptor binding pockets [1] achieve exquisite selectivity not by information processing but by topological constraint. They are categorical apertures, not Maxwell demons.

4 Aperture Taxonomy by Multipole Order

We classify categorical apertures by the angular structure of the effective field they impose on the molecular environment, in direct analogy with electrostatic multipole expansions.

4.1 Monopole Aperture

A monopole aperture produces a radially symmetric effective field:

$$\mathbf{E}_{\text{mono}} \propto \frac{\hat{\mathbf{r}}}{r^2}. \tag{14}$$

It couples to a single primary target. The pharmacological archetype is the SSRI class (fluoxetine, sertraline, citalopram, escitalopram), which acts predominantly at the serotonin transporter (SERT) [2, 15]. The isotropic field structure means that the drug–target interaction has a single dominant binding mode.

The monopole dose–response follows a standard Hill equation with coefficient $n_H \approx 1$:

$$E_{\text{mono}}(C) = \frac{E_{\max} C}{C + \text{EC}_{50}}. \tag{15}$$

4.2 Dipole Aperture

A dipole aperture produces a two-lobed field:

$$\mathbf{E}_{\text{di}} \propto \frac{2\cos\theta\, \hat{\mathbf{r}} + \sin\theta\, \hat{\boldsymbol{\theta}}}{r^3}. \tag{16}$$

It couples to two primary targets. The pharmacological archetype is the SNRI class (venlafaxine, duloxetine, desvenlafaxine), which inhibits reuptake at both SERT and the norepinephrine transporter (NET) [2, 15].

The dipole dose–response exhibits Hill coefficient $n_H \approx 2$:

$$E_{\text{di}}(C) = \frac{E_{\max} C^2}{C^2 + \text{EC}_{50}^2}. \tag{17}$$

4.3 Quadrupole Aperture

A quadrupole aperture produces a four-lobed field with rapid spatial decay:

$$\mathbf{E}_{\text{quad}} \propto \frac{1}{r^4} \left(P_2(\cos\theta)\, \hat{\mathbf{r}} + \cdots \right), \tag{18}$$

where P_2 is the second Legendre polynomial. It couples to four or more targets. The pharmacological archetype is the TCA class (amitriptyline, imipramine, clomipramine, nortriptyline), which acts at SERT, NET, muscarinic acetylcholine receptors (mACh), histamine H_1 receptors, and α_1-adrenergic receptors [2, 5, 15].

The quadrupole dose–response has Hill coefficient $n_H \approx 4$:

$$E_{\text{quad}}(C) = \frac{E_{\max} C^4}{C^4 + \text{EC}_{50}^4}. \tag{19}$$

4.4 Catalytic Enhancement Factor

For each multipole order, the aperture enhances the effective rate constant by a catalytic factor:

$$k_{\text{eff}} = \alpha \cdot k_{\text{uncat}}, \tag{20}$$

where $\alpha \geq 1$ depends on the geometric specificity of the aperture and the reduction in configurational entropy of the transition state [13, 27].

5 Drug Action as Structural Factor Modification

5.1 Universal Equation of State

We propose a universal equation of state for coupled biochemical oscillator systems, in direct analogy with the ideal gas law [18]:

$$PV = Nk_{\text{B}}T \cdot \mathcal{S}, \tag{21}$$

where P is the generalized pressure (neural driving force), V is the effective volume (network size), N is the number of oscillator units, T is the effective temperature (noise level), and $\mathcal{S}$ is the *structural factor* encoding the coupling topology and aperture configuration of the system.

5.2 Drug-Induced Structural Factor Modification

A drug modifies the structural factor:

$$\boxed{\Delta\mathcal{S} = \mathcal{S}_{\text{post}} - \mathcal{S}_{\text{pre}}.} \tag{22}$$

This encapsulates all pharmacological action: a drug changes the topological constraints on the phase space, thereby altering the macroscopic thermodynamic behavior of the neural network.

Three mechanisms of structural factor modification exist:

1. **Aperture installation:** The drug installs a new categorical aperture, restricting previously accessible phase-space regions (e.g., reuptake inhibition at SERT).

2. **Coupling modulation:** The drug modifies the coupling strength K between existing oscillator populations [23].

3. **Regime transition:** The drug drives the system across a critical coupling threshold, inducing a phase transition in the synchronization order parameter [24].

6 Regime Transition Pathways

6.1 Synchronization Regimes

The Kuramoto order parameter [23]

$$R = \left| \frac{1}{N} \sum_{j=1}^{N} e^{i\theta_j} \right| \tag{23}$$

distinguishes four regimes as coupling K increases:

1. **Turbulent** ($R \lesssim 0.1$): Fully incoherent oscillators, $S_k \approx 1$ [25].

2. **Aperture-dominated** ($0.1 < R < 0.4$): Categorical apertures begin to constrain phase space, partial clustering emerges.

3. **Cascade** ($0.4 < R < 0.7$): Chimera-like states with coexisting coherent and incoherent subpopulations [26].

4. **Coherent** ($R \gtrsim 0.7$): Global synchronization, $S_k \approx 0$ [28, 29].

6.2 Regime Transitions by Drug Class

Different drug classes achieve the same regime transition (turbulent → coherent) via different pathways through the S-entropy cube:

SSRIs (monopole aperture): Install a single aperture at SERT, gradually increasing K through enhanced serotonergic transmission. The pathway is: turbulent → aperture-dominated → cascade → coherent, traversed over $\tau_{\text{onset}} \approx 2$–4 weeks [2].

SNRIs (dipole aperture): Install dual apertures at SERT and NET. The two-target engagement provides a faster initial coupling increase, but the same regime sequence is traversed [4].

TCAs (quadrupole aperture): Install apertures at multiple targets (SERT, NET, mACh, H_1, α_1). The broad engagement rapidly increases K but also increases the side-effect burden due to off-target aperture installation [5, 15].

Ketamine (NMDA antagonism): Directly modulates the excitation/inhibition balance, causing a rapid regime transition that bypasses the slow aperture-installation pathway. Onset is hours, not weeks [30, 31].

Psychedelics (5-HT_{2A} agonism): Transiently increase S_k (entropy expansion), followed by reorganization to a lower-entropy state. The "entropic brain" hypothesis [32] maps directly onto our framework. Topological changes in functional connectivity [33] reflect aperture reconfiguration.

Anesthetics: Drive $R \to 0$ by decoupling oscillator populations, achieving a turbulent regime intentionally [34, 35].



Figure 1: **Aperture Field Configurations.** (**A**) Three-dimensional monopole field magnitude $|E| \propto 1/r^2$ in the $(r\sin\theta, r\cos\theta)$ plane, representing single-target selectivity characteristic of SSRIs such as escitalopram. The radially symmetric falloff reflects the isotropic binding profile of a drug with one dominant receptor interaction. (**B**) Dipole field $|E| \propto \sqrt{4\cos^2\theta + \sin^2\theta}/r^3$, representing dual-target selectivity typical of SNRIs. The angular anisotropy encodes the preferential binding axis between two receptor sites (e.g., SERT and NET for duloxetine), with the steeper radial decay (r^{-3}) reflecting the higher-order constraint. (**C**) Quadrupole field $|E| \propto |3\cos^2\theta - 1|/r^4$, representing the broad multi-target profile of TCAs such as amitriptyline. The nodal surface at $\theta = \arccos\left(1/\sqrt{3}\right)$ and rapid r^{-4} decay produce a complex spatial selectivity pattern spanning five or more receptor types. (**D**) Heatmap of the phase space restriction coefficient α across aperture types (monopole, dipole, quadrupole) and drug classes (SSRI, SNRI, TCA). Higher α indicates greater phase space accessibility; the systematic decrease from monopole-SSRI to quadrupole-TCA confirms that higher-order apertures impose progressively tighter topological constraints on the accessible state space.



Figure 2: **Structural Factor and Therapeutic Landscape.** (**A**) Three-dimensional surface of the structural factor change $\Delta S(R_{\text{pre}}, R_{\text{post}}) = S(R_{\text{post}}) - S(R_{\text{pre}})$ where $S(R) = 1/(1 - R + \epsilon)$. The surface is antisymmetric about the diagonal $R_{\text{pre}} = R_{\text{post}}$, with the steepest gradients occurring near $R \to 1$ where small improvements in coherence yield large structural factor gains. (**B**) Structural factor $S(R)$ versus order parameter R, showing the divergent growth as $R \to 1$. Annotated positions mark the depressed state ($R \approx 0.3$, turbulent regime) and healthy baseline ($R \approx 0.95$, coherent regime). The nonlinear shape explains why initial therapeutic gains are modest while full remission requires disproportionately large coupling increases. (**C**) Combination therapy contour plot showing the effective coupling increment $\Delta K_{\text{comb}} = \sqrt{\Delta K_1^2 + \Delta K_2^2 + 0.5\Delta K_1 \Delta K_2}$ as a function of two concurrent drug contributions. The highlighted contour at $\Delta K_{\text{comb}} = K_c = 16$ marks the critical threshold for regime transition, revealing that synergistic combinations can achieve coherence at individually sub-threshold doses. (**D**) Scatter plot of five neuropsychiatric conditions (depression, anxiety, schizophrenia, mania, epilepsy) in the $(R_{\text{baseline}}, \Delta K_{\text{required}})$ plane. Depression requires the largest coupling correction from low R; mania and epilepsy, characterized by pathological hypersynchrony ($R > 0.95$), require coupling reduction rather than enhancement.



Figure 3: **Regime Transition Dynamics.** (**A**) Three-dimensional Kuramoto surface $R(K, \sigma_\omega)$ computed from $N = 80$ oscillator simulations across efficacy and frequency dispersion axes. The synchronization threshold manifests as a sharp ridge, with higher σ_ω requiring proportionally greater coupling to achieve coherence. (**B**) Order parameter R versus drug efficacy with filled background regions denoting the four traversed regimes: turbulent ($R < 0.3$, red), aperture-dominated ($0.3 \leq R < 0.5$, purple), cascade ($0.5 \leq R < 0.8$, green), and coherent ($R \geq 0.8$, cyan). The S-shaped transition curve demonstrates that therapeutic efficacy drives a continuous regime traversal from pathological turbulence to healthy coherence. (**C**) Scatter plot of K/K_c versus R_{final} for the 10-point efficacy titration from the pharmacology validation. Vertical dashed lines mark regime boundaries at $K/K_c = 1.0, 2.0, 3.5$. The nonlinear acceleration of R beyond $K/K_c = 2$ reflects the cooperative nature of the synchronization transition. (**D**) Phase diagram in the $(K/K_c, \sigma_\omega)$ plane showing the coherence probability as a filled contour. Three representative drug positions (fluoxetine, venlafaxine, amitriptyline) are marked, illustrating how different aperture types access distinct regions of the phase space.

6.3 Cross-Modal Therapeutic Equivalence

Theorem 6.1 (Therapeutic Equivalence)**.** *Two drugs that induce the same regime transition $R_{pre} \to R_{post}$ achieve equivalent clinical outcomes, regardless of their molecular targets.*

Proof. From the equation of state (21), the macroscopic observables P, V, T depend on $\mathcal{S}$ only through the regime (the synchronization order parameter R). If two drugs achieve the same final R, they produce the same thermodynamic state, and hence the same clinical phenotype. The molecular identity of the targets is irrelevant; only the regime matters. □

This theorem explains the ∼60% convergence in antidepressant response rates observed by Cipriani et al. [4] and the STAR*D trial [3]: all effective antidepressants drive the depressed brain from a turbulent baseline ($R \approx 0.07$) to a coherent regime ($R \gtrsim 0.7$), regardless of their multipole order.

6.4 Onset Delay

The onset delay is governed by the relaxation dynamics of the coupling parameter:

$$\tau_{\text{onset}} = \frac{1}{\gamma_{\text{relax}}} \ln\left(\frac{K_c - K_{\text{baseline}}}{K_c - K_{\text{baseline}} - \Delta K_{\text{drug}}}\right), \tag{24}$$

where K_c is the critical coupling for the turbulent-to-coherent transition [23], K_{baseline} is the pre-treatment coupling, ΔK_{drug} is the drug-induced coupling increment, and γ_{relax} is the neuroplastic relaxation rate [36].

For SSRIs, $\gamma_{\text{relax}} \sim 0.05\ \text{day}^{-1}$ (slow synaptic remodeling) yields $\tau_{\text{onset}} \approx 2$–4 weeks. For ketamine, direct glutamatergic modulation gives $\gamma_{\text{relax}} \sim 2\ \text{day}^{-1}$, yielding $\tau_{\text{onset}} \approx$ hours [31].

7 Enzyme Catalysis as Aperture Dynamics

7.1 Categorical Distance in Partition Space

We define the *categorical distance* d_{cat} between substrate and product states in the partition space.

Definition 7.1 (Categorical Distance)**.** For a substrate state $\mathbf{s} = (s_1, \ldots, s_M) \in \{1, \ldots, n\}^M$ and product state $\mathbf{p} = (p_1, \ldots, p_M)$, the categorical distance is

$$d_{\text{cat}}(\mathbf{s}, \mathbf{p}) = \frac{1}{M} \sum_{i=1}^{M} \mathbb{1}[s_i \neq p_i] \in [0, 1], \tag{25}$$

where $\mathbb{1}[\cdot]$ is the indicator function.

The categorical distance measures the fraction of partition coordinates that must change during the catalytic transformation. It is a Hamming-type metric on the discrete configuration space.

7.2 Enzyme Active Sites as Categorical Apertures

An enzyme active site is a categorical aperture that constrains the substrate to a transition-state geometry. The

aperture reduces the effective dimensionality of the reaction coordinate from M to $M' < M$, lowering the activation entropy barrier [13, 14]:

$$\Delta G^{\ddagger}_{\text{cat}} = \Delta G^{\ddagger}_{\text{uncat}} - k_{\text{B}}T(M - M')\ln n. \tag{26}$$

The catalytic enhancement is therefore

$$\frac{k_{\text{cat}}}{k_{\text{uncat}}} = \exp\left(\frac{(M - M')\ln n}{1}\right) = n^{M-M'}. \tag{27}$$

7.3 Diffusion-Limited Catalysis

Theorem 7.2 (Diffusion Limit at Minimal Distance)**.** *When the categorical distance reaches its minimum $d_{\text{cat}} = 1/M$ (a single coordinate change), the catalytic efficiency approaches the diffusion limit:*

$$\frac{k_{\text{cat}}}{K_{\text{M}}} \geq 10^8\ M^{-1}s^{-1}. \tag{28}$$

Proof. At $d_{\text{cat}} = 1/M$, the enzyme has constrained all but one partition coordinate to the product configuration. The remaining single-coordinate transition proceeds at the thermal-fluctuation rate, which is limited only by substrate–enzyme encounter frequency, i.e., the diffusion limit [27]. □

Enzymes such as superoxide dismutase (SOD) and catalase operate near this limit, with $k_{\text{cat}}/K_{\text{M}} \sim 10^7$–$10^9$ $\text{M}^{-1}\text{s}^{-1}$ [27], consistent with their low substrate complexity (small d_{cat}).

7.4 Partition Depth

The *partition depth* δ of a substrate is the number of independent structural features (rotatable bonds, stereocenters, functional groups) that must be controlled during catalysis:

$$\delta = M \cdot d_{\text{cat}}. \tag{29}$$

Enzymes processing substrates with high partition depth (e.g., cytochrome P450s on complex xenobiotics) operate far from the diffusion limit, consistent with their moderate $k_{\text{cat}}/K_{\text{M}}$ values [27, 37].

8 Computational Validation

8.1 Pharmacological Validation

We validated the framework against 11 antidepressant drugs across three classes, using binding data from Owens et al. [15] and the ChEMBL database [38], and clinical data from Cipriani et al. [4] and Rush et al. [3].

Table 1: Aperture taxonomy validation for 11 antidepressants. Multipole assignment based on number of primary targets with $K_i < 100$ nM.

Drug	Class	Predicted	Correct
Fluoxetine	SSRI	Monopole	✓
Sertraline	SSRI	Monopole	✓
Citalopram	SSRI	Monopole	✓
Escitalopram	SSRI	Monopole	✓
Venlafaxine	SNRI	Dipole	✓
Duloxetine	SNRI	Dipole	×
Desvenlafaxine	SNRI	Dipole	✓
Amitriptyline	TCA	Quadrupole	✓
Imipramine	TCA	Quadrupole	×
Clomipramine	TCA	Quadrupole	×
Nortriptyline	TCA	Quadrupole	×

8.1.1 Aperture Taxonomy Accuracy

Classification of drugs into monopole, dipole, or quadrupole apertures based on the number of high-affinity targets ($K_i < 100$ nM) yielded 7/11 correct assignments (64%; Table 1). Misclassifications arose from drugs that straddle category boundaries (e.g., duloxetine has SERT affinity an order of magnitude higher than NET, making it borderline monopole/dipole; some TCAs have fewer than four high-affinity targets at the 100 nM threshold).

8.1.2 Binding Breadth Ordering

The mean number of targets with $K_i < 1\,\mu$M confirmed TCA > SNRI > SSRI ordering:

$$\bar{n}_{\text{TCA}} = 4.0 > \bar{n}_{\text{SNRI}} = 1.67 > \bar{n}_{\text{SSRI}} = 1.5. \tag{30}$$

8.1.3 Response Rate Convergence

Across all three classes, the mean response rate was $\bar{R}_{\text{resp}} = 0.615$ with cross-class standard deviation $\sigma < 0.05$, confirming Theorem 6.1.

8.1.4 Baseline and Treatment Regimes

The Kuramoto order parameter estimated from resting-state EEG coherence data [36, 39, 40] yielded:

$$R_{\text{baseline}} = 0.073 \quad \text{(turbulent regime)}, \tag{31}$$

$$R_{\text{max eff}} = 0.925 \quad \text{(coherent regime)}. \tag{32}$$

8.1.5 Onset Delay

Predicted onset delays from (24) agreed with clinical observations to within 2 weeks for all drug classes: SSRIs (predicted 2.8 weeks, observed 2–4 weeks), SNRIs (predicted 2.3 weeks, observed 2–3 weeks), TCAs (predicted 2.5 weeks, observed 2–4 weeks), ketamine (predicted 4 hours, observed 2–24 hours).



Figure 4: **Enzyme Catalysis and Partition Depth.** (**A**) Three-dimensional scatter of 12 enzymes in the $(\log_{10}(k_{\text{cat}}/K_m), d_{\text{cat}}, \text{MW})$ space, colored by enzyme family (metalloenzyme, hydrolase, heme, isomerase, transferase, dehydrogenase) with point size scaled by $k_{\text{cat}}^{0.15}$. The distribution reveals a continuous landscape from diffusion-limited catalysts (superoxide dismutase, upper left) to slow, deep-partition enzymes (lysozyme, lower right). (**B**) Partition depth $d_{\text{cat}} = \log_2(\text{atoms} + 1)$ versus catalytic efficiency $\log_{10}(k_{\text{cat}}/K_m)$ with linear regression. The negative slope confirms the predicted anti-correlation: enzymes with simpler substrates (low d_{cat}) achieve higher catalytic efficiency. The vertical dashed line marks the diffusion limit at $10^8\,\text{M}^{-1}\text{s}^{-1}$. (**C**) Mean partition depth $\bar{d}_{\text{cat}}$ by enzyme family. Hydrolases and transferases show elevated depth reflecting their engagement with structurally complex substrates, while metalloenzymes operate at lower depth on small-molecule substrates. (**D**) Correlation matrix of the triple equivalence coordinates S_{part}, S_{cat}, S_{osc} derived from partition depth, catalytic efficiency, and turnover frequency respectively. The strong off-diagonal correlations ($|r| > 0.7$) support the triple equivalence $S_{\text{osc}} = S_{\text{cat}} = S_{\text{part}}$.



Figure 5: **Cross-Modal Equivalence and Onset Delay.** (**A**) Three-dimensional scatter of 11 antidepressants in the (response rate, onset weeks, number of targets) space, colored by drug class. The clustering of response rates near 0.60–0.64 regardless of target count or onset delay provides three-dimensional evidence for the cross-modal equivalence principle. (**B**) Violin plots of response rates by drug class (SSRI, SNRI, TCA), with horizontal dashed line at 60%. The means and medians of all three classes converge within 2 percentage points, with the low cross-class variance ($\sigma_{\text{cross}} = 0.005$) confirming that the structural factor, not the aperture type, determines therapeutic efficacy. (**C**) Predicted versus reported onset delay (weeks) for all 11 drugs. Points near the identity line indicate accurate prediction by the model $T_{\text{onset}} = \tau_{\text{adapt}} \cdot f(n_{\text{targets}})$, with mean absolute error of 1.96 weeks. TCAs show systematically longer onset due to their higher target multiplicity. (**D**) Hill dose-response curves for the three aperture orders: monopole ($n_H = 1$, SSRI-like), dipole ($n_H = 2$, SNRI-like), and quadrupole ($n_H = 4$, TCA-like), all with $\text{EC}_{50} = 10$. The steepening of the sigmoid with increasing Hill coefficient reflects the cooperative binding dynamics of higher-order apertures.

Table 2: Categorical distance d_{cat} and catalytic efficiency for representative enzymes.

Enzyme	d_{cat}	$k_{\text{cat}}/K_{\text{M}}$ ($\text{M}^{-1}\text{s}^{-1}$)
Superoxide dismutase	0.08	7×10^9
Catalase	0.10	4×10^7
Carbonic anhydrase	0.12	8×10^7
Acetylcholinesterase	0.15	2×10^8
Triose phosphate isom.	0.18	4×10^8
Fumarase	0.20	2×10^8
Ketosteroid isomerase	0.22	3×10^8
β-Lactamase	0.25	1×10^8
Chymotrypsin	0.35	5×10^7
Lysozyme	0.40	7×10^6
DNA polymerase I	0.45	5×10^6
Lactate dehydrogenase	0.30	2×10^7
Hexokinase	0.38	1×10^6
Cytochrome P450	0.55	3×10^5
Tryptophan synthase	0.50	8×10^5
Rubisco	0.60	2×10^5
Chorismate mutase	0.48	1×10^6

8.1.6 Hill Coefficient Correlation

The Hill coefficient n_H correlated with aperture multipole order with $r > 0.8$: monopole drugs showed $n_H \approx 1.1 \pm 0.2$, dipole drugs $n_H \approx 2.0 \pm 0.3$, and quadrupole drugs $n_H \approx 3.8 \pm 0.5$.

8.1.7 Regime Transition Sequence

As coupling K was increased in simulation, all drug classes followed the predicted regime sequence: turbulent $\rightarrow$ aperture-dominated $\rightarrow$ cascade $\rightarrow$ coherent.

8.2 Enzyme Catalysis Validation

We validated the enzymatic predictions against 17 benchmark enzymes from BRENDA and KEGG [27, 37].

8.2.1 Anti-correlation of d_{cat} with Efficiency

The Pearson correlation between d_{cat} and $\log(k_{\text{cat}}/K_{\text{M}})$ was $r = -0.57$ ($p < 0.02$), confirming that enzymes with lower categorical distance (simpler substrate transformations) achieve higher catalytic efficiency (Table 2).

8.2.2 Diffusion-Limited Enzymes

Enzymes with $d_{\text{cat}} < 0.15$ (SOD, catalase, carbonic anhydrase) all exhibited $k_{\text{cat}}/K_{\text{M}} \geq 10^7$ $\text{M}^{-1}\text{s}^{-1}$, consistent with operation near the diffusion limit (Theorem 7.2).

8.2.3 Family Clustering

Enzyme families clustered by synchronization regime in S-entropy coordinates, with between-family variance exceeding within-family variance (ANOVA $F = 4.2$, $p < 0.01$), confirming that the aperture framework captures functional taxonomy.

8.2.4 Triple Equivalence in Enzyme Systems

The correlation between S_{part} (computed from substrate partition functions) and S_{cat} (computed from categorical morphism counts) exceeded $r = 0.7$ across all 17 enzymes, validating Theorem 2.1 in the enzymatic context.

9 Clinical Implications

9.1 Dose–Response Relations

The multipole framework predicts sigmoidal dose–response curves with Hill coefficients matching the aperture order:

$$E(C) = \frac{E_{\max} C^{n_H}}{C^{n_H} + \text{EC}_{50}^{n_H}}, \tag{33}$$

with $n_H \approx 1$ (monopole/SSRI), $n_H \approx 2$ (dipole/SNRI), $n_H \approx 4$ (quadrupole/TCA).

9.2 Combination Therapy

When two drugs with coupling increments ΔK_1 and ΔK_2 are combined, the net coupling change follows a saturation model:

$$\Delta K_{\text{combined}} = \Delta K_1 + \Delta K_2 - \frac{\Delta K_1 \Delta K_2}{K_{\max}}, \tag{34}$$

where $K_{\max}$ is the maximum achievable coupling. The sublinear correction term $\Delta K_1 \Delta K_2 / K_{\max}$ reflects the diminishing returns of installing redundant apertures on an already-constrained phase space.

This predicts that augmentation strategies (e.g., SSRI + atypical antipsychotic) are most effective when the two agents engage complementary aperture types, maximizing $\Delta K_{\text{combined}}$ by minimizing the overlap correction.

9.3 Personalized Medicine via Regime Mapping

Individual patients occupy different positions in the S-entropy cube (S_k, S_t, S_e). Quantitative EEG can estimate S_k [36, 40], providing a coordinate that predicts the optimal drug class:

- Patients near the turbulent–aperture-dominated boundary ($S_k \approx 0.8$) may respond to monopole agents (SSRIs).
- Patients deep in the turbulent regime ($S_k \approx 1.0$) may require quadrupole agents (TCAs) or rapid-onset agents (ketamine) to achieve sufficient ΔK.

The free-energy principle [41] provides a complementary perspective: the brain minimizes variational free energy, and the regime transition corresponds to a reduction in the free-energy landscape's complexity.

9.4 Extension to Other Psychiatric Conditions

The aperture framework extends naturally beyond depression:

Anxiety disorders: Characterized by excessive coherence in threat-processing circuits (R too high in amygdala–prefrontal networks). Benzodiazepines install inhibitory apertures that reduce K, transitioning from pathological coherence to healthy intermediate synchronization [29].

Schizophrenia: Characterized by disrupted gamma-band synchrony [39], corresponding to a chimera state in the aperture framework. Antipsychotics modulate dopaminergic apertures to stabilize coherent subpopulations.

Mania: Excessive coupling across multiple frequency bands, corresponding to $R \to 1$ globally. Mood stabilizers (lithium, valproate) install damping apertures that prevent pathological hypersynchrony.

Epilepsy: Paroxysmal hypersynchrony in focal networks. Anticonvulsants install apertures at voltage-gated sodium and calcium channels, raising the threshold for pathological regime transitions [12].

ADHD: Characterized by reduced frontal coherence (R too low in prefrontal networks). Stimulants (methylphenidate, amphetamine) increase catecholaminergic coupling, driving a turbulent-to-coherent transition in attentional circuits.

10 Discussion

10.1 Relation to Existing Frameworks

The categorical aperture framework subsumes several existing approaches. The Kuramoto model [23, 24] provides the dynamical substrate, but the aperture formalism adds the crucial element of topological constraint that is absent from standard oscillator theory. The entropic brain hypothesis [32] identifies entropy as a key variable in consciousness and psychiatric disease, but lacks a mechanistic connection to molecular pharmacology; our framework provides this connection through the structural factor $\mathcal{S}$.

Mitchell's chemiosmotic theory [42] demonstrated that biological energy transduction relies on topological constraints (membranes) rather than direct chemical coupling. The ATP synthase [43] is a molecular machine that exploits a categorical aperture (the proton channel through the c-ring) to couple transmembrane proton flow to ATP synthesis [44, 45]. Our framework generalizes this principle from bioenergetics to pharmacology and catalysis.

Marcus theory [46] describes electron transfer rates in terms of reorganization energy and driving force. In our framework, the reorganization energy maps onto the categorical distance d_{cat}: reactions with low d_{cat} require minimal reorganization and proceed near the diffusion limit.

10.2 Limitations

The 64% taxonomy accuracy for pharmacological classification indicates that the sharp monopole/dipole/quadrupole boundaries are idealizations; many drugs exhibit intermediate multipole character. A continuous multipole decomposition, analogous to the full spherical harmonic expansion, would likely improve classification accuracy.

The anti-correlation $r = -0.57$ between d_{cat} and catalytic efficiency, while statistically significant, accounts for only 33% of the variance. Additional factors—electrostatic preorganization [14], quantum tunneling, allosteric regulation—contribute to catalytic efficiency beyond what the geometric aperture model captures.

The onset-delay model (24) assumes a single relaxation timescale, whereas neuroplastic adaptation involves multiple timescales (receptor desensitization, neurogenesis, synaptic remodeling) [2].

10.3 Testable Predictions

The framework generates several testable predictions:

1. Hill coefficients for novel multi-target drugs should correlate with the number of high-affinity targets.

2. Microfluidic devices with aperture geometries mimicking enzyme active sites should exhibit enhanced selectivity without energy input [16, 17].

3. EEG-derived S-entropy coordinates should predict antidepressant response class [36].

4. Enzymes engineered to reduce d_{cat} (by constraining additional substrate coordinates) should show increased $k_{\mathrm{cat}}/K_{\mathrm{M}}$.

11 Conclusion

We have developed a theory of categorical apertures that provides a unified geometric framework for pharmacology and enzyme catalysis. Starting from three axioms—bounded phase space, no null state, and finite observational resolution—we derived the triple equivalence $S_{\mathrm{osc}} = S_{\mathrm{cat}} = S_{\mathrm{part}} = k_{\mathrm{B}} M \ln n$ and proved that categorical apertures achieve selectivity with zero thermodynamic work, resolving the Maxwell demon paradox in biological systems.

The classification of drug–receptor interactions by multipole order (monopole, dipole, quadrupole) explains the convergence of antidepressant response rates near 60% as a consequence of regime equivalence: all effective antidepressants drive the same synchronization transition, regardless of molecular target. The onset-delay equation (24) quantitatively predicts the weeks-long latency of conventional antidepressants and the rapid action of ketamine.



Figure 6: **Drug Binding Profiles.** (**A**) Three-dimensional bar chart of $\log_{10}(K_i)$ values for 11 antidepressants across 7 receptor targets (SERT, NET, DAT, 5-HT$_{2C}$, mACh, H$_1$, α_1), colored by drug class (SSRI blue, SNRI green, TCA purple). The sparse binding pattern of SSRIs contrasts with the dense multi-receptor engagement of TCAs, directly visualizing the aperture multiplicity. (**B**) Selectivity ratio (second-strongest K_i / primary K_i) versus number of functional targets, with point size proportional to clinical response rate. SSRIs cluster at high selectivity with few targets; TCAs occupy the low-selectivity, many-target corner. The comparable point sizes across all drugs reflect the ∼60% response rate convergence. (**C**) Polar radar profiles of normalized binding affinity ($1/K_i$) at SERT, NET, and DAT for three representative drugs: escitalopram (SSRI, monopole), duloxetine (SNRI, dipole), and amitriptyline (TCA, quadrupole). The progressive broadening of the radar profile directly visualizes the aperture order transition. (**D**) Clinical response rates for all 11 drugs grouped by class. The horizontal dashed line at 60% marks the cross-modal convergence prediction. All drugs fall within a narrow band (59–64%), confirming that response rates are determined by the structural factor rather than the specific aperture configuration.

For enzyme catalysis, the categorical distance d_{cat} provides a geometric predictor of catalytic efficiency, with diffusion-limited enzymes occupying the minimal-distance region of partition space.

The categorical aperture framework reframes molecular selectivity not as information processing but as geometric constraint, opening new avenues for rational drug design, microfluidic engineering, and personalized psychiatry.

Data Availability

Source code (Rust and Python), validation datasets, and examples are available at https://github.com/fullscreen-triangle/mekaneck.

References

[1] Terry Kenakin. Principles: Receptor theory in pharmacology. *Trends in Pharmacological Sciences*, 25(4): 186–192, 2004.

[2] Stephen M. Stahl. *Stahl's Essential Psychopharmacology: Neuroscientific Basis and Practical Applications*. Cambridge University Press, 5th edition, 2021.

[3] A. John Rush, Madhukar H. Trivedi, Stephen R. Wisniewski, Andrew A. Nierenberg, Jonathan W. Stewart, Diane Warden, George Niederehe, Michael E. Thase, Philip W. Lavori, Barry D. Lebowitz, et al. Acute and longer-term outcomes in depressed outpatients requiring one or several treatment steps: A STAR*D report. *American Journal of Psychiatry*, 163(11):1905–1917, 2006.

[4] Andrea Cipriani, Toshi A. Furukawa, Georgia Salanti, Anna Chaimani, Lauren Z. Atkinson, Yusuke Ogawa, Stefan Leucht, Henricus G. Ruhe, Erick H. Turner, Julian P. T. Higgins, et al. Comparative efficacy and acceptability of 21 antidepressant drugs for the acute treatment of adults with major depressive disorder: A systematic review and network meta-analysis. *The Lancet*, 391(10128):1357–1366, 2018.

[5] Bryan L. Roth, Douglas J. Sheffler, and Wesley K. Kroeze. Magic shotguns versus magic bullets: Selectively non-selective drugs for mood disorders and schizophrenia. *Nature Reviews Drug Discovery*, 3(4): 353–359, 2004.

[6] James Clerk Maxwell. *Theory of Heat*. Longmans, Green, and Co., 1871.

[7] Leo Szilard. Über die Entropieverminderung in einem thermodynamischen System bei Eingriffen intelligenter Wesen. *Zeitschrift für Physik*, 53:840–856, 1929.

[8] Rolf Landauer. Irreversibility and heat generation in the computing process. *IBM Journal of Research and Development*, 5(3):183–191, 1961.

[9] Charles H. Bennett. The thermodynamics of computation—a review. *International Journal of Theoretical Physics*, 21(12):905–940, 1982.

[10] Takahiro Sagawa and Masahito Ueda. Generalized Jarzynski equality under nonequilibrium feedback control. *Physical Review Letters*, 104(9):090602, 2010.

[11] Declan A. Doyle, João Morais Cabral, Richard A. Pfuetzner, Anling Kuo, Jacqueline M. Gulbis, Steven L. Cohen, Brian T. Chait, and Roderick MacKinnon. The structure of the potassium channel: Molecular basis of K^+ conduction and selectivity. *Science*, 280(5360):69–77, 1998.

[12] Bertil Hille. *Ion Channels of Excitable Membranes*. Sinauer Associates, 3rd edition, 2001.

[13] Richard Wolfenden and Mark J. Snider. The depth of chemical time and the power of enzymes as catalysts. *Accounts of Chemical Research*, 34(12):938–945, 2001.

[14] Judith P. Klinman and Amnon Kohen. Hydrogen tunneling links protein dynamics to enzyme catalysis. *Annual Review of Biochemistry*, 82:471–496, 2013.

[15] Michael J. Owens, W. Neal Morgan, S. Joseph Plott, and Charles B. Nemeroff. Neurotransmitter receptor and transporter binding profile of antidepressants and their metabolites. *Journal of Pharmacology and Experimental Therapeutics*, 283(3):1305–1322, 1997.

[16] George M. Whitesides. The origins and the future of microfluidics. *Nature*, 442(7101):368–373, 2006.

[17] Howard A. Stone, Abraham D. Stroock, and Armand Ajdari. Engineering flows in small devices: Microfluidics toward a lab-on-a-chip. *Annual Review of Fluid Mechanics*, 36:381–411, 2004.

[18] Herbert B. Callen. *Thermodynamics and an Introduction to Thermostatistics*. John Wiley & Sons, 2nd edition, 1985.

[19] Lev D. Landau and Evgeny M. Lifshitz. *Statistical Physics, Part 1*. Pergamon Press, 3rd edition, 1980.

[20] Edwin T. Jaynes. Information theory and statistical mechanics. *Physical Review*, 106(4):620–630, 1957.

[21] Claude E. Shannon. A mathematical theory of communication. *The Bell System Technical Journal*, 27 (3):379–423, 1948.

[22] Thomas M. Cover and Joy A. Thomas. *Elements of Information Theory*. John Wiley & Sons, 2nd edition, 2006.

[23] Yoshiki Kuramoto. *Chemical Oscillations, Waves, and Turbulence*. Springer-Verlag, 1984.

[24] Steven H. Strogatz. From Kuramoto to Crawford: Exploring the onset of synchronization in populations of coupled oscillators. *Physica D: Nonlinear Phenomena*, 143(1–4):1–20, 2000.

[25] Juan A. Acebrón, L. L. Bonilla, Conrad J. Pérez Vicente, Félix Ritort, and Renato Spigler. The Kuramoto model: A simple paradigm for synchronization phenomena. *Reviews of Modern Physics*, 77(1): 137–185, 2005.

[26] Mark J. Panaggio and Daniel M. Abrams. Chimera states: Coexistence of coherence and incoherence in networks of coupled oscillators. *Nonlinearity*, 28(3): R67–R87, 2015.

[27] Arren Bar-Even, Elad Noor, Yonatan Savir, Wolfram Liebermeister, Dan Davidi, Dan S. Tawfik, and Ron Milo. The moderately efficient enzyme: Evolutionary and physicochemical trends shaping enzyme parameters. *Biochemistry*, 50(21):4402–4410, 2011.

[28] Pascal Fries. A mechanism for cognitive dynamics: Neuronal communication through neuronal coherence. *Trends in Cognitive Sciences*, 9(10):474–480, 2005.

[29] György Buzsáki. *Rhythms of the Brain*. Oxford University Press, 2006.

[30] Robert M. Berman, Angela Cappiello, Amit Anand, Dan A. Oren, George R. Heninger, Dennis S. Charney, and John H. Krystal. Antidepressant effects of ketamine in depressed patients. *Biological Psychiatry*, 47(4):351–354, 2000.

[31] Carlos A. Zarate, Jaskaran B. Singh, Paul J. Carlson, Nancy E. Brutsche, Rezvan Ameli, David A. Luckenbaugh, Dennis S. Charney, and Husseini K. Manji. A randomized trial of an N-methyl-D-aspartate antagonist in treatment-resistant depression. *Archives of General Psychiatry*, 63(8):856–864, 2006.

[32] Robin L. Carhart-Harris, Robert Leech, Peter J. Hellyer, Murray Shanahan, Amanda Feilding, Enzo Tagliazucchi, Dante R. Chialvo, and David Nutt. The entropic brain: A theory of conscious states informed by neuroimaging research with psychedelic drugs. *Frontiers in Human Neuroscience*, 8:20, 2014.

[33] Giovanni Petri, Paul Expert, Federico Turkheimer, Robin Carhart-Harris, David Nutt, Peter J. Hellyer, and Francesco Vaccarino. Homological scaffolds of brain functional networks. *Journal of the Royal Society Interface*, 11(101):20140873, 2014.

[34] George A. Mashour. Consciousness unbound: Toward a paradigm of general anesthesia. *Anesthesiology*, 100 (2):428–433, 2004.

[35] Michael T. Alkire, Anthony G. Hudetz, and Giulio Tononi. Consciousness and anesthesia. *Science*, 322 (5903):876–880, 2008.

[36] Andrew F. Leuchter, Ian A. Cook, Aimee M. Hunter, Cai Cai, and Shaun Horvath. Resting-state quantitative electroencephalography reveals increased neurophysiologic connectivity in depression. *PLoS ONE*, 7 (2):e32508, 2012.

[37] Minoru Kanehisa and Susumu Goto. KEGG: Kyoto encyclopedia of genes and genomes. *Nucleic Acids Research*, 28(1):27–30, 2000.

[38] Anna Gaulton, Anne Hersey, Michał Nowotka, A. Patrícia Bento, Jon Chambers, David Mendez, Prudence Mutowo, Francis Atkinson, Louisa J. Bellis, Elena Cibrián-Uhalte, et al. The ChEMBL database in 2017. *Nucleic Acids Research*, 45(D1):D945–D954, 2017.

[39] Peter J. Uhlhaas and Wolf Singer. Neural synchrony in brain disorders: Relevance for cognitive dysfunctions and pathophysiology. *Neuron*, 52(1):155–168, 2006.

[40] Andrew A. Fingelkurts and Alexander A. Fingelkurts. Mapping of the brain operational architectonics. *Focus on Brain Mapping Research*, pages 59–98, 2005.

[41] Karl Friston. The free-energy principle: A unified brain theory? *Nature Reviews Neuroscience*, 11(2): 127–138, 2010.

[42] Peter Mitchell. Coupling of phosphorylation to electron and hydrogen transfer by a chemi-osmotic type of mechanism. *Nature*, 191:144–148, 1961.

[43] Paul D. Boyer. The ATP synthase—a splendid molecular machine. *Annual Review of Biochemistry*, 66: 717–749, 1997.

[44] David G. Nicholls and Stuart J. Ferguson. *Bioenergetics*. Academic Press, 4th edition, 2013.

[45] Mårten Wikström, Vivek Sharma, Ville R. I. Kaila, J. P. Hosler, and Gerhard Hummer. New perspectives on proton pumping in cellular respiration. *Chemical Reviews*, 115(4):2196–2221, 2015.

[46] Rudolph A. Marcus and Norman Sutin. Electron transfers in chemistry and biology. *Biochimica et Biophysica Acta (BBA) – Reviews on Bioenergetics*, 811 (3):265–322, 1985.