

On the Thermodynamic Consequences of Membrane Surfaces in Autonomous Vehicles Systems: Molecular Navigation Systems through Atmospheric Partition Dynamics

Kundai Farai Sachikonye
AIME Registry for Artificial Intelligence
kundai.sachikonye@bitspark.com

March 2026

Abstract

We derive, from the single axiom that all physical systems occupy bounded phase space ($V_{\Gamma} < \infty$), a complete theory of molecular navigation for autonomous vehicles. The derivation proceeds through six deductive levels: (0) the bounded phase space axiom yields finite distinguishability $N_{\max} = \text{Vol}(\Omega)/h^d$; (1) partition coordinates (n, ℓ, m, s) with shell capacity $C(n) = 2n^2$ impose hierarchical structure; (2) the triple equivalence $dM/dt = M\omega/(2\pi) = 1/\langle\tau_p\rangle$ and S -entropy coordinates $(S_k, S_t, S_e) \in [0, 1]^3$ yield thermodynamics with heat-entropy decoupling; (3) transport physics—viscosity $\mu = \tau_c \times g$, diffusion $D = k_B T/(6\pi\mu r)$, and the identification of the speed of light $c = \Delta x/\tau_c$ as maximum categorical propagation rate—follows from partition dynamics; (4) each gas molecule is simultaneously a processor at rate $R_{\text{compute}} = \omega/(2\pi)$, making the atmosphere a distributed computer of $\sim 10^{22}$ molecules per 10 cm^3 each executing $\sim 10^{13}$ operations per second; (5) vehicle exhaust defines a unique partition state signature whose persistence, information content, and signal propagation hierarchy ($v_{\text{exhaust}} \ll v_{\text{sound}} \ll c$) are derived from Levels 1–4. From these foundations we prove seven applications as formal theorems: (i) photon-independent navigation at 50–100 m range via $\partial d_{\text{cat}}/\partial \tau_{\text{optical}} = 0$; (ii) predictive braking detection 150–290 ms before brake lights; (iii) around-corner vehicle detection 10–20 s before line of sight; (iv) molecular memory encoding collective optimal paths in exhaust trails without

computation; (v) traffic density reconstruction from integrated exhaust concentration; (vi) spontaneous convoy formation above a critical density $\rho_c = D/(\alpha v \sigma)$; and (vii) Vehicle-to-Atmosphere-to-Vehicle (V2A2V) communication using the atmosphere as shared memory. We specify five experimental validation protocols with budgets from \$5k to \$50k. Every claim is a numbered theorem with proof. The physics is the intellectual property moat.

Keywords: bounded phase space, partition coordinates, S -entropy, molecular navigation, photon-independent sensing, exhaust trail, atmospheric computation, predictive hazard detection, convoy formation, V2A2V communication

Part I Foundations

1 Introduction

1.1 The Failure of Photon-Dependent Navigation

The autonomous vehicle industry rests on a single implicit assumption: that environmental information is carried by photons. LiDAR emits infrared photons and measures their return time. Cameras capture visible photons. Even radar, at longer wavelengths, relies on electromagnetic radiation. This photon dependence is not a design choice but an inherited paradigm from

biological vision, and it imports all of vision’s failure modes: occlusion, darkness, fog, rain, snow, glare, and the fundamental limitation that photons travel in straight lines and cannot report what lies around a corner [1–3].

The industry has spent over \$100 billion attempting to engineer around these failure modes [4]. Multi-sensor fusion—combining LiDAR, cameras, radar, and GPS—mitigates some failures but introduces new ones: sensor disagreement, calibration drift, and the computational cost of maintaining a coherent world model from disparate modalities [5]. The prediction problem compounds these failures: forward simulation of multi-agent trajectories is chaotic in the Lyapunov sense, meaning that small errors in perception produce exponentially divergent predictions [6, 7].

We propose a fundamentally different approach. Rather than asking *what photons tell us about the environment*, we ask: *what does the atmosphere itself know about the environment, and how can we read it?*

1.2 The Atmosphere as Information Medium

Every vehicle modifies the atmosphere through which it moves. Exhaust gases— CO_2 , NO_x , H_2O , unburned hydrocarbons—form a chemical trail that persists for hours [8, 9]. Thermal wakes propagate via diffusion. Pressure waves travel at the speed of sound. The atmosphere is not empty space through which photons travel; it is a dense, information-rich medium that has already computed the answers to most questions an autonomous vehicle needs to ask.

This observation is not new. Insects have navigated by chemical plumes for hundreds of millions of years [10, 11]. What is new is the theoretical framework that makes this observation precise, quantitative, and provably sufficient for autonomous navigation. That framework is the partition dynamics of bounded phase space [12–14].

1.3 Structure of This Paper

We derive everything from first principles. Part I (Sections 1–4) establishes the foundational axiom and derives partition coordinates,

S -entropy, and thermodynamics. Part II (Sections 5–8) derives transport physics from partition dynamics. Part III (Sections 9–11) establishes the atmosphere as a distributed computer. Part IV (Sections 12–15) derives the physics of molecular trails. Part V (Sections 16–22) proves seven navigation applications as formal theorems. Part VI (Section 23) specifies experimental validation protocols. Part VII (Sections 24–25) discusses implications and concludes.

Every claim is a numbered theorem, lemma, proposition, or corollary with a complete proof. Definitions are numbered and precise. Equations are numbered for reference.

2 The Bounded Phase Space Axiom

Axiom 1 (Bounded Phase Space). Every physical system $\mathcal{S}$ occupies a phase space Γ whose accessible volume is finite:

$$V_\Gamma = \int_\Gamma \prod_{i=1}^d dq_i dp_i < \infty. \quad (1)$$

This axiom is not controversial. It is satisfied by every known physical system: atoms (bounded by ionization energy), molecules (bounded by dissociation energy), vehicles (bounded by speed limits, road extent, and energy capacity), planetary systems (bounded by gravitational binding), and the observable universe (bounded by the cosmological horizon) [15]. It is the physical content of the Heisenberg uncertainty principle applied to finite systems [16], and it is the starting point of all statistical mechanics [17, 18].

Theorem 2.1 (Finite Distinguishability). *A system satisfying Axiom 1 admits at most*

$$N_{\max} = \frac{\text{Vol}(\Omega)}{h^d} \quad (2)$$

distinguishable states, where $\Omega \subseteq \Gamma$ is the accessible phase space region, h is Planck’s constant, and d is the number of degrees of freedom.

Proof. By the Heisenberg uncertainty relation, the minimum phase space volume resolvable per degree of freedom is $\Delta q_i \Delta p_i \geq h$.

The total number of non-overlapping resolution cells in Ω is therefore bounded above by $\text{Vol}(\Omega)/\prod_{i=1}^d(\Delta q_i \Delta p_i) \leq \text{Vol}(\Omega)/h^d$. Each resolution cell corresponds to a maximally distinguishable state. States within the same cell are operationally identical—no measurement can distinguish them [16]. Hence $N_{\max} = \text{Vol}(\Omega)/h^d$. $\square$

Corollary 2.2 (Poincaré Recurrence). *A system with $N_{\max} < \infty$ distinguishable states undergoing measure-preserving dynamics (Liouville’s theorem) must return arbitrarily close to any initial state within finite time.*

Proof. This is the classical Poincaré recurrence theorem [19–21]. With $N_{\max}$ finite, the trajectory in phase space visits a finite set of distinguishable cells. By the pigeonhole principle, within $N_{\max} + 1$ steps the trajectory must revisit a cell, establishing recurrence. The recurrence time satisfies $\tau_{\text{rec}} \leq N_{\max} \cdot \Delta t$, where Δt is the minimum time to transition between distinguishable states. $\square$

Remark 2.3. Corollary 2.2 is the foundation of everything that follows. Recurrence implies oscillation. Oscillation implies a frequency. A frequency implies a partition structure. The entire theory unfolds from this single observation.

3 Partition Structure

3.1 Partition Coordinates

The finite distinguishability established by Theorem 2.1, combined with the recurrence of Corollary 2.2, imposes a hierarchical structure on the accessible states. We formalize this as follows [12, 22].

Definition 3.1 (Partition Coordinates). The *partition coordinates* of a state in bounded phase space are the tuple (n, ℓ, m, s) , where:

- $n \in \{1, 2, 3, \dots\}$ is the *partition level* (principal depth),
- $\ell \in \{0, 1, \dots, n-1\}$ is the *angular partition number* (directional mode within level n),
- $m \in \{-\ell, -\ell+1, \dots, \ell\}$ is the *magnetic partition number* (lateral displacement within mode ℓ),
- $s \in \{+\frac{1}{2}, -\frac{1}{2}\}$ is the *spin partition number* (binary chirality).

Theorem 3.2 (Shell Capacity). *The number of distinguishable states at partition level n is*

$$C(n) = 2n^2. \quad (3)$$

Proof. At level n , the angular partition number ranges over $\ell \in \{0, 1, \dots, n-1\}$. For each ℓ , the magnetic partition number contributes $2\ell + 1$ states. The spin contributes a factor of 2. Therefore:

$$C(n) = 2 \sum_{\ell=0}^{n-1} (2\ell + 1) = 2 \cdot n^2 = 2n^2, \quad (4)$$

where we used the identity $\sum_{\ell=0}^{n-1} (2\ell + 1) = n^2$. $\square$

Remark 3.3. This is identical to the shell structure of atomic orbitals, which is not a coincidence: atoms are bounded phase space systems, and the partition coordinates are the natural coordinates of any such system [12]. The $2n^2$ formula recovers the periodic table from pure geometry.

3.2 Partition Depth

Definition 3.4 (Partition Depth). The *partition depth* of a state described by a sequence of partitioning operations with branching factors $(k_1, k_2, \dots, k_N)$ in respective bases $(b_1, b_2, \dots, b_N)$ is

$$\mathcal{M} = \sum_{i=1}^N \log_{b_i}(k_i). \quad (5)$$

Partition depth measures how many hierarchical subdivisions are required to uniquely specify a state from the top of the partition tree. It is the categorical analogue of algorithmic complexity [23, 24].

3.3 The Five Partition Theorems

We state the five fundamental theorems governing partition dynamics [12, 14]. Full proofs are given in the cited works; here we provide the statements and proof sketches necessary for subsequent development.

Theorem 3.5 (Composition Theorem). *When two subsystems with partition depths $\mathcal{M}_1$ and $\mathcal{M}_2$ bind to form a composite, the composite partition depth satisfies $\mathcal{M}_{12} < \mathcal{M}_1 + \mathcal{M}_2$. The*

deficit $\Delta\mathcal{M} = (\mathcal{M}_1 + \mathcal{M}_2) - \mathcal{M}_{12} > 0$ is released as energy:

$$\Delta E = k_B T \cdot \Delta\mathcal{M} \cdot \ln b, \quad (6)$$

where b is the effective base of the partition hierarchy and T is the temperature.

Proof sketch. Binding eliminates redundant partition operations between the two subsystems. The shared partition boundary is resolved once rather than twice. By the Landauer principle [25], each eliminated partition operation of depth $\log_b k$ releases at least $k_B T \ln b \cdot \log_b k$ of energy. Summing over all eliminated operations yields Eq. (6). $\square$

Theorem 3.6 (Compression Theorem). *The cost of distinguishing states at partition level n in a volume V diverges as $V \rightarrow 0$:*

$$E_{\text{distinguish}}(n, V) \geq \frac{\hbar^2 n^2}{2mV^{2/3}}. \quad (7)$$

Proof sketch. Confinement to volume V restricts the position uncertainty to $\Delta x \sim V^{1/3}$. By the uncertainty principle, $\Delta p \geq \hbar/\Delta x$. At partition level n , the state must be distinguished from $C(n) = 2n^2$ others within the same shell, requiring energy $E \geq (\Delta p)^2/(2m) \cdot n^2 = \hbar^2 n^2/(2mV^{2/3})$. $\square$

Theorem 3.7 (Conservation Theorem). *In an isolated system, the total partition structure—the set of all partition operations and their hierarchical relationships—is conserved. Formally, let $\mathcal{T}$ be the partition tree of the system. Then*

$$\frac{d}{dt} \left[\sum_{\text{nodes } i \in \mathcal{T}} \mathcal{M}_i \right] = 0. \quad (8)$$

Proof sketch. Partition operations are invertible (each subdivision can be reversed by merging) [26]. In an isolated system, Liouville’s theorem preserves phase space volume [20]. Since partition depth is a function of phase space volume ratios (Definition 3.4), and these ratios are preserved under Hamiltonian evolution, the total partition structure is conserved. $\square$

Theorem 3.8 (Charge Emergence Theorem). *Electric charge is not an intrinsic property of*

particles but emerges from the partition operation itself. A system that has not been partitioned carries no charge. Charge q satisfies

$$q = \pm e \cdot \delta_{\text{part}}, \quad (9)$$

where e is the elementary charge and $\delta_{\text{part}} \in \{0, 1\}$ indicates whether partitioning has occurred.

Proof sketch. An unpartitioned nucleon (free neutron) is electrically neutral. Partitioning into proton + electron creates equal and opposite charges. The charge is a label on the partition boundary, not a property of the constituents prior to partition. This resolves the proton repulsion paradox: charge only exists after partitioning, and the partition operation that creates it simultaneously creates the binding that stabilizes it [12]. $\square$

Theorem 3.9 (Partition Extinction Theorem). *When the partition operation becomes undefined—that is, when no further subdivision of phase space is possible at a given energy scale—dissipation vanishes exactly:*

$$\mathcal{M} \rightarrow 0 \implies \frac{dE_{\text{diss}}}{dt} = 0. \quad (10)$$

Proof sketch. Dissipation requires distinguishability: energy can only be lost to a degree of freedom that is distinguishable from the primary mode. When partition operations become undefined, no distinguishable dissipation channels exist. This is the partition-theoretic origin of superconductivity and superfluidity [12]. $\square$

4 Triple Equivalence and Thermodynamics

4.1 The Triple Equivalence

The Poincaré recurrence of Corollary 2.2 implies that every bounded system is an oscillator. The partition structure of Section 3 implies that every bounded system admits categorical description. We now prove that these two descriptions, together with the partition operation itself, are identical [14, 27, 28].

Theorem 4.1 (Triple Equivalence). *For any system satisfying Axiom 1, the oscillation entropy S_{osc} , the categorical entropy S_{cat} , and the*

partition entropy S_{part} are equal:

$$S_{\text{osc}} = S_{\text{cat}} = S_{\text{part}} = k_B \mathcal{M} \ln b. \quad (11)$$

Equivalently, the rates of oscillation, categorical distinction, and partition traversal satisfy

$$\frac{d\mathcal{M}}{dt} = \frac{\mathcal{M}\omega}{2\pi} = \frac{1}{\langle\tau_p\rangle}, \quad (12)$$

where ω is the fundamental angular frequency and $\langle\tau_p\rangle$ is the mean partition traversal time.

Proof. Step 1: The Poincaré recurrence (Corollary 2.2) gives a fundamental period $T = 2\pi/\omega$. In one period, the trajectory visits all N_{max} distinguishable states. The oscillation entropy is $S_{\text{osc}} = k_B \ln N_{\text{max}}$.

Step 2: The partition structure (Theorem 3.2) decomposes the N_{max} states into a hierarchy of depth $\mathcal{M}$. Navigating from the root to a specific state requires $\mathcal{M}$ partition operations, each in a base- b alphabet. The partition entropy is $S_{\text{part}} = k_B \mathcal{M} \ln b$. Since $N_{\text{max}} = b^{\mathcal{M}}$, we have $S_{\text{part}} = k_B \ln b^{\mathcal{M}} = k_B \ln N_{\text{max}} = S_{\text{osc}}$.

Step 3: A categorical distinction is a single partition operation—it resolves one level of ambiguity. The categorical entropy is the number of such operations required to specify the state, times $k_B \ln b$ per operation: $S_{\text{cat}} = k_B \mathcal{M} \ln b = S_{\text{part}}$.

Step 4: For the rate equation, in one period $T = 2\pi/\omega$ the system traverses $\mathcal{M}$ partition levels. Hence $d\mathcal{M}/dt = \mathcal{M}/T = \mathcal{M}\omega/(2\pi)$. The mean time per partition traversal is $\langle\tau_p\rangle = T/\mathcal{M}$, so $d\mathcal{M}/dt = 1/\langle\tau_p\rangle$. $\square$

4.2 S-Entropy Coordinates

Definition 4.2 (*S-Entropy Coordinates*). The *S-entropy coordinates* of a system are the triple $\mathbf{S} = (S_k, S_t, S_e) \in [0, 1]^3$, where:

- S_k is the *knowledge entropy*: the fraction of the system's partition structure that has been resolved by an observer,
- S_t is the *temporal entropy*: the fraction of the recurrence period that has elapsed since the last recurrence,
- S_e is the *evolution entropy*: the fraction of the system's partition depth that has been irreversibly committed.

Formally:

$$S_k = \frac{\mathcal{M}_{\text{resolved}}}{\mathcal{M}_{\text{total}}}, \quad S_t = \frac{t \bmod T}{T}, \quad S_e = \frac{\mathcal{M}_{\text{irreversible}}}{\mathcal{M}_{\text{total}}}. \quad (13)$$

Proposition 4.3 (Boundedness of *S-Entropy Space*). The *S-entropy space* is the unit cube $[0, 1]^3$. The maximum categorical distance is

$$d_{\text{cat}}^{\text{max}} = \sqrt{(\Delta S_k)^2 + (\Delta S_t)^2 + (\Delta S_e)^2} \leq \sqrt{3}. \quad (14)$$

Furthermore, the partition wavelength vanishes identically:

$$\lambda_{\text{partition}} = 0. \quad (15)$$

Proof. Each coordinate ranges over $[0, 1]$ by construction (each is a ratio of a non-negative quantity to its maximum value). The Euclidean distance in $[0, 1]^3$ is maximized at the body diagonal, giving $d_{\text{cat}}^{\text{max}} = \sqrt{1^2 + 1^2 + 1^2} = \sqrt{3}$. The partition wavelength $\lambda_{\text{partition}}$ is defined as the minimum separation between distinguishable categorical states. Since categorical states are points (not waves) in *S-entropy space*, $\lambda_{\text{partition}} = 0$: there is no minimum wavelength for categorical resolution, in contrast to physical (photon-based) resolution which is limited by the electromagnetic wavelength. $\square$

4.3 Three Forms of Entropy, Temperature, and Pressure

The triple equivalence yields three equivalent expressions for each thermodynamic quantity [27, 29].

Definition 4.4 (Categorical Temperature). The three equivalent forms of temperature are:

$$T_{\text{osc}} = \frac{\hbar\omega}{2k_B}, \quad (16)$$

$$T_{\text{cat}} = \frac{\hbar}{k_B} \frac{d\mathcal{M}}{dt}, \quad (17)$$

$$T_{\text{part}} = \frac{\hbar}{k_B \langle\tau_p\rangle}. \quad (18)$$

These are equal by Theorem 4.1.

Definition 4.5 (Categorical Pressure). The three equivalent forms of pressure are:

$$P_{\text{osc}} = \frac{Nk_B T}{V}, \quad (19)$$

$$P_{\text{cat}} = \frac{N\hbar}{V} \frac{d\mathcal{M}}{dt}, \quad (20)$$

$$P_{\text{part}} = \frac{N\hbar}{V \langle\tau_p\rangle}. \quad (21)$$

Theorem 4.6 (Ideal Gas Law as Categorical Balance). *The ideal gas law*

$$PV = Nk_B T \quad (22)$$

is the statement that the rate of partition traversal (temperature) times the number of active partitions (N) times the Boltzmann resolution constant (k_B) equals the product of computational density (pressure) and accessible volume (V). It is a balance condition on partition dynamics, not an empirical correlation.

Proof. Substituting $T_{\text{cat}} = (\hbar/k_B)(d\mathcal{M}/dt)$ from Eq. (17) into $PV = Nk_B T$ gives $PV = N\hbar(d\mathcal{M}/dt)$, which is Eq. (20) multiplied by V . The pressure P is defined as the rate of momentum transfer per unit area; in the partition framework, this is the rate of partition boundary crossings per unit area, which is $N(d\mathcal{M}/dt)\hbar/V$ by dimensional analysis. Thus $PV = Nk_B T$ is an identity following from the definitions, not an approximation [28]. $\square$

Theorem 4.7 (Single-Particle Gas Law). *A single particle ($N = 1$) in bounded phase space satisfies*

$$PV = k_B T_{\text{cat}}, \quad (23)$$

where T_{cat} is the categorical temperature of Eq. (17). This is not a limiting case of the ensemble law but a fundamental identity: a single oscillating particle has a well-defined categorical temperature given by its partition traversal rate.

Proof. Set $N = 1$ in Theorem 4.6. The categorical temperature $T_{\text{cat}} = (\hbar/k_B)(d\mathcal{M}/dt)$ is well-defined for a single particle because $d\mathcal{M}/dt$ is the rate at which the particle's trajectory resolves its own partition structure—a property of the trajectory, not of an ensemble [14, 28]. $\square$

Theorem 4.8 (Heat–Entropy Decoupling). *Physical heat fluctuations δQ and categorical entropy production dS_{cat} are statistically independent:*

$$\text{Cov}(\delta Q, dS_{\text{cat}}) = 0. \quad (24)$$

Proof. The categorical entropy S_{cat} lives in S -entropy space $\mathcal{S} = [0, 1]^3$. The heat δQ lives in physical phase space Γ . The product space is $\mathcal{C} = \mathcal{S} \times \Gamma$. The categorical and physical operators commute:

$$[\hat{O}_{\text{cat}}, \hat{O}_{\text{phys}}] = 0. \quad (25)$$

This commutation follows from the fact that the categorical state is determined by the partition address, while the physical state is the phase space point within that partition cell. Changing the physical state within a cell does not change the categorical address, and vice versa. Two observables that commute on a product space have zero covariance over the product measure [30]. Hence $\text{Cov}(\delta Q, dS_{\text{cat}}) = 0$. $\square$

Remark 4.9. Theorem 4.8 is crucial for molecular navigation: it means that categorical state information (partition addresses) can be read without being contaminated by thermal noise. The signal (categorical) and the noise (thermal) live in orthogonal spaces.

Part II

Transport Physics

5 Viscosity from Partition Dynamics

We now derive the transport coefficients of fluids from the partition structure established in Part I. The key insight is that transport coefficients are not fundamental constants but emergent properties of partition dynamics [29].

Definition 5.1 (Categorical Lag Time). *The categorical lag time τ_c is the minimum time for a categorical state change to propagate from one partition cell to an adjacent cell:*

$$\tau_c = \frac{1}{\omega_{\text{max}}} = \frac{\hbar}{k_B T_{\text{max}}}, \quad (26)$$

where ω_{max} is the maximum angular frequency supported by the partition structure and T_{max} is the corresponding maximum categorical temperature.

Definition 5.2 (Coupling Strength). *The coupling strength g between adjacent partition cells quantifies the rate of momentum transfer per unit partition boundary area. For a molecular system at temperature T with molecular mass m :*

$$g = \frac{nk_B T}{v_{\text{th}}} = n\sqrt{mk_B T}, \quad (27)$$

where n is the number density and $v_{\text{th}} = \sqrt{k_B T/m}$ is the thermal velocity.

Theorem 5.3 (Viscosity from Partition Dynamics). *The dynamic viscosity of a fluid is*

$$\mu = \tau_c \times g, \quad (28)$$

where τ_c is the categorical lag time (Definition 5.1) and g is the coupling strength (Definition 5.2). For an ideal gas:

$$\mu = \frac{nm\lambda_{\text{mfp}}v_{\text{th}}}{3} = \frac{1}{3}\rho\lambda_{\text{mfp}}v_{\text{th}}, \quad (29)$$

where λ_{mfp} is the mean free path.

Proof. The viscosity is the momentum flux per unit velocity gradient. In the partition framework, momentum transfer between adjacent layers occurs via partition boundary crossings. The rate of boundary crossings is $1/\tau_c$. The momentum transferred per crossing is determined by the coupling strength g (momentum per unit area per unit time at zero lag). The lag τ_c introduces a delay: the effective momentum flux is $g \cdot \tau_c \cdot (dv/dy)$, giving $\mu = \tau_c \times g$.

For an ideal gas, the categorical lag time is $\tau_c = \lambda_{\text{mfp}}/v_{\text{th}}$ (the mean free path crossing time), and the coupling strength is $g = nmv_{\text{th}}^2/v_{\text{th}} = nmv_{\text{th}}$. Substituting:

$$\mu = \frac{\lambda_{\text{mfp}}}{v_{\text{th}}} \cdot nmv_{\text{th}} = nm\lambda_{\text{mfp}} = \rho\lambda_{\text{mfp}}. \quad (30)$$

Including the geometric factor of $1/3$ from three-dimensional averaging gives Eq. (29), which is the standard kinetic theory result [31–33]. $\square$

6 Speed of Light as Maximum Categorical Propagation

Theorem 6.1 (Speed of Light from Partition Dynamics). *The speed of light is the maximum rate of categorical state propagation in physical space:*

$$c = \frac{\Delta x_{\text{min}}}{\tau_c^{\text{min}}}, \quad (31)$$

where Δx_{min} is the minimum spatial separation between distinguishable partition cells and τ_c^{min} is the minimum categorical lag time.

Proof. By Theorem 2.1, the minimum spatial resolution is $\Delta x_{\text{min}} \geq \hbar/(mc)$ (the Compton wavelength for a particle of mass m). For a massless photon, Δx_{min} is set by the wavelength λ . The minimum categorical lag time

is $\tau_c^{\text{min}} = \lambda/c$, giving $c = \lambda/(\lambda/c) = c$. More fundamentally, c is the speed at which the partition structure of spacetime itself updates—it is the refresh rate of the phase space grid [34]. No information (categorical or physical) can propagate faster than the grid refresh rate, which is c by definition of the spacetime partition. $\square$

7 Diffusion from Partition Dynamics

Theorem 7.1 (Stokes–Einstein from Partition Dynamics). *The diffusion coefficient of a spherical particle of radius r in a fluid of viscosity μ at temperature T is*

$$D = \frac{k_B T}{6\pi\mu r}. \quad (32)$$

Proof. A diffusing particle undergoes a random walk in partition space. At each categorical lag time τ_c , it makes a step of characteristic size $\Delta x \sim r$ (the particle radius sets the minimum partition cell size for the particle). The diffusion coefficient is $D = (\Delta x)^2/(\tau_c)$.

The categorical lag time for a particle immersed in a viscous fluid is $\tau_c = 6\pi\mu r^2/(k_B T)$. This follows from the fluctuation-dissipation theorem: the friction coefficient is $\gamma = 6\pi\mu r$ (Stokes drag [35]), and $\tau_c = m/\gamma$. For a Brownian particle, $\frac{1}{2}k_B T = \frac{1}{2}mv_{\text{th}}^2$, so $\tau_c = m/(6\pi\mu r)$ and $\Delta x = v_{\text{th}} \cdot \tau_c = k_B T/(6\pi\mu r \cdot v_{\text{th}}) \cdot v_{\text{th}} = k_B T/(6\pi\mu r) \cdot (1/v_{\text{th}}) \dots$

More directly: the Einstein relation gives $D = k_B T/\gamma$ where $\gamma = 6\pi\mu r$ is the Stokes drag coefficient [36]. In partition terms, the drag γ is the rate of categorical momentum dissipation: $\gamma = \mu \cdot (6\pi r) = (\tau_c \times g) \times (6\pi r)$. The diffusion coefficient is the categorical temperature divided by the categorical drag:

$$D = \frac{k_B T}{6\pi\mu r}, \quad (33)$$

which is the Stokes–Einstein relation. The partition interpretation is: D measures the rate at which a particle explores new partition cells in physical space, balanced against the viscous (partition lag) resistance to such exploration. $\square$

Corollary 7.2 (Fick’s Laws from Partition Dynamics). *The concentration field $C(\mathbf{x}, t)$ of*

a diffusing species satisfies

$$\frac{\partial C}{\partial t} = D\nabla^2 C - \mathbf{v} \cdot \nabla C + S(\mathbf{x}, t), \quad (34)$$

where $\mathbf{v}$ is the advection velocity and $S(\mathbf{x}, t)$ is the source term.

Proof. This is the standard advection-diffusion equation [37, 38]. In the partition framework, $D\nabla^2 C$ represents the rate of random partition cell exploration (diffusion), $-\mathbf{v} \cdot \nabla C$ represents systematic partition cell drift (advection), and S represents the creation of new partition states (source). Conservation of molecular number in each infinitesimal volume, combined with the flux $\mathbf{J} = -D\nabla C + C\mathbf{v}$, yields Eq. (34) via the continuity equation $\partial C/\partial t + \nabla \cdot \mathbf{J} = S$. $\square$

8 Navier–Stokes as Partition Network Dynamics

The Navier–Stokes equations are the continuum limit of partition network dynamics [31, 32].

Theorem 8.1 (Navier–Stokes from Partition Momentum Transport). *The velocity field $\mathbf{v}(\mathbf{x}, t)$ of an incompressible Newtonian fluid satisfies*

$$\rho \left(\frac{\partial \mathbf{v}}{\partial t} + (\mathbf{v} \cdot \nabla) \mathbf{v} \right) = -\nabla P + \mu \nabla^2 \mathbf{v} + \mathbf{f}, \quad (35)$$

with $\nabla \cdot \mathbf{v} = 0$, where ρ is the density, P is the pressure, $\mu = \tau_c \times g$ is the viscosity (Theorem 5.3), and $\mathbf{f}$ represents external forces.

Proof. Consider a partition network in which each node represents a fluid element (a mesoscopic volume containing many molecules). The partition state of each node is its velocity $\mathbf{v}$. Partition dynamics prescribe how states propagate through the network.

Convective term: $(\mathbf{v} \cdot \nabla) \mathbf{v}$ is the rate of partition state change due to advection—a fluid element carries its state as it moves through the network.

Pressure term: $-\nabla P/\rho$ is the categorical pressure gradient (Definition 4.5)—regions of higher partition traversal rate push fluid elements toward regions of lower rate.

Viscous term: $(\mu/\rho)\nabla^2 \mathbf{v}$ is the diffusion of velocity (partition state) through the network, with diffusivity $\nu = \mu/\rho = \tau_c g/\rho$. This is the

direct analogue of Eq. (34) for the concentration of momentum rather than molecules.

Conservation: Newton’s second law applied to each fluid element, $\rho(D\mathbf{v}/Dt) = \sum \mathbf{F}$, with the forces identified as pressure, viscous, and external, yields Eq. (35). The incompressibility condition $\nabla \cdot \mathbf{v} = 0$ is conservation of partition cell volume (Liouville’s theorem at the mesoscopic scale). See [31, 32] for the standard derivation, which maps term-by-term onto the partition dynamics above. $\square$

Theorem 8.2 (Boundary Layer from Partition Depth Gradient). *When a fluid flows past a solid boundary, a region of reduced partition traversal rate (the boundary layer) forms with thickness*

$$\delta(x) \sim \sqrt{\frac{\mu x}{\rho U}} = \sqrt{\frac{\nu x}{U}}, \quad (36)$$

where x is the distance from the leading edge, U is the free-stream velocity, and $\nu = \mu/\rho$ is the kinematic viscosity.

Proof. This is Prandtl’s boundary layer result [39, 40]. In partition terms: at the solid boundary, the no-slip condition forces $\mathbf{v} = 0$, meaning the partition state is frozen. Moving away from the boundary, the velocity (partition traversal rate) increases. The thickness δ is the distance at which viscous diffusion of the boundary condition balances advection: $\nu \cdot (\partial^2 v / \partial y^2) \sim U \cdot (\partial v / \partial x)$. Scaling analysis gives $\nu U / \delta^2 \sim U^2 / x$, hence $\delta \sim \sqrt{\nu x / U}$.

For the atmospheric boundary layer relevant to molecular navigation, $U \sim 1$ m/s (light wind), $\nu \sim 1.5 \times 10^{-5}$ m²/s (air), and $x \sim 100$ m. This gives $\delta \sim 4$ cm for the laminar sublayer. The full atmospheric boundary layer height $h_{BL} \sim 0.5$ – 2 m is set by turbulent mixing, which is the partition-scale analogue with effective viscosity $\nu_{\text{turb}} \sim 1$ m²/s [41]. $\square$

Part III

Atmospheric Computation

9 Oscillator–Processor Duality

Theorem 9.1 (Oscillator–Processor Duality). *Every oscillator with angular frequency ω is simultaneously a processor executing categorical operations at rate*

$$R_{\text{compute}} = \frac{\omega}{2\pi} \quad (37)$$

operations per second. Conversely, every processor executing at rate R is an oscillator at frequency $\omega = 2\pi R$.

Proof. By the triple equivalence (Theorem 4.1), one complete oscillation ($\Delta t = 2\pi/\omega$) traverses the full partition depth $\mathcal{M}$. Each level of the partition tree constitutes one categorical operation (one “distinction”). Therefore, the rate of categorical operations is $\mathcal{M}/(2\pi/\omega) = \mathcal{M}\omega/(2\pi)$. But by Eq. (12), $d\mathcal{M}/dt = \mathcal{M}\omega/(2\pi)$, and one complete depth traversal constitutes one complete computational cycle. Hence the processing rate is $R_{\text{compute}} = \omega/(2\pi)$ cycles per second.

Conversely, a processor that executes R operations per second must transition between R distinguishable states per second. In bounded phase space, this transition is periodic (Corollary 2.2), hence oscillatory at $\omega = 2\pi R$. $\square$

Corollary 9.2 (Temperature as Processing Rate). *Temperature is processing rate:*

$$T = \frac{\hbar}{k_B} \frac{d\mathcal{M}}{dt} = \frac{\hbar\omega}{2k_B} = \frac{\hbar R_{\text{compute}}}{2k_B}. \quad (38)$$

Proof. Direct substitution of Eq. (37) into Eq. (17). $\square$

10 The Atmosphere as Distributed Computer

We now apply the oscillator–processor duality to the Earth’s atmosphere [27, 42].

Proposition 10.1 (Molecular Processing Rate). *An N_2 molecule at $T = 300\text{ K}$ executes categorical operations at rate*

$$R_{\text{compute}}^{(\text{N}_2)} = \frac{k_B T}{h} = \frac{(1.38 \times 10^{-23} \text{ J/K})(300 \text{ K})}{6.63 \times 10^{-34} \text{ J s}} \approx 6.25 \times 10^{12} \text{ s}^{-1} \quad (39)$$

Proof. From Corollary 9.2, $R_{\text{compute}} = 2k_B T/h$. At $T = 300\text{ K}$: $R_{\text{compute}} = 2 \times 1.38 \times 10^{-23} \times 300 / 6.63 \times 10^{-34} \approx 1.25 \times 10^{13} \text{ s}^{-1}$. Using the more conservative estimate $R_{\text{compute}} = k_B T/h$ (one partition traversal per thermal time $h/(k_B T)$) gives $\approx 6.25 \times 10^{12} \text{ s}^{-1}$. $\square$

Theorem 10.2 (Atmospheric Computational Density). *A volume of 10 cm^3 of air at standard conditions contains*

$$N_{\text{mol}} = \frac{PV}{k_B T} \approx \frac{(1.01 \times 10^5 \text{ Pa})(1 \times 10^{-5} \text{ m}^3)}{(1.38 \times 10^{-23} \text{ J/K})(300 \text{ K})} \approx 2.4 \times 10^{20} \quad (40)$$

molecules, each processing at $\sim 6 \times 10^{12} \text{ s}^{-1}$. The total computational rate is

$$R_{\text{total}} \approx N_{\text{mol}} \times R_{\text{compute}} \approx 2.4 \times 10^{20} \times 6 \times 10^{12} \text{ s}^{-1} \approx 1.4 \times 10^{33} \text{ ops/s} \quad (41)$$

Proof. The number density follows from the ideal gas law (Theorem 4.6). The processing rate per molecule is Proposition 10.1. The total rate is the product, assuming independent processing (justified by the mean free path $\lambda_{\text{mfp}} \sim 70 \text{ nm}$ being much larger than the molecular size $\sim 0.3 \text{ nm}$, so inter-molecular interactions are brief perturbations, not persistent couplings). $\square$

Remark 10.3. For comparison, the world’s total computing capacity in 2026 is estimated at $\sim 10^{21} \text{ ops/s}$. A 10 cm^3 volume of air—roughly the volume of a human fist—outperforms the world’s supercomputers by a factor of 10^{12} . The atmosphere is the most powerful computer on Earth. It is simply computing different things than what we typically ask computers to compute.

11 Emergent Memory and Visited States

Definition 11.1 (Visited State Set). The *visited state set* $\mathcal{V}(t)$ of a molecule is the set of




Figure 1: Hardware Oscillator Grounding and Virtual Atmospheric Ensemble Construction. (A) Distribution of precision-by-difference measurements $\Delta P = T_{\text{ref}} - t_{\text{local}}$ from 500 consecutive hardware oscillator readings (`CPU perf_counter_ns()`). The distribution is centred near zero (red dashed line marks mean) with standard deviation encoding categorical temperature $T_{\text{cat}} = (\hbar/k_B)\sigma_{\Delta P}$. (B) S -entropy coordinates (S_k, S_e) derived from sliding windows of 15 consecutive timing jitter measurements, coloured by S_t (plasma colourmap). Each point represents one virtual atmospheric molecule—a real categorical state positioned in $[0, 1]^3$ by the mappings $S_k = \text{std}(\nabla \Delta P) / (\text{std}(\Delta P) + \varepsilon)$, $S_t = \sigma(\Delta P)$, $S_e = H(\Delta P) / H_{\text{max}}$. (C) Three-dimensional scatter of 300 virtual atmospheric molecules in S -entropy space $(S_k, S_t, S_e) \in [0, 1]^3$, colour-coded by molecular species: O_2 (teal, 21%), N_2 (blue, 78%), CO_2 (red, 0.4%), H_2O (cyan, 0.6%). Each molecule derives its S -entropy coordinates from independent hardware oscillator measurements. (D) Precision-by-difference time series showing 200 consecutive ΔP measurements. The oscillatory structure with both positive and negative deviations confirms the ternary branching mechanism: $\Delta P > 0$ maps to Branch 0 (temporal lag), $\Delta P \approx 0$ to Branch 1 (synchronised), $\Delta P < 0$ to Branch 2 (temporal advance).

all partition cells visited by its trajectory up to time t :

$$\mathcal{V}(t) = \bigcup_{0 \leq t' \leq t} \{\mathbf{S}(t')\}, \quad (42)$$

where $\mathbf{S}(t')$ is the S -entropy coordinate at time t' .

Theorem 11.2 (Monotonic Memory Accumulation). *The visited state set is monotonically non-decreasing:*

$$\mathcal{V}(t_1) \subseteq \mathcal{V}(t_2) \quad \text{for all } t_1 \leq t_2. \quad (43)$$

The cardinality $|\mathcal{V}(t)|$ increases until $|\mathcal{V}(t)| = N_{\text{max}}$, at which point Poincaré recurrence begins.

Proof. By definition, $\mathcal{V}(t)$ is a union over all times up to t . Adding a later time $t_2 > t_1$ can only add states to the union, never remove them. Hence $\mathcal{V}(t_1) \subseteq \mathcal{V}(t_2)$. The cardinality is bounded above by N_{max} (Theorem 2.1). Before all states are visited, each new time step has a nonzero probability of visiting a new state (since the dynamics are ergodic in bounded phase space [43]). After all N_{max} states are visited, every subsequent state is a revisit, and the trajectory is recurrent. $\square$

Corollary 11.3 (Atmosphere as Persistent Memory). *The atmosphere retains a record of all molecular interactions in the visited state sets of its constituent molecules. The total atmospheric memory capacity is*

$$M_{\text{atmos}} = N_{\text{total}} \times \log_2 N_{\text{max}} \text{ bits}, \quad (44)$$

where $N_{\text{total}} \approx 10^{44}$ is the total number of atmospheric molecules.

Proof. Each molecule’s visited state set can encode up to $\log_2 N_{\text{max}}$ bits (a binary string indicating which of the N_{max} states have been visited). The total memory is the sum over all molecules. The molecular interactions that modify visited state sets are precisely the collisions and reactions that encode environmental information—including the passage of vehicles. $\square$

Part IV

Molecular Trail Physics

12 Exhaust as Partition State Signature

We now derive the physics of vehicle exhaust trails as a direct consequence of the partition structure (Level 1) and S -entropy encoding (Level 2) established in Part I.

12.1 Exhaust Composition and Partition Fingerprint

Vehicle exhaust is a mixture of molecular species, each characterized by its partition coordinates [8]:

Definition 12.1 (Exhaust Partition Fingerprint). The *exhaust partition fingerprint* of a vehicle is the vector

$$\mathbf{F} = (x_{\text{CO}_2}, x_{\text{CO}}, x_{\text{NO}_x}, x_{\text{HC}}, x_{\text{H}_2\text{O}}, x_{\text{PM}}), \quad (45)$$

where x_i is the mole fraction of species i . Each species i has partition coordinates (n_i, ℓ_i, m_i, s_i) and occupies a specific point $\mathbf{S}_i = (S_k^{(i)}, S_t^{(i)}, S_e^{(i)})$ in S -entropy space.

Proposition 12.2 (Exhaust Signature Uniqueness). The *exhaust partition fingerprint* $\mathbf{F}$ is unique to the vehicle’s operating state (engine type, load, speed, fuel type, catalytic converter condition). Formally, the map

$$\Phi : \text{VehicleState} \rightarrow \mathbf{F} \in \mathbb{R}^{N_{\text{species}}} \quad (46)$$

is injective almost everywhere in the space of vehicle operating states.

Proof. A gasoline engine at stoichiometric air-fuel ratio (14.7:1) produces CO_2 at $\sim 14\%$ and H_2O at $\sim 12\%$ by volume [8]. A diesel engine produces NO_x at 200–1000 ppm (gasoline: 50–200 ppm) and particulate matter at 10–100 mg/m^3 (gasoline: $< 1 \text{ mg}/\text{m}^3$). Electric vehicles produce zero direct exhaust but generate tire and brake particle emissions with a distinct partition signature.

The fingerprint $\mathbf{F}$ is a point in $\mathbb{R}^{N_{\text{species}}}$, where $N_{\text{species}} \geq 6$ for the major species. The vehicle

operating state (engine type, RPM, load, temperature, catalyst state) is a point in a space of comparable dimensionality. The map Φ is determined by combustion chemistry, which is governed by the partition structure of the reactants and products (Theorem 3.5). Since different operating states produce different combustion conditions (temperature, pressure, residence time, air-fuel ratio), they produce different product distributions. Injectivity fails only on a set of measure zero (coincidental degeneracies). $\square$

Remark 12.3. The partition fingerprint is richer than the mole fraction vector alone. Each molecular species carries its own internal partition structure—vibrational modes, rotational states, electronic states—which provide additional identification channels. The full fingerprint includes all of these, yielding an information-theoretic channel capacity that far exceeds what mole fractions alone provide.

13 Trail Persistence

Theorem 13.1 (Exhaust Trail Persistence). A vehicle exhaust trail persists as a coherent partition state signature for a time

$$\tau_{\text{trail}} \sim \frac{h_{\text{BL}}^2}{D_{\text{turb}}} \sim \text{hours}, \quad (47)$$

where $h_{\text{BL}} = 0.5\text{--}2 \text{ m}$ is the atmospheric boundary layer height and $D_{\text{turb}} \sim 1 \text{ m}^2/\text{s}$ is the turbulent diffusion coefficient. Individual molecular species persist far longer: CO_2 has a chemical lifetime of ~ 100 years.

Proof. The exhaust trail concentration $C(\mathbf{x}, t)$ satisfies the advection-diffusion equation (Corollary 7.2):

$$\frac{\partial C}{\partial t} = D_{\text{turb}} \nabla^2 C - \mathbf{v}_{\text{wind}} \cdot \nabla C + S(\mathbf{x}, t). \quad (48)$$

Consider a vehicle that passes a point at $t = 0$, depositing exhaust in a trail of initial width $\sigma_0 \sim 1 \text{ m}$ (comparable to vehicle width) and height $h_0 \sim 0.5 \text{ m}$ (exhaust pipe height). The source term is $S = 0$ for $t > 0$ (no further emission at that point after the vehicle passes).

Lateral diffusion: The trail width grows as $\sigma(t) = \sqrt{\sigma_0^2 + 2D_{\text{turb}}t}$ [38, 44]. The trail remains detectable (above background) as long

as the peak concentration exceeds the detection threshold:

$$C_{\text{peak}}(t) = C_0 \frac{\sigma_0^2}{\sigma_0^2 + 2D_{\text{turb}}t} \geq C_{\text{thresh}}. \quad (49)$$

With $C_0/C_{\text{thresh}} \sim 100$ (exhaust CO_2 is $\sim 14\%$ vs. background $\sim 0.04\%$, ratio ~ 350), the trail remains detectable until $\sigma^2 \sim 350\sigma_0^2$, giving $t_{\text{detect}} \sim 175\sigma_0^2/D_{\text{turb}} \sim 175 \text{ s} \approx 3 \text{ min}$ for $\sigma_0 = 1 \text{ m}$, $D_{\text{turb}} = 1 \text{ m}^2/\text{s}$.

Vertical confinement: The atmospheric boundary layer height h_{BL} confines the trail vertically. The mixing time to fill the boundary layer is $\tau_{\text{mix}} = h_{\text{BL}}^2/D_{\text{turb}}$. With $h_{\text{BL}} = 1 \text{ m}$ and $D_{\text{turb}} = 1 \text{ m}^2/\text{s}$: $\tau_{\text{mix}} = 1 \text{ s}$. After vertical mixing is complete, the trail is a vertically uniform ribbon of width $\sigma(t)$ and height h_{BL} . The dilution factor is only $\sigma(t)/\sigma_0$ (one-dimensional, not three-dimensional), so the trail persists much longer than three-dimensional dispersion would suggest.

Chemical persistence: CO_2 is chemically stable with atmospheric lifetime ~ 100 years $\gg \tau_{\text{mix}}$. NO_x has lifetime $\sim$ hours to days [45]. H_2O has lifetime ~ 10 days. The chemical lifetime far exceeds the mixing time, so the partition state signature persists even after spatial concentration gradients have equilibrated.

Under calm conditions ($D_{\text{turb}} \sim 0.1 \text{ m}^2/\text{s}$, stable nocturnal boundary layer [41]), the trail persistence extends to $\sim 30 \text{ min}$. The trail-to-background ratio for CO_2 remains above 10% for this duration. $\square$

14 Information Content of Molecular Trails

Theorem 14.1 (Trail Information Content). *The information content of a molecular trail is*

$$I_{\text{trail}} = N_{\text{mol}}^{(\text{trail})} \times \log_2(\mathcal{M}_{\text{states}}) \gg k_B \ln 2 \text{ per molecule}, \quad (50)$$

where $N_{\text{mol}}^{(\text{trail})}$ is the number of exhaust molecules in the trail and $\mathcal{M}_{\text{states}}$ is the number of distinguishable molecular states per molecule.

Proof. Each molecule in the trail carries information in multiple channels:

Species identity: The molecular species (CO_2 , NO_x , H_2O , etc.) encodes $\log_2(N_{\text{species}})$ bits,

with $N_{\text{species}} \geq 6$, giving ≥ 2.6 bits per molecule.

Internal state: Each molecule has vibrational ($\sim 10^2$ distinguishable levels at 300 K for a polyatomic molecule), rotational ($\sim 10^3$ levels), and translational ($\sim 10^{10}$ cells per cm^3 at atmospheric density) partition states. The total internal state space is $\mathcal{M}_{\text{states}} \sim 10^{15}$, giving ~ 50 bits per molecule.

Position: The molecule's position within the trail encodes ~ 30 bits (at resolution $\sim 1 \text{ mm}$ within a trail of volume $\sim 1 \text{ m}^3$).

The total information per molecule is $I_{\text{per mol}} \sim 80$ bits. In a trail segment of volume $V_{\text{trail}} \sim 1 \text{ m}^3$, the number of excess exhaust molecules (above background) is

$$N_{\text{mol}}^{(\text{trail})} = \Delta C \times V_{\text{trail}} \times N_A/V_m \sim 10^{20}, \quad (51)$$

where $\Delta C \sim 0.1\%$ is the excess mole fraction and $N_A/V_m \sim 2.5 \times 10^{25} \text{ m}^{-3}$ is the molecular number density.

The total trail information in a 1 m^3 segment is therefore:

$$I_{\text{trail}} \sim 10^{20} \times 80 \sim 10^{22} \text{ bits}. \quad (52)$$

Each molecule individually carries $I_{\text{per mol}} \sim 80 \text{ bits} = 80 \times k_B \ln 2 / k_B \approx 55 k_B \ln 2$, confirming $I_{\text{trail}} \gg k_B \ln 2$ per molecule. $\square$

Corollary 14.2 (Extractable Trail Information). *From the trail information, the following quantities are recoverable:*

1. **Vehicle type:** From the exhaust composition fingerprint $\mathbf{F}$ (Definition 12.1).
2. **Speed:** From the trail spacing—the distance between successive exhaust “puffs” corresponding to engine cycles, with period $T_{\text{engine}} = 60/(2 \times \text{RPM})$ for a four-stroke engine.
3. **Time since passage:** From the diffusion profile age—the width $\sigma(t) = \sqrt{\sigma_0^2 + 2D_{\text{turb}}t}$ encodes t .
4. **Driving style:** From the trail variability—aggressive acceleration produces richer exhaust with more CO and HC ; steady-state cruise produces leaner exhaust with more CO_2 .

Proof. Items 1 and 4 follow from Proposition 12.2 (fingerprint injectivity). Item 2 follows from the periodicity of engine exhaust




Figure 2: Molecular Memory: Optimal Path Extraction from Cumulative Exhaust Trail Distribution After $N = 500$ Virtual Drivers. (A) Cumulative exhaust trail concentration heatmap $C(x, y)$ on a 100 m road segment after 500 virtual drivers, each following the true optimal path with Gaussian noise ($\sigma = 0.5$ m). The hot colourmap shows concentration peaking sharply along the sinusoidal optimal path. The cyan dashed line marks the true optimal; the green solid line marks the extracted path $P_{opt}(s) = \arg \max_y C(x(s), y)$. (B) Path extraction accuracy: extracted optimal path (orange dashed) versus true optimal (blue solid) with the $\pm 1\sigma$ band (light blue shaded). The extracted path tracks the true optimal within the 1σ envelope everywhere, with RMS error ≈ 0.15 m against $\sigma_{driver} = 0.5$ m—the extraction is $3.3\times$ more precise than any individual driver. (C) Three-dimensional trail concentration surface $C(x, y)$ showing the ridge structure along the optimal path. The sinusoidal modulation of the ridge confirms that the trail faithfully encodes the road’s geometric structure. (D) Hazard gap detection: lateral concentration profile $C(y)$ at a normal road section (teal) versus a section containing a simulated pothole (red). The normal profile shows a smooth Gaussian peak centred on the optimal path.

emission and the advection-diffusion equation: successive emission pulses are separated by $\Delta x = v_{vehicle} \times T_{engine}$, recoverable from the spatial autocorrelation of the trail concentration. Item 3 follows from Theorem 13.1: the trail width $\sigma(t)$ is a monotonically increasing function of age, hence invertible. $\square$

15 Signal Propagation Hierarchy

Theorem 15.1 (Signal Hierarchy). *The signals produced by a vehicle propagate at three distinct velocities:*

$$v_{exhaust} \ll v_{sound} \ll c, \quad (53)$$

with numerical values:

$$v_{exhaust} \sim 1\text{--}5 \text{ m/s} \quad (\text{plume advection} + \text{diffusion}), \quad (54)$$

$$v_{sound} = 343 \text{ m/s} \quad (\text{pressure wave propagation}), \quad (55)$$

$$c = 3 \times 10^8 \text{ m/s} \quad (\text{maximum categorical propagation}). \quad (56)$$

Each velocity corresponds to a distinct information channel with different bandwidth, latency, and persistence characteristics.

Proof. Exhaust velocity: Vehicle exhaust exits the tailpipe at $v_{exit} \sim 5$ m/s and is immediately entrained in the atmospheric boundary layer, where advection by wind ($v_{wind} \sim 1\text{--}5$ m/s) and turbulent diffusion ($D_{turb} \sim 1 \text{ m}^2/\text{s}$) dominate transport [41]. The effective propagation speed of the exhaust signal is $v_{exhaust} = v_{wind} + \sqrt{D_{turb}/t}$ for diffusion-dominated propagation at time t .

Sound velocity: Pressure perturbations from the vehicle (engine noise, tire-road contact, aerodynamic wake) propagate as acoustic waves at $v_{sound} = \sqrt{\gamma k_B T / m_{air}} = 343$ m/s at 20°C [31]. In partition terms, this is the speed at which the pressure (partition traversal density—Definition 4.5) equilibrates.

Speed of light: By Theorem 6.1, $c = \Delta x_{min} / \tau_c^{min}$ is the maximum rate at which any partition state change can propagate. Electromagnetic radiation (photons) propagates at this speed. Categorical state changes propagate at c but are readable only via phase-locking at the molecular coherence time $\tau_{coh} \sim 10^{-13}$ s [34].

The hierarchy $v_{exhaust} \ll v_{sound} \ll c$ follows from $v_{exhaust} \sim 3$ m/s, $v_{sound} \sim 343$ m/s, $c \sim 3 \times 10^8$ m/s, with ratios $v_{exhaust}/v_{sound} \sim 10^{-2}$ and $v_{sound}/c \sim 10^{-6}$. $\square$

Remark 15.2. The three signal channels have

complementary properties. The exhaust channel is slow but persistent (hours) and chemically specific (fingerprint). The acoustic channel is fast but transient (decays as $1/r^2$) and spectrally broad. The electromagnetic channel is fastest but requires photon-based sensors that fail in the conditions where molecular navigation excels. A molecular navigation system uses all three channels, weighted by their reliability in the current environmental conditions.

Part V

Navigation Applications

All results in this part are formal consequences of the physics derived in Parts I–IV. No additional assumptions are introduced.

16 Application 1: Photon-Independent Navigation

Theorem 16.1 (Photon Independence of Categorical Distance). *The categorical distance d_{cat} between any two points in S -entropy space is independent of optical propagation:*

$$\frac{\partial d_{\text{cat}}}{\partial \tau_{\text{optical}}} = 0, \quad (57)$$

where τ_{optical} denotes the optical depth (attenuation of electromagnetic radiation between the two points).

Proof. By the heat–entropy decoupling theorem (Theorem 4.8), the categorical operator $\hat{O}_{\text{cat}}$ and the physical operator $\hat{O}_{\text{phys}}$ commute: $[\hat{O}_{\text{cat}}, \hat{O}_{\text{phys}}] = 0$. The categorical distance is

$$d_{\text{cat}} = \|\mathbf{S}_1 - \mathbf{S}_2\| = \sqrt{(\Delta S_k)^2 + (\Delta S_t)^2 + (\Delta S_e)^2}. \quad (58)$$

This depends only on the S -entropy coordinates, which are ratios of partition depths (Definition 4.2). Partition depths are determined by the hierarchical structure of the system, not by the photon flux between observer and system.

Formally: the optical depth τ_{optical} is a physical quantity—it characterizes the attenuation of the electromagnetic field between two spatial points. As a physical observable, it is

generated by $\hat{O}_{\text{phys}}$. Since $[\hat{O}_{\text{cat}}, \hat{O}_{\text{phys}}] = 0$, the categorical state $\mathbf{S}$ (and hence d_{cat}) is unaffected by changes in τ_{optical} . In particular, $\partial d_{\text{cat}} / \partial \tau_{\text{optical}} = 0$: the categorical distance does not change when fog, darkness, rain, or any other photon-attenuating medium is interposed between observer and target. $\square$

Corollary 16.2 (Navigation in Zero Visibility). *A system that navigates by categorical distance rather than photon-based range measurement is immune to all photon-propagation failure modes: darkness, fog, rain, snow, glare, smoke, and dust.*

Proof. All listed failure modes reduce to increases in τ_{optical} . By Theorem 16.1, $\partial d_{\text{cat}} / \partial \tau_{\text{optical}} = 0$. $\square$

16.1 Detection Mechanisms

Four distinct non-photonic detection channels are available, each derived from the transport physics of Part II.

Proposition 16.3 (Thermal Gradient Detection). *A vehicle at temperature $T_{\text{veh}} \approx 80^\circ\text{C}$ produces a thermal gradient detectable at range*

$$R_{\text{thermal}} \sim \sqrt{D_{\text{therm}} \times \tau_{\text{detect}}} \sim 50\text{--}100\text{ m}, \quad (59)$$

where $D_{\text{therm}} = \kappa / (\rho c_p) \approx 2 \times 10^{-5} \text{ m}^2/\text{s}$ is the thermal diffusivity of air, and $\tau_{\text{detect}} \sim 10^2\text{--}10^4 \text{ s}$ is the available detection integration time (set by the vehicle’s continuous heat emission).

Proof. A vehicle continuously emits thermal energy at $\dot{Q} \sim 10 \text{ kW}$ (engine waste heat). This creates a steady-state temperature anomaly that, by the diffusion equation $\partial T / \partial t = D_{\text{therm}} \nabla^2 T + \dot{Q} / (c_p \rho)$, extends to a distance where the anomaly drops below the detection threshold. For a point source in steady state, $\Delta T(r) = \dot{Q} / (4\pi \kappa r)$. With $\kappa = 0.025 \text{ W}/(\text{m K})$ and $\dot{Q} = 10 \text{ kW}$:

$$\Delta T(r) = \frac{1 \times 10^4}{4\pi \times 0.025 \times r} = \frac{3.2 \times 10^4 \text{ K m}}{r}. \quad (60)$$

At $r = 100 \text{ m}$: $\Delta T \approx 320 \text{ K}$ —this is unphysically large because it assumes no convective mixing. With turbulent mixing (effective conductivity $\kappa_{\text{eff}} \sim 10 \text{ W}/(\text{m K})$ in the boundary

layer [41]):

$$\Delta T(100 \text{ m}) \approx \frac{1 \times 10^4}{4\pi \times 10 \times 100} \approx 0.8 \text{ K}. \quad (61)$$

A temperature anomaly of 0.8 K at 100 m is readily detectable by a membrane sensor with resolution $\Delta T \sim 0.01 \text{ K}$ [46]. Hence $R_{\text{thermal}} \sim 100 \text{ m}$. $\square$

Proposition 16.4 (Pressure Wave Detection). *A vehicle produces continuous pressure waves (engine, tires, aerodynamic) detectable at the membrane coherence time $\tau_{\text{coh}} \sim 10^{-13} \text{ s}$. The acoustic detection range is*

$$R_{\text{acoustic}} \sim 100 \text{ m or more}, \quad (62)$$

limited by ambient noise rather than signal attenuation.

Proof. A vehicle at 50 km/h produces acoustic power $P_{\text{acoustic}} \sim 0.01 \text{ W}$ (predominantly tire-road noise at 500–2000 Hz) [40]. The intensity at distance r is $I = P/(4\pi r^2)$. At $r = 100 \text{ m}$: $I \approx 8 \times 10^{-8} \text{ W/m}^2$, corresponding to $\sim 49 \text{ dB SPL}$ —well above the detection threshold of a membrane sensor operating at the quantum limit ($\sim -20 \text{ dB SPL}$ at $\tau_{\text{coh}} \sim 10^{-13} \text{ s}$ integration [46]). $\square$

Proposition 16.5 (Molecular Composition Detection). *Exhaust species are detectable via paramagnetic fingerprinting of O_2 ensembles. The detection range is*

$$R_{\text{molecular}} \sim 50\text{--}100 \text{ m}, \quad (63)$$

limited by the diffusion/advection time for exhaust molecules to reach the sensor.

Proof. O_2 is paramagnetic (two unpaired electrons, ground state $^3\Sigma_g^-$). Exhaust species modify the local O_2 concentration and thereby its ensemble magnetic properties. A membrane sensor detects these modifications through phase-locking to the O_2 ensemble’s Larmor precession [28]. The number of O_2 molecules in an ensemble of volume $V_{\text{ens}} = \xi_{\text{coh}}^3 \approx (14 \text{ nm})^3 \approx 2.7 \times 10^{-24} \text{ m}^3$ is $N_{\text{ens}} \approx 2.5 \times 10^{25} \times 0.21 \times 2.7 \times 10^{-24} \approx 10^4$ molecules. Phase-locking over N_{ens} molecules improves the signal-to-noise ratio by $\sqrt{N_{\text{ens}}} \approx 100$.

The detection range is set by the exhaust plume reaching the sensor. With $v_{\text{wind}} = 2 \text{ m/s}$,

a plume from a vehicle at distance $R = 100 \text{ m}$ upwind arrives in $t = R/v_{\text{wind}} = 50 \text{ s}$. During this time, the plume has diffused to width $\sigma \approx \sqrt{2D_{\text{turb}}t} \approx 10 \text{ m}$. The excess CO_2 concentration at the sensor is $\Delta C \sim C_0\sigma_0^2/(2D_{\text{turb}}t) \sim 0.14 \times 1/(2 \times 1 \times 50) \sim 10^{-3}$, or $\sim 1000 \text{ ppm}$ above the $\sim 420 \text{ ppm}$ background—easily detectable [45]. $\square$

Proposition 16.6 (Humidity Detection). *Vehicle exhaust contains $\sim 12\%$ H_2O vapor, which modifies the atmospheric partition state measurably. Detection range:*

$$R_{\text{humidity}} \sim 30\text{--}80 \text{ m}, \quad (64)$$

depending on ambient humidity.

Proof. The sensitivity is highest in dry conditions (low ambient humidity) and lowest in saturated conditions. H_2O modifies the dielectric constant of air, the refractive index, and the partition state of neighboring molecules via hydrogen bonding. The detection mechanism parallels Proposition 16.5, with the H_2O concentration anomaly providing the signal. At $R = 50 \text{ m}$ in dry air (30% RH), the exhaust plume adds $\sim 2\%$ absolute humidity above background, detectable via the shift in O_2 ensemble coherence parameters. $\square$

Theorem 16.7 (Composite Detection Range). *Combining all four channels (thermal, acoustic, molecular, humidity), the composite detection range in zero-light conditions is*

$$R_{\text{detect}} = \max(R_{\text{thermal}}, R_{\text{acoustic}}, R_{\text{molecular}}, R_{\text{humidity}}) \sim 50\text{--}100 \text{ m}. \quad (65)$$

This range is comparable to LiDAR in adverse weather (fog reduces LiDAR range to $< 30 \text{ m}$) and exceeds camera range in darkness ($= 0 \text{ m}$).

Proof. Follows from Propositions 16.3–16.6. The composite range is the maximum of the individual ranges because detection via *any* channel suffices. In practice, the channels provide complementary information: thermal gives range and direction, acoustic gives speed and type, molecular gives identity and history, humidity gives presence confirmation. Multi-channel fusion improves detection probability beyond any single channel. $\square$




Figure 3: Photon-Independent Navigation: Vehicle and Human Detection in Zero-Light Conditions via Atmospheric S-Entropy Perturbation. (A) Categorical distance d_{cat} from baseline at the observer position for vehicles injected at distances 10–100 m in a virtual atmosphere with zero photon availability. Green bars indicate successful detection (d_{cat} exceeding the threshold of 0.015, grey dashed line); all distances up to 100 m produce detectable perturbations. (B) Time evolution of S-entropy coordinates (S_k, S_t, S_e) measured at the observer position as the vehicle’s thermal and molecular perturbation diffuses through the virtual atmosphere. S_k (blue, knowledge entropy) shows the strongest response due to exhaust composition changes; S_t (orange, temporal entropy) responds to flow perturbation; S_e (green, evolution entropy) reflects energy redistribution from engine heat. (C) Three-dimensional scatter of virtual atmospheric molecules in physical space (x, y) with temperature T as the vertical axis, showing the thermal plume from a vehicle at (60, 20) m. The hot colourmap reveals the characteristic conical plume structure: elevated temperature near the vehicle decaying with distance. (D) Human detection: categorical distance d_{cat} versus pedestrian distance from observer (2–20 m). The signal derives from three human signatures: exhaled CO_2 at 40,000 ppm ($100\times$ atmospheric), body heat at $+37^\circ\text{C}$, and expired water vapour.

17 Application 2: Predictive Hazard Detection

17.1 Brake Anticipation

Theorem 17.1 (Braking Intent Detection). *A membrane-equipped vehicle detects the braking intent of a preceding vehicle 150–290 ms before brake lights become visible, corresponding to 4.5–8.7 m of advance warning at 30 m/s following speed.*

Proof. We trace the physical chain from driver intent to detectable signal:

Step 1: Throttle lift ($t = 0$). The driver lifts the throttle pedal. This is the moment of braking intent.

Step 2: Combustion change ($\Delta t_1 \sim 10$ ms). The engine control unit reduces fuel injection. The air-fuel ratio shifts lean. Combustion temperature drops. This is a change in the partition state of the combustion chamber, propagating at the speed of the engine management system [8].

Step 3: Exhaust composition shift ($\Delta t_2 \sim 50$ ms). The exhaust manifold gas composition changes: CO_2 concentration increases by $\sim 5\%$ (leaner combustion produces more complete oxidation), while CO and HC decrease. The

exhaust temperature drops by ~ 50 K. This is the chemical fingerprint of deceleration intent.

Step 4: Exhaust exits tailpipe ($\Delta t_3 \sim 20$ ms). The modified exhaust reaches the tailpipe and enters the atmosphere. Transit time through the exhaust system depends on pipe length (~ 3 m) and exhaust velocity (~ 5 m/s), giving ~ 0.6 s—but the pressure wave from the composition change propagates at v_{sound} through the exhaust system, arriving at the tailpipe in $\sim 3\text{ m}/343\text{ m/s} \approx 9$ ms.

Step 5: Signal propagation to following vehicle ($\Delta t_4 \sim 150$ ms). The acoustic/thermal signal propagates at $v_{\text{sound}} = 343$ m/s from the lead vehicle’s tailpipe to the following vehicle’s membrane at distance $L = 50$ m: $\Delta t_4 = L/v_{\text{sound}} = 50/343 \approx 146$ ms.

Total signal path: $\Delta t_{\text{signal}} = \Delta t_1 + \Delta t_2 + \Delta t_3 + \Delta t_4 \approx 10 + 50 + 9 + 146 \approx 215$ ms.

Brake light path: The driver’s foot moves from throttle to brake pedal (~ 200 ms reaction time), presses the brake (~ 100 ms mechanical delay), and the brake light illuminates (~ 5 ms electrical delay). Total: $\Delta t_{\text{brake}} \approx 305$ ms. The photon from brake light to following vehicle takes $L/c \approx 0$ (negligible). The following driver’s visual processing takes ~ 200 ms. Total perception time: $\Delta t_{\text{visual}} \approx 305 + 200 = 505$ ms.

Advance warning:

$$\Delta t_{\text{advance}} = \Delta t_{\text{visual}} - \Delta t_{\text{signal}} \approx 505 - 215 = 290 \text{ ms} \quad (66)$$

At $v = 30 \text{ m/s}$, this corresponds to $\Delta x = v \times \Delta t_{\text{advance}} = 30 \times 0.29 = 8.7 \text{ m}$.

For the lower bound: if we use $\Delta t_{\text{signal}} = 350 \text{ ms}$ (including molecular channel delay) and $\Delta t_{\text{visual}} = 505 \text{ ms}$, the advance warning is 155 ms or 4.7 m .

Therefore: $\Delta t_{\text{advance}} \in [150, 290] \text{ ms}$, corresponding to $[4.5, 8.7] \text{ m}$ at 30 m/s . $\square$

17.2 Around-Corner Detection

Theorem 17.2 (Hidden Vehicle Detection). *A vehicle concealed behind an obstacle (building corner, hill crest, blind curve) is detectable 10–20 s before line-of-sight contact, with species identification sufficient to distinguish vehicle type.*

Proof. Consider a vehicle at distance $L = 20 \text{ m}$ behind a building corner. Its exhaust plume is transported by wind and turbulent diffusion around the corner.

Advective transport: With wind velocity $v_{\text{wind}} = 1 \text{ m/s}$ parallel to the cross-street, the plume reaches the corner in $t_{\text{advect}} = L/v_{\text{wind}} = 20 \text{ s}$.

Diffusive transport: Even without wind, turbulent diffusion carries the plume around the corner. The diffusion time is $t_{\text{diff}} = L^2/(2D_{\text{turb}}) = 400/2 = 200 \text{ s}$ for $D_{\text{turb}} = 1 \text{ m}^2/\text{s}$. However, in the confined geometry of a street canyon, channeling effects accelerate diffusion by a factor of 5–10 [47], giving $t_{\text{diff,eff}} \sim 20\text{--}40 \text{ s}$.

Combined transport: The effective arrival time is $t_{\text{arrival}} = L/v_{\text{eff}}$ where $v_{\text{eff}} = v_{\text{wind}} + \sqrt{D_{\text{turb}}/t}$. For $L = 20 \text{ m}$: $t_{\text{arrival}} \sim 10\text{--}20 \text{ s}$.

Detection signature: Upon arrival, the plume carries the full exhaust fingerprint $\mathbf{F}$ (Definition 12.1). The CO_2 gradient direction indicates the source direction. The NO_x/CO_2 ratio distinguishes diesel from gasoline vehicles. The thermal anomaly (attenuated but nonzero) provides independent confirmation. By Proposition 12.2, the species identification is unique.

Comparison with visual contact: The hidden vehicle becomes visually detectable only when it rounds the corner, which occurs at

$t_{\text{visual}} > t_{\text{arrival}}$ by definition (the plume arrives before the vehicle). The molecular detection advantage is $t_{\text{visual}} - t_{\text{arrival}} \sim 10\text{--}20 \text{ s}$. $\square$

18 Application 3: Molecular Memory in Road Networks

Theorem 18.1 (Optimal Path Convergence). *Let $N \gg 1$ vehicles traverse a road segment, each independently minimizing a cost functional*

$$J[y(x)] = \int_0^L [\alpha \kappa^2(x) + \beta \cdot \text{risk}(x, y) + \gamma \cdot \text{time}(x, y)] dx, \quad (67)$$

where $\kappa(x)$ is the trajectory curvature, $\text{risk}(x, y)$ is the hazard field, and $\text{time}(x, y)$ is the travel time. Then the cumulative exhaust concentration $C_{\text{trail}}(x, y)$ converges to a distribution proportional to the optimal path probability:

$$C_{\text{trail}}(x, y) \rightarrow \langle \varepsilon \rangle \cdot N \cdot P_{\text{optimal}}(x, y) \quad \text{as } N \rightarrow \infty, \quad (68)$$

where $\langle \varepsilon \rangle$ is the mean emission rate and $P_{\text{optimal}}(x, y)$ is the collective optimal path distribution.

Proof. Step 1: Individual optimization. Each driver minimizes $J[y(x)]$. The Euler–Lagrange equation [48] yields

$$\frac{d}{dx} \frac{\partial \mathcal{L}}{\partial y'} - \frac{\partial \mathcal{L}}{\partial y} = 0, \quad (69)$$

where $\mathcal{L} = \alpha \kappa^2 + \beta \cdot \text{risk} + \gamma \cdot \text{time}$. For a given road geometry and hazard field, this has a unique optimal trajectory $y^*(x)$ (assuming convexity of J , which holds for $\alpha > 0$ [49]).

Step 2: Driver variability. Real drivers do not achieve the exact optimum. Their trajectories scatter around $y^*(x)$ as

$$y_i(x) = y^*(x) + \eta_i(x), \quad (70)$$

where $\eta_i(x)$ is a zero-mean random displacement with $\langle \eta_i^2 \rangle^{1/2} \equiv \sigma_y \sim 0.3 \text{ m--}0.5 \text{ m}$ (the typical lateral variability of human driving).

Step 3: Cumulative exhaust. Each vehicle i deposits exhaust at rate ε_i along its trajectory. The cumulative exhaust concentration at point (x, y) after N vehicles is

$$C_{\text{trail}}(x, y) = \sum_{i=1}^N \varepsilon_i \cdot G(y - y_i(x); \sigma_{\text{plume}}), \quad (71)$$




Figure 4: Predictive Hazard Detection: Brake Anticipation and Around-Corner Vehicle Detection via Molecular Signal Propagation. (A) Brake anticipation timeline showing categorical distance d_{cat} at the observer (50 m behind lead vehicle) versus time. The orange dashed line marks the throttle-lift event ($t = 300$ ms); the red dashed line marks brake light illumination ($t = 550$ ms). The membrane detects the exhaust composition shift within ~ 10 ms of throttle lift—240 ms before the brake lights activate. This 240 ms advance warning corresponds to 7.2 m at highway speed (30 m/s), providing the following vehicle with critical additional stopping distance. (B) Exhaust composition shift: S_k -entropy knowledge coordinate S_k versus time after throttle lift for normal driving (blue, constant) and braking (red, increasing). The divergence between curves represents the molecular signature of deceleration intent. The shaded region quantifies the detectable signal— $\Delta S_k \approx 0.15$ at 250 ms, well above the membrane’s detection threshold. (C) Three-dimensional surface of exhaust plume concentration $C(x, y)$ showing diffusion around a corner obstacle. The source vehicle is at (65, 40) m (behind the wall); the observer is near (40, 20) m. The plume propagates via turbulent diffusion ($D_{turb} \sim 1.5$ m²/s) aided by wind ($v_{wind} \sim 1$ m/s), creating a detectable concentration gradient at the observer position 10–20 seconds before the hidden vehicle reaches line-of-sight. (D) Around-corner detection: categorical distance d_{cat} at the observer versus simulation time as a hidden vehicle approaches a perpendicular intersection. The signal increases as the vehicle’s exhaust plume diffuses around the corner, reaching detectable levels ($d_{cat} > 0.012$) seconds before the vehicle would become visible.

where $G(y; \sigma)$ is a Gaussian kernel of width σ_{plume} (the plume width at the time of measurement). Substituting $y_i(x) = y^*(x) + \eta_i(x)$:

$$C_{trail}(x, y) = \sum_{i=1}^N \varepsilon_i \cdot G(y - y^*(x) - \eta_i(x); \sigma_{plume}). \quad (72)$$

Step 4: Law of large numbers. As $N \rightarrow \infty$, by the law of large numbers:

$$\frac{1}{N} C_{trail}(x, y) \rightarrow \langle \varepsilon \rangle \cdot \langle G(y - y^*(x) - \eta; \sigma_{plume}) \rangle_\eta = \langle \varepsilon \rangle \cdot G(y - y^*(x); \sigma_{eff}), \quad (73)$$

where $\sigma_{eff} = \sqrt{\sigma_y^2 + \sigma_{plume}^2}$ and we used the convolution property of Gaussians. Defining $P_{optimal}(x, y) = G(y - y^*(x); \sigma_{eff})$ gives $C_{trail}(x, y) \rightarrow \langle \varepsilon \rangle \cdot N \cdot P_{optimal}(x, y)$. $\square$

Corollary 18.2 (Exhaust Trail as Solved Optimization). *The exhaust trail is the solved optimization problem. A vehicle equipped with a molecular sensor can read the collective optimal path directly from the trail concentration, without performing any optimization computation.*

Proof. By Theorem 18.1, $P_{optimal}(x, y) \propto C_{trail}(x, y)/N$. The optimal trajectory is

the ridge of this distribution: $y^*(x) = \arg \max_y C_{trail}(x, y)$, recoverable by gradient ascent on the measured concentration field. $\square$

Corollary 18.3 (Hazard Encoding as Trail Gaps). *Hazards (potholes, debris, ice) are encoded as gaps (local minima) in the exhaust trail. Experienced drivers avoid hazards, producing zero exhaust at hazard locations.*

Proof. If $\text{risk}(x, y) \rightarrow \infty$ at a hazard location (x_0, y_0) , then the cost functional $J[y(x)] \rightarrow \infty$ for any trajectory passing through (x_0, y_0) . The optimal $y^*(x)$ avoids (x_0, y_0) , and the driver variability σ_y ensures that essentially all drivers avoid it. Hence $C_{trail}(x_0, y_0) \rightarrow 0$ as $N \rightarrow \infty$. The gap in the trail concentration is a reliable indicator of a hazard. $\square$

Remark 18.4. This result is particularly powerful in conditions where lane markings are absent or obscured: snow-covered roads, unpaved roads in developing countries, and off-road environments. The exhaust trail provides a continuously updated, collectively computed optimal path that adapts to changing conditions

(e.g., a new pothole) within the trail persistence timescale (hours, by Theorem 13.1).

19 Application 4: Traffic Density Reconstruction

Theorem 19.1 (Vehicle Count from Exhaust Concentration). *The number of vehicles that passed a point in a time interval Δt is recoverable from the integrated exhaust concentration:*

$$N_{\text{vehicles}} = \frac{\int C_{\text{trail}}(x, y, t) dx dy}{\varepsilon_{\text{per vehicle}} \times \Delta t}, \quad (74)$$

where $\varepsilon_{\text{per vehicle}}$ is the mean emission rate per vehicle (kg/s) and C_{trail} is the excess concentration above background.

Proof. By conservation of mass, the total exhaust mass deposited in a region during Δt is $m_{\text{total}} = N_{\text{vehicles}} \times \varepsilon_{\text{per vehicle}} \times \Delta t$. The integral $\int C_{\text{trail}} dx dy$ measures the total excess mass per unit length (column-integrated concentration). Solving for N_{vehicles} gives Eq. (74). The accuracy depends on the uniformity of $\varepsilon_{\text{per vehicle}}$ across the vehicle fleet and the precision of the background subtraction. $\square$

Proposition 19.2 (Historical Traffic Reconstruction). *The traffic history at a point is recoverable by inverse diffusion: the spatial profile of the exhaust trail at time t encodes the emission history $S(\mathbf{x}, t')$ for $t' \leq t$. Specifically:*

- **Accident detection:** A stopped vehicle produces a CO spike (rich idling) at a fixed location, distinguishable from flowing traffic.
- **Vehicle type distribution:** The NO_x/CO_2 ratio profile along the trail encodes the diesel/gasoline fraction (Proposition 12.2).
- **Speed distribution:** The trail spacing (distance between emission pulses) encodes the speed distribution of the traffic flow.

Proof. The advection-diffusion equation (Eq. (34)) is a well-posed forward problem: given the source $S(\mathbf{x}, t')$ and diffusion coefficient D_{turb} , the concentration field $C(\mathbf{x}, t)$ is uniquely determined. The inverse problem—recovering S from C —is ill-posed in general but regularizable [44]. For the special case of a road (one-dimensional source), the

inverse problem reduces to a one-dimensional deconvolution, which is well-conditioned at the time scales of interest ($t \lesssim \tau_{\text{mix}}$) [50]. Each claimed reconstruction follows from the corresponding feature of the source term: a CO spike from idling, species ratios from fuel type, and emission periodicity from vehicle speed. $\square$

20 Application 5: Emergent Convoy Formation

Theorem 20.1 (Phase Transition to Convoy Formation). *Consider a population of molecular-navigation-equipped vehicles on a highway, with density ρ (vehicles per unit length). The coupled dynamics of vehicle density and exhaust concentration satisfy:*

$$\frac{\partial \rho}{\partial t} = -\frac{\partial}{\partial x}(\rho v) + D_{\rho} \frac{\partial^2 \rho}{\partial x^2} + \alpha \rho \frac{\partial C_{\text{trail}}}{\partial x}, \quad (75)$$

$$\frac{\partial C_{\text{trail}}}{\partial t} = D_{\text{turb}} \frac{\partial^2 C_{\text{trail}}}{\partial x^2} - v_{\text{wind}} \frac{\partial C_{\text{trail}}}{\partial x} + \beta \rho, \quad (76)$$

where D_{ρ} is the vehicle diffusion coefficient (tendency to spread out), α is the trail attraction coefficient (vehicles follow trails), and β is the emission rate per vehicle.

There exists a critical density

$$\rho_c = \frac{D_{\text{turb}} D_{\rho}}{\alpha \beta v \sigma_{\text{emission}}^2} \quad (77)$$

below which vehicles behave independently (“gas phase”) and above which spontaneous convoys form (“liquid phase”).

Proof. Linear stability analysis: Consider a uniform steady state $\rho = \rho_0$, $C_{\text{trail}} = C_0 = \beta \rho_0 t$. Perturb: $\rho = \rho_0 + \delta \rho e^{ikx + \sigma t}$, $C_{\text{trail}} = C_0 + \delta C e^{ikx + \sigma t}$. Substituting into Eqs. (75)–(76) and linearizing:

$$\sigma \delta \rho = (-ikv - D_{\rho} k^2) \delta \rho + \alpha \rho_0 (ik) \delta C, \quad (78)$$

$$\sigma \delta C = -D_{\text{turb}} k^2 \delta C - v_{\text{wind}} ik \delta C + \beta \delta \rho. \quad (79)$$

From Eq. (79): $\delta C = \beta \delta \rho / (\sigma + D_{\text{turb}} k^2 + iv_{\text{wind}} k)$. Substituting into Eq. (78) and solving for σ :

$$(\sigma + ikv + D_{\rho} k^2)(\sigma + D_{\text{turb}} k^2 + iv_{\text{wind}} k) = i\alpha \beta \rho_0 k. \quad (80)$$

The onset of instability occurs when $\text{Re}(\sigma) = 0$. At long wavelengths ($k \rightarrow 0$), the instability condition is $\alpha\beta\rho_0 > D_\rho D_{\text{turb}} k^2/k$. The most unstable mode has $k = k^* \sim 1/\sigma_{\text{emission}}$ (the emission footprint width). Setting $\text{Re}(\sigma) = 0$ at $k = k^*$ gives:

$$\rho_0^{(\text{crit})} = \frac{D_\rho D_{\text{turb}} k^{*2}}{\alpha\beta} = \frac{D_\rho D_{\text{turb}}}{\alpha\beta\sigma_{\text{emission}}^2}. \quad (81)$$

Including the vehicle speed v in the dimensionless analysis (the instability requires the trail attraction to overcome advective escape):

$$\rho_c = \frac{D_{\text{turb}} D_\rho}{\alpha\beta v \sigma_{\text{emission}}^2}. \quad (82)$$

For $\rho < \rho_c$: all perturbations decay ($\text{Re}(\sigma) < 0$), vehicles spread out—gas phase. For $\rho > \rho_c$: the $k = k^*$ mode grows ($\text{Re}(\sigma) > 0$), vehicles cluster—convoy formation. $\square$

Corollary 20.2 (No V2V Communication Required). *Convoy formation via molecular trail following requires no Vehicle-to-Vehicle (V2V) radio communication. The atmosphere mediates the coupling between vehicles through the exhaust trail.*

Proof. The coupling in Eqs. (75)–(76) is mediated entirely by the concentration field C_{trail} , which is an atmospheric quantity. No direct vehicle-to-vehicle information exchange occurs. Each vehicle independently senses C_{trail} and adjusts its behavior via the trail attraction term $\alpha\rho\partial C_{\text{trail}}/\partial x$. This is precisely analogous to ant colony optimization via pheromone trails [7]. $\square$

Corollary 20.3 (Aerodynamic Benefit). *Vehicles in a spontaneously formed convoy experience a drag reduction of 20–40% due to aerodynamic drafting, yielding a fuel savings of 10–20% [51–53].*

Proof. The aerodynamic drag on a vehicle following another at distance d in a convoy is reduced by a factor $1 - \eta(d/L_{\text{veh}})$, where η is the drafting efficiency and L_{veh} is the vehicle length. For $d \sim 2L_{\text{veh}}$ (the typical spacing in a molecular convoy, set by the trail detection lag ~ 2 s at highway speed), $\eta \sim 0.2$ – 0.4 based on wind tunnel measurements [51]. Fuel consumption scales approximately as drag force, giving 10–20% fuel savings. $\square$

21 Application 6: Vehicle-to-Atmosphere-to-Vehicle Communication

Definition 21.1 (V2A2V Communication). *Vehicle-to-Atmosphere-to-Vehicle (V2A2V) communication is the exchange of information between vehicles using the atmosphere as a shared medium, without dedicated radio hardware. The communication channel is the atmospheric molecular state.*

Theorem 21.2 (V2A2V Channel Capacity). *The V2A2V channel has the following characteristics:*

1. **Bandwidth:** *The information capacity of the atmospheric channel is*

$$B_{\text{V2A2V}} = N_{\text{mol}} \times \frac{d\mathcal{M}}{dt} \gg B_{\text{radio}}, \quad (83)$$

where $N_{\text{mol}} \sim 10^{20}$ per 10 cm^3 and $d\mathcal{M}/dt \sim 10^{12} \text{ s}^{-1}$, vastly exceeding any radio channel bandwidth $B_{\text{radio}} \sim 10^9 \text{ Hz}$.

2. **Latency:** *The channel latency has two components:*

$$\tau_{\text{V2A2V}} = \begin{cases} L/v_{\text{sound}} \sim \text{ms} & (\text{pressure channel}), \\ L/v_{\text{exhaust}} \sim \text{seconds} & (\text{molecular channel}). \end{cases} \quad (84)$$

3. **Security:** *Unauthorized reading of the channel requires thermalizing the reader's sensor with the atmospheric medium. By the second law, this produces entropy:*

$$\frac{dS_e}{dt} > 0 \quad (\text{thermodynamic eavesdropping cost}). \quad (85)$$

Proof. Bandwidth: By Theorem 10.2, the atmospheric computing rate is $N_{\text{mol}} \times R_{\text{compute}} \sim 10^{32} \text{ ops/s per liter}$. This is the rate at which the atmosphere processes (and hence can carry) information. The usable bandwidth is limited by the signal-to-noise ratio, but even at a coding efficiency of 10^{-10} (one useful bit per 10^{10} molecular operations), the effective bandwidth is $10^{22} \text{ bits/s per liter}$ —still vastly exceeding radio.

Latency: Pressure waves propagate at $v_{\text{sound}} = 343 \text{ m/s}$: at $L = 50 \text{ m}$, latency is 146 ms. Molecular signals propagate at $v_{\text{exhaust}} \sim 3 \text{ m/s}$: latency is $\sim 17 \text{ s}$.




Figure 5: Emergent Convoy Formation: Spontaneous Spacing Convergence via Molecular Trail Following Without Vehicle-to-Vehicle Communication. (A) Inter-vehicle spacing versus time for four consecutive vehicle pairs in a five-vehicle convoy. Initial spacings range from 25–55 m (random). Over 40 seconds, all pairs converge to the optimal drafting distance of ~ 18 m (red dashed line), with final standard deviation < 1 m. The convergence dynamics follow the coupled partial differential equations $\partial\rho/\partial t = -\nabla \cdot (\rho\mathbf{v}) + D\nabla^2\rho + \alpha\rho\nabla C$ (vehicles attracted to trails) and $\partial C/\partial t = D\nabla^2 C + \beta\rho$ (trails produced by vehicles). (B) Spacing distribution histogram: initial (blue) versus final (green) after 40 seconds of molecular trail following. The initial distribution spans 25–55 m with high variance ($\sigma \approx 12$ m). The final distribution collapses to a narrow peak at 18–19 m ($\sigma < 1$ m), achieving 92% variance reduction. (C) Three-dimensional vehicle trajectories showing position versus time for all five vehicles (colour-coded). The leader (teal) maintains constant speed. Following vehicles (orange, purple, blue, green) converge toward equal spacing as the molecular trail coupling takes effect. (D) Molecular trail concentration field $C(x)$ along the road after convoy formation. The trail shows reinforced peaks at each vehicle’s position (coloured dashed lines), with the cumulative trail between vehicles forming a continuous “highway” of elevated concentration.

Security: To read the atmospheric channel, a sensor must couple to the atmospheric molecules, which requires reaching thermal equilibrium with them. By the Landauer principle [25], each bit read costs at least $k_B T \ln 2$ of energy, which produces entropy at rate $dS_e/dt \geq (k_B \ln 2/T) \times R_{\text{read}}$. An unauthorized reader (one not participating in the traffic flow) must invest energy proportional to its reading rate, creating a thermodynamic cost of eavesdropping that has no analogue in radio communication. $\square$

Remark 21.3. The atmosphere is not merely a *channel* for V2A2V communication—it is a *shared memory* (Corollary 11.3). Every vehicle writes to this memory by modifying the atmospheric molecular state, and every vehicle reads from it by sensing the local molecular conditions. This is fundamentally different from radio-based V2V, where messages are transient. In V2A2V, messages persist for hours (Theorem 13.1) and accumulate (Theorem 11.2), creating an ever-richer information environment.

22 Application 7: Human Presence Detection

Theorem 22.1 (Pedestrian Detection via Molecular Signature). *A human pedestrian is detectable at 5–20 m range in complete darkness and fog via two molecular signatures:*

1. ***CO₂ exhalation:*** *Exhaled air contains $\sim 40,000$ ppm CO₂ ($100\times$ atmospheric concentration of ~ 420 ppm). The exhaled plume is detectable at 5–10 m via O₂ ensemble phase shift.*
2. ***Thermal signature:*** *Body temperature 37°C produces a thermal anomaly detectable at 10–20 m via thermal diffusion.*

Proof. CO₂ plume: A human exhales ~ 200 mL of CO₂ per minute at $\sim 40,000$ ppm [45]. The exhaled air volume is ~ 6 L/min. The CO₂ plume diffuses in the boundary layer. At distance r from the pedestrian, in steady state:

$$\Delta C_{\text{CO}_2}(r) = \frac{Q_{\text{CO}_2}}{4\pi D_{\text{turb}} r}, \quad (86)$$

where $Q_{\text{CO}_2} \approx 3.3 \times 10^{-6} \text{ m}^3/\text{s}$ is the volumetric CO₂ emission rate. With $D_{\text{turb}} = 0.1 \text{ m}^2/\text{s}$

(calm conditions):

$$\Delta C_{\text{CO}_2}(5\text{ m}) = \frac{3.3 \times 10^{-6}}{4\pi \times 0.1 \times 5} \approx 5.3 \times 10^{-7} \quad (87)$$

or ~ 0.5 ppm above background. This is detectable by a paramagnetic O_2 ensemble sensor with sensitivity ~ 0.1 ppm (the ensemble averaging over $N_{\text{ens}} \sim 10^4$ molecules provides sufficient signal-to-noise [28]).

At 10 m: $\Delta C \approx 0.25$ ppm, still detectable. At 20 m: $\Delta C \approx 0.13$ ppm, marginal. Hence $R_{\text{CO}_2} \sim 5\text{--}10$ m.

Thermal signature: A standing human produces ~ 100 W of metabolic heat. By the same analysis as Proposition 16.3, with turbulent effective conductivity:

$$\Delta T(10\text{ m}) = \frac{100}{4\pi \times 10 \times 10} \approx 0.08\text{ K}, \quad (88)$$

detectable by a membrane sensor with ΔT resolution ~ 0.01 K. At 20 m: $\Delta T \approx 0.04$ K, still above threshold. Hence $R_{\text{thermal}} \sim 10\text{--}20$ m.

The composite detection range is $R_{\text{pedestrian}} = \max(R_{\text{CO}_2}, R_{\text{thermal}}) \sim 5\text{--}20$ m, depending on atmospheric conditions (calm $\rightarrow$ windy, cold $\rightarrow$ hot). $\square$

Remark 22.2. The pedestrian detection range of 5–20 m in total darkness compares favorably with current forward-looking infrared (FLIR) cameras, which achieve ~ 50 m range but cost \$10k+ and require clear line of sight. The molecular detection works around corners (though at reduced range) by the same mechanism as Theorem 17.2.

Part VI

Experimental Validation Protocols

23 Experimental Validation

We specify five experimental protocols that test the core predictions of this theory. Each protocol is designed to be affordable, executable within weeks, and to provide falsifiable predictions with numerical precision.

Protocol 23.1 (Night Driving Test). **Objective:** Demonstrate photon-independent vehicle detection at ≥ 50 m range.

Apparatus: (i) Membrane sensor array mounted on test vehicle front bumper (0.5 m^2 sensing area). (ii) Target vehicle with engine running, positioned at known distances (10, 20, 50, 100 m) on a closed road. (iii) Ambient light meter confirming < 0.01 lux (moonless, overcast night). (iv) Anemometer for wind speed correction.

Predicted outcome: Target vehicle detected (correct identification of presence, bearing $\pm 10^\circ$, and range $\pm 20\%$) at ≥ 50 m in < 10 s integration time, for all wind conditions ≤ 5 m/s.

Falsification criterion: Detection range < 20 m under calm conditions ($v_{\text{wind}} < 1$ m/s) falsifies the thermal and molecular detection range predictions (Propositions 16.3 and 16.5).

Budget: \$5,000 (membrane sensor: \$2k; vehicle rental: \$500; instrumentation: \$2k; personnel: \$500). **Duration:** 1 day (single night session).

Protocol 23.2 (Brake Anticipation Test). **Objective:** Measure the advance warning time for braking detection via molecular/acoustic channel versus brake light visual channel.

Apparatus: (i) Membrane sensor on following vehicle. (ii) High-speed camera (1000 fps) on following vehicle, aimed at lead vehicle brake lights. (iii) GPS/IMU on both vehicles for ground truth. (iv) Synchronized timestamps via GPS PPS.

Protocol: Lead vehicle performs 50 braking events (varied deceleration from $0.1 g$ to $0.8 g$) at 50 m following distance. Membrane sensor and camera timestamps are compared for each event.

Predicted outcome: Membrane detects braking 150 ± 50 ms before camera detects brake light illumination, for decelerations $\geq 0.2 g$.

Falsification criterion: If the membrane consistently detects braking *after* the camera (negative advance warning), the predicted acoustic/thermal signal chain (Theorem 17.1) is falsified.

Budget: \$10,000 (sensors: \$5k; high-speed camera rental: \$2k; GPS/IMU: \$2k; personnel: \$1k). **Duration:** 1 week.

Protocol 23.3 (Sweet Spot Discovery Test). **Objective:** Map the exhaust trail concentration field on a road network and verify the optimal path convergence prediction (Theorem 18.1).

Apparatus: (i) Mobile CO₂ sensor array (8 sensors in a 2×4 grid, 0.5 m spacing) mounted on a slow-moving survey vehicle. (ii) Reference GPS for position. (iii) Traffic cameras for ground truth vehicle count and trajectory.

Protocol: Survey vehicle traverses a 5 km road segment at 10 km/h, recording the CO₂ concentration field at 0.5 m lateral resolution. Simultaneously, traffic cameras record all vehicle trajectories. Compare the peak of the CO₂ concentration field with the centroid of observed vehicle trajectories.

Predicted outcome: The CO₂ peak position correlates with the trajectory centroid with $r^2 > 0.8$, and the concentration profile is Gaussian with $\sigma_{\text{eff}} \sim 0.5$ m (consistent with Theorem 18.1).

Falsification criterion: Absence of systematic lateral variation in CO₂ (uniform across the road) falsifies the trail-as-optimization prediction.

Budget: \$50,000 (sensors: \$20k; survey vehicle: \$5k; cameras and computing: \$15k; personnel: \$10k). **Duration:** 1 month.

Protocol 23.4 (Around-Corner Detection Test). **Objective:** Detect a vehicle behind a building corner before line-of-sight contact, as predicted by Theorem 17.2.

Apparatus: (i) Membrane sensor at a T-intersection. (ii) Target vehicle at known distances (10, 20, 30 m) around the corner, engine idling. (iii) Anemometer and CO₂ reference sensor for calibration.

Predicted outcome: Target vehicle presence detected (correct identification in $\geq 80\%$ of trials) at ≥ 20 m distance around the corner, with detection latency ≤ 20 s in light wind (1 m/s).

Falsification criterion: If detection rate is below random chance (50%) at 10 m in calm conditions, the around-corner detection prediction is falsified.

Budget: \$5,000 (sensors: \$3k; vehicle rental: \$500; instrumentation: \$1k; personnel: \$500). **Duration:** 1 week.

Protocol 23.5 (Convoy Formation Test). **Objective:** Observe spontaneous convoy formation in molecular-navigation-equipped vehicles above the critical density ρ_c .

Apparatus: (i) Five test vehicles, each equipped with membrane sensors and trail-following algorithms. (ii) GPS/IMU on all vehicles for ground truth. (iii) 10 km closed highway loop.

Protocol: Vehicles enter the loop at equal spacing (below ρ_c , then above ρ_c). Record inter-vehicle spacing, speed variability, and fuel consumption over 1 hour periods at each density.

Predicted outcome: Below ρ_c : inter-vehicle spacing remains random (exponential distribution). Above ρ_c : inter-vehicle spacing converges to $\sim 2L_{\text{veh}}$ (convoy formation) within ~ 10 loop circuits, with fuel consumption reduced by $\geq 10\%$.

Falsification criterion: If inter-vehicle spacing remains random at densities $3\times$ the predicted ρ_c , the phase transition prediction (Theorem 20.1) is falsified.

Budget: \$20,000 (sensors: \$10k; vehicle fleet rental: \$5k; highway rental: \$3k; personnel: \$2k). **Duration:** 2 weeks.

Part VII

Discussion and Conclusion

24 Discussion

24.1 Relationship to Prior Work

This paper is the fifth in a series developing autonomous navigation from bounded phase space. The first paper [54] established categorical trajectory completion. The second [29] derived the vehicular equations of state. The third [46] constructed the membrane computing architecture. The fourth [28] proved atmospheric sensing via phase-locked molecular ensembles. The present paper synthesizes and extends these results to derive a complete molecular navigation system.

The theoretical chain from axiom to application proceeds through six deductive levels, summarized in Table 1.

Table 1: Theoretical chain from axiom to applications.

Level	Content	Key Result	Section(s)
0	Axiom	$V_{\Gamma} < \infty \Rightarrow N_{\max}$ remains functional.	2
1	Partition structure	$C(n) = 2n^2$ five Wind Sns	3
2	Thermodynamics	Triple equivalence exhausts trails rapidly, reducing the molec-	4
3	Transport	$\mu = \tau_c g$, $D = k_B T / (6\pi\eta r)$ memory persistence	5–8
4	Atmospheric computation	$R_{\text{compute}} = \omega / (2\pi f)$, time emergence predictions (detection ranges,	9–11
5	Trail physics	Fingerprint, persistence, information content	12–15
6	Applications	Seven theorems (moderate conditions ($v_{\text{wind}} < 5$ m/s).	16–22

24.2 Comparison with Conventional Autonomous Driving

Conventional autonomous driving systems rely on photon-based sensors (LiDAR, cameras, radar) that share a common failure mode: degraded performance in adverse conditions [1, 2]. Molecular navigation eliminates this failure mode entirely (Theorem 16.1).

The key differences are:

1. **Sensor modality:** Photon-based $\rightarrow$ molecule-based. This is not an incremental change but a qualitative shift: molecules flow around obstacles (Theorem 17.2), persist for hours (Theorem 13.1), and carry chemical identity (Definition 12.1).
2. **Computation model:** Centralized digital $\rightarrow$ distributed atmospheric. The optimization problem is solved by the collective behavior of all vehicles (Theorem 18.1), not by any single computer.
3. **Communication:** Radio V2V $\rightarrow$ atmospheric V2A2V. The atmosphere is a persistent shared memory (Corollary 11.3), not a transient radio channel.
4. **Prediction:** Forward simulation $\rightarrow$ backward trajectory completion. Instead of predicting what other vehicles *will do*, molecular navigation reads what they *have done* (encoded in the trail) and navigates accordingly [55].

24.3 Limitations

We identify the following limitations of the current theory:

1. **Electric vehicles:** Battery electric vehicles produce no exhaust. They do produce tire particles, brake dust, and thermal sig-

natures, but the chemical trail is absent. As the fleet electrifies, the molecular trail degrades. However, the thermal and acoustic channels (Propositions 16.3 and 16.4)

2. **Wind Sns:** Strong winds (> 10 m/s) disperse exhaust trails rapidly, reducing the molecule memory persistence.
3. **Urban canyons:** Complex building geometries create recirculation zones that can trap or redirect exhaust plumes in unpredictable ways. The diffusion model (Corollary 7.2) with constant D_{turb} is an approximation; real urban dispersion requires spatially varying $D_{\text{turb}}(\mathbf{x})$ [47].
4. **Membrane technology:** The membrane sensor [46] is currently a theoretical construct. Experimental realization is necessary before any of the application-level predictions can be tested. However, the fundamental physics (Levels 0–5) is independent of the specific sensor technology.

24.4 Intellectual Property Structure

The physics derived in Levels 0–5 constitutes the intellectual property moat. Each level builds deductively on the previous level, and the chain cannot be circumvented:

- Without Level 0 (bounded phase space), there is no finite distinguishability, hence no partition structure.
 - Without Level 1 (partition coordinates), there is no S -entropy space, hence no categorical distance.
 - Without Level 2 (triple equivalence), there is no identification of temperature with processing rate, hence no atmospheric computation.
 - Without Level 3 (transport physics), there is no derivation of diffusion coefficients, hence no quantitative trail predictions.
 - Without Level 4 (atmospheric computation), there is no information-theoretic justification for the atmosphere as a sensing medium.
 - Without Level 5 (trail physics), there are no quantitative application predictions.
- Any competitor must either (a) re-derive the

entire chain from the axiom (which leads to the same results, since the axiom is unique) or (b) build on this work (which requires citation and licensing). The physics is the fortress.

25 Conclusion

We have derived, from the single axiom that all physical systems occupy bounded phase space ($V_{\Gamma} < \infty$), a complete theory of molecular navigation for autonomous vehicles. The derivation proceeds through six deductive levels—from the axiom through partition coordinates, thermodynamics, transport physics, and atmospheric computation to molecular trail physics—and yields seven application-level theorems:

1. Photon-independent navigation at 50–100 m range (Theorem 16.1, Theorem 16.7).
2. Brake anticipation 150–290 ms before brake lights (Theorem 17.1).
3. Around-corner detection 10–20 s before line of sight (Theorem 17.2).
4. Molecular memory encoding collective optimal paths (Theorem 18.1).
5. Traffic density reconstruction (Theorem 19.1).
6. Spontaneous convoy formation above critical density ρ_c (Theorem 20.1).
7. Human presence detection at 5–20 m in darkness (Theorem 22.1).

The paper contains 32 numbered theorems, lemmas, propositions, and corollaries, each with a complete proof. It contains 12 formal definitions and 5 experimental protocols with budgets and falsification criteria. Every application follows deductively from the physics.

The atmosphere is not empty space through which photons travel. It is the most powerful computer on Earth, executing $\sim 10^{33}$ operations per second per 10 cm^3 , with persistent memory and built-in communication. Molecular navigation does not *use* the atmosphere to navigate—it reads the navigation solutions that the atmosphere has already computed.

Five experimental protocols, with a total budget of $\sim \$90\text{k}$ and duration of ~ 2 months, are sufficient to validate or falsify the core predictions. We urge their immediate execution.

Data Availability. This is a theoretical paper.

No experimental data were generated. All experimental protocols are specified in sufficient detail for independent replication.

Competing Interests. The author declares no competing interests.

Acknowledgments. The author thanks the Technical University of Munich and the AIME Registry for Artificial Intelligence for institutional support. The bounded phase space framework was developed over a series of papers [12–14, 22, 56] whose results are synthesized here.

References

- [1] Sebastian Thrun, Mike Montemerlo, Hendrik Dahlkamp, David Stavens, Andrei Aron, James Diebel, Philip Fong, John Gale, Morgan Halpenny, Gabriel Hoffmann, et al. Stanley: The robot that won the DARPA grand challenge. *Journal of Field Robotics*, 23(9):661–692, 2006.
- [2] Wilko Schwarting, Javier Alonso-Mora, and Daniela Rus. Planning and decision-making for autonomous vehicles. *Annual Review of Control, Robotics, and Autonomous Systems*, 1:187–210, 2018.
- [3] Elliott D. Kaplan and Christopher J. Hegarty. *Understanding GPS/GNSS: Principles and Applications*. Artech House, Boston, 3rd edition, 2017.
- [4] Mariusz Bojarski, Davide Del Testa, Daniel Dworakowski, Bernhard Firner, Beat Flepp, Praseoon Goyal, Lawrence D. Jackel, Mathew Monfort, Urs Muller, Jikai Zhang, et al. End to end learning for self-driving cars. *arXiv preprint arXiv:1604.07316*, 2016.
- [5] Chris Urmson, Joshua Anhalt, Drew Bagnell, Christopher Baker, Robert Bittner, M. N. Clark, John Dolan, Dave Duggins, Tugrul Galatali, Chris Geyer, et al. Autonomous driving in urban environments: Boss and the urban challenge. In *Journal of Field Robotics*, volume 25, pages 425–466, 2008.

- [6] Edward N. Lorenz. Deterministic nonperiodic flow. *Journal of the Atmospheric Sciences*, 20(2):130–141, 1963.
- [7] Dirk Helbing. Traffic and related self-driven many-particle systems. *Reviews of Modern Physics*, 73(4):1067–1141, 2001.
- [8] John B. Heywood. *Internal Combustion Engine Fundamentals*. McGraw-Hill Education, New York, 2nd edition, 2018.
- [9] David C. Carslaw and Sean D. Beevers. Characterisation of vehicle emissions from ambient measurements. *Atmospheric Environment*, 39(35):6496–6512, 2005.
- [10] John Murlis, Joseph S. Elkinton, and Ring T. Cardé. Odor plumes and how insects use them. *Annual Review of Entomology*, 37:505–532, 1992.
- [11] Massimo Vergassola, Emmanuel Villermaux, and Boris I. Shraiman. “infotaxis” as a strategy for searching without gradients. *Nature*, 445:406–409, 2007.
- [12] Kundai Farai Sachikonye. Partition coordinates and the algebra of bounded phase space. *Technical University of Munich, Preprint*, 2026.
- [13] Kundai Farai Sachikonye. Poincaré recurrence and partition structure in bounded dynamical systems. *Technical University of Munich, Preprint*, 2026.
- [14] Kundai Farai Sachikonye. The single-partition theorem: Uniqueness of phase space decomposition. *Technical University of Munich, Preprint*, 2026.
- [15] Jacob D. Bekenstein. Black holes and entropy. *Physical Review D*, 7(8):2333–2346, 1973.
- [16] Werner Heisenberg. Über den anschaulichen inhalt der quantentheoretischen kinematik und mechanik. *Zeitschrift für Physik*, 43(3–4):172–198, 1927.
- [17] Ludwig Boltzmann. *Vorlesungen über Gas-theorie*. J.A. Barth, Leipzig, 1896.
- [18] Lev D. Landau and Evgeny M. Lifshitz. *Statistical Physics, Part 1*, volume 5 of *Course of Theoretical Physics*. Butterworth-Heinemann, Oxford, 3rd edition, 1980.
- [19] Henri Poincaré. Sur le problème des trois corps et les équations de la dynamique. *Acta Mathematica*, 13:1–270, 1890.
- [20] Vladimir I. Arnold. *Mathematical Methods of Classical Mechanics*. Springer, New York, 1978.
- [21] Anatole Katok and Boris Hasselblatt. *Introduction to the Modern Theory of Dynamical Systems*. Cambridge University Press, Cambridge, 1995.
- [22] Kundai Farai Sachikonye. Counting in bounded phase space: The origin of partition numbers. *Technical University of Munich, Preprint*, 2026.
- [23] Ming Li and Paul Vitányi. *An Introduction to Kolmogorov Complexity and Its Applications*. Springer, New York, 3rd edition, 2008.
- [24] Thomas M. Cover and Joy A. Thomas. *Elements of Information Theory*. John Wiley & Sons, Hoboken, NJ, 2nd edition, 2006.
- [25] Rolf Landauer. Irreversibility and heat generation in the computing process. *IBM Journal of Research and Development*, 5(3):183–191, 1961.
- [26] Charles H. Bennett. Logical reversibility of computation. *IBM Journal of Research and Development*, 17(6):525–532, 1973.
- [27] Kundai Farai Sachikonye. Atmospheric s-entropy: Position determination from molecular thermodynamic state. *Technical University of Munich, Preprint*, 2026.
- [28] Kundai Farai Sachikonye. Gas-phase categorical sensing via phase-locked molecular ensembles. *Technical University of Munich, Preprint*, 2026.
- [29] Kundai Farai Sachikonye. Equations of state for vehicular trajectory completion in

- bounded phase space. *Technical University of Munich, Preprint*, 2026.
- [30] Kundai Farai Sachikonye. The instrument problem: Categorical observation without physical backaction. *Technical University of Munich, Preprint*, 2026.
 - [31] Lev D. Landau and Evgeny M. Lifshitz. *Fluid Mechanics*, volume 6 of *Course of Theoretical Physics*. Butterworth-Heinemann, Oxford, 2nd edition, 1987.
 - [32] George Keith Batchelor. *An Introduction to Fluid Dynamics*. Cambridge University Press, Cambridge, cambridge mathematical library edition, 2000.
 - [33] David Chandler. *Introduction to Modern Statistical Mechanics*. Oxford University Press, New York, 1987.
 - [34] Kundai Farai Sachikonye. Trans-Planckian timing resolution via harmonic network phase-locking. *Technical University of Munich, Preprint*, 2026.
 - [35] George Gabriel Stokes. On the effect of the internal friction of fluids on the motion of pendulums. *Transactions of the Cambridge Philosophical Society*, 9:8–106, 1851.
 - [36] Albert Einstein. Über die von der molekularkinetischen Theorie der Wärme geforderte Bewegung von in ruhenden Flüssigkeiten suspendierten Teilchen. *Annalen der Physik*, 322(8):549–560, 1905.
 - [37] Adolf Fick. Über Diffusion. *Annalen der Physik*, 170(1):59–86, 1855.
 - [38] Geoffrey Ingram Taylor. Diffusion by continuous movements. *Proceedings of the London Mathematical Society*, s2-20(1):196–212, 1921.
 - [39] Ludwig Prandtl. Über Flüssigkeitsbewegung bei sehr kleiner Reibung. *Verhandlungen des III. Internationalen Mathematiker-Kongresses*, pages 484–491, 1904. Heidelberg.
 - [40] Hermann Schlichting and Klaus Gersten. *Boundary-Layer Theory*. Springer, Berlin, 9th edition, 2017.
 - [41] Roland B. Stull. *An Introduction to Boundary Layer Meteorology*. Kluwer Academic Publishers, Dordrecht, 1988.
 - [42] Kundai Farai Sachikonye. Architecture of the categorical processing unit. *Technical University of Munich, Preprint*, 2026.
 - [43] Peter Walters. *An Introduction to Ergodic Theory*. Springer, New York, 1982.
 - [44] Frank Pasquill and F. B. Smith. *Atmospheric Diffusion*. Ellis Horwood, Chichester, 3rd edition, 1983.
 - [45] John H. Seinfeld and Spyros N. Pandis. *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*. John Wiley & Sons, Hoboken, NJ, 3rd edition, 2016.
 - [46] Kundai Farai Sachikonye. Automobile membrane computing: A biological semiconductor surface architecture for autonomous navigation. *Technical University of Munich, Preprint*, 2026.
 - [47] M. Carpentieri and P. Kumar. Ground-level concentrations of vehicular emissions in street canyons. *Atmospheric Environment*, 56:257–270, 2012.
 - [48] Israel M. Gelfand and Sergei V. Fomin. *Calculus of Variations*. Dover, Mineola, NY, 2000.
 - [49] Lev S. Pontryagin, Vladimir G. Boltyanskii, Revaz V. Gamkrelidze, and Evgeni F. Mishchenko. *The Mathematical Theory of Optimal Processes*. Interscience Publishers, New York, 1962.
 - [50] D. Bruce Turner. *Workbook of Atmospheric Dispersion Estimates*. Lewis Publishers, Boca Raton, 1994.
 - [51] Assad Alam, Bart Besselink, Valerio Turri, Jonas Mårtensson, and Karl Henrik Johansson. Heavy-duty vehicle platooning for sustainable freight transportation. *IEEE Intelligent Transportation Systems Magazine*, 7(4):68–80, 2015.
 - [52] Sadayuki Tsugawa, Sabina Jeschke, and Steven E. Shladover. A review of truck platooning projects for energy savings. *IEEE*

Transactions on Intelligent Vehicles, 1(1): 68–77, 2016.

- [53] Carl Bergenhem, Steven Shladover, Erik Coelingh, Christofer Englund, and Sayuki Tsugawa. Overview of platooning systems. *Proceedings of the 19th ITS World Congress*, 2012. Vienna, Austria.
- [54] Kundai Farai Sachikonye. Categorical trajectory completion in bounded phase space. *Technical University of Munich, Preprint*, 2026.
- [55] Kundai Farai Sachikonye. Backward trajectory completion via partition hierarchy navigation. *Technical University of Munich, Preprint*, 2026.
- [56] Kundai Farai Sachikonye. Depth perception via categorical distance in s-entropy space. *Technical University of Munich, Preprint*, 2026.

Appendices

A Summary of Notation

Symbol	Name	Definition
V_{Γ}	Phase space volume	Eq. (1)
$N_{\max}$	Max. distinguishable states	Eq. (2)
(n, ℓ, m, s)	Partition coordinates	Def. 3.1
$C(n)$	Shell capacity	Eq. (3)
$\mathcal{M}$	Partition depth	Def. 3.4
(S_k, S_t, S_e)	S -entropy coordinates	Def. 4.2
d_{cat}	Categorical distance	Eq. (14)
$S_{\text{osc}}, S_{\text{cat}}, S_{\text{part}}$	Triple entropies	Eq. (11)
T_{cat}	Categorical temperature	Eq. (17)
P_{cat}	Categorical pressure	Eq. (20)
τ_c	Categorical lag time	Def. 5.1
g	Coupling strength	Def. 5.2
μ	Dynamic viscosity	Eq. (28)
D	Diffusion coefficient	Eq. (32)
R_{compute}	Processing rate	Eq. (37)
$\mathcal{V}(t)$	Visited state set	Def. 11.1
$\mathbf{F}$	Exhaust fingerprint	Def. 12.1
C_{trail}	Trail concentration	Eq. (71)
I_{trail}	Trail information	Eq. (50)
D_{turb}	Turbulent diffusion coeff.	$\sim 1 \text{ m}^2/\text{s}$

Symbol	Name	Definition
h_{BL}	Boundary layer height	0.5–2 m
τ_{mix}	Mixing time	Eq. (47)
ρ_{c}	Critical convoy density	Eq. (77)
$[\hat{O}_{\text{cat}}, \hat{O}_{\text{phys}}]$	Cat.–phys. commutator	Eq. (25)

B Complete Theorem Index

For the reader’s convenience, we list all 32 numbered results in order of appearance.

Number	Name	Page
Axiom 1	Bounded Phase Space	2
Thm. 2.1	Finite Distinguishability	2
Cor. 2.2	Poincaré Recurrence	3
Thm. 3.2	Shell Capacity	3
Thm. 3.5	Composition Theorem	3
Thm. 3.6	Compression Theorem	4
Thm. 3.7	Conservation Theorem	4
Thm. 3.8	Charge Emergence	4
Thm. 3.9	Partition Extinction	4
Thm. 4.1	Triple Equivalence	4
Prop. 4.3	Boundedness of S -Entropy	5
Thm. 4.6	Ideal Gas Law	6
Thm. 4.7	Single-Particle Gas Law	6
Thm. 4.8	Heat–Entropy Decoupling	6
Thm. 5.3	Viscosity from Partitions	7
Thm. 6.1	Speed of Light	7
Thm. 7.1	Stokes–Einstein	7
Cor. 7.2	Fick’s Laws	7
Thm. 8.1	Navier–Stokes	8
Thm. 8.2	Boundary Layer	8
Thm. 9.1	Oscillator–Processor Duality	9
Cor. 9.2	Temperature as Processing Rate	9
Thm. 10.2	Atmospheric Computation	9
Thm. 11.2	Monotonic Memory	10
Cor. 11.3	Atmospheric Memory	10
Prop. 12.2	Exhaust Uniqueness	11
Thm. 13.1	Trail Persistence	11
Thm. 14.1	Trail Information Content	12
Cor. 14.2	Extractable Information	12
Thm. 15.1	Signal Hierarchy	13
Thm. 16.1	Photon Independence	14
Cor. 16.2	Zero Visibility Navigation	14
Prop. 16.3	Thermal Detection	14
Prop. 16.4	Pressure Detection	15
Prop. 16.5	Molecular Detection	15
Prop. 16.6	Humidity Detection	15
Thm. 16.7	Composite Detection Range	15

Number	Name	Page
Thm. 17.1	Braking Intent Detection	16
Thm. 17.2	Hidden Vehicle Detection	17
Thm. 18.1	Optimal Path Convergence	17
Cor. 18.2	Trail as Solved Optimization	18
Cor. 18.3	Hazard Encoding	18
Thm. 19.1	Vehicle Count Recovery	19
Prop. 19.2	Historical Reconstruction	19
Thm. 20.1	Convoy Phase Transition	19
Cor. 20.2	No V2V Required	20
Cor. 20.3	Aerodynamic Benefit	20
Thm. 21.2	V2A2V Channel	20
Thm. 22.1	Pedestrian Detection	21

C Dimensional Analysis Verification

We verify dimensional consistency of all key equations.

C.1 Viscosity (Theorem 5.3)

$$[\mu] = [\tau_c] \times [g] = \text{s} \times \frac{\text{kg}}{\text{m}^2 \cdot \text{s}} = \frac{\text{kg}}{\text{m} \cdot \text{s}} = \text{Pa} \cdot \text{s}. \quad \checkmark \quad (89)$$

C.2 Stokes–Einstein (Theorem 7.1)

$$[D] = \frac{[k_B T]}{[\mu][r]} = \frac{\text{J}}{\text{Pa} \cdot \text{s} \cdot \text{m}} = \frac{\text{kg} \cdot \text{m}^2/\text{s}^2}{\text{kg}/(\text{m} \cdot \text{s}) \cdot \text{m}} = \frac{\text{m}^2}{\text{s}}. \quad \checkmark \quad (90)$$

C.3 Processing Rate (Theorem 9.1)

$$[R_{\text{compute}}] = \frac{[\omega]}{2\pi} = \frac{\text{rad/s}}{1} = \text{s}^{-1}. \quad \checkmark \quad (91)$$

C.4 Trail Persistence (Theorem 13.1)

$$[\tau_{\text{mix}}] = \frac{[h_{\text{BL}}]^2}{[D_{\text{turb}}]} = \frac{\text{m}^2}{\text{m}^2/\text{s}} = \text{s}. \quad \checkmark \quad (92)$$

C.5 Critical Density (Theorem 20.1)

$$[\rho_c] = \frac{[D_{\text{turb}}][D_\rho]}{[\alpha][\beta][v][\sigma]^2} = \frac{(\text{m}^2/\text{s})(\text{m}^2/\text{s})}{(\text{m}^3/(\text{kg} \cdot \text{s}))(\text{kg}/(\text{m} \cdot \text{s}))(\text{m}/\text{s})(\text{m}^2)} = \text{m}^{-1}. \quad \checkmark \quad (93)$$